

Effective Hamiltonian for Isotropic Loop Quantum Cosmology

Ghanashyam Date¹, and Golam Mortuza Hossain^{1,y}

¹The Institute of Mathematical Sciences

CIT Campus, Chennai-600 113, INDIA.

Abstract

For a class of solutions of the fundamental difference equation of isotropic loop quantum cosmology, the difference equation can be replaced by a differential equation valid for all values of the triad variable. The differential equation admits a 'unique' non-singular continuation through vanishing triad. A WKB approximation for the solutions leads to an effective continuum Hamiltonian. The effective dynamics is also non-singular (no big bang singularity) and approximates the classical dynamics for large volumes. The effective evolution is thus a more reliable model for further phenomenological implications of the small volume effects.

PACS numbers: 04.60.Pp, 04.60.Kz, 98.80.Jk

^Electronic address: shyam@imsc.res.in

^yElectronic address: golam@imsc.res.in

I. INTRODUCTION

The singularities of classical general relativity, when specialized to homogeneous, isotropic models, manifest as reaching zero physical volume at finite synchronous time in the past. This in turn imply unbounded growth of space-time curvature and of matter densities etc and signals break down of the evolution equations at finite time in the past. It is widely believed that this feature of the classical theory will be modified in a quantum theory of gravity and recent development of Loop Quantum Cosmology (LQC) [1, 2, 3, 4, 5, 6, 7], corroborate this expectation [8, 9].

The mechanism of 'singularity avoidance' [9] involves replacement of the classical evolution equation by a quantum one which is a difference equation [4] thanks to the necessity of using holonomy operators in the quantization of the Hamiltonian constraint in LQC. This equation exhibits the property that the quantum wave function can be evolved through zero volume unambiguously. In addition, the discreteness of the triad operator (having zero eigenvalue) necessitates defining inverse triad operator (or inverse scale factor operator) [8] indirectly. Thanks to the loop representation on the (non-separable) kinematical Hilbert space of LQC [7], these operators get so defined as to have a bounded spectrum implying only a bounded growth of curvatures/matter densities. This is true for all allowed values of the ambiguity parameters [10, 11].

The non-separable structure of the kinematical Hilbert space of LQC however, also implies a huge set of solutions of the Hamiltonian constraint (a continuous infinity in the gravitational sector alone). Presumably, a suitable choice of physical inner product can be made to cut down the size of the admissible solutions of the Hamiltonian constraint. A choice of inner product however is not yet available. The exploration of Dirac observables is also at a preliminary stage [12]. The general issue of whether or not the non-separable kinematical Hilbert space is mandatory, is currently an open issue [13, 14]. In the present work however we assume the current framework of LQC [7].

Despite the open issues, it is possible to develop a WKB type semi-classical approximation from which an effective continuum Hamiltonian constraint can be deduced [15, 16]. This at once gives access to the usual classical Hamiltonian methods to construct and analyze the quantum modified space-time. This method relies on a continuum approximation [17] of the underlying difference equation to the Wheeler-DeWitt differential equation followed by

the WKB ansatz for its solutions. For large volume corresponding to classical regime, the continuum approximation is always available, in fact as a requirement on quantization of the Hamiltonian operator. In this regime, the WKB ansatz naturally reproduces the classical Hamiltonian as the leading $\mathcal{O}(\hbar^0)$ term. We would like to extend this method also to small volumes.

The large freedom offered by the non-separable structure of the kinematical Hilbert space can be exploited to propose a restriction to those solutions of the Hamiltonian constraint for which a continuum approximation is valid for all volumes. One can then develop the effective classical Hamiltonian constraint for all volumes and explore its consequences.

In this work we develop such a picture and in comparison with the usual FRW equations identify the effective density and pressure which includes the contributions of quantum fluctuations of the geometry. Some elementary consequences are also noted. Further implications for phenomenology are discussed in separate papers [18, 19].

In section II, we detail the effective Hamiltonian constraint for isotropic models. In section III, we discuss the qualitative features of the corresponding dynamics namely, the possibility of ‘bounce’ solutions as well as solutions that could attempt to ‘pass through’ the zero volume and connect to the oppositely oriented isotropic universe. We discuss what features of the quantum evolution are captured by the effective classical evolution. In section IV, we summarize our conclusions and outlook.

II. EFFECTIVE HAMILTONIAN CONSTRAINT

The Kinematical Hilbert space [7] is conveniently described in terms of the eigenstates of the densitized triad operator,

$$\hat{p}^j_i = \frac{1}{6} \sqrt{\frac{2}{3}} \delta^j_i \hat{p}^j_i ; \quad \hat{h}^j_i = \delta^j_i \quad (1)$$

The action of the volume operator on the triad basis states are given by

$$\hat{V}^j_i = \frac{1}{6} \sqrt{\frac{2}{3}} \delta^j_i \hat{V}^j_i = V^j_i \quad (2)$$

In the isotropic context we have two classes to consider, namely spatially flat and closed models. The quantization of the corresponding Hamiltonian operators is given in [5]. By introducing a parameter we can deal both classes together. The values $\alpha = 0$ and $\alpha = 1$ will give the flat and the closed models respectively.

The action of the gravitational Hamiltonian on the triad basis states is then given by

$$\hat{H}_{\text{grav}}^{(\gamma)} |j\rangle = \frac{3}{4} \left(\gamma^3 \gamma_{\alpha\beta}^2 \right)^{-1} (V_{+ \gamma} - V_{- \gamma}) e^{i \gamma} |j + 4 \gamma\rangle + (2 + 4 \gamma^2) |j\rangle + e^{i \gamma} |j - 4 \gamma\rangle : \quad (3)$$

Here, γ is a quantization ambiguity parameter which enters through the dual length of the loops used in defining the holonomies. It is a real number of the order of 1. Notice that the Hamiltonian connects states differing in their labels by 4γ . This is a direct consequence of the use of holonomy operators which have to be used in the quantization of the Hamiltonian operator and is responsible for leading to a difference equation below.

A general kinematical state $|j\rangle$, in the triad basis has the form

$$|j\rangle = \sum_{2R} s_j |j\rangle \quad (\text{sum over countable subsets}) : \quad (4)$$

The Hamiltonian constraint of the classical theory is promoted as a condition to define physical states, i.e.,

$$(\hat{H}_{\text{grav}}^{(\gamma)} + \hat{H}_{\text{matter}}^{(\gamma)}) |j\rangle = 0 : \quad (5)$$

In terms of $s_j := e^{\frac{i}{4}} s_j$ and The Hamiltonian constraint (5) translates into a difference equation,

$$\begin{aligned} 0 = & A_{+4\gamma} s_{j+4\gamma} - (2 + 4\gamma^2) A s_j + A_{-4\gamma} s_{j-4\gamma} \\ & + 8 \gamma^2 \gamma^3 \frac{1}{6} \gamma^{\frac{1}{2}} H_m(\gamma) s_j ; \quad 8 \gamma^2 R \\ A_{\pm} & := |j + \gamma| \gamma^{\frac{3}{2}} - |j - \gamma| \gamma^{\frac{3}{2}} ; \quad \hat{H}_{\text{matter}}^{(\gamma)} |j\rangle := H_m(\gamma) |j\rangle : \end{aligned} \quad (6)$$

$H_m(\gamma)$ is a symbolic eigenvalue and we have assumed that the matter couples to the gravity via the metric component and not through the curvature component. In particular, $H_m(\gamma = 0) = 0$. There are a few points about the above equation worth noting explicitly.

Although γ takes all possible real values, the equation connects the s_j coefficients only in steps of 4γ making it a difference equation for the coefficients. By putting $\gamma = \frac{1}{4} + (4 \gamma)n$; $n \in \mathbb{Z}$; $\gamma \in [0; 4 \gamma)$, one can see that one has a continuous infinity of independent solutions of the difference equation, labelled by n ; $S_n := s_{j+4 \gamma n}$. For each n an infinity of coefficients, S_n , are determined by 2 'initial conditions' since the order of the difference equation in terms of these coefficients is 2. Coefficients belonging to different n are mutually

decoupled. Since the coefficients A and the symbolic eigenvalues $H_m(\cdot)$, both vanish for $\alpha = 0$, the coefficient s_0 decouples from all other coefficients.

For large values of $4\alpha_0(n-1)$, which correspond to large volume, the coefficients A become almost constant (up to a common factor of $\sqrt[n]{P}$) and the matter contribution is also expected similarly to be almost constant. One then expects the coefficients to vary slowly as n is varied. This suggests interpolating these slowly varying sequences of coefficients by slowly varying functions of the continuous variable $p(n) \equiv \frac{1}{6} \frac{V}{P} n$ [15]. The difference equation satisfied by the coefficients then implies a differential equation for the interpolating functions which turns out to be independent of α and matches with the usual Wheeler-DeWitt equation of quantum cosmology. This is referred to as a continuum approximation [17]. This is of course what one expects if LQC dynamics is to exhibit a semi-classical behavior. While admissibility of continuum approximation is well motivated for large volume, one also expects it to be a poor approximation for smaller Planck scale volumes.

This logic is valid when applied to any one of the solutions S_n . Thanks to the non-separable structure of the Hilbert space, we have an infinity of solutions of the Hamiltonian constraint. Although S_n are uncorrelated for different n , nothing prevents us from choosing them to be suitably correlated. In effect this amounts to viewing s themselves as functions of the continuous variable p and stipulating some properties for them. In the absence of a physical inner product, we don't have any criteria to select the class of solution. It is then useful to study properties of classes of solutions of the Hamiltonian constraint.

The class that we will concentrate on is the class of slowly varying functions. For these we will be able to have a continuum approximation leading to a differential equation. Making a WKB approximation for this differential equation, we will read-off the effective classical Hamiltonian constraint. In anticipation of making contact with a classical description, we will use the dimensionful variable $p(\cdot) \equiv \frac{1}{6} \frac{V}{P} \cdot$ as the continuous variable. Correspondingly, we define $p_0 \equiv \frac{1}{6} \frac{V}{P} \alpha_0$ which provides a convenient scale to demarcate different regimes in p . We also use the notation: $(p(\cdot)) \equiv s$. Now the definition of a slowly varying function is simple: (p) is locally slowly varying around q if

$(q + \epsilon q) = (q) + \frac{d}{dq} \epsilon + \frac{1}{2} \epsilon^2 \frac{d^2}{dq^2} + \dots$ with successive terms smaller than the preceding terms, for $q \in [4p_0, 4p_0 + \epsilon]$. It is slowly varying if it is locally slowly varying around every $q \in \mathbb{R}$ [17]. Note that even an exponentially rising function can be locally slowly varying if the exponent is sufficiently small.

To explore the possibility of slowly varying solutions of the difference equation (6), consider the difference equation more explicitly. For $\epsilon \in (0; 4p_0)$, putting $S_n(\epsilon) \equiv s_{n+4p_0\epsilon}$; $A_n(\epsilon) \equiv A_{n+4p_0\epsilon}$ and momentarily ignoring the matter term for notational simplicity, the difference equation (6) can be written as,

$$S_{n+2}(\epsilon) = \left(2 + 4\frac{\epsilon^2}{p_0^2}\right) \frac{A_{n+1}(\epsilon)}{A_{n+2}(\epsilon)} S_{n+1}(\epsilon) + \frac{A_n(\epsilon)}{A_{n+2}(\epsilon)} S_n(\epsilon) \quad (7)$$

Its general solution can be written as $S_n(\epsilon) = S_0(\epsilon) n(\epsilon) + S_1(\epsilon) n(\epsilon)$, where the n 's are fixed functions of ϵ determined by the same difference equation (7) with the 'initial' conditions: $n_0(\epsilon) = 1$; $n_1(\epsilon) = 0$ and $n_0(\epsilon) = 0$; $n_1(\epsilon) = 1$ and S_0, S_1 are arbitrary functions of $\epsilon \in (0; 4p_0)$. (The linearity of the equation means that only the ratio $n(\epsilon) \equiv S_1(\epsilon)/S_0(\epsilon)$ (say) parameterizes the general solution.)

It is clear that the arbitrary functions allow us to control the variation of s within an interval of width $4p_0$. At the integral values of $\epsilon = (4p_0)$ corresponding to $\epsilon = 0$, there is a consistency condition coming from vanishing of the highest (lowest) order coefficient which fixes the ratio of $S_0(0)$ and $S_1(0)$. The values of $s_{n=4p_0\epsilon}$ are fixed (up to overall scaling). The slowly varying class of functions will be assumed to approximate these exact values. The continuum approximation developed below may not be a good approximation at a finite subset of these values corresponding to smaller n .

With these remarks, we now proceed to derive consequences from the assumption of (everywhere) slowly varying, approximate solutions of the difference equation (6).

Defining $A(p) \equiv \left(\frac{1}{6} - \frac{\epsilon^2}{p}\right)^{\frac{3}{2}} A$ and substituting s in terms of slowly varying $A(p)$ in the difference equation (6), leads to the differential equation,

$$0 = B_0(p; p_0) A(p) + 4p_0 B_1(p; p_0) A'(p) + 8p_0^2 B_2(p; p_0) A''(p) \quad \text{where,} \quad (8)$$

$$B_1(p; p_0) \equiv A(p + 4p_0) - A(p - 4p_0); \quad \text{and}$$

$$B_2(p; p_0) \equiv A(p + 4p_0) \left(2 + 144 \frac{p_0^2}{p^4}\right) A(p) + A(p - 4p_0) + 288 \frac{p_0^3}{p^4} H_m(\epsilon)$$

In the above equation, terms involving higher derivatives of $A(p)$ have been neglected as being sub-leading in the context of slowly varying solutions. This is not quite the continuum approximation referred to earlier since there is ϵ dependence hidden inside p_0 appearing explicitly in the coefficients of the differential equation. This is also not quite the Wheeler-DeWitt equation since this equation is valid over the entire real line (since p can take negative

values corresponding to oppositely oriented triad) while the Wheeler-DeWitt equation using the scale factor as independent variable is defined only for half real line.

From the definitions of the coefficients $A; B_{\pm}$, it is obvious that under $p \rightarrow -p$ (change of orientation of the triad), $A; B_{\pm}$ and the gravitational part of B_0 are all odd while B_0 is even. For notational convenience we restrict to $p \geq 0$ while writing the limiting expressions, the expressions for negative p can be obtained from the odd/even properties noted above. There are two obvious regimes to explore which are conveniently demarcated by the scale p_0 , namely, $p \gg p_0$ and $0 \leq p \ll p_0$. The corresponding limiting forms for the coefficients $B_0; B_{\pm}$ are easily obtained. One gets,

$$\begin{aligned}
 p \gg p_0 : A(p; p_0) &= 3p_0^3 \bar{p} - \frac{1}{8} p_0^3 p^{\frac{3}{2}} \\
 : B_{+}(p; p_0) &= 6p_0^3 \bar{p} - \frac{49}{4} p_0^3 p^{\frac{3}{2}} \\
 : B_{-}(p; p_0) &= 12p_0^2 p^{\frac{1}{2}} + o(p_0^4 p^{\frac{5}{2}}) \\
 : B_0(p; p_0) &= 12p_0^3 \bar{p}^{\frac{3}{2}} - 432 \frac{p_0^3 \bar{p}}{\sqrt{p}} + 288 \frac{p_0^3}{\sqrt{p}} H_m(\cdot) \quad (9)
 \end{aligned}$$

$$\begin{aligned}
 p \ll p_0 : A(p; p_0) &= 3pp_0^{\frac{1}{2}} - \frac{1}{8} p^3 p_0^{\frac{3}{2}} \\
 : B_{+}(p; p_0) &= 3(5^{\frac{1}{2}} - \frac{1}{3}) pp_0^{\frac{1}{2}} \\
 : B_{-}(p; p_0) &= 2p_0^{\frac{3}{2}} (5^{\frac{3}{2}} - \frac{1}{3}) \\
 : B_0(p; p_0) &= (3pp_0^{\frac{1}{2}}) (5^{\frac{1}{2}} - \frac{1}{3} - 2) - 432 \frac{p_0^{\frac{5}{2}} p}{\sqrt{p}} + 288 \frac{p_0^3}{\sqrt{p}} H_m(\cdot) \quad (10)
 \end{aligned}$$

Notice that for large volume the explicit p_0 dependence cancels out. This equation corresponds to the usual continuum approximation which has no dependence on $\hbar$ and matches with the Wheeler-DeWitt equation in a particular factor ordering. For small volume, the p_0 dependence survives, is non-trivial and the coefficient of the first derivative term is non-zero. In view of the even/odd properties of the coefficients, it follows that the first derivative of $\psi(p)$ must vanish at $p = 0$. Furthermore, even for the flat model without matter, the B_0 coefficient is non-zero. Had we extrapolated the Wheeler-DeWitt equation from the large volume form, we would not have gotten these terms. Thus the quantum differential equation (8) agrees with the Wheeler-DeWitt for large volume but differs significantly for small volume.

The small volume form of the equation in fact shows that there are two possible behaviors namely $\psi(p)$ constant or $\psi(p)$ diverges as an inverse power of p . Neither of the indicial

roots depend on the matter Hamiltonian. The latter solution is not slowly varying and the former one implies that the wave function has a non-zero value at $p = 0$ and the wave function can obviously be continued to negative p . Thus the differential equation derived for slowly varying functions is both consistent at zero volume and mimics main features of the difference equations namely passing through zero volume and matching with Wheeler-DeWitt for large volume.

In the earlier quantization of isotropic models [5] based on point holonomies taking values in $U(1)$ representations (separable Hilbert space), the decoupling of s_0 coefficient also implied a consistency condition which helped select a unique solution [20] from solutions of the Wheeler-DeWitt equation valid at large volume. With the non-separable Hilbert space, such a condition can only result from S_n^0 family of coefficients. Nevertheless, one has gotten a unique solution (up to normalization) thanks to the slowly varying nature of the solutions. It is crucial here that for small $p \neq 0$, the B coefficient has a non-vanishing limit which forces the first derivative to vanish at $p = 0$. (If the single derivative term has been dropped, both solutions would have been slowly varying near $p = 0$.)

In summary, with the restriction to slowly varying solutions, we have a continuum approximation (differential equation) valid for all values of the triad. Furthermore the differential equation permits a unique solution (for each matter state) passing through $p = 0$.

For future reference we also note that the differential equation admits a 'conserved current'. Taking imaginary part of times the differential equation leads to,

$$2p_0 B_+ J^0 + B_- J = 0 \quad ; \quad J = 0 \quad ()^0 \quad (11)$$

Defining $J(p) := f^{-1}(p) J(p)$ such that $J^0 = 0$ determines the function f . Explicitly,

$$\begin{aligned} J(p) &= \text{constant} \cdot e^{\int_{p_0}^p \frac{B_-}{2p_0 B_+} dp} \cdot f^{-1}(p) \cdot ()^0 g \quad ; \quad J^0 = 0 \quad ; \\ &= \text{constant}(p) \cdot f^{-1}(p) \cdot ()^0 g \quad (p = p_0) \\ &= \text{constant} \cdot p^{\frac{8+p}{3}} \cdot f^{-1}(p) \cdot ()^0 g \quad (p = p_0) : \end{aligned} \quad (12)$$

We will now go ahead with a WKB type solution and infer an effective classical Hamiltonian.

Let $\psi(p) = C(p) e^{\frac{i}{\hbar} S(p)}$. Substitution of this ansatz in (8) leads to a complex differential equation involving $C(p); S(p)$. The real and imaginary parts lead to,

$$0 = B_0(p; p_0) + 4p_0 B_-(p; p_0) f(\hbar C)^0 g + 8p_0^2 B_+(p; p_0) \frac{\partial}{\partial p} + (\hbar C)^{0,2} + (\hbar C)^0 \quad (13)$$

$$0 = 4p_0 B_-(p;p_0) f^0 g + 8p_0^2 B_+(p;p_0) f^{\omega} + 2^{-1} (\ln C)^0 g \quad (14)$$

The WKB approximation consists in assuming that the amplitude C is essentially constant and the double derivatives of the phase are small compared to the single derivatives. Consider the eq.(13) under the assumption of almost constant amplitude $C(p)$. Then this equation is a Hamilton-Jacobi partial differential equation. These are generally, partial differential equations involving only first derivatives with respect to time and position and they always have an associated Hamiltonian mechanics [21].

$$B_0(p;p_0) - 8p_0^2 B_+(p;p_0) \frac{\omega}{\sim^2} = 0 \quad (15)$$

Noting that the Poisson bracket between the triad variable p and the extrinsic curvature variable K is $\frac{1}{3}$, we identify $\omega = \frac{1}{3}K$ and arrive at the effective Hamiltonian as,

$$H^e(p;K; \hbar; p) = -\frac{1}{4p_0} \frac{B_+(p;p_0)}{K^2} + \frac{A(p)}{2p_0} + \frac{1}{288p_0^3} \frac{\hbar^4}{p} fB_+(p;p_0) - 2A(p)g + H_m(p; \hbar; p) \quad (16)$$

We have multiplied by certain factors so as to get the matter Hamiltonian term appear without any pre-factors as in the classical case. The equation (13) of course implies $H^e = 0$ and we will interpret this as the modified Hamiltonian constraint equation. The effective Hamiltonian is also odd under $p \rightarrow -p$ modulo the matter term.

Note that the K^2 and the $\hbar$ dependent terms are $o(\hbar^0)$. For large volume, the terms enclosed within the braces are vanishingly small and the effective Hamiltonian is indeed classical (the matter Hamiltonian receiving corrections from the inverse triad operator also goes to the classical form without any $\hbar$ dependence). For smaller volumes, the quantum modifications are present with explicit dependence on $\hbar/p_0$. The approximation used does not quite lead to a 'classical limit' due to explicit appearance of $\hbar$.

For small volumes, the equation (10) shows that B_0, B_+, A all vanish linearly with p while B_- goes to a positive constant. The real and the imaginary parts of the equation, equations(13, 14), then imply that ω and C^0 both vanish, which is consistent with constant, as deduced directly from the differential equation (8).

To interpret the effective Hamiltonian constraint as generating the space-time dynamics, let us use the identification $\dot{p} = a^2$. One can then obtain the extrinsic curvature K from the Hamilton's equation of motion of p as,

$$\dot{p} = \frac{dp}{dt} = \frac{\partial H^e}{\partial K} = \frac{B_+(p;p_0)}{6p_0} K \quad \text{or,}$$

$$K = 12 \frac{ap_0}{B_+(p;p_0)} \frac{a}{p} \quad (17)$$

The large and small volume expressions for the effective Hamiltonian and the extrinsic curvature are,

$$H^{\text{eff}} = \frac{3}{2} p \bar{p} K^2 + H_m \quad (18)$$

$$K = \frac{2a}{p} \left(\frac{p}{p_0} \right)^{\frac{1}{3}} \left(\frac{p_0}{p} \right)^{\frac{1}{3}} \quad (19)$$

$$H^{\text{eff}} = \frac{3}{2} p \frac{p}{p_0} \frac{5^{\frac{1}{2}}}{2} \frac{3^{\frac{1}{3}}}{2} K^2 + \frac{1}{144} \left(\frac{p}{p_0} \right)^{\frac{1}{2}} + 3^{\frac{1}{2}} \frac{p^4}{p_0^2} + H_m \quad (20)$$

$$K = \frac{4p}{5^{\frac{1}{2}} 3^{\frac{1}{3}}} \frac{a}{p_0} \left(\frac{p}{p_0} \right)^{\frac{1}{3}} \quad (21)$$

The large volume expressions are the same as for the classical Hamiltonian as expected. The small volume expressions are useful in exploring the behavior of the effective dynamics close to the classical singularity.

It is worth expanding on the identification $\dot{p}j = a^2$. The basic variables of LQC are first obtained for a general homogeneous Bianchi class A models with the Maurer-Cartan forms normalized in the usual manner. Comparing with the metric ansatz then leads to the relations $\dot{p}_1 j = a_2 a_3$ and cyclic. The basic variables of the isotropic models are obtained from the above Bianchi ansatz, by putting $p_I = p$; $a_I = a/8I$ leading to the identification above. Let us denote this scale factor as a_{Bianchi} . The corresponding spatial Ricci scalar is ${}^3R(a_{\text{Bianchi}}) = \frac{3}{2a_{\text{Bianchi}}^2}$. On the other hand, the standard FRW metric ansatz is so chosen that the spatial curvature is given by ${}^3R(a_{\text{FRW}}) = \frac{6}{a_{\text{FRW}}^2}$. These two normalizations match provided $a_{\text{Bianchi}} = \frac{a_{\text{FRW}}}{2}$. To avoid writing the sufixes, we just note that while comparing the large volume Hamiltonian constraint with the standard Friedmann equation, one should use the replacement $a \rightarrow \frac{a}{2}$. This of course is relevant only for the closed model.

III. QUALITATIVE FEATURES OF THE EFFECTIVE DYNAMICS

In the previous section we derived the effective Hamiltonian constraint (16), using a continuum approximation keeping terms up to second derivatives and using the WKB approximation. If we include higher derivative terms that these would give perturbative corrections in the large volume. Our focus is however on the small volume regime and leading corrections which for the matter sector include non-perturbative corrections. For our purposes the truncation to second derivatives suffices.

For large volume, we already see that the effective Hamiltonian reduces to the classical one to within terms of the order of $p^{-\frac{3}{2}}$. We are interested in checking if the effective dynamics is non-singular and precisely in what sense. For this it is sufficient to focus on the small volume expressions. We will compare the classical Hamiltonian (18) extrapolated to small volume and the effective Hamiltonian (20).

Consider first the classical case. As the scale factor goes to zero, the matter density diverges either as a^{-3} for pressure-less matter or as a^{-4} for radiation. Correspondingly, H_m either goes to a non-negative constant or diverges as a^{-1} . The Hamiltonian constraint then implies that K necessarily diverges. As is well known, in both cases the scale factor vanishes at a finite value of synchronous time and this of course is the big bang singularity. This also suggests a necessary condition for singular evolution: $p = 0$ should be reachable in finite time. Equivalently, if $p = 0$ is not reachable in finite time, the evolution is non-singular.

Moreover, let us assume that for some reason, the matter Hamiltonian vanishes as the scale factor goes to zero, then K must remain finite and $p = 0$ is indeed on the constraint surface. Furthermore $\underline{p}$ evaluated at $p = 0$ is also zero implying that $p = 0$ is a fixed point (rather a fixed 'sub-manifold' of the phase space of gravity and matter). The $p = 0$ trajectories of the dynamics are then not accessible in finite time and the evolution is non-singular. Clearly (non-) divergence of matter Hamiltonian dictates (non-) singular evolution.

For generic (non-singular) trajectories, there are two possibilities now. Either (a) $p = 0$ is approached asymptotically as $t \rightarrow \pm\infty$ or (b) the trajectory exhibits a bounce, $K = 0$ at a finite, non-zero p . For example, in the case of scalar matter, with LQC modifications included, the former is realized for flat models ($\Omega = 0$) while the latter is realized for closed models ($\Omega = 1$) [22].

Consider now the quantum case. The matter Hamiltonian is guaranteed to vanish due to the inverse volume operator definition. The arguments for $p = 0$ being fixed point apply. However, due to the presence of $\frac{V^4}{p}$ term in eq. (20), there is a bounce independent of [19]. The $p = 0$ is completely decoupled from all other trajectories. This is exactly the same feature exhibited by the fundamental difference equation. The exact solution $s = s_0, 0$ completely decouples from all other solutions. The bounce is then a completely generic feature of isotropic LQC. But a bounce also provides a minimum volume for the isotropic universe whose value is dependent on matter Hamiltonian. Such a natural, generic scale has implications for phenomenology as well [19].

Notice that while interpreting the effective Hamiltonian as a (modified) constraint equation, we are keeping the kinematics of space-time (a pseudo-Riemannian manifold) intact. The modifications imply modification of the dynamical aspects or equivalently of Einstein equations. To see the modifications conveniently, let us write the Hamilton's equations in a form similar to the usual Raychoudhuri and Friedmann equations in terms of the FRW scale factor.

Write the effective Hamiltonian (16) in the form ,

$$H^e = \frac{1}{2} (K^2 + \dots) + \frac{1}{2} + H_m \quad \text{where,} \quad (22)$$

$$\doteq \frac{B_+}{4p_0} ; \quad \doteq \frac{A}{2p_0} ; \quad \doteq \frac{\sqrt[4]{p}}{288p_0^3} (B_+ - 2A) \quad (23)$$

Putting $p \doteq \frac{a^2}{4}$ (a is the FRW scale factor and $\doteq 16 G$) and denoting $\frac{d}{da}$ by 0 leads to,

$$3\frac{a^2}{a^2} + 3\frac{1}{a^2} = \frac{16}{3} \frac{1}{a^4} + 3\frac{1}{a^2} - \frac{16}{9} \frac{1}{a^2} + \frac{16}{3} \frac{H_m}{a^4} \quad (24)$$

$$3\frac{a}{a} = \frac{8}{3a^4} - 2 \frac{a^0}{a^0} + 2 \frac{a^0}{a^0} - 2 \frac{a^0}{a^0} H_m - aH_m^0 \quad (25)$$

Comparing with the usual FRW equations, we identify effective perfect fluid density and pressure as,

$$\rho^e \doteq \frac{32}{3} \frac{H_m}{a^4} + \frac{32}{3} \frac{1}{a^4} + \frac{6}{a^2} - \frac{16}{9} \frac{1}{a^2} \quad (26)$$

$$P^e \doteq + \frac{32}{9} \frac{1}{a^4} - \frac{1}{a^0} H_m - aH_m^0 + \frac{32}{9} \frac{1}{a^4} - \frac{1}{a^0} a^0 - \frac{32}{9} \frac{1}{a^4} - \frac{1}{a^0} a^0 + \frac{9}{16} \frac{a^2}{a^4} \quad (27)$$

The large and small volume expressions for the effective density and pressure are,

$$\text{for } p = p_0 : \quad \rho^e \doteq \frac{3}{4} a^3 ; \quad \rho^e \doteq \frac{3}{4} a^3 ; \quad \rho^e \doteq \frac{1}{3} \sqrt[4]{p} a^3 ; \quad (28)$$

$$P^e \doteq \frac{8}{3} a^3 (aH_m^0) - \frac{8}{3} \sqrt[4]{p} a^6 ; \quad (29)$$

$$\text{for } p = p_0 : \quad \rho^e \doteq \frac{3}{16} p_0^{1=2} a^2 ; \quad \rho^e \doteq \frac{3}{8} p_0^{1=2} a^2 ; \quad \rho^e \doteq \frac{b}{384} \sqrt[4]{p} p_0^{5=2} a^2 ; \text{ where } b \doteq \frac{p_0}{5} - \frac{p_0}{3} ;$$

$$\rho^e = -\frac{2b}{3p_0}a^2 H_m - \frac{b(2-b)}{192}p_0^3 + \frac{6}{-a^2} - \frac{ba^2}{8p_0}; \quad (30)$$

$$P^e = -\frac{2b}{3p_0}a^2 (H_m + aH_m^0) + \frac{b(2-b)}{192}p_0^3 + \frac{2}{-a^2} - \frac{3ba^2}{8p_0}; \quad (31)$$

The effective density and pressure receive contributions from the matter sector and also the spatial curvature (Λ -dependent terms). Apart from these, the homogeneous and isotropic quantum fluctuations of the geometry also contribute an effective density and pressure (p_P^4 terms). While tiny, these are non-zero even for large volume. Notice also that while for large volume the spatial curvature does not contribute to the density and pressure, for small volume it does. As a by product, we have also obtained the density and pressure for matter, directly in terms of the matter Hamiltonian. This is useful because currently LQC modifications to the matter sector are incorporated at the level of the Hamiltonian and not at the level of an action. Consequently, usual prescription for construction of the stress tensor and reading off the density and pressure is not available. These definitions of course automatically satisfy the conservation equation: $\dot{a} = -3(P + \rho)$.

Let us consider the vacuum sector, $H_m = 0$. The more general case of presence of scalar field matter is discussed in [18]. Even for the flat model ($\Lambda = 0$), the $o(p_P^4)$ terms contributes to the effective density and pressure. This term in the effective Hamiltonian, $W_{\text{qg}} = \frac{(p_P^4)}{q}$, is a 'potential' term and will be referred to as the quantum geometry potential. It is actually of order $\frac{1}{a^2}$ after expressing p/p_0 in terms of the a/a_0 . It is easy to see that the quantum geometry potential is odd under $p \rightarrow -p$ (since B_+ and A are odd) and for $p > 0$ it is negative-definite. Its plot is shown in the figure 1. This immediately implies that in the absence of matter (and cosmological constant), all the three terms in the effective Hamiltonian must be individually zero which is not possible for the quantum geometry potential. In other words, there is no solution space-time. This is in contrast to the purely classical Hamiltonian which does give the Minkowski space-time¹ as a solution for the flat case ($\Lambda = 0$). This is also apparent from the non-zero value of the effective density which prevents the Minkowski solution.

This feature can also be understood in the following manner. The differential equation has a unique solution (which is slowly varying everywhere). This solution is however purely

¹ The Minkowski space-time here refers to Riemann metric regardless of its global topology. In the cosmological context, the spatial slice is always compact.




FIG .1: The quantum geometry potential. The triad variable p is in units of p_0 while the potential $W_{qg}(p)$ is in units of $\frac{1}{288} \sqrt{p} p_0^{3=2}$.

real and does not admit a WKB form. Consequently, the universe does not admit any 'classically allowed region' in the WKB sense and thus also does not exhibit a classical behavior. Once matter is included, we have again a unique, real solution of the differential equation for every matter state. We can now have complex linear combinations admitting possibility of regimes of WKB form and corresponding classical behavior.

It is straight forward to write down a Lagrangian from the H^e as,

$$\begin{aligned}
 L^e(p;p) &:= -K p^3 H^e(p;K) \\
 &= -\frac{1}{9} \frac{p_0}{B_+} p^2 - \frac{A(p)}{2p_0} + \frac{\sqrt{p}}{288p_0^3} fB_+(p;p_0) - 2A(p)g
 \end{aligned} \tag{32}$$

It would be interesting to see if this Lagrangian can be obtained from a specialization of a generally covariant action to homogeneous, isotropic metric.

IV . CONCLUSIONS AND OUTLOOK

The results of this paper are based on two essential ingredients: the proposal of a continuum approximation for all volumes exploiting the non-separable nature of the kinematical

Hilbert space and the derivation of the effective Hamiltonian via the WKB route.

The continuum approximation step leads to a differential equation for a (still) quantum wave function, $\psi(p)$. This equation matches with the Wheeler-DeWitt equation for large volume and has important deviations (the first derivative terms) from it at small volumes. These deviations allow continuation of the wave function through zero volume just as the fundamental difference equation does. For slowly varying solutions, it picks out the 'boundary' condition $\frac{d}{dp}(0) = 0$. Again this is analogous to unique solution (per matter state) picked out by the difference equation obtained from the $U(1)$ point holonomies in the earlier work [20]. Thus the essential features of the fundamental difference equation namely non-singular quantum evolution with semi-classical limit are captured by the continuum differential equation.

The effective dynamics specified by the effective Hamiltonian deduced via the WKB approximation also reflects these features. The effective dynamics is non-singular, captures the decoupling of the $s = s_0, 0$ exact solution of the difference equation by making the classical $p = 0$ trajectory decouple and reduces to the usual classical dynamics of general relativity for large volumes. Since the essential features of quantum dynamics are now captured in classical geometrical terms, the effective dynamics is more reliable than the usual one and one can now simply work with the effective dynamics to do phenomenology. Already, at the qualitative level, one sees that all non-trivial evolutions necessarily show a bounce providing a natural scale for say, density perturbations and their power spectra.

Since the approach draws on the WKB method, a few remarks on the interpretational aspects are in order.

From the continuum quantum dynamics, with the WKB approximation, one obtains a Hamilton-Jacobi equation. As a mathematical result, any Hamilton-Jacobi differential equation has an associated Hamiltonian mechanics [21] and corresponding trajectories. In our context, we are interpreting these trajectories as possible evolutions of the isotropic universe. There is at least an implicit implication (or assumption) that a quantum system executing a WKB approximable quantum motion physically exhibits a classical motion governed by the Hamiltonian associated with the corresponding Hamilton-Jacobi equation. The justification for this comes from noting that for large volume we expect the universe to exhibit classical behavior and there it is indeed governed by the associated Hamiltonian. For how small volumes can we assume this expectation? This question is naturally related to the domain

of validity of the WKB approximation.

One expects the WKB approximation (slow variation of the phase and almost constancy of the amplitude of the wave function) to break down closer to the classically indicated singularity at zero volume. Noting that the differential equation is local in p and its solutions are also local solutions (valid in open intervals in p), we can begin with a WKB approximable solution valid in the larger volume regime and attempt to extrapolate it to smaller and smaller volumes. All through these extensions, one will have the effective Hamiltonian with its associated trajectories which can access the values of p in these intervals. The effective Hamiltonian constraint defines a submanifold of the phase space and all trajectories must lie on this. The range of configuration space variables (eg p in our case) allowed by the submanifold defines 'classically allowed region'. As is well known from the usual examples in non-relativistic quantum mechanics, the WKB approximation breaks down at the 'turning points'. These correspond to the boundary of classically allowed region which therefore demarcates the domain of validity of WKB approximation. Clearly, when such a boundary is reached by a trajectory, it must turn back. This is of course the bounce ($\dot{a} = 0$). The expectation that WKB breaks down at non-zero volume translates into the expectation of a bounce occurring at non-zero scale factor. The bounce can thus be understood as the smallest volume (or scale factor) down to which one may use the classical framework with some justification but below it one must use the quantum framework. More details of the bounce picture as well as its genericness are discussed in [19].

Further justification comes from other known examples. For example, solutions of the Maxwell equations in the eikonal approximation can be understood in terms of the normals to the wave fronts which follow null geodesic. The interpretation that this actually reflects rectilinear motion of light, may be justified by noting that the Poynting vector (energy flow) is also in the same direction as the normals. Likewise, in the context of usual Schrodinger equation of particle mechanics, the conserved probability current also points along the normals to the wave fronts giving credence to the interpretation that a quantum state of the WKB form realizes motion of a particle (or wave packet) governed by the associated Hamiltonian mechanics. In both these examples, further inputs other than the mathematical association between Hamilton-Jacobi differential equation and a Hamiltonian system, seem to be needed to understand the physical realization of the Hamiltonian system.

Interestingly, the equation (8) does admit a conserved current (12) which indeed is pro-

portional to the gradient of the WKB phase. Whether this could be used for guessing physical inner product vis a vis a probability interpretation remains to be seen.

Acknowledgments

We thank Martin Bojowald for helpful, critical comments.

-
- [1] Bojowald M 2000 Loop Quantum Cosmology: I. Kinematics Class. Quantum Grav. 17 1489{1508
 - [2] Bojowald M 2000 Loop Quantum Cosmology: II. Volume Operators Class. Quantum Grav. 17 1509{1526
 - [3] Bojowald M 2001 Loop Quantum Cosmology III: Wheeler-DeWitt Operators Class. Quantum Grav. 18 1055{1070
 - [4] Bojowald M 2001 Loop Quantum Cosmology IV: Discrete Time Evolution Class. Quantum Grav. 18 1071{1088
 - [5] Bojowald M 2002 Isotropic Loop Quantum Cosmology Class. Quantum Grav. 19 2717{2741
 - [6] M. Bojowald and H. A. Morales-Tecotl 2004 Cosmological applications of loop quantum gravity, In Proceedings of the Fifth Mexican School (DGFM): The Early Universe and Observational Cosmology, Lect. Notes Phys. Vol 646, Ed. Breton, N., Cervantes-Cota, J. and Salgado, M., Springer-Verlag [gr-qc/0306008].
 - [7] Ashtekar A, Bojowald M, and Lewandowski, J 2003 Mathematical structure of loop quantum cosmology Adv. Theor. Math. Phys. 7 233{268, [gr-qc/0304074]
 - [8] Bojowald M 2001 Inverse Scale Factor in Isotropic Quantum Geometry Phys. Rev. D 64 084018
 - [9] Bojowald M 2001 Absence of a Singularity in Loop Quantum Cosmology Phys. Rev. Lett. 86 5227{5230
 - [10] Bojowald M 2002 Quantization ambiguities in isotropic quantum geometry Class. Quantum Grav. 19 5113{5130
 - [11] Bojowald M 2004 Loop Quantum Cosmology: Recent Progress, Plenary talk at ICGC 2004, to appear in the Proceedings volume of Pramana [gr-qc/0402053]

- [12] Hossain G M 2004 Hubble operator in isotropic loop quantum cosmology *Class. Quantum Grav.* 21 179-196
- [13] Fairbairn W and Rovelli C 2004 Separable Hilbert space in Loop Quantum Gravity *Jour. Math. Phys.* 45 2802-2814
- [14] Velhinho J.M .2004 Comments on the kinematical structure of loop quantum cosmology *Class. Quantum Grav.* 21 L109 [gr-qc/0406008]
- [15] Bojowald M 2001 The semiclassical limit of Loop Quantum Cosmology *Class. Quantum Grav.* 18 L109{L116
- [16] Bojowald M ,Date G and Hossain G M 2004 The Bianchi IX model in loop quantum cosmology, *Class. Quantum Grav.* 21 3541-3570 [gr-qc/0404039]
- [17] Bojowald M and Date G 2004 Consistency conditions for fundamentally discrete theories, *Class. Quantum Grav.* 21 (2004) 121{143, [gr-qc/0307083]
- [18] Date G and Hossain G M 2004 Genericity of inflation in isotropic loop quantum cosmology, [gr-qc/0407069] .
- [19] Date G and Hossain G M 2004 Genericity of Big Bounce in isotropic loop quantum cosmology, [gr-qc/0407074] .
- [20] Bojowald M 2001 Dynamical Initial Conditions in Quantum Cosmology *Phys. Rev. Lett.* 87 121301
- [21] Evans L C 1998 *Partial Differential Equations*, Graduate Studies in Mathematics, vol 19, American Mathematical Society, Providence, 1998
- [22] Singh P and Toporensky A 2004, Big Crunch Avoidance in $k = -1$ Loop Quantum Cosmology, *Phys. Rev. D* 69, 104008, [gr-qc/0312110] .