

Sudden future singularities in FLRW cosmologies

Kayll Lake [*]

Department of Physics and Department of Mathematics and Statistics,
Queen's University, Kingston, Ontario, Canada, K7L 3N6
(Dated: March 23, 2024)

The standard energy conditions of classical general relativity are applied to FLRW cosmologies containing sudden future singularities. Here we show, in a model independent way, that although such cosmologies can satisfy the null, weak and strong energy conditions, they always fail to satisfy the dominant energy condition. They require a divergent spacelike energy flux in all but the comoving frame.

Recently Barrow [1] has shown that a singularity can occur at a finite future time in an expanding Friedmann-Lemaître-Robertson-Walker (FLRW) universe even when $\rho > 0$ and $+3p > 0$ (and p the comoving energy density and isotropic pressure respectively). This study was motivated by the many recent investigations (see references in [1]) into "big rip" singularities which are precipitated by the "phantom" equation of state

$$+p < 0; \quad (1)$$

Barrow emphasized the fact that (1) is a sufficient but not necessary condition for a singularity in the future of a non-contracting FLRW cosmology by way of introducing sudden future singularities. Because of the possible importance of such singularities they are examined here within the context of the standard energy conditions of classical general relativity. In particular we show that sudden future singularities always fail to satisfy the dominant energy condition. The slightest deviation from the comoving frame results in a divergent spacelike energy flux sufficiently close to any sudden future singularity.

The standard energy conditions of classical general relativity are well known [2] [3] [4]. We pay special attention to the treatment of the cosmological constant in these conditions here since the recent observations of Riess et al. [5] not only confirm earlier reports that we live in an accelerating universe [6] [7], but also sample the transition from deceleration to acceleration. To begin we recall a remarkable feature of four dimensions: the fact that the only divergence free two index tensor A derivable from the metric tensor and its first two derivatives is (up to a disposable multiplicative constant) [8]

$$A = G + \Lambda g; \quad (2)$$

where G is the Einstein tensor and Λ is a finite constant. We write Einstein's equations in the form [9]

$$G + \Lambda g = 8\pi T + 8\pi \Lambda g; \quad (3)$$

where the total energy-momentum tensor consists of the "matter" contribution T and the vacuum contribution Λg . Here we simply assume that the latter is of the form $8\pi \Lambda g$, Λ a finite constant. Since the source of the Robertson-Walker metric can always be

taken (formally) to be a perfect mathematical fluid with comoving velocity field, care must be taken as to how the constants ρ and p are interpreted. Let us write the bare field equations in the form

$$G + C g = 8\pi T; \quad (4)$$

where C is a constant. In terms of the flow (congruence of unit timelike vectors u) and unit normal field n (in the tangent space of the associated Lorentzian two-space) the energy density and isotropic pressure (including bulk viscosity) are given by T_{uu} and $p = T_{nn}$ respectively. Writing the Robertson-Walker metric in the form

$$ds^2 = a^2 \left(\frac{dr^2}{1 - kr^2} + r^2 d\Omega^2 \right) dt^2; \quad (5)$$

where $a = a(t)$ and $d\Omega^2$ is the metric of a unit 2-sphere ($d\Omega^2 = d\theta^2 + \sin^2\theta d\phi^2$), the Friedmann equations follow as

$$8\pi \rho = \frac{3}{a^2} (k + \dot{a}^2) - C; \quad (6)$$

and

$$8\pi p = \frac{1}{a^2} (k + \dot{a}^2 + 2a\ddot{a}) + C; \quad (7)$$

where $\dot{a} = da/dt$.

There are three possible choices for C . (i) $C = \rho - p$: In this procedure the vacuum contribution is extracted from the energy-momentum tensor and considered part of the geometry. The apparent smallness of the effective cosmological constant arises out of the almost perfect cancellation of ρ and p an explanation of which constitutes a standard form of one of the "cosmological constant problems" [10]. This procedure is the common one and, for example, is the procedure used to construct the observers Σ plane [11]. We write the associated energy density and isotropic pressure of the perfect fluid source as ρ and p respectively. These are given by (6) and (7). (ii) $C = \rho$: In this procedure the vacuum contribution is not extracted from the energy-momentum tensor but rather considered part of it. The perfect fluid source now has an associated energy density and isotropic pressure given by $\tilde{\rho} = \rho - \frac{p}{8}$

and $\bar{p} = p + \frac{v}{8}$ respectively. These are given by (6) and (7) with ρ and p replaced by $\bar{\rho}$ and $\bar{p}$. (iii) $C = 0$: In this procedure the geometrical contribution from ρ_c and the contribution from the vacuum ρ_v are both considered part of the energy-momentum tensor. The perfect fluid source now has an associated energy density and isotropic pressure given by $\bar{\rho} = \rho + \frac{v}{8}$ and $\bar{p} = p + \frac{v}{8}$ respectively. Again these are given by (6) and (7) with ρ and p replaced by $\bar{\rho}$ and $\bar{p}$.

We now impose standard energy conditions on the mathematical fluid. The local energy conditions considered here are the null energy condition (NEC), weak energy condition (WEC), strong energy condition (SEC) and dominant energy condition (DEC). In the present context we are always dealing with a formal perfect fluid and in that case the conditions are given in case (i) by

$$\text{NEC} \quad (\rho + p \geq 0); \quad (8)$$

$$\text{WEC} \quad (\rho \geq 0) \text{ and } (\rho + p \geq 0); \quad (9)$$

$$\text{SEC} \quad (\rho + 3p \geq 0) \text{ and } (\rho + p \geq 0); \quad (10)$$

and

$$\text{DEC} \quad (\rho \geq 0) \text{ and } (p \geq 0); \quad (11)$$

where ρ and p are replaced by $\bar{\rho}$ and $\bar{p}$ in case (ii) and by $\bar{\rho}$ and $\bar{p}$ in case (iii). From (6) and (7) for case (i) we find

$$\rho + 3p \geq 0 \quad (\rho \geq 0) \quad C \geq \frac{3a}{a}; \quad (12)$$

$$\rho + p \geq 0 \quad (\rho \geq 0) \quad k + \frac{a^2}{a^2} \geq a; \quad (13)$$

$$p \geq 0 \quad (\rho \geq 0) \quad C \geq \frac{2}{a^2} (k + \frac{a^2}{a^2}) + \frac{a}{a}; \quad (14)$$

and

$$0 \leq (\rho \geq 0) \quad C \geq \frac{3}{a^2} (k + \frac{a^2}{a^2}); \quad (15)$$

again with ρ and p are replaced by $\bar{\rho}$ and $\bar{p}$ in case (ii) and by $\bar{\rho}$ and $\bar{p}$ in case (iii).

A sudden future singularity is defined informally as follows [1]:

$$\begin{aligned} t_s &> 0; \quad 0 < a(t_s) < 1; \\ 0 < \frac{a}{a}(t_s) < 1; \quad a(t \rightarrow t_s) &\rightarrow 1; \end{aligned} \quad (16)$$

(A formal definition can be given in any spacetime which is not Ricci-flat in terms of Ricci invariants [12].) Equations (11) and (14) now make it clear that there is no choice of C for which the DEC is satisfied at a sudden future singularity. We now examine this failure more closely.

Recall that the dominant energy condition states that the energy-momentum tensor $T_{\mu\nu}$ must satisfy

$$T_{\mu\nu} W^\mu W^\nu \geq 0; \quad T_{\mu\nu} W^\mu T^\nu{}_\mu = 0 \quad (17)$$

over the set of all possible timelike 4-vectors W^μ . That is, the local energy density appears non-negative and the local energy flow vector $(T_{\mu\nu} W^\mu)$ is not spacelike. Note that if $T_{\mu\nu}$ decomposes into a perfect fluid (of energy density ρ and isotropic pressure p) with respect to a unit timelike vector field u^μ , then with respect to any unit timelike vector field W^μ we have $T_{\mu\nu} W^\mu W^\nu = (u \cdot W)^2 (\rho + p) + p$ and $T_{\mu\nu} W^\mu T^\nu{}_\mu = (u \cdot W)^2 (\rho^2 - p^2) - p^2$.

For an arbitrary unit timelike 4-vector W^μ and comoving u^μ it follows from (5) that $(W \cdot u)^2 = 1 + w^2$ where $w^2 = \hat{W}_a \hat{W}^a$ and $\hat{W}^a$ is the proper spatial three velocity $(\vec{r}; \vec{v})$, $\vec{v} = \frac{d\vec{r}}{dt}$, the proper time). In case (i) we have $T_{\mu\nu} W^\mu W^\nu = w^2 (\rho + p) + p$ and $T_{\mu\nu} W^\mu T^\nu{}_\mu = w^2 (\rho^2 - p^2) - p^2$ with cases (ii) and (iii) treated in the manner explained above. In all three cases it follows from (17) that for $w^2 \neq 0$ as $t \rightarrow t_s$

$$T_{\mu\nu} W^\mu W^\nu \rightarrow w^2 \left(\frac{a}{4a}\right) \rightarrow +1 \quad (18)$$

and

$$T_{\mu\nu} W^\mu T^\nu{}_\mu \rightarrow +w^2 \left(\frac{a}{4a}\right)^2 \rightarrow +1; \quad (19)$$

We conclude that sudden future singularities defined by (17) violate the dominant energy condition as strongly as possible. The slightest deviation from the comoving frame (any $w^2 \neq 0$) results in a divergent spacelike energy flux as $t \rightarrow t_s$.

Acknowledgments

This work was supported by a grant from the Natural Sciences and Engineering Research Council of Canada. It is a pleasure to thank M. Visser and E. Poisson for helpful remarks.

[*] Electronic Address: lake@astro.queensu.ca

[1] J. D. Barrow, Class. Quantum Grav. 21:179 (2004). (gr-

- qc/0403084) For a somewhat more general consideration and updated references see S.Nojiri and S.D.Odintsov, "The final state and thermodynamics of dark energy universe" (2004). (hep-th/0408170)
- [2] S.W.Hawking and G.F.R.Ellis, *The Large Scale Structure of Space-Time* (Cambridge University Press, Cambridge, 1973)
- [3] M.Visser, *Lorentzian Wormholes* (Springer-Verlag, New York, 1996)
- [4] E. Poisson, *A Relativist's Toolkit: The Mathematics of Black-hole Mechanics* (Cambridge University Press, Cambridge, 2004).
- [5] A. G. Riess et al., *ApJ*, 607, 665 (2004). (astro-ph/0402512)
- [6] A. G. Riess et al., *A. J.*, 116 1009 (1998). (astro-ph/9805201)
- [7] S. Perlmutter et al., *Ap. J.*, 517, 565 (1999). (astro-ph/9812133)
- [8] D.Lovelock, *J.Math.Phys.* 13, 874 (1972).
- [9] We use geometrical units and a signature of + 2.
- [10] See, for example, S.Weinberg astro-ph/9610044 and citations thereto and T.Padmanabhan, *Phys.Rept.* 380, 235 (2003). (hep-th/0212290)
- [11] See, for example, K.Lake, "Cosmological observations and their representation in the M plane" (2004). (astro-ph/0404319)
- [12] In any spacetime which is not Ricci-flat the Ricci invariants of lowest degree are usually taken to be the Ricci scalar R ($R = R^{\mu}_{\mu}$, R the Ricci tensor of the spacetime) and the first Ricci invariant R_1 ($R_1 = S^{\mu\nu} S_{\mu\nu} = 4 S^2$, S the trace-free Ricci tensor $R^{\mu\nu} - \frac{1}{4} R g^{\mu\nu}$). Without any loss in generality we can also take two independent invariants to be I_1 and I_2 defined as follows:
- $$8 I_1 = R = 4 + \text{sign}(S) \sqrt{3 S^2} + C; \quad (20)$$
- and
- $$8 I_2 = R = 4 + \text{sign}(S) \sqrt{S^2 = 3} + C; \quad (21)$$
- where C is a constant. In the coordinates of (5) these evaluate to (6) and (7) respectively, that is $I_1 = \rho$ and $I_2 = p$. A sudden future singularity can be defined formally by the divergence of I_2 while I_1 remains finite. Rather than "sudden" these singularities might be referred to more generally as "pressure dominated".