Hamiltonian and physical Hilbert space in polymer quantum mechanics

Alejandro Corichi, 1, 2, 3, * Tatjana Vukašinac, 4, † and José A. Zapata^{1, ‡}

¹ Instituto de Matemáticas, Unidad Morelia,
Universidad Nacional Autónoma de México, UNAM-Campus Morelia,
A. Postal 61-3, Morelia, Michoacán 58090, Mexico

²Departamento de Gravitación y Teoría de Campos,
Instituto de Ciencias Nucleares, Universidad Nacional Autónoma de México,
A. Postal 70-543, México D.F. 04510, Mexico

³Institute for Gravitational Physics and Geometry, Physics Department,
Pennsylvania State University, University Park PA 16802, USA

⁴Facultad de Ingeniería Civil, Universidad Michoacana
de San Nicolas de Hidalqo, Morelia, Michoacán, Mexico

Abstract

In this paper, a version of polymer quantum mechanics, which is inspired by loop quantum gravity, is considered and shown to be equivalent, in a precise sense, to the standard, experimentally tested, Schrödinger quantum mechanics. The kinematical cornerstone of our framework is the so called polymer representation of the Heisenberg-Weyl (H-W) algebra, which is the starting point of the construction. The dynamics is constructed as a continuum limit of effective theories characterized by a scale, and requires a renormalization of the inner product. The result is a physical Hilbert space in which the continuum Hamiltonian can be represented and that is unitarily equivalent to the Schrödinger representation of quantum mechanics. As a concrete implementation of our formalism, the simple harmonic oscillator is fully developed.

PACS numbers: 04.60.Pp, 04.60.Ds, 04.60.Nc 11.10.Gh.

^{*}Electronic address: corichi@matmor.unam.mx †Electronic address: tatjana@shi.matmor.unam.mx

[‡]Electronic address: zapata@matmor.unam.mx

I. INTRODUCTION

A non-standard and 'exotic' representation of the canonical commutation relations, recently introduced in the context of non-perturbative quantizations of gravity and known as polymer quantum mechanics has been used to explore both mathematical and physical issues in background independent theories such as quantum gravity [1]. A notable example of this type of quantization, when applied to minisuperspace models has given way to what is known as loop quantum cosmology [2]. As in any toy model situation, one hopes to learn about the subtle technical and conceptual issues that are present in full quantum gravity by means of simple, finite dimensional examples. This formalism is not an exception in this regard. Such a quantization is made of several steps. The first one is to build a representation of the Weyl algebra (sometimes also known as the Heisenberg-Weyl algebra) on a Kinematical Hilbert space that is "background independent', and it is sometimes referred to as the polymeric Hilbert space. The second part, the dynamics, deals with the implementation of a Hamiltonian (or Hamiltonian constraint) on this space. In the examples studied so far, the first part is fairly well understood, yielding a kinematical Hilbert space $\mathcal{H}_{\text{poly}}$ that is, however, non-separable. Dynamics has proved to be a bit more difficult, given that a direct implementation of the Hamiltonian H on \mathcal{H}_{poly} is not possible since one of the main features of this quantum mechanics is that the operators \hat{q} and \hat{p} cannot be both simultaneously defined (nor its analogues in theories involving more elaborate variables). Thus, any operator that involves (powers of) the not defined variable has to be regulated by a well defined operator. However, the regulator can not be removed and one is left with the ambiguity that is present in any regularization. The freedom in choosing it can be sometimes associated with a length scale, as is the case of a simple harmonic oscillator, where the freedom can be cast in terms a length scale associated with the 'discreteness of space'. In the standard treatment of that system, it has been argued that if this length scale is taken to be 'sufficiently small', one can arbitrarily approximate standard Schrödinger quantum mechanics [1].

In this paper we shall adopt a different viewpoint. The new viewpoint is that any scale that one chooses for defining the Hamiltonian should be seen as providing an effective description at the given scale. A natural question is what happens when we change the scale and go to even smaller 'distances'. Can we define consistency conditions between these scales? Or even better, can we take the limit and find thus a continuum limit? As we shall see in this paper, the answer to both questions is in the affirmative.

In the remainder of the introduction, let us pose the question with a bit of more precision. The kinematical Hilbert space $\mathcal{H}_{\text{poly},x}$ is the completion of the vector space Cyl_x which is finitely generated by the family of functions $\{\delta_{x_0}\}_{x_0\in\mathbb{R}}$ (where $\delta_{x_0}(x)=0$ if $x\neq x_0$ and $\delta_{x_0}(x)=1$ if $x=x_0$) with the inner product that makes this family an orthonormal basis. That is, the "polymer representation" of the Heisenberg-Weyl algebra in $\mathcal{H}_{\text{poly},x}$ [1], is such that the operator \hat{x} acts by multiplication and the generator of translations $\hat{V}(\lambda)$ acts as: $\hat{V}(\lambda) \cdot \Psi(x) = \Psi(x-\lambda)$. The inner product on the basis is given by: $\langle \delta_{x_0}(x), \delta_{x_1}(x) \rangle = \delta_{x_0,x_1}$. A state $\Psi \in \mathcal{H}_{\text{poly},x}$ is of the form $\Psi(x) = \sum_i \psi_i \, \delta_{x_i}(x)$ with $\sum_i |\psi_i|^2 < \infty$. As we have already mentioned, the dynamics is problematic in this framework; not even the Hamiltonian

¹ In loop quantum cosmology, the minimum area gap of the full quantum gravity theory imposes such a scale [2].

² This corresponds to the 'position representation' of the polymeric quantum mechanics. In most treatments, the Hilbert space is described in the 'momentum representation' where the basis is given by quasiperiodic

of the simple harmonic oscillator can be represented inside $\mathcal{H}_{\text{poly},x}$. This is the case given that the operator \hat{p} (the generator of translations in x) is not well defined in $\mathcal{H}_{\text{poly},x}$, so any observable that depends on finite powers of p is not well defined as such. In this regard this simplified quantum mechanical system captures the essence of loop quantum gravity, including the mathematical origin of the non triviality of the dynamics. Recall that in the case of LQG, the curvature of the connection –that appears in the constraints– is not a well defined operator so one has to approximate it by a holonomy.

In this paper we construct a physical Hilbert space and a Hamiltonian operator on it as a continuum limit of effective theories. That is, we import the proposal of constructing the dynamics of loop quantized theories as a continuum limit of effective theories developed in [3] to this model structure for quantum mechanics.

As an example of the continuum limit here presented, the case of the simple harmonic oscillator is fully developed. We make extensive use of previous results by Ashtekar, Fairhurst and Willis [1]. However, the viewpoint here advocated is different from the viewpoint of [1], where the authors already had evidence of "some convergence with standard quantum mechanics" (see also [4] for a different perspective). The question that these papers have tried to address is rather simple: Can polymer quantum mechanics approximate the standard representation of quantum mechanics? This question implicitly comes with the next question: What is the meaning of approximating a quantum theory? In a sense the difference between [1], [4] and this work deals with the different answers given to the later question. The viewpoint here adopted, as we shall elaborate throughout the paper, is based on the construction of a continuum theory as an appropriately defined limit. This theory will turn out to be equivalent to the standard Schrödinger quantum mechanics. Our procedure involves, as intermediate steps, the construction of effective theories at different scales, that on themselves can be seen as an approximation (given certain tolerances) of the continuum theory. When we focus attention to one given scale our treatment bears some resemblances to the studies presented in [1] and [4].

The strategy for constructing the continuum theory starting from the polymer representation will involve three steps. In the first one we choose to view the Hamiltonian as a quadratic form, which implies that it is naturally coarse grained by a map relating different scales, and that its continuum limit will be a projective limit. We should explicitly mention that the matching of effective theories at different scales that is behind the continuum limit involves coarse graining as well as rescaling. The second step corresponds to working at one given scale, but replacing the effective theory at that scale by better theories that know about degrees of freedom from finer scales and that are generated by coarse graining and rescaling the effective theory of a finer scale. If at a given scale these microscopically corrected theories converge, the limit is called a completely renormalized theory at that scale. In the last step the completely renormalized theories at all the scales are pasted together (they satisfy a compatibility condition) to construct the continuum limit. A product of this construction is the physical Hilbert space. This space turns out to be equivalent to the ordinary L^2 space of the Schrödinger theory.

The structure of the paper is as follows. In Sec. II we introduce the necessary steps for defining the notion of scale and the continuum limit at the kinematical level. Section III is devoted to the dynamics, namely to the Hamiltonian. In this section the physical Hilbert space is constructed. In Sec. IV we show in detail how the program outlined in the previous

functions and the inner-product is given by the Haar measure on the Bohr compactification of \mathbb{R} [2].

sections is implemented in the case of the harmonic oscillator. We conclude the paper with a discussion in Sec. V.

II. EFFECTIVE THEORIES, COARSE GRAINING AND CONTINUUM LIMIT: KINEMATICS

A central object of the kinematics is the Hilbert space $\mathcal{H}_{\text{poly},x}$; it will be the starting point for our formalism. In this section we shall see how our notions of effective theories at a given scale, the coarse graining maps that relate the different effective theories and the continuum limit are compatible with $\mathcal{H}_{\text{poly},x}$.

The role of scales will be played by decompositions of \mathbb{R} as a disjoint union of closed-open intervals.

Definition 1 (Scale) In our context, a scale C is a decomposition of the real line of the form

$$\mathbb{R} = \bigcup_{\alpha_i \in C} \, \alpha_i,$$

where $\alpha_i = [L(\alpha_i), R(\alpha_i))$, and the vertex set $\{L(\alpha_{i+1}) = R(\alpha_i)\}_{\alpha_i \in C}$ is required not to have any accumulation point.

Recall that $\mathcal{H}_{\text{poly},x}$ is the completion of Cyl_x , which is finitely generated by the basis given by $\{\delta_{x_0}(x)\}$, consisting of all Kronecker deltas for $x_0 \in \mathbb{R}$. A scale C defines an equivalence relation \sim_C in Cyl_x ; we define this relation on one basis and extend it by linearity to the whole space. We declare

$$\delta_{x_0} \sim_C \delta_{y_0} \iff \exists \alpha_i \in C : x_0, y_0 \in \alpha_i.$$

To every scale C we associate a space of states at scale C defined as the vector space of equivalence classes

$$Cyl(C) = Cyl_x/\sim_C$$
.

This vector space is finitely generated by an orthonormal basis $\{e_{\alpha}\}_{{\alpha}\in C}$ labeled by the cells of C.³ The completion of Cyl(C) is the Hilbert space \mathcal{H}_C .

We will also work with the dual space \mathcal{H}_{C}^{\star} . Since the space of states at any given scale is a Hilbert space, its dual is isomorphic to it, but for us the dual space will be of special interest. Notice that its elements have a natural \sim_{C} -preserving action on $Cyl(\mathbb{R})_{x}$. This action is particularly simple to see for the elements of the dual basis $\{\omega_{\alpha}\}_{{\alpha}\in C}$; for them we have

$$\omega_{\alpha}(\delta_{x_0}) = \chi_{\alpha}(x_0)$$

where χ_{α} is the characteristic function of the set $\alpha \subset \mathbb{R}$. Thus, we write

$$\mathcal{H}_C^{\star} \subset Cyl(C)^{\star}$$

where we have noticed that $Cyl(C)^*$ can be thought of as the \sim_C -preserving subspace of $Cyl(\mathbb{R})_x^*$.

³ There is a natural projection $P_C: Cyl_x \to Cyl(C)$ and the inner product in \mathcal{H}_C can be pulled back from $\mathcal{H}_{poly,x}$ by any section. The end result is that the basis $\{e_{\alpha}\}$ becomes orthonormal: $(e_{\alpha}, e_{\beta})_C = \delta_{\alpha,\beta}$.

Definition 2 (Coarse graining) Given two scales we write $C_a \leq C_b$ and say that C_a is a coarse graining of C_b (or C_b is a refinement of C_a) if any interval $\alpha_i \in C_a$ is a finite union of intervals of C_b .

Our coarse graining maps work by decimation. If we have two scales related by refinement $C_a \leq C_b$ our decimation map will be defined to be the injective isometry $d: \mathcal{H}_{C_a} \to \mathcal{H}_{C_b}$ characterized by

$$d(e_{\alpha}) = e_{\beta} \iff L(\alpha) = L(\beta).$$

It is important to notice that if $C_a \leq C_b \leq C_c$ the corresponding d-triangle diagram commutes.

On the other hand, $d^*: \mathcal{H}_{C_b}^* \to \mathcal{H}_{C_a}^*$ sends part of the elements of the dual basis to zero while keeping the information of the rest: $d^*(\omega_\beta) = \omega_\alpha$ if $L(\alpha) = L(\beta)$; if there is no interval $\alpha \in C_a$ with that property $d^*(\omega_\beta) = 0$.




FIG. 1: (a)
$$d(\sum_{i=1}^{3} e_{\alpha_i}) = e_{\beta_1} + e_{\beta_3} + e_{\beta_4}$$
 (b) $d^*(\sum_{i=1}^{5} \omega_{\beta_i}) = \sum_{j=1}^{3} \omega_{\alpha_j}$

Now we mention two properties that will be central when defining a continuum limit.

- 1. Apart from being partially ordered, the set of scales is directed towards refinement. Any two scales C_a , C_b have a common refinement, $C_c \ge C_a$ and $C_c \ge C_b$. (In particular, the decomposition consisting of the intersection of the intervals of C_a and C_b defines a scale that is finer than the original pair of scales.)
- 2. The set of scales satisfies the following property of *infinite refinement*. Given any open set $U \subset \mathbb{R}$ there is a scale C which has an interval completely contained in it, $\alpha_i \in C$ and $\alpha_i \subset U$.

Our objective now is to introduce a construction that will let us define the limit of \mathcal{H}_C as the scale C gets infinitely fine. At a given scale C the corresponding Hilbert space

 \mathcal{H}_C contains "states that come from coarser scales" $C_0 \leq C$ using the decimation map $d: \mathcal{H}_{C_0} \to \mathcal{H}_C$. It is clear that if C_0 is not equal to C then there are states in \mathcal{H}_C that do not come from states in \mathcal{H}_{C_0} . Also those states of the form $d(\phi_{C_0})$ may "come from scales even coarser than C_0 ", since it could happen that there is a scale $C_0' \leq C_0$ and a state at this scale such that $\phi_{C_0} = d(\phi_{C_0'})$. That is, the state $\phi_{C_0} \in \mathcal{H}_{C_0}$ and the state $\phi_{C_0'} \in \mathcal{H}_{C_0'}$ (such that $\phi_{C_0} = d(\phi_{C_0'})$) agree when they are brought to scale C by the corresponding decimation maps. The systematic construction that organizes this repeated information is the following. Given a state at any scale coarser than C, say $\phi_{C_0'} \in \mathcal{H}_{C_0'}$ with $C_0' \leq C$, we assign to it a collection consisting of one state at every scale C_0 that is finer than C_0' coarser than C, $\{\phi_{C_0} = d(\phi_{C_0'}) \in \mathcal{H}_{C_0}\}_{C_0' \leq C_0 \leq C}$. In addition we must regard two such collections as equivalent if they agree at the finer scales; that is, the collection written above is equivalent to $\{\psi_{C_0} = d(\psi_{C_0''}) \in \mathcal{H}_{C_0}\}_{C_0' \leq C_0 \leq C}$ if and only if there is a scale C_1 such that $C_0' \leq C_1$, $C_0'' \leq C_1$ and for any scale $C_2 \geq C_1$ we have $d(\psi_{C_0''}) = d(\phi_{C_0'}) \in \mathcal{H}_{C_2}$. The set of such equivalence classes of collections can be endowed with the structure of a vector space and this vector space is isomorphic to \mathcal{H}_C itself. The corresponding map, I_C is given by

$$I_C(e_{\alpha_i,C}) = \{(C_0, e_{\beta_j,C_0}) | C_0 \le C\}$$

where the scale C_0 contains an interval β_j such that $L(e_{\alpha_i,C}) = L(e_{\beta_j,C_0})$. Moreover, since the decimation maps d are isometries the set of equivalence classes of collections can be made a Hilbert space.

One can notice that the scale C on the collections described above only plays the role of a cut-off allowing only scales coarser than it. We can work with collections that do not have any cut-off by simply replacing the scale C by the symbol \mathbb{R} . Again we can define an equivalence relation and it can be given the structure of a Hilbert space. This is formalized in the following definition.

Definition 3 (Continuum limit of \mathcal{H}_C) Our decimation map d gives a meaning to the limit of \mathcal{H}_C as the scale gets finer and finer. In this case the continuum limit is called a direct limit and it is denoted by

$$\overrightarrow{\mathcal{H}}_{\mathbb{R}}:= \stackrel{\longrightarrow}{\lim}_{C \to \mathbb{R}} \mathcal{H}_{C}.$$

It is defined to be the complete vector space generated by collections of the type

$$\phi_{\mathbb{R}} := \{ (C, \phi_C \in \mathcal{H}_C) | C \ge C_0 \text{ and } \phi_C = d \phi_{C_0} \},$$

for some scale C_0 , and where two such collections are identified if they coincide for all $C \geq C'_0$ for some scale C'_0 . Addition and multiplication by scalars is defined "component by component", and the inner product is inherited from any of the involved scales because the decimation maps are isometries. ⁴

Notice that due to the absence of a cut-off, that allows only certain scales, we have to include a clause to guarantee the completeness of the Hilbert space. The effect of such a clause is to add states that have the form of infinite linear combinations of states that "have their origin at infinitely many distinct given scales". Let us then pose the following result:

⁴ This is exactly the same mathematical structure appearing in the definition of cylindrical functions. In the collection written above C_0 plays the role of the graph Γ labelling the cylindrical function ψ_{Γ} .

Theorem 1 $\overrightarrow{\mathcal{H}}_{\mathbb{R}}$ is naturally isomorphic to $\mathcal{H}_{\mathrm{poly},x}$.

To prove this theorem we simply exhibit the isomorphism $I: \mathcal{H}_{\text{poly},x} \to \overrightarrow{\mathcal{H}}_{\mathbb{R}}$,

$$I(\delta_{x_0}) = \{ (C, e_{\alpha_{x_0}} \in \mathcal{H}_C) | C \ge C_{x_0} \},$$

where the decomposition C_{x_0} contains an interval for which x_0 is its left extreme point, and where the interval $\alpha_{x_0} \in C$ is the one that contains x_0 .

Definition 4 (Continuum limit of \mathcal{H}_C^*) Also d^* gives a meaning to the continuum limit of \mathcal{H}_C^* . Since in this case the limit goes in the direction opposite to d^* the limit is called an inverse limit (or projective limit) and denoted by

$$\stackrel{\longleftarrow}{\mathcal{H}}_{\mathbb{R}}^{\star} := \stackrel{\longleftarrow}{\lim}_{C \to \mathbb{R}} \mathcal{H}_{C}^{\star}.$$

It is defined to be the complete vector space generated by collections

$$\Phi_{\mathbb{R}} := \{ \Phi_C \in \mathcal{H}_C^{\star} \}$$

which satisfy the following two properties:

- 1. It is d^{\star} -compatible, $d_{a,b}^{\star}(\Phi_{C_b}) = \Phi_{C_a}$ whenever $C_a \leq C_b$.
- 2. The norm of its elements converges, $\lim_{C\to\mathbb{R}} \|\Phi_C\|_C$ exists.

Notice that the first property guarantees that the elements of $\mathcal{H}_{\mathbb{R}}^{\star}$ have a natural action on $Cyl(\mathbb{R})$ given by

$$\Phi_{\mathbb{R}}(\delta_{x_0}) := \lim_{C \to \mathbb{R}} \Phi_C(\delta_{x_0}).$$

The space that we have just described is naturally isomorphic to $\mathcal{H}^{\star}_{\text{poly},x}$. The natural isomorphism $I^{\star}: \stackrel{\leftarrow}{\mathcal{H}}_{\mathbb{R}}^{\star} \to \mathcal{H}^{\star}_{\text{poly},x}$ is given by

$$I^{\star}(\omega_{x_0}) = (\delta_{x_0}, \cdot)_{\text{poly}},$$

where the symbol ω_{x_0} is a short hand for the collection

$$\omega_{x_0} := \{ \sum_{\alpha \in C} \delta_{x_0}(L(\alpha)) \omega_{\alpha} \}.$$

To summarize we can write

$$\stackrel{\longleftarrow}{\mathcal{H}_{\mathbb{R}}^{\star}} \simeq \mathcal{H}_{\mathrm{poly},x}^{\star} \subset Cyl(\mathbb{R})_{x}^{\star}.$$

This concludes our construction of the kinematical structure needed for the second step, namely for the definition of dynamics.

III. EFFECTIVE THEORIES, COARSE GRAINING AND CONTINUUM LIMIT: DYNAMICS

As mentioned in the introduction we will import the proposal of constructing the dynamics of loop quantized theories as a continuum limit of effective theories developed in Ref [3] to this model structure for quantum mechanics. The basic ideas of this approach are presented and developed for the quantum field theories in the mentioned article, here we give a self-contained treatment for the case of quantum mechanics. As expected in the case of quantum mechanics, the procedure is considerably simplified; only the inner product will have to be renormalized, but not the coupling constants.

For convenience we will use a smaller auxiliary set of scales defined by regular decompositions $\{C_n\}$. We start with a regular decomposition C_0 with intervals of length a_0 , and generate a totally ordered family of decompositions $\{C_n\}$ from it by regularly subdividing its intervals repeatedly; then C_n 's intervals have length $\frac{a_0}{2^n}$.

The first ingredient is a regularization of the Hamiltonian as a self-adjoint operator at each scale. With the aid of the inner product, these Hamiltonians will be treated as quadratic forms; the quadratic forms that give its expectation value on normalized states $\lambda_{C_n}^2(\psi, h_n\psi)_{C_n} = h_n(\psi)$

$$h_n:\mathcal{H}_{C_n}\to\mathbb{R}.$$

The normalization factors in the inner product $\lambda_{C_n}^2 \in \mathbb{R}^+$ have to be adjusted in such a way that, at least in the continuum limit, the Hamiltonians of different scales are pasted correctly by the decimation maps⁵.

At a given scale C_m we can "include the effects of more microscopic degrees of freedom" by using our decimation map. When $C_m \leq C_n$ we define

$$h_{m(n)} := d_{m,n}^{\star} h_n. \tag{1}$$

That is, $h_{m(n)}(e_{\alpha_i}) := h_n(d_{m,n}e_{\alpha_i})$. If these microscopically corrected Hamiltonians converge, their limit will be called a completely renormalized Hamiltonian at the given scale h_m^{ren} : $\mathcal{H}_{C_m} \to \mathbb{R}$,

$$h_m^{\text{ren}} := \lim_{C_n \to \mathbb{R}} h_{m(n)}. \tag{2}$$

By construction, when the completely renormalized Hamiltonians exist, they are compatible with each other in the sense that

$$d_{m,n}^{\star}h_n^{\text{ren}} = h_m^{\text{ren}}.$$

A collection of compatible Hamiltonians defines in itself a continuum limit Hamiltonian. We will construct a Hilbert space of physical states after we study the issue of convergence stated in Eq. (2) in some detail.

We will work with effective Hamiltonians that have a purely discrete spectrum (labelled by ν) $h_n \cdot \Psi_{\nu,C_n} = E_{\nu,C_n} \Psi_{\nu,C_n}$ (if we see the effective Hamiltonians h_n as quadratic forms

⁵ In standard introductions the continuum limit in the renormalization group framework includes a wave function renormalization; here we choose the equivalent action of renormalizing the inner product instead. We choose to absorb the normalization factors in the inner product to find a non trivial action of the completely renormalized Hamiltonian in $\mathcal{H}_{\text{poly},x}$ (6).

the condition is that they have isolated critical points Ψ_{ν,C_n}). We shall also introduce, as an intermediate step, a cut-off in the energy levels. Thus, we can write

$$h_m^{\nu_{\text{cut-off}}} = \sum_{\nu=0}^{\nu_{\text{cut-off}}} E_{\nu,C_m} \Psi_{\nu,C_m} \otimes \Psi_{\nu,C_m} , \qquad (3)$$

where the eigen covectors $\Psi_{\nu,C_m} \in \mathcal{H}_{C_m}^{\star} \subset Cyl(\mathbb{R})_x^{\star}$ are normalized according to the inner product rescaled by $\frac{1}{\lambda_{C_n}^2}$, and the cut-off can vary up to a scale dependent bound, $\nu_{\text{cut-off}} \leq \nu_{\text{max}}(C_m)$. The origin of this cut-off is in the regularization which models the Hamiltonian of our system at a given scale with a Hamiltonian of a periodic system in a regime of small energies. Note that in the polymer representation the operator of momentum \hat{p} is not defined, so it has to be approximated using the translation operator, that is well defined. As a result of this approximation the standard kinetic term $p^2/2m$ is replaced by a periodic function of p.

In the presence of a cut-off, the convergence of the microscopically corrected Hamiltonians, equation (2) is equivalent to the existence of the following two limits. The first one is the convergence of the energy levels,

$$\lim_{C_n \to \mathbb{R}} E_{\nu, C_n} = E_{\nu}^{\text{ren}} \,. \tag{4}$$

Second is the existence of the completely renormalized eigen covectors,

$$\lim_{C_n \to \mathbb{R}} d_{m,n}^{\star} \Psi_{\nu,C_n} = \Psi_{\nu,C_m}^{\text{ren}} \in \mathcal{H}_{C_m}^{\star} \subset Cyl_x^{\star}. \tag{5}$$

When the completely renormalized eigen covectors exist, they form a collection that is d^* -compatible.

In addition one could ask whether the eigen covectors $\Psi_{\nu,C_m}^{\text{ren}}$ converge in the continuum limit as elements of $Cyl(\mathbb{R})_x^*$. We can choose the normalization factors $\lambda_{C_n}^2$ in a way that that the renormalized eigen covectors converge point-wise when acting on cylindrical functions δ_{x_0} when the point x_0 is a vertex of one of the intervals at some scale C_n . In most cases the dynamics leads to completely renormalized eigen covectors that are continuous in the sense that they also converge point-wise when acting on other cylindrical functions. In these cases, the collection of d^* -compatible Hamiltonians can be extended to act on $\mathcal{H}_{\text{poly},x}$. This Hamiltonian (with cut-off) $h_{\mathbb{R}}^{\nu_{\text{cut-off}}\text{ren}}: \mathcal{H}_{\text{poly},x} \to \mathbb{R}$ in the continuum is defined by,

$$h_{\mathbb{R}}^{\nu_{\text{cut-offren}}}(\delta_{x_0}) := \lim_{C_n \to \mathbb{R}} h_n^{\nu_{\text{cut-offren}}}([\delta_{x_0}]_{C_n}). \tag{6}$$

Notice that the domain of this Hamiltonian is $\mathcal{H}_{\text{poly},x}$ and not only Cyl_x ; this happens even when the domain of the $\Psi_{\nu,C_m}^{\text{ren}}$ is only $Cyl_x \subsetneq \mathcal{H}_{\text{poly},x}$ because the quadratic nature of the Hamiltonian improves convergence, so we can extend the action from Cyl_x to $\mathcal{H}_{\text{poly},x}$. From now on, we will assume that the limit in equation (6) exists for any $x_0 \in \mathbb{R}$.

This continuum limit Hamiltonian can be coarse grained to any scale $d^*h^{\nu_{\text{cut-off}ren}}_{\mathbb{R}}:\mathcal{H}_C\to\mathbb{R}$. Clearly, for a regular decomposition we have $d^*h^{\nu_{\text{cut-off}ren}}_{\mathbb{R}}=h^{\nu_{\text{cut-off}ren}}_n$.

⁶ In the case of the harmonic oscillator we shall see in detail that the system can be described in terms of a periodic potential, namely, that of a pendulum.

Now we return to the construction of a Hilbert space of physical states. Recall that an inner product renormalization was used to obtain our results concerning the continuum limit. Had we insisted on using normalized eigen covectors $\{\frac{\Psi_{\nu,C_n}}{||\cdot||}\}$ according to an inner product which did not have the normalization factors, we would have found that the limit in equation (5) did not exist. In this case the norm of the eigen covectors diverge as $C_n \to \mathbb{R}$.

A choice of normalization factors that leads to convergence in (2) for "standard systems" is $\lambda_{C_n}^2 = 2^n$, which means that the renormalized inner product in \mathcal{H}_n^* is

$$(\omega_{\alpha_i}, \omega_{\alpha_j})_{C_n}^{\text{ren}} = \frac{1}{2^n} \delta_{ij}. \tag{7}$$

The Hilbert space of covectors together with such inner product will be called $\mathcal{H}_{C_n}^{\star \mathrm{ren}}$.

The sequence of compatible renormalized eigen covectors $\{\Psi^{\text{ren}}_{\nu,C_n}\}$ does define an element of $\mathcal{H}_{\mathbb{R}}^{\star \text{ren}}$, which is the projective limit of the renormalized spaces of covectors

$$\{\Psi_{\nu,C_n}^{\text{ren}}\} \in \mathcal{H}_{\mathbb{R}}^{\star \text{ren}} = \lim_{C_n \to \mathbb{R}} \mathcal{H}_{C_n}^{\star \text{ren}}.$$
 (8)

The inner product in this space is defined by

$$(\{\Psi_{C_n}\}, \{\Phi_{C_n}\})_{\mathbb{R}}^{\text{ren}} := \lim_{C_n \to \mathbb{R}} (\Psi_{C_n}, \Phi_{C_n})_{C_n}^{\text{ren}}.$$

This inner product is degenerate. Collections of the form

$$\omega_{x_0} := \{ \sum_{\alpha_i \in C_r} \delta_{x_0}(L(\alpha_i)) \omega_{\alpha_i} \}$$

have zero norm, since there is at most one term contributing. While collections corresponding to characteristic functions of intervals that fit inside some decomposition $\alpha_i \in C_m$

$$\tilde{\chi}_{\alpha_i} := \{0 \text{ if } C_n \leq C_m, \sum_{\alpha_j \in C_n} \chi_{\alpha_i}(L(\alpha_j))\omega_{\alpha_j} \text{ if } C_m \leq C_n \}$$

now belong to the Hilbert space and their squared norm is the length of α_i in units of a_0 .

Given that the inner product identifies many states, the natural definition of the Hilbert space of physical states is to take the quotient by the zero norm states, namely:

Definition 5 (Hilbert space of physical states)

$$\begin{split} \mathcal{H}^{\star}_{\mathrm{phys}} := & \stackrel{\longleftarrow}{\mathcal{H}}^{\star \mathrm{ren}}_{\mathbb{R}} / \ker(\cdot, \cdot)^{\mathrm{ren}}_{\mathbb{R}} \\ \mathcal{H}_{\mathrm{phys}} := & \mathcal{H}^{\star \star}_{\mathrm{phys}} \end{split}$$

Theorem 2 The Hilbert space of physical states is naturally unitarily isomorphic to the space of square integrable functions with respect to the Lebesgue measure

$$\mathcal{H}_{\text{phys}} \simeq L^2(\mathbb{R}, \mathrm{d}x)$$

⁷ "Standard systems" include those with classical Hamiltonian of the form $H = p^2/2m + V(x)$.

Proof. We will define a unitary anti-isomorphism (invertible antilinear map that preserves the inner product) from $L^2(\mathbb{R}, dx) \to \mathcal{H}^{\star}_{phys}$.

Consider in $L^2(\mathbb{R}, dx)$ elements of the form $\Psi = \sum_{\alpha} \Psi(\alpha) [\chi_{\alpha}]_{L^2}$, where the sum runs over finitely many closed-open intervals and χ_{α} is the characteristic function of the corresponding interval. The subset of elements of this form is dense in L^2 . Thus, our unitary anti-isomorphism is defined by assigning $\Psi = \sum_{\alpha} \bar{\Psi}(\alpha) [\chi_{\alpha}]_{\text{phys}}$ to those elements. \square

Notice that we could have worked with the set of all the scales C (not necessarily corresponding to regular decompositions) using the corresponding renormalized inner products for them and we would have obtained the same result. The use of our auxiliary set of scales corresponding to regular decompositions simply selects a dense subset of the physical Hilbert space.

Now we return to the definition of the Hamiltonian in the continuum limit. We can use the renormalized inner product to induce an action of the cut-off Hamiltonians on $\mathcal{H}_{\mathbb{R}}$

$$h_{\mathbb{R}}^{\nu_{\text{cut-off}}\text{ren}}(\{\Psi_{C_n}\}) := \lim_{C_n \to \mathbb{R}} h_n^{\nu_{\text{cut-off}}\text{ren}}((\Psi_{C_n}, \cdot)_{C_n}^{\text{ren}}),$$

where we have used the fact that $(\Psi_{C_n}, \cdot)_{C_n}^{\text{ren}} \in \mathcal{H}_{C_n}$. The existence of this limit is trivial because the renormalized Hamiltonians are finite sums and the limit exists term by term.

These cut-off Hamiltonians descend to the physical Hilbert space

$$h_{\mathbb{R}}^{\nu_{\mathrm{cut-off}}\mathrm{ren}}([\{\Psi_{C_n}\}]) := h_{\mathbb{R}}^{\nu_{\mathrm{cut-off}}\mathrm{ren}}(\{\Psi_{C_n}\})$$

for any representative $\{\Psi_{C_n}\}\in [\{\Psi_{C_n}\}]\in \mathcal{H}_{phys}^{\star}$. The context will prevent any confusion with the notation.

Finally we can address the issue of removal of the cut-off. The Hamiltonian $h_{\mathbb{R}}^{\mathrm{ren}} : \stackrel{\longleftarrow}{\mathcal{H}_{\mathbb{R}}}^{\mathrm{ren}} \to \mathbb{R}$ is defined by the limit

$$h_{\mathbb{R}}^{\mathrm{ren}} := \lim_{\nu_{\mathrm{cut-off}} \to \infty} h_n^{\nu_{\mathrm{cut-off}}}$$
ren

when the limit exists. Its corresponding Hermitian form in \mathcal{H}_{phys} is defined whenever the above limit exists.

Moreover, we can construct $h_{\mathbb{R}}^{\text{ren}} : \mathcal{H}_{\mathbb{R}}^{\text{ren}} \to \mathbb{R}$ without introducing an auxiliary cut-off. We can simply define

$$h_{\mathbb{R}}^{\text{ren}}(\{\Psi_{C_n}\}) := \lim_{C_n \to \mathbb{R}} h_n((\Psi_{C_n}, \cdot)_{C_n}^{\text{ren}})$$

$$\tag{9}$$

and arrive to the same Hamiltonian in the continuum as the previous equation. On the other hand, the cut-off can not be removed when working on $\mathcal{H}_{\text{poly},x}$, or at a given scale; the completely renormalized Hamiltonians can not be liberated from their cut-off given that, as we explained earlier, at any given scale the cut-off is nedeed in order to approximate the given quantum mechanical system by a periodic one. There is no solution to this "problem" but, as we will show, it is really not a problem, but simply an issue related to the domain of the Hamiltonian.

With the use of the decimation map, the elements of \mathcal{H}_{C_m} have an action as linear forms on $\mathcal{H}_{\mathbb{R}}$

$$e_{\alpha_i}(d_{m,n}^{\star}(\{\Psi_{C_n}\})) := e_{\alpha_i}(\Psi_{C_m}).$$

Also, the elements of $\mathcal{H}_{\text{poly},x}$ act on $\mathcal{H}_{\mathbb{R}}$

$$\delta_{x_0}(\{\Psi_{C_n}\}) := \lim_{C_n \to \mathbb{R}} [\delta_{x_0}]_{C_n}(\Psi_{C_n}).$$

If any such action is restricted to smooth functions with compact support, $C_0^{\infty} \subset \mathcal{H}_{\mathbb{R}}^{-}$, it becomes the standard action of Dirac's delta functional. The natural inclusion of C_0^{∞} in $\mathcal{H}_{\mathbb{R}}^{-}$ is by an antilinear map which assigns to any $\Psi \in C_0^{\infty}$ the d^* -compatible collection $\Psi_{C_n}^{\mathrm{shad}} := \sum_{\alpha_i} \omega_{\alpha_i} \bar{\Psi}(L(\alpha_i)) \in \mathcal{H}_{C_n}^{\mathrm{*ren}} \subset Cyl_x^{\mathrm{*}}; \ \Psi_{C_n}^{\mathrm{shad}}$ will be called the shadow of Ψ at scale C_n and acts in Cyl_x as a piecewise constant function. Clearly other types of test functions like Schwartz functions are also naturally included in $\mathcal{H}_{\mathbb{R}}^{-}$. This natural inclusion map is compatible with the isomorphism of Theorem 2.

When \mathcal{H}_{C_m} or $\mathcal{H}_{\text{poly},x}$ are seen from point of view of the natural home of the dynamics in the continuum they correspond to highly distributional objects which are not in the domain of the Hamiltonians with the standard kinetic term. For example, the calculation of equation (6) corresponds, in the Schrödinger representation, to the expectation value of the cut-off Hamiltonian on $\delta_{x_0}^{\text{Dirac}}$.

Notice that equation (9) leads to a very compact version of the dynamics of polymeric quantum mechanics. We can say that a dynamics in this framework is characterized by a collection of effective Hamiltonians such that the limit $h_{\mathbb{R}}^{\text{ren}}$ in (9) exists for all the elements of a dense domain, $[\{\Psi_{C_n}\}]_{\text{phys}} \in D_{\text{phys}} \subset \mathcal{H}_{\text{phys}}^{\star}$.

Now we can compare with the standard quantum mechanics resulting from the Schrödinger representation, where the dynamics is taken to be defined by a Hamiltonian quadratic form h_{Schr} defined on a dense domain $[\Psi]_{L^2} \in D_{L^2} \subset L^2(\mathbb{R}, dx)$. These two dynamics are equivalent if the unitary anti-isomorphism of Theorem 2 identifies the domains of definition and the Hamiltonian quadratic forms. In the next section we will explicitly see that this is indeed the case for the simple harmonic oscillator.

We have completed the main result of this paper, namely to construct a physical Hilbert space which turns out to be isomorphic to that of the standard Schrödinger representation. In the next section, we shall examine a paradigmatic system that has been the subject of extensive study, namely the simple harmonic oscillator.

IV. SIMPLE HARMONIC OSCILLATOR

In this section, let us consider the example of a Simple Harmonic Oscillator (SHO) with parameters m and ω , classically described by the following Hamiltonian

$$H = \frac{1}{2m} p^2 + \frac{1}{2} m \omega^2 x^2.$$

Recall that from these parameters one can define a length scale $D=\sqrt{\hbar/m\omega}$. In the standard treatment one uses this scale to define a complex structure J (and an inner product from it) that uniquely selects the standard Schrödinger representation. In our case, since the operator

$$\hat{V}(\mu) = e^{i\mu\hat{p}/\hbar}$$

is not weakly continuous in μ in the polymeric representation, the corresponding would-be self-adjoint momentum operator \hat{p} does not exist. As a starting point we take the suggestion of Ashtekar, Fairhurst and Willis who define (regulate) \hat{p} , at a given scale C_n , as [1],

$$\hat{p} := \frac{i\hbar \, 2^{n-1}}{a_0} \left[\hat{V} \left(\frac{a_0}{2^n} \right) - \hat{V} \left(-\frac{a_0}{2^n} \right) \right] \,. \tag{10}$$

However, the operator \hat{p}^2 that appears in the Hamiltonian, is then defined not as the square of (10) but rather as,

$$\hat{p^2} = \frac{\hbar^2 \, 2^{2n}}{a_0^2} \left[2 - \hat{V} \left(\frac{a_0}{2^n} \right) - \hat{V} \left(-\frac{a_0}{2^n} \right) \right] \,. \tag{11}$$

As they note this election is not unique and is guided by the requirement of non-degenerate energy levels (that makes it preferred compared with $(\hat{p})^2$). Then, the Hamiltonian \hat{H}_{C_n} : $\mathcal{H}_{C_n} \to \mathcal{H}_{C_n}$ at scale C_n , is given by,

$$\hat{H}_{C_n} = \frac{\hbar^2}{2m} \left(\frac{2^n}{a_0}\right)^2 \left[2 - \hat{V}\left(\frac{a_0}{2^n}\right) - \hat{V}\left(-\frac{a_0}{2^n}\right)\right] + \frac{1}{2} m \omega^2 \hat{x}^2$$

where the operators are represented as

$$\hat{V}\left(\frac{a_0}{2^n}\right) e_{\alpha_i, C_n} = e_{\alpha_{i+1}, C_n} ,$$

and

$$\hat{x} \, e_{\alpha_i, C_n} = L(\alpha_i) \, e_{\alpha_i, C_n} \,,$$

with $L(\alpha_i) = \frac{i a_0}{2^n}$. Let us now discuss what is being done. The original system is rather symmetric for x and p, being quadratic in both coordinates. This means that one can alternatively think of x or p as the configuration variable. Then, what one is doing is to replace the 'potential' p^2 for a periodic function, for each n as

$$p^2 \approx 2 \left(\frac{2^n \hbar}{a_0}\right)^2 \left[1 - \cos\left(\frac{a_0 p}{2^n \hbar}\right)\right].$$

This is precisely the potential of a pendulum with frequency ω and n-dependent length $\ell_n = \frac{\hbar 2^n}{m\omega a_0}$. That is, we are approximating, for each scale C_n the SHO by a pendulum that is longer (and with 'stronger gravity' since the effective gravitational constant is given by $g = \frac{\hbar \omega^2}{ma_0}$) as we go to smaller distances. In the limit we do recover the SHO. There is, however, an important difference. From our knowledge of the pendulum, we know that the quantum system will have a spectrum for the energy that has two different asymptotic behaviors, the SHO for low energies and the planar rotor in the higher end, corresponding to oscillating and rotating solutions respectively⁸. As we refine our scale and both the length of the pendulum and the height of the periodic potential increase, we expect to have an increasing number of oscillating states (for a given pendulum system, there is only a finite number of such states). Thus, it is justified to consider the cut-off in the energy eigenvalues, as discussed in the last section, given that we only expect a finite number of states of the pendulum to approximate SHO eigenstates. With these consideration in mind, the next question is whether the conditions (4) and (5) are satisfied for the SHO.

We denote the corresponding eigen-covectors as $\Psi_{\nu,C_n} = \sum_k \bar{\psi}_{\nu,C_n} \left(\frac{ka_0}{2^n}\right) \omega_{\alpha_k,C_n} \in \mathcal{H}_{C_n}^{*ren}$. Let us take into account the microscopic corrections and define the renormalized eigen-covectors, at the scale C_m as

$$\Psi_{\nu,C_m}^{ren} = \lim_{C_n \to \mathbb{R}} d_{m,n}^* \Psi_{\nu,C_n} .$$

⁸ Note that both types of solutions are, in the phase space, closed. This is the reason behind the purely discrete spectrum. The distinction we are making is between those solutions inside the separatrix, that we call oscillating, and those that are above it that we call rotating.

We can now prove the following result

$$\Psi_{\nu,C_m}^{\text{ren}} = \Psi_{\nu,C_m}^{\text{shad}}, \tag{12}$$

where $\Psi_{\nu,C_m}^{\text{shad}}(e_{\alpha_i,C_m}) = \bar{\psi}_{\nu}^{\text{Schr}}(ia_0/2^m)$ is (the complex conjugate of) the eigenfunction in the representation of Schrödinger calculated at the extreme left side of the interval α_i of the partition C_m (see FIG 2). The proof is based on the following numerical result in [1],




FIG. 2: The solid continuous line represents the graph of the vacuum state $\Psi_{0,\text{SHO}}$ and the piecewise constant function represents $\Psi_{0,C_m}^{\text{ren}}$.

obtained for the case of a fixed regular lattice (for our considerations are relevant only the results referring to $x_0 = 0$ in their notation) and modified by using the renormalized inner product,

$$2^{n/2} \parallel \Psi_{\nu,C_n} - \Psi_{\nu,C_n}^{\text{shad}} \parallel^{\text{ren}} \sim (\nu + 1)^{1.35} \left(\frac{a_0}{2^n D}\right)^{1.10} , \qquad (13)$$

where $D^2 = \frac{\hbar}{m\omega}$. This result is obtained for $10^{-6} < \frac{a_0}{2^n D} < 1$. Using the properties of d^* we can easily prove that

$$\parallel d_{m,n}^{\star} \Psi_{\nu,C_n} - \Psi_{\nu,C_m}^{\mathrm{shad}} \parallel^{\mathrm{ren}} := \parallel d_{m,n}^{\star} (\Psi_{\nu,C_n} - \Psi_{\nu,C_n}^{\mathrm{shad}}) \parallel^{\mathrm{ren}} \leq 2^{(n-m)/2} \parallel \Psi_{\nu,C_n} - \Psi_{\nu,C_n}^{\mathrm{shad}} \parallel^{\mathrm{ren}},$$

which, by means of (13), leads us to the wanted convergence result. Note that the numerical result of [1] is valid in a range of values that implies a maximum number n'. However, there is strong evidence to assume that the desired result holds valid (see [1] and [5] for further analytic arguments).

Note that the $\Psi^{\text{ren}}_{\nu,C_n}$ are compatible in the sense that

$$d^*\Psi_{\nu,C_n}^{\text{ren}}(e_{\alpha_i,C_{n-1}}) = \Psi_{\nu,C_{n-1}}^{\text{ren}}(e_{\alpha_i,C_{n-1}})$$
.

For any $\delta_x \in \mathcal{H}_{\text{poly},x}$ we have

$$\lim_{C_n \to \mathbb{R}} \Psi_{\nu, c_n}^{\text{ren}}(\delta_x) \equiv \Psi_{\nu, \mathbb{R}}(\delta_x) = \bar{\psi}_{\nu}^{\text{Schr}}(x) ,$$

so that in this limit we recover all the information about the eigenstates from the Schrödinger quantum mechanics.

Regarding the energy levels we take the corresponding result from [1] and rewrite it in our notation leading to

$$E_{\nu,C_n} \sim (2\nu + 1)\frac{\hbar\omega}{2} + \mathcal{O}\left(\frac{a_0}{2^n D}\right)$$
,

valid for $(a_0/2^nD) \ll 1$, so that

$$\lim_{C_n \to \mathbb{R}} E_{\nu, C_n} = E_{\nu, SHO} \,,$$

where $E_{\nu, \text{SHO}}$ are the energy levels for the harmonic oscillator in the Schrödinger representation.

The convergence of the eigen covectors and of the discrete energy levels imply that we have completely renormalized Hamiltonians for any allowed value of the cut-off

$$h_m^{\nu_{\mathrm{cut-off}},\mathrm{ren}} := \lim_{C_n \to \mathbb{R}} d_{m,n}^{\star} h_n^{\nu_{\mathrm{cut-off}}}.$$

Also the Hamiltonian $h_{\mathbb{R}}^{\text{ren}}: \stackrel{\longleftarrow}{\mathcal{H}}_{\mathbb{R}}^{\text{ren}} \to \mathbb{R}$ as defined in equation (9) (or equivalently by taking the limit $\nu_{\text{cut-off}} \to \infty$) can be calculated. The calculation is especially simple when we consider the states $\{\Psi_{\nu,C_n}^{\text{shad}}\} \in \stackrel{\longleftarrow}{\mathcal{H}}_{\mathbb{R}}^{\text{ren}}$. After using the convergence result (12) it is easy to find

$$h_{\mathbb{R}}^{\text{ren}}(\{\Psi_{\nu,C_n}^{\text{shad}}\}) = E_{\nu,\text{SHO}}.\tag{14}$$

Since the L^2 -classes corresponding to states of this type form a dense subset of L^2 , we have shown that the unitary anti-isomorphism of Theorem 2 identifies our Hamiltonian and the Hamiltonian of the SHO written in the Schrödinger representation. We can thus conclude that we have recovered the standard treatment of the harmonic oscillator when applying our formalism.

Let us end this section with a remark. As we have seen in the system considered in this section, there is no intrinsic length scale that one can consider as fundamental and below which the theory should not be considered. One could imagine that there is such a minimum scale, the Planck length, which would provide a cut-off in the scales C_n , namely with a maximum number $N_{\rm Pl}$ for n, but as our results here suggest, we do not need to take such a cut-off. Furthermore, even if one chose to have such a cut-off $N_{\rm Pl}$ near the Planck scale, one might not be able to describe in an accurate manner, arbitrarily large, macroscopic systems, such as a stone attached to a coil. The larger the system, the higher the amplitude of the pendulum potential needed to approximate the system, so that one could even need a scale C_n well beyond the Planck scale. How could one then justify the minimum length in that case? Could one consider, for instance, the polymer description as fundamental, but only for elementary systems, and not for composite systems such as a large stone? We shall leave these questions for future research.

V. DISCUSSION AND CONCLUSIONS

Let us summarize our results. We have shown that by starting with the usual polymer representation of a quantum mechanical system, we can implement a renormalization procedure that allows us to define a continuum theory where the physical Hilbert space and the Hamiltonian are well defined. We proceeded in three steps. The first one was to consider an equivalent formulation of the polymer representation, where we could define the notion of scale refinement and the continuum limit at the kinematical level. In order to implement a Hamiltonian, that is regarded as a quadratic form, we showed that, under very mild assumptions, one could perform a renormalization of the inner product and arrive at a physical Hilbert space. This space is equivalent to the standard L^2 space of Schrödinger quantum mechanics. In order to make contact with other approaches, we have shown that if certain conditions are fulfilled, the shadows of the Schrödinger system can be approximated by the continuum polymeric system, in such a way that in the limit, both systems coincide. We have analyzed in detail the case of a simple harmonic oscillator, were all the steps can successfully be completed. The program is general enough that it can, in principle, be implemented to other physically motivated systems where the polymer representation is also the starting point, and to models defined by a bounding potential and a discrete spectrum. Work in this direction is underway, as well as in extending the formalism to systems that in the standard representation have also a continuum energy spectrum and for totally constrained systems.

An important issue regarding loop quantization and the applicability of these methods to different physical systems has to do with the adequacy of loop quantization for dealing with all class of systems. That is, are loop methods tailored to background independent theories only? And, why should we expect that these methods are useful for background dependent theories at all? Our considerations here have been motivated by our expectations for loop quantization, namely, that it should be applicable to theories that depend on a metric background as well as to theories that are independent of such structure. Thus we expect that, when applied to the case of metric background dependent theories, it should recover known physics while giving new theories when applied in the background independent category. Such a goal can only be obtained if loop quantization itself knows of no preferred background metric and the dynamics of each particular theory (that might itself depend on extra structures) brings in the knowledge of the background structure that it needs. This is precisely what happens with the polymer representation of quantum mechanics.

As we have seen, the model of polymeric quantum mechanics is a concrete realization of our goal. At the kinematical level we started with the polymer representation of the Weyl algebra on \mathcal{H}_{poly} . The algebra itself does not know about any extra structure such as a metric on the configuration space (nor phase space). A remarkable fact about \mathcal{H}_{poly} is that it hosts a faithful unitary representation of the group of homeomorphisms of \mathbb{R} . Later on, we continued with our quantization process and found that the dynamics called for a renormalization of the inner product which lead to the physical Hilbert space \mathcal{H}_{phys} which now knows of a metric structure in \mathbb{R} ; only the group of rigid translations can be represented unitarily and faithfully on it. The moral that this model is trying to teach us is that, as already mentioned, it is the system under consideration that introduces extra structure and thus reduces the symmetries of the final, dynamical theory.

We do not want to claim that our implementation of Wilson's Renormalization Group is the only possible one. In fact, the general procedure of [3] (applicable to loop quantized field theories), calls for modeling a given scale by a general decomposition of the space (or spacetime) manifold. In the case of \mathbb{R} , cell decompositions are built out of open intervals (1-d cells) and points (0-d cells). The procedure that we followed here is more economical and it has the useful property of being consistent with only one decimation map. A procedure of this type is available for \mathbb{R}^n (simply considering the cartesian product of our construction n times) and for T^n . If the procedure is not unique, then one must ask if the results can be trusted in any way. The existence of the limits defining the continuum limit imply that,

as the scale is refined, the relevant objects acquire some degree of regularity which justifies, to some extent, our initial expectation for robustness of the formalism. In cases (like the SHO) where not only the continuum limit exists but the relevant wave functions coincide with the shadows of the corresponding wave functions on the Schrödinger representation which in addition happen to be continuous, we know that as the scale is refined the wave functions become continuous. In these cases we can expect that other similar procedures would produce the same continuum limit.

The particular example we have chosen to analyze, namely the simple harmonic oscillator, has received much attention lately, mainly in connection with a proposed loop quantization of strings (See for instance [6] and [7]). Even when it might seem that our results contradict in a sense the results of [7], this is not the case. This is because we are looking at different physical questions. In our case, we are concerned with a procedure to implement the Hamiltonian operator in the continuum, while [7] analyze a closely related representation where the finite time evolution operator is well defined but neither \hat{p} nor \hat{q} exist as well defined operators. What the articles in Ref. [7] show is that, by coupling the oscillator to an external field, one might get very different answers for physically motivated quantities. It would be interesting though to apply the procedure here presented to that particular quantization and see whether this behavior remains in the limit.

What lessons for loop quantum gravity can we learn from our study? The first immediate 'application' of our formalism that comes to mind is loop quantum cosmology. The classical system possesses a finite number of degrees of freedom, so it falls within the quantum mechanical category. The relevant operator, namely the Hamiltonian constraint, does not have the standard form with a kinetic and a potential term, such as a standard particle on a potential, but nevertheless, the methods of polymer quantization are employed for them [2]. It is also remarkable that our case study, namely the SHO, has an intrinsic length scale $(D = \sqrt{\hbar/m\omega})$ that can play a parallel role to that of the the Planck length in loop quantum cosmology. One could imagine that the intrinsic length scale determines a minimum scale, which would provide a 'physical cut-off' of the type $C_n \leq C_{N_{\max}}$, with $\frac{a_0}{2^{N_{\max}}} = L_{\text{intrinsic}}$. Our results here are not conclusive for general systems, but they suggest that stopping at any given scale might be physically incorrect.

On the other hand, if loop quantum cosmology is to be seen as a symmetric sector of loop quantum gravity, then one is allowed in principle to impose such a cut-off, coming from the minimum area gap of the full theory [2]. The question remains open, however, as to whether one should take the Planck scale $C_{N_{\rm Pl}}$ as fundamental and keep the truncated theory, or to perform the continuum limit and take the resulting description at face value. Would in that case the continuum theory coincide with the Wheeler-DeWitt quantization? or, do the main features of the truncated theory prevail such as a a bounce near the singularity? Can one define, for instance, physical observables that implement the quantization of geometry at the physical Hilbert space level? Can, therefore, a minimum scale be recovered dynamically? Work in this direction is underway.

Let us end this paper by asking what might be the relevance of our results for the program of loop quantum gravity at large. That is, can we apply what we have learned to full loop quantum gravity? In this respect, we will give one technical comment and later a more general one. First, we will elaborate on the issue of "summing over discrete structures versus refining discrete structures" to define the continuum limit (or to get rid

⁹ Some issues related to the existence of this fundamental cut-off in LQC have also been raised in [8].

of the discretization introduced as a mere auxiliary regularization ingredient). Our point of view uses the discrete structures C to define effective theories; for example, the states of \mathcal{H}_C are related to equivalence classes of states of \mathcal{H}_{poly} . Then when we consider $C' \geq C$, the equivalence classes $[\cdot]_C$ become unions of $[\cdot]_{C'}$ equivalence classes. Thus, in our formulation "refinement" and "summing over" can not be separated. Now we turn to comment on a more general issue that calls for attention after our work. A possible strategy would be to advocate the viewpoint that we should take seriously the lessons from this simple model as well as from actual field theories where a more nontrivial Wilsonian renormalization is needed. Examples of these are 2d Yang-Mills, 3d quantum gravity and the much more non trivial (scalar) example that is an interacting relativistic quantum field theory defined as the continuum limit of 2d Ising field theory [3]. Also, when we apply the formalism of [3] to gauge theories on a flat metric background, our effective theories and decimation maps lead us to Lattice Gauge Theory, where the renormalization group flow has been extensively studied numerically. These studies suggest that one might need to take, in loop quantum gravity, the Wilsonian viewpoint and perform a nontrivial renormalization. These and other questions certainly deserve further investigation.

Acknowledgments

We thank Abhay Ashtekar, Raymundo Bautista, Homero Díaz, Robert Helling, Michael Hrusak, Jorma Louko, Seth Major, Gerardo Raggi and José Velhinho for discussions, suggestions and advise. This work was in part supported by CONACyT U47857-F and 40035-F grants, by NSF PHY04-56913, by the Eberly Research Funds of Penn State, by the AMC-FUMEC exchange program and by funds of the CIC-Universidad Michoacana de San Nicolás de Hidalgo.

^[1] Ashtekar A, Fairhurst S and Willis J L 2003 Quantum gravity, shadow states, and quantum mechanics *Class. Quant. Grav.* **20** 1031 [arXiv:gr-qc/0207106].

^[2] Bojowald M Loop quantum cosmology 2005 Living Rev. Rel. 8, 11 [arXiv:gr-qc/0601085]; Ashtekar A, Bojowald M and Lewandowski L 2003 Mathematical structure of loop quantum cosmology Adv. Theor. Math. Phys. 7 233 [arXiv:gr-qc/0304074]; Ashtekar A, Pawlowski T and Singh P 2006 Quantum nature of the big bang: Improved dynamics Phys. Rev. D 74 084003 [arXiv:gr-qc/0607039]

^[3] Manrique E, Oeckl R, Weber A and Zapata J A 2006 Loop quantization as a continuum limit Class. Quant. Grav. 23 3393 [arXiv:hep-th/0511222]; Manrique E, Oeckl R, Weber A and Zapata J A 2006 Effective theories and continuum limit for canonical loop quantization (in preparation)

^[4] Fredenhagen K and Reszewski F 2006 Polymer state approximations of Schrödinger wave functions Class. Quant. Grav. 23 6577 [arXiv:gr-qc/0606090]

^[5] Aunola M 2003 The discretized harmonic oscillator: Mathieu functions and a new class of generalized Hermite polynomials J. Math. Phys. 44 1913

^[6] Thiemann T 2006 The LQG string: Loop quantum gravity quantization of string theory. I: Flat target space *Class. Quant. Grav.* **23** 1923 [arXiv:hep-th/0401172]; Thiemann T 2006 Loop quantum gravity: An inside view [arXiv:hep-th/0608210].

- [7] Helling R C and Policastro G 2004 String quantization: Fock vs. LQG representations [arXiv:hep-th/0409182]; Helling R C 2006 Lessons from the LQG string [arXiv:hep-th/0610193].
- [8] Velhinho J M 2004 Comments on the kinematical structure of loop quantum cosmology Class. Quant. Grav. 21 L109 [arXiv:gr-qc/0406008].