PROBABILITIES DON'T MATTER

D on N. Page ^y

CIAR Cosm ology Program, Institute for Theoretical Physics Department of Physics, University of Alberta Edmonton, Alberta, Canada T 6G 2J1

(1994 Nov. 25)

Abstract

It is suggested that probabilities need not apply at all to matter in the physical world, which may be entirely described by the amplitudes given by the quantum mechanical state. Instead, probabilities may apply only to conscious perceptions in the mental world. Such perceptions may not form unique sequences that one could call individual minds.

1 Probabilities in Quantum Mechanics

One of the most mysterious aspects of quantum mechanics is its usual probabilistic interpretation. There is not the uncertainty of which amplitudes should be squared to get probabilities. Then there is the even deeper puzzle of what the resulting probabilities mean.

For example, one view point on the rst question is that whenever a measurement is made, the amplitude for each macroscopically-distinct outcome should be squared to get a probability. More precisely, one takes a complete set of orthogonal projection operators, each representing one of the macroscopically-distinct outcomes. One projects the wavefunction by one of these projection operators to get a reduced wavefunction. Then one takes the inner product of this reduced wavefunction with itself i.e., \squares the amplitude" to get the probability of the corresponding outcome. This probability is the same as the expectation value of the projection operator in the quantum state given by the original wavefunction.)

A lberta-T hy-28-94, gr-qc/9411004, to be published in P roceedings of the 7th M arcelG rossm ann M eeting on G eneral Relativity, eds. M .K eiser and R.T. Jantzen (W orld Scienti c, Singapore 1995).

^yInternet address: don@ phys.ualberta.ca

A broader view point is that the expectation value of any projection operator is a probability for the corresponding \event." An even broader view point is that one can square the amplitude given by projecting the wavefunction not just by one, but by a whole sequence of (possibly noncommuting) projection operators representing a \history" or sequence of \events." (For the resulting probabilities to obey the sum rules under a coarse-graining of the projection operators, the sequences must obey certain consistency conditions [1, 2].) One can extend this view point, of assigning probabilities to \consistent histories," yet further to the view point that one can project the wavefunction by sum s of sequences of projection operators that represent coarser-grained histories. (Then one needs \decoherence" conditions for the resulting probabilities to obey the sum rules for \decohering histories" [3, 4, 5].) An even further extension is the view point that probabilities are the real parts of the expectation values of sum s of sequences of projection operators, whenever these obey a \linear positivity" condition of being nonnegative, giving probabilities for \linearly positive histories" that autom atically obey the sum rules [6].

Besides this sam ple of the wide variety of view points of what quantities should be assigned probabilities in quantum mechanics, there is the enigm a of how to interpret the resulting probabilities. One attitude is that a unique one of the possible \events" or \histories" actually occurs, with the probability assigned by quantum mechanics, and that the other possibilities do not. This attitude still leaves open the question of whether the \probability" is a \frequency" in an ensemble of actual worlds, or whether it is a \propensity" in a single world and what that could mean. If the latter, what is it that actually makes the choice of the actual \event" or \history" from those potentially possible?

A nother attitude is that all of the possible \events" or \histories" with nonzero probabilities actually occur, but with measures proportional to the probabilities. This \m any-worlds" interpretation [7] is very sim ilar to the frequency interpretation in a ensemble of actual worlds mentioned above, but it need not have the implication that our present world is a single member of the ensemble that has a de nite (albeit unpredictable) future.

In any of these approaches in which one or all of the possibilities are actualized, there is the further question of which set of possibilities is singled out. In general, there are m any di erent allowed sets of possibilities (e.g., di erent sets of orthogonal projection operators, or of sequences of projection operators, that each add up to the identity operator). If only one \event" or only one \history" actually occurs, there m ust be mysterious choices both of the set of possibilities and of the single actual element within that chosen set. If, on the other hand, all of the possible \events" or \histories" with nonzero probabilities in a given set of possibilities actually occur, there m ust apparently still be a mysterious choice of this set of possibilities out of the fam ily of all such sets of possibilities.

One conceivable answer is that all possibilities in all sets of possibilities actually occur, with measures given by the probabilities deduced from quantum mechanics

in one of the ways discussed above. Since, for a norm alized quantum state, these probabilities are designed to add up to unity for a single set of possibilities, the measures will sum to more than unity when one adds up all the sets. In fact, typically the number of allowed sets of possibilities is in nite (e.g., even for a twostate spin-half system, there is a rank-one projection operator for each direction in space and hence an in nite number of such projection operators). This means that the sum of the measures for all possibilities typically is unnorm alizable, which may lead to problem s. (These problems may be avoidable. For example, in the spin-half system, the sum of the quantum probabilities for all of the in nite number of possibilities for the spin direction is in nite, but instead of simply adding these quantum probabilities, one can divide by the total solid angle of the unit sphere of spin directions to get a probability density which can then be integrated over a nonzero solid angle to get a norm alized second-order probability that the spin is in a direction within that solid angle.)

2 Sensible Quantum Mechanics

Here I shallpropose instead that probabilities not be applied at all to the physical world (the \m atter" of the title), which is instead to be described entirely by the quantum amplitudes of its wavefunction (or by the elements of its density m atrix, or possibly by a more general description, such as a C -algebra state). I suggest instead that probabilities apply only to the mental world of conscious perceptions. In this view point, and in a certain bose manner of speaking in which \m ind" is this whole mental world rather than a sequence of perceptions, probabilities are only in the mind.

Consider the set of all possible perceptions p, which I shall call the m ental world M. By a perception, I m ean all that one is consciously aware or consciously experiencing at one m om ent. This is what Lockwood, in a book expressing what seems to be ideas mostly concordant with m ine [8], calls a \phenom enal perspective" or \m axim al experience." In another way of putting it, m y p denotes a total \raw feel" that one has at once. It can include com ponents such as a visual sensation, an auditory sensation, a pain, a conscious m em ory, a conscious in pression of a thought or belief, etc., but it does not include a sequence of m ore than one im m ediate perception that in other proposals m ight be considered to be strung together to form a stream of consciousness of an individual m ind.

Suppose that there is a basic measure on the mental world which weights each perception equally, which I shall denote by dp without implying a choice between the conceptual possibilities that the space of perceptions is discrete (in which case dp simply counts the number of perceptions) or continuous (in which case dp m ight represent some basic multi-dimensional integration measure or volume element). Now I shall postulate that our actual world does not have all possible perceptions occurring equally, but instead that there is a nontrivial measure m (p)dp, the non-

negative real scalar function or measure density m (p) times the basic measure, for the perceptions in our actual world. Thus I get the following basic assumption:

M easure A xiom for P erceptions: There is a nontrivial measure on the space of (m axim al) perceptions p, namely m (p) times the basic measure dp that weights each perception equally.

Because of the fact that our perceptions seem to be more simply explained by assuming that they are related to a physical world, I shall assume a principle of psycho-physical parallelism, that the nontrivial part m (p) of the measure for perceptions is a functional of the properties of the physical world.

For example, if the physical world were represented by a single classical trajectory in some phase space, it might be natural to assume that m (p) has a form of a sum of K ronecker deltas or an integral of a discrete set of D irac delta function distributions over sequences of perceptions that a set of conscious beings has as the point m oves along the trajectory in phase space. In other words, each point in phase space might naturally be assigned a discrete set of conscious perceptions, one for each conscious being whose point in its phase space is derived from the point in the phase space of the entire system (e.g., the universe). Even in this classical model, there can be m any perceptions at one time, but each is ascribed to a di erent conscious being.

If the physical world is represented by a quantum state that has no preferred classical trajectories (such as do occur in Bohm's version of quantum mechanics [9], but which I shall not further consider here), then it seems unnatural to assume that m (p) is completely concentrated on a discrete sequence of trajectories, one for each conscious observer. Instead, in view of the linearity of quantum mechanics, I propose the following basic assumption for interpreting quantum mechanics 'sensibly,' i.e., in term s of sensations or perceptions:

Sensible Q uantum A xiom : Each m (p) is given by the expectation value of a corresponding positive perception or \mbox{m} axim al experience" operator E (p) in the quantum state of the universe. As a formula,

$$m(p) = hE(p)i = h E(p)ji = Tr(E(p));$$
 (1)

where the third expression applies if the quantum state is represented by the wavefunction or pure state j i, and the fourth expression applies if the quantum state is represented by the statistical operator or density matrix \cdot . (The second expression can apply in more general situations, such as in C -algebra.)

In this fram ework, which I shall call Sensible Q uantum M echanics, the quantum state of the universe is xed, in the H eisenberg picture I am using, and never collapses or changes to another state, so the von N eum ann intervention 1 [10] is assumed never to occur. N either is there assumed to be any nonlinearity in quantum m echanics when consciousness is involved, as W igner proposed [11]. If the quantum state and all the perception operators E (p) are known, one can in principle calculate from Eq. (1) the m easure density m (p) for all perceptions. (O f course, I am not competent to give these essential elements, so the present proposal is a fram ework,

on the level of other fram eworks or interpretations of quantum mechanics, rather than a complete theory.) Since all maximal perceptions p with m(p) > 0 really occur in this fram ework, it is completely determ inistic if the quantum state and the E (p) are determined: there are no random or truly probabilistic elements in this fram ework of Sensible Quantum Mechanics.

Thus Sensible Q uantum M echanics has no need for any axiom of what is typically called \m easurem ent" [10], or what U nruh calls \determ ination" [12] to distinguish this hypothetical process from the physical m easurem ent interactions that are encoded in the quantum state and the structure of the operators. In particular, there are no probabilistic results of such \determ inations."

Nevertheless, because the fram ework has measures for perceptions, one can readily use them to calculate quantities that can be interpreted as conditional probabilities. One can consider sets of perceptions A, B, etc., de ned in term s of properties of the perceptions. For example, A m ight be the set of perceptions in which there is a feeling that the universe is approximately described by a Friedman-Robertson-W alkermodel, and B m ight be the set of perceptions in which there is a feeling that the universe is approximately described by a Friedman-Robertson-W alker model with an age (at the perceived time) between ten and twenty billion years. By sum – ming or integrating m (p)dp over the sets A; B; A \ B (= B in the example here), etc., one can get corresponding measures m (A); m (B); m (A \ B), etc. Then one can interpret

as the conditional probability that the perception is in the set B, given that it is in the set A. In our example, this would be the conditional probability that a perception including the feeling that the universe is approximately described by a Friedman-Robertson-W alker model, also has the feeling that at the time of the perception the age is between ten and twenty billion years.

An analogue of this conditional \probability" is the conditional probability that a person in 1994 is the Q ueen of England. If we consider a model of all the ve to six billion people, including the Q ueen, that we agree to consider as our contemporary hum and on Earth in 1994, then at the basic level of this model the Q ueen certainly exists in it; there is nothing random or probabilistic about her existence. But if the model weights each of the ve to six billion people equally, then one can in a manner of speaking say that the conditional probability that one of these persons is the Q ueen is somewhat less than 2 10¹⁰. I am proposing that it is in the same manner of speaking that one can assign conditional probabilities to sets of perceptions, even though there is nothing truly random about them at the basic level.

W hen one's perceptions include feelings of belief about what is ascribed to be external events or histories (e.g., results of experiments in the physical world rather than in the mind), and when it is believed that these beliefs are an accurate representation of some aspects of those ascribed-to-be-external events or histories, it is tem pting to the theorist to interpret the conditional probabilities of the perceptions as giving conditional probabilities for those aspects of the ascribed-to-be-external events or histories. One may even go further and develop formalisms for directly calculating probabilities of such events or histories. W ith this view point one can say that the historical development of quantum mechanics has been fruitful (or more strictly, I have a feeling that what I perceive to have been its historical development was fruitful), but it has left unexplained which events or histories are to be assigned probabilities and what those probabilities mean.

Thus I am proposing that at the basic level, probabilities (or, more strictly, measures) have meaning only for perceptions in the mental world and should not be assigned to anything (e.g., to events or histories) in the physical world of matter. In this sense probabilities don't matter."

On the other hand, I am not saying that it is forbidden to assign conditional \probabilities" to events and/or histories in the physical world. They can be much simpler to calculate than those in the mental world given by Eqs. (1) and (2), since we don't know what the perception operators E (p) are, and these physical \probabilities" may offen give good approximations for the mental probabilities. Indeed, I have been happy to help play the gam e of broadening the scope of histories to which one can assign mathematically-consistent probabilities [6]. How ever, I am now proposing that these probabilities are not fundamental and need not be added to complicate the basic ontology of a measurable set of perceptions (which I have been calling the mental world) and a quantum state of the universe (which I have been calling the physical world), with the measure and interpretation given entirely by the Measure Axiom for Perceptions and the Sensible Quantum Axiom above.

G oldstein has pointed out [14] that one can simplify the ontology and avoid the assumption of a basic measure dp on the mental world by replacing the measure density m (p) for single perceptions p with a measure (S) on sets S of perceptions and by replacing the perception operators E (p) with a positive-operator-valued measure, say A (S). This alternative formulation is given in [15, 16]. Another way to minimize the number of independent entities is to postulate that the basic measure dp is the volume element of a R iem annian metric

$$g_{ij}dp^{i}dp^{j} = Trf[E(p^{i} + dp^{i}) E(p^{i})][E(p^{j} + dp^{j}) E(p^{j})]g;$$
 (3)

if this is nite and nondegenerate, so that the basic measure is determined by the perception operators them selves.

3 Perceptions rather than M inds

A nother point I should emphasize is that in Sensible Quantum M echanics, the set of all perceptions is basic, but not any higher power of this set. In other words, perceptions and the measure m (p)dp on them are basic, but not n-tuples of perceptions, or measures on n-tuples of perceptions. Thus, for example, there is no

fundam ental notion of a correlation between individual complete perceptions given by any measure. (On the other hand, if a perception can be broken up into component parts, say A and B, there can be a correlation between the parts, in the sense that the measure m (A \ B) for all perceptions containing the part A and the part B need not be the same as m (A) m (B), the measure for all perceptions containing A times the measure for all perceptions containing B. The enorm ous structure in a single perception seem s to suggest that such correlations within perceptions are highly nontrivial, but I see no evidence for a nontrivial correlation between m axim al perceptions, since no two di erent m axim al perceptions can be perceived together.)

Furtherm ore, Sensible Quantum M echanics postulates no fundam ental equivalence relation on the set of perceptions. For example, the measure gives no way of classifying di erent perceptions as to whether they belong to the same conscious being (e.g., at di erent tim es) or to di erent conscious beings. The only such classication would be by the content (including the qualia) of the perceptions them selves, which distinguish the perceptions, so that no two di erent perceptions, $p \in p^0$, have the same content. Based upon my own present perception, I nd it natural to suppose that perceptions that could be put into the classi cation of being alert hum an perceptions have such enorm ous structure that they could easily distinguish between all of the 10¹¹ or so persons that are typically assigned to our history of the hum an race. However, I doubt that in a fundam ental sense there is any absolute classi cation that uniquely distinguishes each person in all circum stances. Therefore, in the present fram ework perceptions are fundam ental, but persons (or individual minds) are not, although they certainly do seem to be very good approximate entities that I do not wish to deny. The concept of persons and minds certainly occurs in some sense as part of the content of my present perception, even if there is no absolute de nition of it in the fram ework of Sensible Quantum Mechanics itself.

In this way the fram ework of Sensible Quantum M echanics proposed here is a particular manifestation of H um e's ideas [13], that \what we call a m ind, is nothing but a heap or collection of di erent perceptions, united together by certain relations, and suppos'd, tho' falsely, to be endow 'd w ith a perfect sim plicity and identity" (p. 207), and that the self is \nothing but a bundle or collection of di erent perceptions" (p. 252). As he explains in the Appendix (p. 634), \W hen I turn m y re exion on m yself, Inever can perceive this self without some one orm ore perceptions; nor can I ever perceive any thing but the perceptions. 'T is the com position of these, therefore, which form s the self." (H ere I should note that what H um e calls a perception m ay be only one com ponent of the \phenom enal perspective" or \m axim al experience" [8] that I have been calling a perception p, so one p can include \one or m ore perceptions" in H um e's sense.)

Furtherm ore, each perception operator need not have any precise location in either space or time associated with it, so there need be no fundamental place or time connected with each perception. Indeed, Sensible Quantum Mechanics can easily survive a replacement of spacetime with some other structure (e.g., superstrings) as m ore basic in the physical world. O focurse, the contents of a perception can include a sense or impression of the time of the perception, just as my present perception at the perceived time of writing this includes a feeling that it is now 1994 A D ., so the set of perceptions p must include perceptions with such beliefs, but there need not be any precise time in the physical world associated with a perception. That is, perceptions are butside' physical spacetime (even if spacetime is a fundam ental element of the physical world, which I doubt).

As a consequence of these considerations, there are no unique time sequences of perceptions to form an individual mind or self in Sensible Quantum M echanics. In this way the present fram ework appears to dier from those proposed by Squires [17], A bert and Loewer [18], and Stapp [19]. (Stapp's also di ers in having the wavefunction collapse at each \Heisenberg actual event," whereas the other two agree with mine in having a xed quantum state, in the Heisenberg picture, which never collapses.) Lockwood's proposal [8] seems to be more similar to mine, though he also proposes (p. 232) \a continuous in nity of parallel such stream s" of consciousness, \di erentiating over tim e," whereas Sensible Quantum Mechanics has no such stream as fundam ental. On the other hand, later Lockwood [20] does explicitly repudiate the Albert-Loewer many-minds interpretation, so there seems to me to be little disagreem ent between Lockwood's view and Sensible Quantum Mechanics except for the detailed form alism and manner of presentation. Thus one might label Sensible Quantum Mechanics as the Hume-Everett-Lockwood-Page (HELP) interpretation, though I do not wish to imply that these other three scholars, on whose work my proposal is heavily based, would necessarily agree with my present form ulation.

Of course, the perceptions them selves can include components that seem to be m em ories of past perceptions or events. In this way it can be a very good approxim ation to give an approxim ate order for perceptions whose content include m em ories that are correlated with the contents of other perceptions. It m ight indeed be that the m easure (or m easure density) m (p) for perceptions including detailed m em ories is rather heavily peaked around approxim ate sequences constructed in this way. But I would doubt that either the content of the perceptions or the m easure on the set of perceptions would give unique sequences of perceptions that one could rigorously identify with individual m inds.

Because the physical state of our universe seems to obey the second law of therm odynamics, with growing correlations in some sense, I suspect that the measure density m (p) may have rather a smeared peak (or better, ridge) along approximately tree-like structures of branching sequences of perceptions, with perceptions further out along the branches having contents that includes memories that are correlated with the present-sensation components of perceptions further back toward the trunks of the trees. This is dierent from what one might expect from a classical model with a discrete number of conscious beings, each of which might be expected to have a unique sharp sequence or non-branching trajectory of perceptions. In the quantum case, I would expect that what are crudely viewed as quantum choices would cause smeared-out trajectories to branch into larger numbers of smeared-out trajectory is viewed as a di erent individualm ind, we do get roughly a \m any-m inds" picture that is analogous to the \m any-worlds" interpretation [7], but in m y fram ework of Sensible Q uantum M echanics, the \m any m inds" are only approxim ate and are not fundam ental as they are in the proposal of A lbert and Loewer [18]. Instead, Sensible Q uantum M echanics is a \m any-perceptions" or \m any-sensations" interpretation. O ne m ight label it philosophically as M indless Sensationalism.

Even in a classical model, if there is one perception for each conscious being at each moment of time in which the being is conscious, the fact that there may be many conscious beings, and many conscious moments, can be said to lead to a m any-perceptions" interpretation. However, in Sensible Quantum Mechanics, there may be vastly more perceptions, since they are not limited to a discrete set of one-parameter sharp sequences of perceptions, but occur for all perceptions p for which m (p) is positive. In this way a quantum model may be said to be even m ore sensible" than a classical model.

One might fear that the present attack on the assumption of any de nite notion of a precise identity for persons or minds as sequences of perceptions would threaten hum an dignity. A lthough I would not deny that I feel that it might, I can point out that on the other hand, the acceptance of the view point of Sensible Q uantum M echanics might increase one's sense of identity with all other hum ans and other conscious beings. Furtherm ore, it might tend to undercut the motivations tow and sel shness that I perceive in myself if I could realize in a deeply psychological way that what I norm ally anticipate as my own future perceptions are in no fundam ental way picked out from the set of all perceptions. (O fcourse, what I norm ally think of as m y own future perceptions are presumably those that contain memory components that are correlated with the content of my present perception, but I do not see logically why I should be any more concerned about trying to make such perceptions happy than about trying to make perceptions happy that do not have such memories: better to do unto others as I would wish they would do unto me.) Lockwood [21] inform s me that Part 1[22] has drawn sim ilar conclusions from a Hum ean view.

4 Properties of Perception Operators

A lthough no one is competent to give the complete set of perception operators E (p), one can speculate about some of their properties. In this speculation, a theoretical physicist such as myself would like to be guided by the principles of simplicity and of agreement with observations. Both are dicult, the form erbecause we do not know all that is logically possible and have a measure of the simplicity of the di erent possibilities, and the latter because we do not have direct access to more than one perception at once.

On the form er principle, it is because of simplicity that I do not stop at the M easure A xiom for Perceptions but also postulate a physical quantum state and a set of perception operators E (p) which give them easure density m (p) by the Sensible Q uantum A xiom. If one has the correct m (p) (as well as the basic measure dp, which is a separate element from the set of perceptions if they are not discrete), the M easure A xiom for Perceptions is logically su cient for describing a measured set of perceptions. It might seem to be complicating the theory to add a physical quantum state and a set of perception operators E (p), but I believe that this structure of a postulated physical world can give a simpler explanation of m (p) than just giving it directly without this explanation in terms of a postulated physical world. In this way the whole of the mental world and the physical world can be simpler than just the mental world considered by itself. (O ne might also raise the reverse question of whether the whole is simpler than the physical world alone, by which I mean an alternative logically possible world in which all E (p) are zero, so that in it the quantum state is the same as in ours, but no conscious perceptions occur.)

On the latter principle, the only agreem ent with observations that one can impose is the assumption that one's perception be not too atypical, i.e., that it have not too low a measure density m (p). For example, if both the basic measure dp and the quantum measure m (p)dp were integrable, over the set of all perceptions p, to nite numbers, say b and q respectively, then one can ask that one's particular perception not have m (p) q=b, the latter being the average of m (p) over all perceptions. Unfortunately, I see no reason why a simple theory should make either of these integrals nite, since alm ost any nite number ism ore complex than in nity. (Perhaps the fact that my present perception seems to have a large but not in nite am ount of inform ation in it is evidence that the simplest complete theory is not extrem ely simple, since I would expect an extrem ely simple theory to make typical perceptions have either an extrem ely sm all or an in nite am ount of inform ation.)

Perhaps a more realistic approach one can make toward agreement with observations is to assume that the measure density m (p) for one's perception is not much lower than the measure density for slightly dierent perceptions. For example, if one has a perception p of having made a certain quantum measurement n times and having gotten m positive results, one can in agine another perception p^0 which is similar except for perceiving m⁰ positive results. Then one would like m (p) to be not much lower than m (p⁰). If the measures (or measure densities) m (p) and m (p⁰) of the perceptions p and p⁰ can to a good approximation be replaced by the quantum derived measures m (r) and m (r⁰) for the respective perceived results r and r⁰ in the physical world, one can check whether m (r) is not much lower than m (r⁰). In such cases there is considerable experimental evidence that the ordinary quantum predictions are consistent with observations ifm (r) is the expectation value of a projection operator P (r) for the physical result r.

Sim ilarly, it m ight be natural to assume that each perception measure density m (p) is given by the expectation value of a projection operator, say one that projects

onto the brain states that cause the perception, if indeed the perception is caused by various brain states. However, before making this speci c assumption, let me make some weaker postulates that one could add to Sensible Quantum Mechanics to make it more restrictive and yet have a more speci c content:

H ypothesis A : The expectation value of each E (p) has a constant maximum value, say unity, in the set of all norm alized quantum states.

A sum ing that the quantum state is normalizable (perhaps an overly restrictive assumption), Hypothesis A and the resulting Sensible Quantum M echanics A would have 0 m (p) 1 for all p, and there would exist a normalizable quantum state for each p such that the corresponding m (p) would be unity in that state.

W ithout some such restriction like that, one could have all the explanation of m(p) in the operators E(p) rather than in the quantum state. For example, one could get m(p) to be any positive function whatsoever simply by choosing E(p) to be that function times the identity operator, which would make m(p) independent of the state (so long as the state is norm alized so that the expectation of the identity operator is unity; other norm alizations would change the scale of the measure but would have the conditional probabilities of Eq. (2) unchanged). We would like to assume that instead the E(p) are restricted so the explanation for the m(p) lies largely in the quantum state.

Now Hypothesis A is still not highly restrictive, so one may wish to look for more restrictions on the operators E (p). For example, one may be motivated by the consistent or decohering histories approaches [1, 2, 3, 4, 5] to assume that perceptions are connected with histories and so perhaps make the following assumption:

H ypothesis B : Each E (p) has the form E (p) = $C^{Y}C = m \operatorname{ax} (C^{Y}C)$, where $C = P^{(n)}P^{(n-1)}$ (2) $PP^{(1)}$ with the integer n and the projection operators $P^{(i)}$ all depending on the perception p, and where m ax ($C^{Y}C$) is the maximum expectation value of $C^{Y}C$ in any normalized quantum state.

The denom inator in the expression for E (p) in Hypothesis B is chosen so that Hypothesis B is a special case of Hypothesis A, but one could also consider an alternative Hypothesis B' in which the denom inator is om itted. One could also consider generalizing Hypothesis B or B' to B* or B*' respectively, in which C is a sum of sequences of projection operators, as is allowed in decohering histories [3].

It is certainly logically possible that perceptions might depend on histories rather than events that one could consider localized on hypersurfaces of constant time if the physical world has such a time. However, as a previous advocate of the in arvelous moment' approach to quantum mechanics in which only quantities on one such hypersurface can be tested [23], I nd it more believable to assume that perceptions are caused by brain states which could be at one moment of time if there are such things in the physical world. The generalization of this hypothesis to the case in which there may not be a well-de ned physical time leads me to make the following restriction of Hypothesis B or B' to the case in which the integer n is always 1:

Hypothesis C: E(p) = P(p), a projection operator that depends on the per-

ception p.

Hypothesis C appears to be a speci c m athem atical realization of part of Lockwood's proposal [8] (p. 215), that \a phenom enal perspective [what I have here been calling simply a perception p] may be equated with a shared eigenstate of som e preferred (by consciousness) set of compatible brain observables." Here I have expressed the \equating" by Eq. (1), and presum ably the \shared eigenstate" can be expressed by a corresponding projection operator P (p).

I should also emphasize that if the same conscious perception is produced by several di erent orthogonal \eigenstates of consciousness" (e.g., di erent states of a brain and surroundings that give rise to the same perception p), then in Hypothesis C the projection operator P (p) would be a sum of the corresponding rank-one projection operators and so would be a projection operator of rank higher than unity (perhaps even in nite), which is what I would expect. On the other hand, if E (p) were a sum of noncommuting projection operators corresponding to nonorthogonal states, or a weighted sum of projection operators with weights di erent from unity, then generically E (p) would not have the form of a projection operator P (p).

If one has a constrained system, such as a closed universe in general relativity, the quantum state m ay obey certain constraint equations, such as the W heeler-D eW itt equations. The projection operators P (p) of perception in Hypothesis C m ay not commute with these constraints, in which case they m ay give technically 'unphysical' states when applied to the quantum state. But so long as their expectation values can be calculated, that is su cient for giving the perception m easure density m (p). W hat it m eans is that in Hypothesis C, the perception operators should be considered as projection operators in the space of unconstrained states, even though the actual physical state does obey the constraints.

A lternatively, if one w ishes to write the perception operators E (p) as operators w ithin the space of constrained states, H ypothesis C could be m odi ed to the follow – ing assumption to give perception operators E (p) that commute with the constraints and so keep the state physical':

H ypothesis $C := (p) = P_C P(p)P_C$, where P_C is the projection operator within the space of unconstrained states that takes any state to the corresponding constrained state, and P(p) is a projection operator in the space of unconstrained states that depends on the perception p.

O ne could obviously alternatively insert the projection operator P_c before and after the perception operators of H ypothesis B, B', B*, or B*' to get H ypothesis B', B', or B*' respectively.

O ne can also get som ething like H ypothesis C, say H ypothesis \hat{C} , even for unconstrained system s if they have sym m etries (e.g., the Poincare sym m etries of quantum eld theory in a classical M inkow ski spacetim e, though one would not expect these sym m etries to survive when one includes gravity), since one m ight then expect that E (p) should be invariant under the sym m etry group with elements g. Then if one starts with a projection operator P (p) that is not invariant under the action of each group element, say P (p) \notin gP (p)g¹, then one might expect E (p) to be proportional to the sum or integral of gP (p)g¹ over the group elements g. Unless all these di erent gP (p)g¹ 's are orthogonal (which does not appear possible for a continuous sym metry group), the resulting E (p) will generically not be a projection operator, but it can be said to have arisen from one, which is what I would propose as the translation of the marvelous moment assumption to Sensible Q uantum M echanics.

If one can learn what the E (p)'s are, one can compare them with the form s given by these hypotheses and thereby distinguish between the consistent or decohering histories approaches and the m arvelous m om ent approach as I here propose they be applied to conscious perceptions (if indeed any of them do). O fcourse, either of these approaches could be applied without inconsistency to m athem atical probabilities that one m ight wish to de ne in the physical world, but in the present fram ework of Sensible Q uantum M echanics, such probabilities are an unnecessary addition to the ontology.

I should emphasize that in no case am I assuming that the E (p)'s commute for di erent perceptions, or that the sum or integral of the E (p)'s over all perceptions is the identity operator. Neither am I assuming that the resulting expectation values m (p) in the particular quantum state of the universe are normalized so that their sum or integral over all perceptions gives a mite number q, or that this number is unity, though in any case the conditional probabilities given by Eq. (2) are automatically normalized to give unity when summed over a complete set of disjoint sets B of perceptions. Of course, if q is nite, one can simply rescale E (p) to e(p) = E(p)=q, which rescales m (p) to m (p) = m (p)=q that is normalized to give unity when summed or integrated over all perceptions. This rescaling obviously leaves the conditional probabilities of Eq. (2) invariant when m (A \ B) and m (A) there are replaced by m (A \ B) and m (A) respectively. On the other hand, I am sceptical that the simplest consistent description of our universe will give a normalizable m (p) (nite q).

If a perception operator E (p) is a projection operator, and the quantum state of the universe is represented by the pure state j i, one can ascribe to the perception p the pure E verett \relative state"

$$\dot{p}i = \frac{E(p)ji}{kE(p)jik} = \frac{E(p)ji}{h \not E(p)E(p)ji^{1-2}};$$
(4)

A lternatively, if the quantum state of the universe is represented by the density matrix , one can associate the perception with a relative density matrix x = 1

$$p_{p} = \frac{E(p) E(p)}{TrE(p) E(p)]};$$
(5)

E ither of these form ulas can be applied when the perception operator is not a projection operator, but then the meaning is not necessarily so clear.

If one has two perceptions p and p^0 , one can calculate an overlap fraction between them as

$$f(p;p^{0}) = \frac{hE(p)E(p^{0})ihE(p^{0})E(p)i}{hE(p)E(p)ihE(p^{0})E(p^{0})i}:$$
(6)

If the quantum state of the universe is pure, this is the same as the overlap probability between the two Everett relative states corresponding to the perceptions: $f(p;p^0) = jppp^0if$. Thus one might in some sense say that if $f(p;p^0)$ is near unity, the two perceptions are in nearly the same one of the Everett \m any worlds," but if $f(p;p^0)$ is near zero, the two perceptions are in nearly orthogonal di erent worlds. However, this is just a manner of speaking, since I do not wish to say that the quantum state of the universe is really divided up into m any di erent worlds. Thus I do not wish to propose that $f(p;p^0)$ be interpreted as a fundam ental element of Sensible Q uantum M echanics. In any case, one can be conscious only of a single perception at once, so there is no way in principle that one can test any properties of joint perceptions such as $f(p;p^0)$.

5 Quantum Field Theory M odel

A lthough Sensible Q uantum M echanics transpends quantum theories in which space and time are fundamental, and although I believe that such theories will need to be transpended to give a good theory of our universe, it might help to get a better feel for the spacetime properties of perceptions by considering the context of quantum eld theory in an unquantized curved globally-hyperbolic background spacetime in which spacetime points are unambiguously distinguished by the spacetime egometry (so that the Poincare symmetries are entirely broken and one need not worry about integrating over gP (p)g¹'s to satisfy superselection rules for energy, momentum, and/or angular momentum [24]). This simplied model might in some sense be a good approximation for part of the entire quantum state of the universe in a correct theory if there is one that does the time work of Sensible Q uantum M echanics and does give a suitable classical spacetime approximation.

In the H eisenberg picture used in this paper, the quantum state is independent of time (i.e., of a choice of C auchy hypersurface in the spacetime), but the H eisenberg equations of evolution for the fundamental elds and their conjugate m omenta can be used to express the operators E (p) in terms of the elds and m omenta on any C auchy hypersurface. The arbitrariness of the hypersurface m eans that even in this quantum eld theory with a well-de ned classical spacetime, and even with a de nite foliation of the spacetime by a one-parameter (time) sequence of C auchy hypersurfaces, there is no unique physical time that one can assign to any of the perceptions p; they are 'butside' time as well as space.

Furtherm ore, the operators E (p) in this sim pli ed m odelare all likely to be highly nonlocal in terms of local eld operators on any C auchy hypersurface, since quantum eld theories that we presently know do not seem to have enough local operators to describe the complexities of an individual perception, unless one considers high spatial derivatives of the eld and conjugate m om entum operators. However, for a given one-parameter (tim e) sequence of C auchy hypersurfaces, one m ight rather arbitrarily choose to de ne a preferred tim e for each perception p as the tim e giving the Cauchy hypersurface on which the corresponding E (p), if expressed in terms of elds and m om enta on that hypersurface, has in some sense the sm allest spatial spread at that time.

For example, to give a tediously explicit ad hoc prescription, on a C auchy hypersurface labeled by the time t one m ight choose a point P and a ball that is the set of all points w ithin a certain geodesic radius r of the point. Then one can de ne an operator $E^{0}(p;t;P;r)$ that is obtained from E (p) written in terms of the elds and conjugate m on enta at points on the hypersurface by throwing away all contributions that have any elds or conjugate m on enta at points outside the ball of radius r from the point P. Now de ne the overlap fraction

$$F (p;t;P;r) = \frac{hE (p)E^{0}(p;t;P;r)ihE^{0}(p;t;P;r)E (p)i}{hE (p)E (p)ihE^{0}(p;t;P;r)E^{0}(p;t;P;r)i}$$
(7)

(If both E (p) and E 0 (p;t;P;r) were projection operators, and the actual quantum state were a pure state, then F would be the overlap probability between the states obtained by projecting the actual quantum state by these projectors and norm alizing.) If E (p) is nonlocal, this fraction F will be small if the radius r is small but will be nearly unity if the radius r is large enough for the ball to encompass alm ost all of the Cauchy hypersurface. For each perception p, time t, and point P, one can nd the smallest r that gives F = 1=2, say, and call that value of the radius r (p;t;P). Then one can nd the point P = P (p;t) on the hypersurface that gives the smallest r (p;t;P) on that hypersurface for the xed perception p and call the resulting radius r (p;t). Finally, de ne the preferred time t_p as the time t for which r (p;t) is the smallest, and label that smallest value of r (p;t) for the xed perception p as r_p.

If the perception operator E (p) for some hum an conscious perception is not unduly nonlocal in the simpli ed model under present consideration, and if the quantum state of the elds in the spacetime has macroscopic structures that at the time t_p of the perception are fairly well localized (e.g., with quantum uncertainties less than a millimeter, say, which would certainly not be a generic state, even am ong states which give a signi cant m (p) for the perception in question), one might expect that at this time the ball within radius r_p of the point P (p; t_p) on the hypersurface labeled by t_p would be inside a hum an brain. It would be interesting if one could learn where the point P (p; t_p) is in a hum an brain, and what the radius r_p is, for various hum an perceptions, and how the location and size of this ball depends on the perception p.

6 Questions and Speculations

O ne can use the fram ework of Sensible Q uantum M echanics to ask questions and m ake speculations that would be di cult without such a fram ework. I shall here give som e examples, without intending to imply that Sensible Q uantum M echanics itself, even if true, would guarantee that these questions and speculations make sense, but it does seem to allow circum stances in which they might.

First, in the model of quantum eld theory on a classical spacetime with no symmetries, and with a quantum state having well-localized hum an brains on some C auchy hypersurface labeled by time t, one might ask whether it is possible to have two quite dierent perceptions, say p and p^0 , in nearly the same E verett world in the sense of having the f $(p;p^0)$ of Eq. (6) near unity, and giving E (p) and E (p^0) both with the same preferred time $t_p = t$ and both localized (by the rather ad hoc prescription above) in balls in the same brain. In other words, can one brain have two dierent (maximal) perceptions in the same world at the same time, each not aware of the other? Unless we are solipsists, we generally believe this is possible for two separate brains, but would one brain be su cient? Furtherm ore, if it is possible, can the two balls (corresponding to p and p^0 respectively) be overlapping spatially, or need they be separate regions in the brain?

Second, one might ask whether and how the sum (or integral) of the measures (or measure densities) m (p) associated with an individual brain region at the time t depends on the brain characteristics. One m ight speculate that it m ight be greater for brains that are in some sense more intelligent, so that in a crude sense brighter brains have more perceptions. This could explain why you do not perceive yourself to be an insect, for example, even though there are farm ore insects than hum ans. To speculate even further, it might imply that your perception is not atypical even if you perceive yourself to be more intelligent than the average person, as I predict that most of my readers do. Of course, my prediction is based on an assumed selection e ect of those who read physics papers, on my own sinful pride that leads me to assume that physicists are brighter than average, and on a psychologically-projected assumption that most people think they are more intelligent than most people. But if you really had good reasons for believing that you were brighter than average, even before reading this paper, you may not really be justiled in any feeling of atypicality; it m ight simply be that your perceptions, like m ost perceptions in the m easure m (p)dp, are associated with brighter-than-average people.

I should emphasize that even if this wild conjecture, which is not an inevitable consequence of Sensible Quantum M echanics (but which can be considered under this fram ework in a way that m ight be di cult under other fram eworks), could be shown to be true, I would not propose that a more intelligent person be assigned more value or be given more political rights or privileges. It would just say that, when weighted by the quantum -m echanical operators E (p) for perceptions, it is conceivable that more intelligent people would have the bulk of the weight rather than being unusual. But being in this new ly-de ned majority (if indeed it is the bulk of the weight) should not confer more individual political rights or privileges, just as with the weighting of num bers the majority of people who are econom ically poor are not given more individual political rights or privileges than the minority of people who are econom ically rich.

Third, one might conjecture that an appropriate measure on perceptions might give a possible explanation of why most of us perceive ourselves to be living on the same planet on which our species developed. This observation might seem surprising when one considers that we may be technologically near the point at which we could leave Earth and colonize large regions of the Galaxy, presum ably greatly increasing the number of hum and beyond the roughly 10^{11} that are believed to have lived on Earth. If so, why don't we have the perceptions of one of the vast num bers of hum ans that may be born away from Earth? One answer is that some sort of doom is likely to prevent this vast colonization of the G alaxy from happening [25, 26, 27, 28], though these arguments are not conclusive [29]. Although I would not be surprised if such a doom were likely, I would na vely expect it to be not so overwhelm ingly probable that the probability of vast colonization would be as smallas is the presum ably very sm all ratio of the total num ber of hum ans who could ever live on Earth to those who could live throughout the G alaxy if the colonization occurs. Then, even though the colonization may be unlikely, it may still produce a higher measure for conscious perceptions of hum ans living o Earth than on it.

However, another possibility is that colonization of the G alaxy is not too in probable, but that it is mostly done by self-replicating computers or machines who do not tolerate m any hum ans going along. If the num ber of these dom inate hum ans as \intelligent" beings, one m ight still have the question of why we perceive ourselves as being hum ans rather than as being one of the vastly greater num bers of such machines. But the explanation m ight simply be that the weight of conscious perceptions (the sum or integral of the m (p)'s corresponding to the type of perceptions under consideration) is dom inated by hum an perceptions, even if the num ber of \intelligent" beings is not. In other words, hum an brains m ay be m uch more e cient in producing conscious perceptions than the kinds of self-replicating com puters or machines are more \intelligent" than hum ans in terms of inform ation-processing cababilities and yet are less e cient in producing conscious perceptions, our perceptions of being hum an would suggest that the measure of perceptions is not merely correlated with intelligence.

It m ight be tempting to take the observations that these speculations m ight explain (your perception of yourself as hum an rather than as insect or even possibly as more intelligent than average hum ans, if my prediction of that is correct, and our perception of ourselves as hum ans on our hom e planet) as evidence tending to support the speculations. For example, if one assigns prior probabilities P (H_i) to an exhaustive set of hypotheses H_i (say each giving a quantum state of the universe and a set of operators E (p) in Sensible Q uantum M echanics), and then if the conditional probability of a perception p, given the hypothesis H_i, were P (pH_i), by Bayes' rule the posterior conditional probability of the speculation H_i would be

$$P (H_{i} \dot{P}) = \frac{P (p H_{i}) P (H_{i})}{p (p H_{i}) q_{i}} = (q_{i} H_{i}) P (H_{i})$$
(8)

Unfortunately, in Sensible Q uantum M echanics one has at most measure densities m (p) for individual perceptions, and so one cannot unam biguously give the probability P (pH_i) without some speci cation of the norm alization of m (p) in the hypothesis H_i. Possibly one can replace the probability for a perception with the 'typicality' of the perception [15, 16]. W ithout some e such solution to this problem, I do not see how to use observations to turn prior probabilities (say given by some function of the sim plicity of the quantum state and perception operators) into posterior probabilities for quantum states and perception operators. O there is it would appear that one could only start with a given quantum state and set of perception operators, and then can one calculate the measure for all perceptions by Eq. (1) and use Eq. (2) to calculate conditional probabilities for sets of perceptions.

7 Conclusions

In conclusion, I am proposing that Sensible Q uantum M echanics, with its M easure A xiom for Perceptions and its Sensible Q uantum A xiom above, is the best fram ework we have at the present level for understanding conscious perceptions and the interpretation of quantum m echanics. Of course, the fram ework would only becom e a complete theory once one had the set of all perceptions p, the basic m easure dp on it, the perception operators E (p), and the quantum state of the universe.

Even such a complete theory of perceptions and the physical world need not be the ultimate simplest complete theory. There might be a simpler set of unifying principles from which one could in principle deduce the perceptions, basic measure, perception operators, and quantum state, or perhaps some simpler entities that replaced them . For example, although in the present fram ework of Sensible Quantum Mechanics, the physical world (i.e., the quantum state), along with the perception operators, determ ines the measure density for perceptions in the mental world, there m ight be a reverse e ect of the mental world a ecting the physical world to give a simpler explanation than we have at present of the correlation between will and action (why my desire to do something I feel am capable of doing is correlated with my perception of actually doing it, i.e., why I \do as I please"). If the quantum state is partially determ ined by an action functional, can desires in the mental world a ect that functional (say in a coordinate-invariant way that therefore does not violate energy-momentum conservation)? Such considerations may call for a more unied fram ework than Sensible Quantum Mechanics, which one might call Sensational Quantum Mechanics. Such a more uni ed fram ework need not violate the limited assumptions of Sensible Quantum Mechanics, though it might do that as well and perhaps reduce to Sensible Quantum Mechanics only in a certain approximate sense.

To explain these fram eworks in term s of an analogy, consider a classical model of spinless massive point charged particles and an electrom agnetic eld in M inkow ski spacetime. Let the charged particles be analogous to the physical world (the quantum state of the universe), and the electrom agnetic eld be analogous to the mental

world (the set of perceptions with its measure m (p)dp). At the level of a sim plistic m aterialist m ind-body philosophy, one m ight m erely say that the electrom agnetic eld is part of, or perhaps a property of, the material particles. At the level of Sensible Quantum Mechanics, the charged particle worldlines are the analogue of the quantum state, the retarded electrom agnetic eld propagator (Coulom b's law in the nonrelativistic approximation) is the analogue of the perception operators, and the electrom agnetic eld determ ined by the worldlines of the charged particles and by the retarded propagator is the mental world. (Here you can see that this analoque of Sensible Quantum Mechanics is valid only if there are no free incoming electrom agnetic waves.) At the level of Sensational Quantum Mechanics, at which the mental world may a ect the physical world, the charged particle worldlines are partially determ ined by the electrom agnetic eld through the change in the action it causes. (This more unied fram ework better explains the previous level but does not violate its description, which simply had the particle worldlines given.) At a yet higher level, there is the possibility of incoming free electrom agnetic waves, which would violate the previous fram eworks that assumed the electrom agnetic eld was uniquely determ ined by the charged particle worldlines. Finally, at a still higher level, there might be an even more unifying fram ework in which both charged particles and the electrom agnetic eld are seen as modes of a single entity (e.g., to take a popular current speculation, a superstring).

Therefore, although it is doubtful that Sensible Q uantum M echanics is the correct fram ework for the nalunifying theory (if one does indeed exist), it seems to me to be a move in that direction that is consistent with what we presently know about consciousness and the physical world.

A cknow ledgm ents

I consciously perceive a gratitude for having had fruitful discussions or correspondence on this general subject with D avid A lbert, Andrei Barvinsky, A rvind Borde, D avid Boulware, Patrick Brady, Howard Brandt, Bruce Campbell, Brandon Carter, D avid D eutsch, Valeri Frolov, M urray G ell-M ann, Farhad G hoddoussi, Shelly G oldstein, R ichard G ott, Jim Hartle, G eo Hayward, Bei-Lok Hu, V igar Husain, Rocky Kolb, Tom as Kopf, Pavel K rtous, W emer Israel, John Leslie, Andrei Linde, M ichael Lockwood, Robb M ann, John Polkinghome, D avid Salopek, Abner Shim ony, Euan Squires, Lenny Susskind, B ill U nruh, A lex V ilenkin, John W heeler, and various others whose nam es aren't in my immediate perception. I am expecially grateful to my wife C athy for helping me becom emore aware of my feelings. Finally, nancial support has been provided by the Natural Sciences and Engineering C ouncil of C anada.

References

- [1] R.B.Griths, J.Stat.Phys. 36, 219 (1984); Phys. Rev. Lett. 70, 2201 (1993).
- [2] R.Omnes, J. Stat. Phys. 53, 893 (1988); 53, 933 (1988); 53, 957 (1988); 57, 357 (1989); 62, 841 (1991); Ann. Phys. (NY) 201, 354 (1990); Rev. M od. Phys. 64, 339 (1992).
- [3] M. Gell-M ann and J. B. Hartle, in Complexity, Entropy and the Physics of Information, Santa Fe Institute Studies in the Sciences of Complexity, Vol. VIII, edited by W. H. Zurek (Addison-Wesley, Redwood City, 1990), p. 425; Phys. Rev. D 47, 3345 (1993).
- [4] J.B.Hartle, in Quantum Cosmology and Baby Universes, Jerusalem W inter School for Theoretical Physics, Vol. 7, edited by S.Coleman, J.B.Hartle, T. Piran, and S.W einberg (W orld Scientic, Singapore, 1991), p. 65.
- [5] M.Gell-M ann and J.B.Hartle, in PhysicalO rigins of T in e A sym m etry, edited by J.J.Halliwell, J.Perez-M ercador, and W.H.Zurek (Cambridge University Press, Cambridge, 1994), p. 311, gr-qc/9304023.
- [6] S. Goldstein and D. N. Page, \Linearly Positive Histories: Probabilities for a Robust Family of Sequences of Quantum Events," Alberta-Thy-43-93, grqc/9403055, submitted to Phys. Rev. Lett.
- [7] H. Everett, III, Rev. M od. Phys. 29, 454 (1957); B. S. DeW itt and N. Graham, eds., The M any-W orlds Interpretation of Q uantum M echanics (Princeton University Press, Princeton, 1973).
- [8] M. Lockwood, M ind, B rain and the Quantum : The Compound 'I' (BasilBlackwell, Oxford, 1989).
- [9] D.Bohm, Phys. Rev. 85, 166 and 180 (1952).
- [10] J. von Neum ann, M athem atische G rundlagen der Quantenm echanik (Springer, Berlin, 1932), pp.184 ; translated by T.Beyer as M athem atical Foundations of Quantum M echanics (Princeton University Press, Princeton, 1955), pp. 347 ; reprinted in Quantum Theory and M easurem ent, edited by J.A.W heeler and W.H.Zurek (Princeton University Press, Princeton, 1983), p. 549.
- [11] E.P.W igner, \Remarks on the M ind-Body Question," in The Scientist Speculates, edited by I.J.Good (Heinem ann, London, 1961; Basic Books, New York, 1962), p. 284; reprinted in E.W igner, Symmetries and Re ections (Indiana University Press, Bloom ington, 1967), p. 171, and in Quantum Theory and Measurem ent, edited by J.A.W heeler and W.H.Zurek (Princeton University Press, Princeton, 1983), p. 168; J.A.W heeler (private communication, 1994) inform sme that W igner later abandoned this view point.
- [12] W.G.Unruh, \Varieties of Quantum Measurement," to be published in Proceedings of the New York Academy of Sciences Conference on Quantum Measurement, Baltimore, Maryland, June 19-22, 1994, edited by D.Greenberger (New York Academy of Sciences, New York, 1985), hep-th/9410168.

- [13] D.Hume, A Treatise of Human Nature, reprinted from the original edition in three volumes and edited by L.A. Selby-Bigge (Clarendon, Oxford, 1888).
- [14] S.Goldstein (private communication).
- [15] D.N.Page, \Information Loss in Black Holes and/or Conscious Beings?" to be published in Heat Kernel Techniques and Quantum Gravity, edited by S.A. Fulling (D iscourses in M athematics and Its Applications, No. 4, Texas A & M University Department of M athematics, College Station, Texas, 1995), hepth/9411193.
- [16] D.N.Page, \Sensible Quantum Mechanics: Are Only Perceptions Probabilistic?" in preparation.
- [17] E.J.Squires, Found. Phys. Lett. 1, 13 ((1987); Conscious M ind in the Physical W orld (A dam H ilger, B ristol and New York, 1990); Synthese 89, 283 (1991); 97, 109 (1993).
- [18] D. A Ibert and B. Loewer, Synthese 77, 195 (1988); 86, 87 (1991); D. Z. A Ibert, Quantum Mechanics and Experience (Harvard University Press, Cambridge, Massachusetts, 1992).
- [19] H.P.Stapp, M ind, M atter, and Quantum M echanics (Springer-Verlag, Berlin, 1993).
- [20] M. Lockwood, in Erwin Schrodinger: Philosophy and the Birth of Quantum M echanics, edited by M. Bitbol and O. D arrigol (Editions Frontieres, G if-sur-Y vette C edex, 1992), p. 363.
- [21] M. Lockwood (private communication).
- [22] D. Part, Phil. Rev. 80 (1971), reprinted in Personal Identity, edited by J. Perry (University of California Press, Berkeley, 1975), p. 199; Reasons and Persons (Clarendon Press, Oxford, 1984).
- [23] D.N.Page, in PhysicalO rigins of T in e A sym m etry, edited by J.J.Halliwell, J.Perez-M ercader, and W.H.Zurek, (C am bridge University Press, C am bridge, 1994), p. 287, gr-qc/9303020.
- [24] D.N.Page and W.K.Wootters, Phys. Rev. D 27, 2885 (1983).
- [25] B.Carter, Phil. Trans. Roy. Soc. Lond. A 310, 347 (1983).
- [26] J. Leslie, Bull. Canad. Nucl. Society 10, 10 (1989); Interchange 21, 49-58 (1990); Universes (Routledge, London and New York, 1989), p. 214; Phil. Quart. 40, 65 (1990); 42, 85 (1992); M ind 101, 521 (1992); 102, 489 (1993); M ath. Intelligencer 14, 48 (1992); 15, 5 (1993); Interchange 23, 289 (1992); J. Appl. Phil. 11, 31 (1994).
- [27] H.B.Nielsen, Acta Physica Polonica B 20, 427 (1989).
- [28] J.R.Gott III, Nature 363, 315 (1993); 368, 108 (1994).

[29] T.Kopf, P.Krtous, and D.N.Page, \Too Soon for Doom G loom ?" A lberta-Thy-17-94, gr-qc/9407002, submitted to Nature.