# D ynam ics of a self-gravitating thin string in scalar-tensor theories of gravitation

B.Boisseau and B.Linet<sup>y</sup>

Laboratoire de M athem atiques et Physique T heorique CNRS/UPRES-A 6083, Universite Francois Rabelais

Faculte des Sciences et Techniques, Parc de Grandmont 37200 TOURS, France

#### A bstract

We exam ine the dynam ics of a self-gravitating string in the scalar-tensor theories of gravitation by considering a thin tube of matter to describe it. For a class of solutions, we obtain in the generic case that the extrinsic curvature of the world sheet of the central line is null in the lim it where the radius of the string tends to zero. However, if we impose a speci c constraint on the behaviour of the solution then we nd that only the mean curvature of the world sheet of the central line vanishes which is just the Nambu-Goto dynamics. This analysis can include the massless dilatonic theories of gravity.

PACS num bers: 11.27, 04.25

#### 1 Introduction

The know ledge of the equations of motion of the cosm ic strings is essential to predict the evolution and the physical consequences of an eventual string network in the early U niverse [1, 2, 3]. In general, the N am bu-G oto dynam ics of the cosm ic strings is postulated in a background curved spacetime. Yet, there is no proof that a self-gravitating cosm ic string obeys the N am bu-G oto dynam ics as the strings in eld theory moving in the M inkow ski spacetime [4, 5]. On the contrary, several authors [6, 7, 8, 9] state in general relativity that a world sheet whose the points are conical singularities is totally geodesic which is a very particular case of a m inim al surface characteristic of the history of a N am bu-G oto string.

However, a recent paper [10] showed in general relativity the existence of self-gravitating thin strings of m atter having their central line governed by the N am bu-G oto dynam ics in the lim it where their radius goes to zero.

E-mail: boisseau@ cel phys.univ-tours.fr

<sup>&</sup>lt;sup>y</sup>E-mail: linet@cel.phys.univ-tours.fr

The purpose of this work is to exam ine the dynam ics of a self-gravitating cosm ic string in the scalar-tensor theories of gravitation [11, 12, 13] which generalize the B rans-D icke theory [14]. There is a renewed interest of these theories of gravitation due to the importance of the scalar eld or dilaton in the low-energy elective string theory presently used to study the early U niverse. In general the dilaton is massive and consequently the theory is practically the general relativity for distances much larger than the range of the dilaton. However, a massless dilaton would be possible [15] and in this case our method can be immediately transposed.

The straight U (1) gauge cosm ic string has been studied in the Brans-D icke theory [16], in the scalar-tensor theories [17] and in the dilatonic theories of gravity [18]. Now the distribution of matter vanishes practically beyond the range  $l_0$  of the massive gauge and Higgs elds, therefore we can substitute to the U (1) gauge cosm ic string a tube of matter describing a cylinder of radius  $l_0$  having as exterior solution the vacuum solution which coincides with the asymptotic solution of the U (1) gauge cosm ic string.

Then, for a thin string describing a tube of matter of arbitrary shape, we adopt the assumption that the exterior metric is locally the one describing a straight string. The interior solution is constrained from this. The aim of the work is to nd the equations of motion of the central line of the thin tube of matter in the limit where its radius goes to zero.

The plan of the work is as follows. In section 2, we recall the basic equations of the scalar-tensor theories and the dilatonic theory. We give in section 3 the form of the interior and exterior metric representing a static cylinder of matter. In section 4, we introduce a self-gravitating thin string of arbitrary shape. We derive the equations of motion in section 5 by taking the limit where its radius tends to zero. We add in section 6 some concluding remarks.

# 2 Gravitational eld equations

The gravitational eld variables of the scalar-tensor theories are the metric  $\hat{g}$  of the spacetime and the scalar eld . Nevertheless, it is more convenient to use a non-physical metric g dened by  $g = A^2$  () $\hat{g}$  where the function A () characterizes each scalar-tensor theory. The matter elds M are coupled as usual to the metric  $\hat{g}$  within the Lagrangian density  $L_M$  [M; $\hat{g}$ ]. The eld equations can be then derived from the action

$$S = \frac{1}{16 \text{ G}}^{Z} d^{4}x^{P} - g (R \quad 2g \ (e \quad ) + d^{4}xL_{M} [M; A^{2}(M)]$$
(1)

where G is the coupling gravitational constant which can be related to the Newtonian constant. By varying action (1) with respect to g and , we get the gravitational eld equations

R 20 0 = 8 G (T 
$$\frac{1}{2}$$
Tg); (2)

2 = 4 G ()T with () = 
$$\frac{1}{A}\frac{dA}{d}$$
 (3)

and by varying with respect to M the matter eld equations. The source T is related to the conserved energy-momentum tensor  $\hat{T}$  by  $T = A^2$  () $\hat{T}$ ; it satisfies the equation

$$r T = ()TQ$$
 : (4)

We emphasize that we cannot give independently g and to deduce from them the source T  $\cdot$  W e have necessarily the condition

R 2g Q Q = 
$$\frac{2}{()}$$
2 (5)

resulting from the trace of eld equations (2). We now turn to the inverse. We consider that g and verify (5). We can calculate T by (2) and then (3) is automatically verified. Moreover it is easy to see that the source T satisfies (4).

We now consider the massless dilatonic theory of gravity. The eld variables are the string metric  $\hat{g}$ , the dilaton and the matter eld <sub>M</sub>. The metric g in the Einstein frame is de ned by g = exp(2) $\hat{g}$ . Following Damour and Polyakov [15], the eld equations can be derived from the following action

$$S = \frac{1}{16 \text{ G}}^{Z} d^{4}x^{P} - g (R \quad 2g \ (e \quad ) + d^{4}xe^{2} L_{M} [_{M};e^{2}g]:$$
(6)

in the simplest case. We obtain the gravitational eld equations

R 20 0 = 8 
$$\overline{G}$$
 (T  $\frac{1}{2}$ Tg); (7)

$$2 = 4 \overline{G} \quad \text{with} \quad = p \frac{1}{g} \frac{(e^2 L_M)}{g}$$
(8)

where  $\overline{G}$  is a bare constant coupling and also the eld matter equations. The quantity is not directly related to the T; it depends on the matter eld <sub>M</sub> as T. We point out that the physical energy-momentum tensor T is not conserved since we have

$$r T = 0$$
 : (9)

As in the previous case, we cannot give independently g and but the condition takes into account the matter eld.

## 3 Static, cylindrically sym m etric solutions

The vacuum solutions to eld equations (2) and (3) with T = 0 which are static and cylindrically symmetric are known as solutions dependent only on one coordinate [19]. By adding the boost symmetry, they have nally the general form

$$ds^{2} = -\frac{\bar{b}^{\prime}}{2k} (dt^{2} + dz^{2}) + d^{2} + B^{2} (\bar{b})^{2} - \frac{\bar{b}^{\prime}}{2k} d'^{2}; \quad (10)$$

$$= _{0} + \ln \frac{-\frac{1}{b}}{k} \quad \text{with} \quad = \quad \frac{s}{k} \frac{-1}{4} k \quad (11)$$

in the coordinate system (t;z; ;') with 0 < 2. For k = 0, we recover the metric with conical singularity of the general relativity. For  $k \in 0$ , the curvature tensor of metric (10) blow up at  $= \bar{l}_0$ . We can perform a rede nition of and a rescaling of t and z to set B = 1. Metric (10) depends only on two arbitrary constants.

M etric (10) and scalar eld (11) are respectively the asymptotic metric and the asymptotic scalar eld of the solution for a straight U (1) gauge cosm is string. Then, the constants appearing in expression (10) can be approximatively determined in this case [16, 17, 18]. The lengths and  $l_0$   $\bar{l}_0$  have the same order of magnitude as the range  $l_0$  of the gauge and H iggs elds... We emphasize that k is quadratic in G and consequently very small but strictly positive.

We substitute a tube of matter representing a cylinder in place of this U (1) gauge cosm ic string. Its energy-momentum tensor will be determined by the data of the interior metric via the gravitational eld equations. The interior metric is regular at the central line = 0 and it matches to the exterior metric (10) at =  $\frac{1}{2}$ , likewise the scalar eld. In the next sections, we will perform a limit process in which  $\frac{1}{2}$  tends to zero but the constants B and k keep the same values and in which and  $\frac{1}{2}$  are dependent on  $\frac{1}{2}$ . To this end, we put the following family of regular metrics parametrized by

$$ds^{2} = \underbrace{\frac{1}{2}}_{k}^{8} \frac{1}{(dt^{2} + dz^{2}) + d^{2} + b^{2} - t^{2}}_{k}^{1} \frac{1}{(dt^{2} + dz^{2}) + d^{2} + b^{2} (-\frac{1}{b})^{2}}_{k} \frac{-\frac{1}{b}}{(dt^{2} + dz^{2}) + d^{2} + b^{2} (-\frac{1}{b})^{2}} \frac{-\frac{1}{b}}{(dt^{2} + b^{2}) + b^{2} +$$

where the assumption of regularity at = 0 is ensured by the behaviours

$$X(x) = 1 + X_2 x^2 +$$
 and  $h(x) = x \frac{1}{3}x^3 +$ 

The requirement X (0) = 1 in (12) xes the rescaling of the coordinates t and z and consequently it justiles to keep the constant B. The matching conditions at  $= 1_0$  require the continuity of the components of the metric and their rst derivative with respect to . A fler som e manipulation, we get

:

It is easy to show that a change of parameter in form (12) corresponds to a rede nition of the functions h and X which is compatible with (13). If  $_1$  and  $_2$  denote two parameters,

we have  $_1h_1(=_1) = _2h_2(=_2)$  and  $X_1(=_1) = X_2(=_2)$ . Thus, we can choose the parameter of the family of metrics (12) to be

In general relativity, another geom etrical choice had been m ade [10].

For the scalar eld, we choose the following form

$$= \underbrace{\begin{array}{cccc} & & & & \\ \underbrace{\$} & _{0} + Z & - & & 0 & < \underbrace{1}_{0} \\ & & & \underbrace{-1}_{0} & & \\ \underbrace{\$} & _{0} + & \underbrace{\ln} & \underbrace{-1}_{0} & & > \underbrace{1}_{0} \end{array}$$
(15)

taking account (14) and with the behaviour

$$Z(x) = Z_2 x^2 + Z_4 x^4 +$$

to ensure the regularity at = 0. Scalar eld (15) and its rst derivative with respect to are continuous at  $= \frac{1}{2}$ ; we get

$$Z = \frac{l_0}{l_0} = \ln \frac{l_0}{l_0} = \ln \frac{l_0}{l_0} = \frac{l_0}$$

In sum marizing, it is possible to nd a family of metrics (12) and scalar elds (15) representing a cylinder of matter of radius  $l_0$ . The exterior solution is characterized by xed constants B and k. Now we can take  $l_0$ ! 0 but without considering a line source which is a di cult concept in non-linear eld equations [20]. For given functions h and X, we see from (13) that the ratio  $l_0$  = remains constant. It is equivalent to take ! 0 in the family of solutions (12) and (15).

W hen we will use these solutions we will furtherm one need to the continuity of the second derivative with respect to of the components of metric (12) and scalar eld (15). It is not necessary to write down them . We con ne ourselves to these solutions. In consequence, T vanishes at  $= \frac{1}{2}$ .

### 4 A model of self-gravitating thin string

We now consider a general tube of matter with a circular section of nite radius  $l_0$  whose the central line has an arbitrary shape. The central line sweeps a timelike world sheet. A local coordinate system (<sup>A</sup>; <sup>a</sup>) can be then attached to the spacetime by taking the two coordinates <sup>A</sup> of the world sheet and the two geodesic coordinates <sup>a</sup> pointing in a direction orthogonal to the world sheet. The polar coordinates and ' are related to <sup>a</sup> by <sup>1</sup> = cos' and <sup>2</sup> = sin'. The world sheet is represented by the equation = 0.

W e in agine that this cylinder of matter describes a cosm ic string. By assuming that the typical variation on which we have a change in the direction of the central line is much

larger than the radius  $l_0$ , we can suppose that its metric can approximatively be identied with metric (12) in a small neighbourhood. We insist that the constants B and k have the same values during all the history of the tube of matter. We restrict ourselves to metric having no cross terms and so we put in the coordinates (<sup>A</sup>; ;')

$$ds^{2} = \iint_{i}^{8} X - g_{AB} (^{c}; ;') d^{A} d^{B} + d^{2} + {}^{2}h^{2} - d'^{2} 0 < \frac{1}{2}$$

$$= \iint_{i}^{1} g_{AB} (^{c}; ;') d^{A} d^{B} + (17)$$

$$= d^{2} + B^{2} (\overline{1})^{2} - \frac{1}{2} d'^{2} > \frac{1}{2}$$

where the typical variation of components  $g_{AB}$  is much larger than . The matching conditions at  $= \frac{1}{2}$  are given by (13) as in the previous section. Instead of polar coordinates, we prefer express metric (17) in coordinates <sup>a</sup> and we denote  $g_{ab}$  the metric of the surface  $^{A}$  = constant. These components are functions of  $^{a}$  = . We remark that  $g_{ab} = {}_{ab} + 0$  (<sup>2</sup>).

A coording to metric (17), the metric of the world sheet is

$$_{AB} = g_{AB} \dot{j}_{=0} \tag{18}$$

and the extrinsic and mean curvature of the world sheet are

0

$$K_{aAB} = \frac{1}{2} \varrho_a g_{AB} j_{=0} \text{ and } K_a = K_{aAB}$$
<sup>AB</sup>
<sup>(19)</sup>

since X (0) = 1 and X  $^{0}(0) = 0$ .

Further calculations, that we do not reproduce here, show that the scalar eld of such a solution is very particular. We anticipate and we take directly

$$= \begin{array}{c} & & & & & \\ & & & \\ & & & &$$

The matching conditions at = 1 are given by (16). Since h, X and Z are supposed known from the straight string, we obtain from (5) that the components  $g_{AB}$  giving the evolution of the string of arbitrary shape must verify

$$R = 2g^{ab} \mathcal{Q}_{a} Z \mathcal{Q}_{b} Z + \frac{2}{(_{0} + Z)} p \frac{1}{\det g_{ab} \det g_{AB} X} \mathcal{Q}_{a} \stackrel{q}{\det g_{ab} \det g_{AB} X} \mathcal{Q}^{ab} \mathcal{Q}_{b} Z : (21)$$

We add moreover that the second derivative with respect to of the metric and the scalar eld are continuous. We think that the cylinder of matter described by metric (17) and scalar eld (20) is reasonably general in despite of (21) and it allows us to obtain results about the equations of motion of the central line having an arbitrary shape.

### 5 Equations of motion of the string

W e proceed as in the previous paper [10]. The method is based on an expansion in powers of 1= of the geometrical quantities and of the scalar eld associated to the family of solutions (17) and (20). We use the fact that

$$g_{ab} j_{=b} = O(\frac{1}{0}); \quad @_{c}g_{ab} j_{=b} = O(\frac{1}{0}); \quad X j_{=b} = O(\frac{1}{0}); \quad @_{c}X j_{=b} = O(\frac{1}{0}); \quad :::$$

On the contrary, we use

$$g_{AB} j_{=b} = O(\frac{1}{0}); \quad Q_{C}g_{AB} j_{=b} = O(\frac{1}{0}); \quad :::$$

since the typical variation of  $g_{AB}$  is much larger than . The variations of the di erent quantities are thus characterized by . We now expand in powers of 1= the gravitational eld equations on the circle =  $l_0$ . By our choice of matching conditions, we have T  $j_{=l_0} = 0$  and therefore we obtain the following form of the equations

R 
$$j_{=b}$$
 20 0  $j_{=b}$  L  $\frac{1}{2}$  + L  $\frac{1}{-1}$  + L  $\frac{1}{-1}$  = 0; (22)

The term s in  $1=2^{\circ}$  in (22) and (23) vanish identically because they contain only  $g_{ab}$ , X and Z and their rst and second derivatives at  $= 1_0$  which satisfy the vacuum eld equations of the straight string (12) and (15). For instance, we have

$$L_{AB} \quad \frac{1}{2} = \frac{1}{2^{p} \frac{1}{\det g_{ab}}} e_{a} \quad \frac{q}{\det g_{ab}} g^{ab} e_{b} X \quad g_{AB} \quad j_{ab} = 0$$

as we can easily veried it, likewise for  $L_{ab}$  (1= <sup>2</sup>),  $L_{aA}$  (1= <sup>2</sup>) and L (1= <sup>2</sup>).

We now full a limit process in which ! 0, i.e.  $\frac{1}{2}!$  0, the ratio  $\frac{1}{2}$  remaining constant. To analyse the now remaining 1= order equations, we point out that we have

$$AB = \lim_{l_0 \neq 0} g_{AB} j_{=l_0} \text{ and } K_{aAB} = \frac{1}{2} \lim_{l_0 \neq 0} \theta_a g_{AB} j_{=l_0}$$

from (18) and (19) since the typical variation of  $g_{AB}$  is much larger than . The leading terms in 1= in (22) and (23) must go to zero as tends to zero. It is straightforward to carry out the calculation of the (A;B) components

$$\lim_{i \to 0} L_{AB} = F_{1}^{b} - \frac{a}{0} K_{bAB} - F_{2}^{b} - \frac{a}{0} A_{B} K_{b} = 0$$
(24)

where

$$F_{1}^{b} \stackrel{a}{\longrightarrow} = \lim_{! 0} \frac{1}{2}^{h} X \quad (a_{a}g^{ab} + a_{ad}g^{db} \quad g^{ab}(a_{a}X^{i}j_{=b};$$

$$F_{2}^{b} \stackrel{a}{\longrightarrow} = \lim_{! 0} \frac{1}{2}g^{ab}(a_{a}X^{j}j_{=b};$$
(25)

the (a; b) components

$$\lim_{i \to 0} L_{ab} = F_{ab}^{d} - K_{d} = 0$$
 (26)

where

$$F_{ab}^{d} \stackrel{a}{\longrightarrow} = \lim_{\substack{i=0\\ j \in \mathcal{A}}} \frac{d}{ab} \frac{1}{2} \frac{\theta_{b} X}{X} \stackrel{d}{a} \frac{1}{2} \frac{\theta_{a} X}{X} \stackrel{\pi}{b} \stackrel{\pi}{j}_{=b}; \qquad (27)$$

the (a;A) components

$$L_{aA} \quad \frac{1}{2} = 0 \tag{28}$$

#

and the scalar eld equation

$$\lim_{i \to 0} L \frac{1}{i} = F_3^{a} \frac{a}{0} K_a = 0$$
 (29)

where

$$F_{3}^{a} - \frac{a}{0} = \lim_{l \to 0} g^{ab} Q_{b} \quad j_{=l_{0}} :$$
 (30)

The functions  $F_1^{b}$ ,  $F_2^{b}$ ,  $F_{ab}^{c}$  and  $F_3^{b}$  are functions of  ${}_0^{a}=l_0$ , i.e. they depend only on the polar angle '. They are proportional to  ${}^a=l_0$ . We now turn to equation (24); since it must be veried for all polar angles, we have in the generic case

$$K_{aAB} = 0$$
: (31)

Then, the other equations (26) and (29) are obviously satis ed. So, the world sheet swept by the central line of the string is totally geodesic.

However, another possibility exists as in general relativity [10]. We can explicitly calculate  $F_1^{b}$ ; we get "

$$F_{1}^{b} = \frac{b}{2l_{0}} \frac{l_{0}}{h^{2}} \frac{X}{h^{2}} \frac{1}{l_{0}}hh^{0} 1 + X^{0}$$
(32)

for h,  $h^0$  and X , X  $^0$  at  $l_0$  = . By in posing the following constraint on the functions h and X at the value  $l_0$  =

$$h \quad \frac{l_0}{2} \quad h^0 \quad \frac{l_0}{2} \quad + \quad h^2 \quad \frac{l_0}{2} \quad \frac{X^0(l_0 = 1)}{X(l_0 = 1)} = \frac{l_0}{2};$$
(33)

we annihilate the coe cient of the extrinsic curvature in (24). So, equation (31) disappears and the remaining equations give

$$K_{a} = 0$$
: (34)

This expresses that the world sheet of the central line is m inimal which is the behaviour of the N am bu-G oto string. We point out that constraint (33) is invariant under the choice of parametrization of the family of solutions (17) and (20).

In order to obtain the Nambu-G oto dynamics, we must nd an interior metric of the straight string whose the functions h and X satisfy the matching conditions (13), the second order junction conditions and constraint (33) at  $= \frac{1}{2}$ . This corresponds to specify the behaviour of the matter.

#### 6 Conclusion

In the problem of the equations of motion of a self-gravitating thin cosm is string in the scalar-tensor theories of gravitation, we have found that in generic case the world sheet of the central line is totally geodesic i.e.  $K_{aAB} = 0$  for the metric g and in consequence for the physical metric  $\hat{g}$  since the rst derivative of the scalar eld vanishes at = 0. However by in posing constraint (33) on the behavior of metric (17), the scalar eld being given by (20), we have seen that only the mean curvature  $K_a$  vanishes, characteristic of the N am bu-G oto dynam ics. This result coincides with the one in general relativity [10].

However, we have obtained this result only for a class of solutons with a particular scalar eld. Instead of form (17) of the metric, we have attempted more general forms as

$$ds^{2} = \begin{cases} \begin{cases} x & -g_{AB} (c; c) d^{A} d^{B} + (c; c) g_{ab} d^{a} d^{b} & 0 < b \\ & & & \\ &$$

but the fact that the eld equations yield a scalar eld independent on A remains.

To obtain this, we have assumed that the scalar eld does not interact directly with the matter within the framework of the scalar-tensor theories. Our method can be also applied for the massless dilatonic theory whose gravitational eld equations (7) and (8) are similar. We obtain the same result. In the case of a massive dilaton, with eventually a specie conteraction with the energy per unit of length for the string [21], the problem of the equations of motion should be taken up.

In our m ethod for nding the equations of m otion, we have replaced the cosm ic string de ned in eld theory by a tube of m atter. An open question is to determ ine the equations of m otion w ithin the coupled theory of the eld theory and the gravity when the range of the m assive gauge and Higgs elds tends to zero.

#### A cknow ledgm ent

W e would like to thank N.M oham m edi for helpful discussions.

#### References

- Vilenkin A and Shellard E P S 1994 Cosm ic String and Other Topological Defects (Cambridge: Cambridge University Press)
- [2] Brandenberger R H 1994 Int. J. M od. Phys. A 9 2117

- [3] Hindmarsh M B and Kibble T W B 1995 Rep. Prog. Phys. 58 477
- [4] Nielsen H B and Olesen P 1973 Nucl. Phys. B 61 45
- [5] Forster D 1974 Nucl. Phys. B 81 84
- [6] Vickers JA G 1987 Class. Quant. Grav. 4 1
- [7] V P.Frolov V P, Israel W and Unruh W G 1989 Phys. Rev. D 39 1084
- [8] Unruh W G, Hayward G, Israel W and M dM anus D 1989 Phys. Rev. Lett. 62 2897
- [9] Clarke C J S, Ellis G F R and Vickers J A 1990 Class. Quant. Grav. 7 1
- [10] Boisseau B, Charmousis C and Linet B 1997 Phys. Rev. D 55 616
- [11] Bergm ann P G 1968 Int. J. Theor. Phys. 1 25
- [12] W agoner R V 1970 Phys. Rev. D 1 3209
- [13] Nordtverdt Jr K 1970 A strophys. J. 161 1059
- [14] Brans C and Dicke R H 1961 Phys. Rev. 124 925
- [15] Damour T and Polyakov A M 1994 Nucl. Phys. B 423 532
- [16] Gundlach C and Ortiz M E 1990 Phys. Rev D 42 2521
- [17] Guimanaes M E X 1997 Class. Quant. Grav. 14 435
- [18] G regory R and Santos C 1997 Phys. Rev. D 56 to appear
- [19] Ruban V A and Finkelstein A M 1972 Lett. Nuovo Cim. 5 289
- [20] Geroch R and Traschen J 1987 Phys. Rev. D 36 1017
- [21] Damour T and Vilenkin A 1997 Phys. Rev. Lett. 78 2288