$Re(A_0), Re(A_2)$ and RG evolution for $N_f = 3^*$

Keunsu Choi^a and Weonjong Lee^a

^aSchool of Physics, Seoul National University, Seoul, 151-747, South Korea

We present results of $Re(A_0)$ and $Re(A_2)$ calculated using HYP staggered fermions on the lattice of $16^3 \times 64$ at $\beta = 6.0$. These results are obtained using leading order chiral perturbation in quenched QCD. Buras's original RG evolution matrix develops a removable singularity for $N_f = 3$. This subtlety is resolved by finding a finite solution to RG equation and the results are presented.

In the standard model, we can express $Re(A_I)$ in terms of the matrix elements of the effective weak Hamiltonian between hadronic states as follows:

$$Re(A_{I}) = \frac{G_{F}}{\sqrt{2}} |V_{ud}V_{us}| \left[\sum_{i=1,2} z_{i}(\mu) \langle Q_{i}(\mu) \rangle_{I} + Re(\tau) \sum_{i=3}^{10} y_{i}(\mu) \langle Q_{i}(\mu) \rangle_{I} \right]$$
(1)

where V_{ij} is an element of CKM matrix. Here, $z_i(\mu)$ and $y_i(\mu)$ are the Wilson coefficients which are obtained analytically. The hadronic matrix elements $\langle Q_i(\mu) \rangle_I$ are defined as

$$\langle Q_i(\mu) \rangle_I = \langle \pi \pi_I | Q_i(\mu) | K \rangle \tag{2}$$

Since these matrix elements are in a highly nonperturbative regime of QCD, it is necessary to calculate them using non-perturbative tools such as a numerical method based on lattice gauge theory. In fact, we calculate $\langle \pi | Q_i | K \rangle$ and $\langle 0 | Q_i | K \rangle$ on the lattice using HYP staggered fermions and we construct $\langle \pi \pi_I | Q_i | K \rangle$ out of them using chiral perturbation in its leading order [1]. The $Re(\tau)$ in Eq. (1) is defined as

$$Re(\tau) = -Re(\lambda_t/\lambda_u) = 0.002$$
$$\lambda_f = V_{fs}^* V_{fd}$$

In Eq. (1), Q_i $(i = 3, 4, \dots, 10)$ represents the QCD and electroweak penguin operators. Since

 $z_i, y_i \approx 1$, the contribution from the penguin operators to $Re(A_I)$ is suppressed by $Re(\tau)$. Therefore, in practice, only the current-current operators $\langle Q_i \rangle$ (i = 1, 2) can contribute dominantly to $Re(A_I)$ whereas the contribution from penguin operators is negligible. In the case of the direct CP violation, only penguin operators can contribute to ϵ'/ϵ [2] and current-current operators do not play any role in it. In this regards, we can say that ϵ'/ϵ and $Re(A_I)$ are completely different physics.

In Fig. 1, we show individual channels contributing to $Re(A_0)$ with the total sum in the last column. The dominant contribution comes from $\langle Q_i \rangle$ (i = 1, 2) as expected. Each channel is obtained by fitting the data to a function: $f_1(m_K) = c_0 + c_1 m_K^2$ and extrapolating to the chiral limit. We use the matching formula given in [3,4] to convert lattice results into continuum values in the NDR scheme at the scale of $\mu = 1/a$. By assuming that these values are close enough to those for $N_f = 3$, we run them from $\mu = 1/a$ down to $\mu = m_c$, using the RG evolution equation for $N_f = 3$.

There are two alternative methods to transcribe the Q_6 and Q_5 operators to the lattice: the standard (STD) method and the Golterman & Pallante (GP) method [5]. In the case of ϵ'/ϵ , the dominant contribution to the $\Delta I = 1/2$ amplitude comes from $\langle Q_6 \rangle$ and so there is a big difference in ϵ'/ϵ between the STD and GP methods [2]. In the case of $Re(A_0)$, however, the contribution from $\langle Q_6 \rangle$ is extremely suppressed by $Re(\tau)$ and so both methods do not make much difference to $Re(A_0)$.

^{*}Presented by W. Lee. Research supported in part by BK21, by Interdisciplinary Research Grant of Seoul National University, and by KOSEF contract R01-2003-000-10229-0.




Figure 1. $Re(A_0)$, linear fitting for chiral extrapolation, combined at $\mu = m_c$.

In Fig. 2 we show the same kind of plot as in Fig. 1 except for the scale at which the Wilson coefficient and the matrix elements are combined. We set the scale to $\mu = 1/a$ in Fig. 2 and to $\mu = m_c$ (charm quark mass) in Fig. 1. The contribution from each individual channel can fluctuate depending on this matching scale but the total sum should be invariant.

In Fig. 3, we show the same kind of plot as in Fig. 1 except for the fitting function. Each channel in Fig. 3 is obtained by fitting the data to a function: $f_2(m_K) = c_0 + c_1 m_K^2 + c_2 m_K^4$ and extrapolating it to the chiral limit. Comparing Fig. 1 and 3, we observe that the dependence of the chiral extrapolation on the fitting function is large. These results are consistent with the experimental value of $Re(A_0) = 33.3 \times 10^{-8}$ GeV within the systematic and statistical uncertainty.

In Fig. 4, we present individual channels contributing to $Re(A_2)$ and the total sum in the last column. Each channel is obtained by fitting the data to a linear function $f_1(m_K)$ and extrapolating it to the chiral limit. The result of Fig. 4 is slightly lower than the experimental value of $Re(A_2) = 1.5 \times 10^{-8}$ GeV.

Now, let us address a subtle issue on the RG evolution for $N_f = 3$. The $\Delta S = 1$ effective Hamiltonian can be written as follows:

$$\mathcal{H} = \frac{G_F}{\sqrt{2}} \sum_{i=1}^{10} C_i(\mu) Q_i(\mu)$$
(3)




Figure 2. $Re(A_0)$, linear fitting for chiral extrapolation, combined at $\mu = 1/a$.

The RG (renormalization group) equation for the Wilson coefficient C_i is

$$\left[\mu \frac{\partial}{\partial \mu} + \beta(g) \frac{\partial}{\partial g}\right] \vec{C} = \gamma^T(g, \alpha) \vec{C}$$
(4)

where $\beta(g)$ is the QCD beta function:

$$\beta(g) = -\beta_0 \frac{g^3}{16\pi^2} - \beta_1 \frac{g^5}{(16\pi^2)^2} - \beta_{1e} \frac{e^2 g^3}{(16\pi^2)^2}$$
(5)

and $\gamma(g, \alpha)$ is the anomalous dimension matrix:

$$\gamma(g,\alpha) = \gamma_s(g^2) + \frac{\alpha}{4\pi}\Gamma(g^2) + \cdots$$
 (6)

$$\gamma_s(g^2) = \gamma_s^{(0)} \frac{\alpha_s}{4\pi} + \gamma_s^{(1)} \frac{\alpha_s^2}{(4\pi)^2} + \cdots$$
 (7)

A solution to the RG equation can be expressed in terms of the evolution matrix.

$$\vec{C}(\mu) = U(\mu, \mu', \alpha)\vec{C}(\mu') \tag{8}$$

Here the RG evolution matrix is, in general,

$$U(m_1, m_2, \alpha) = T_g \exp\left(\int_{g(m_2)}^{g(m_1)} dg' \frac{\gamma^T(g', \alpha)}{\beta(g')}\right) (9)$$

In the perturbative expansion, we can express the RG evolution matrix as follows:

$$U(m_1, m_2, \alpha) = U(m_1, m_2) + \frac{\alpha}{4\pi} R(m_1, m_2) \quad (10)$$

where $U(m_1, m_2)$ represents the pure QCD evolution and $R(m_1, m_2)$ describes the additional evolution in the presence of the electromagnetic interaction. The pure QCD evolution matrix which




Figure 3. $Re(A_0)$, quadratic fitting for chiral extrapolation, combined at $\mu = m_c$.

Buras, et al. provided originally [6,7], is

$$U(m_1, m_2) = \left(1 + \frac{\alpha_s(m_1)}{4\pi}J\right)U^{(0)}(m_1, m_2) \\ \cdot \left(1 - \frac{\alpha_s(m_2)}{4\pi}J\right)$$
(11)

where $U^{(0)}(m_1, m_2)$ denotes the evolution in the leading logarithmic approximation and J corresponds to the next-to-leading correction.

The J matrix can be expressed as follows:

$$J = VSV^{-1}, \quad G = V^{-1}\gamma_s^{(1)T}V$$
(12)

$$S_{ij} = \delta_{ij} \gamma_i^{(0)} \frac{\beta_1}{2\beta_0^2} - \frac{G_{ij}}{2\beta_0 + \gamma_i^{(0)} - \gamma_j^{(0)}}$$
(13)

where V is a matrix which diagonalize the $\gamma_s^{(0)T}$ matrix and $\gamma_i^{(0)}$ represents corresponding diagonal elements. Here, note that the J or S matrix is singular if $2\beta_0 + \gamma_i^{(0)} - \gamma_j^{(0)} = 0$. In fact, for $N_f = 3$, $2\beta_0 + \gamma_i^{(0)} - \gamma_j^{(0)} = 0$ when i = 8, j = 7. However, since $U(m_1, m_2)$ is not singular, this singularity must be removable at the nextto-leading order. After we make a correct combination at the next leading order and remove the singularity, we can make the $U(m_1, m_2)$ finite.

$$U(m_1, m_2) = U_0(m_1, m_2) + \frac{1}{4\pi} V A(m_1, m_2) V^{-1}(14)$$

where, if $a_j \neq a_i + 1$ (note that $a_i \equiv \gamma_i^{(0)}/(2\beta_0)$),

$$A(m_1, m_2) = S_{ij} \left[\alpha_s(m_1) \left(\frac{\alpha_s(m_2)}{\alpha_s(m_1)} \right)^a \right]$$




Figure 4. $Re(A_2)$, linear fitting for chiral extrapolation, combined at $\mu = m_c$.

$$-\alpha_s(m_2) \left(\frac{\alpha_s(m_2)}{\alpha_s(m_1)}\right)^{a_i} \right] \qquad (15)$$

and if $a_j = a_i + 1$,

$$A(m_1, m_2) = \frac{G_{ij}}{2\beta_0} \alpha_s(m_2) \left(\frac{\alpha_s(m_2)}{\alpha_s(m_1)}\right)^{a_i} \\ \cdot \ln\left(\frac{\alpha_s(m_2)}{\alpha_s(m_1)}\right)$$
(16)

Here, note that the A matrix is finite while J matrix is singular.

In the case of the $R(m_1, m_2)$, the same kind of removable singularity makes the RG evolution much more complicated, which we will present in Ref. [8].

REFERENCES

- C. Bernard, et al., Phys. Rev. D32 (1985) 2343.
- 2. T. Bhattacharya, et al., hep-lat/0409046.
- W. Lee and S. Sharpe, Phys. Rev. D68 (2003) 054510.
- 4. K. Choi and W. Lee, hep-lat/0309070.
- M. Golterman and E. Pallante, Phys. Rev. D69 (2004) 074503; JHEP 0110 (2001) 037.
- A. Buras, M. Jamin and M. Lautenbacher, Nucl. Phys. **B408** (1993) 209.
- 7. A. Buras, hep-ph/9806471.
- 8. W. Lee *et al.*, in preparation.