

Exploring Lattice Methods for Cold Fermionic Atoms

Matthew Wingate^a

^aInstitute for Nuclear Theory, University of Washington, Seattle, WA 98195-1550

There has been a surge of experimental effort recently in cooling trapped fermionic atoms to quantum degeneracy. By varying an external magnetic field, interactions between atoms can be made arbitrarily strong. When the S wave scattering length becomes comparable to and larger than the interparticle spacing, standard mean field analysis breaks down. In this case the system exhibits a type of universality, and J-W. Chen and D.B. Kaplan [1] recently showed how this system can be studied from first principles using lattice field theory. This poster presents the first results of exploratory simulations. The existence of a continuum limit is checked and the pairing condensate is studied as a function of the external source strength over a range of temperatures. Preliminary results show simulations can locate the critical temperature.

1. INTRODUCTION

At $T = 0$ the physics of a dilute gas is characterized by just two length scales: the interparticle spacing $n^{-1/3} \equiv (\frac{V}{N})^{1/3}$ and the S wave scattering length a . The range of interactions is very small compared to the average interparticle spacing $R \ll n^{-1/3}$, so only 2-body S wave scattering is important. By varying an external magnetic field experimentalists can get a trapped Fermi gas with any scattering length. (*e.g.* [2]).

Depending on the scattering length, the ground state of cold Fermi gases can be very different. In two extreme cases, theoretical expansions in $|a|n^{1/3} \ll 1$ are valid. If there is a deep 2-fermion bound state (*i.e.* if $0 < an^{1/3} \ll 1$) then the fermion molecules form a Bose Einstein Condensate (BEC) at low temperatures. If there is no 2-body bound state ($-1 \ll an^{1/3} < 0$) then delocalized Cooper pairs condense in a Bardeen-Cooper-Schrieffer (BCS) superfluid. What is the ground state for fermions when $|a|$ is large? No perturbative expansion converges for $|a|n^{1/3} > 1$. Ref. [1] is the only first principles approach for theoretical calculation with $|a|n^{1/3} > 1$. The case where $|a| \rightarrow \infty$ is especially interesting since systems should then exhibit universal behavior.

2. LATTICE THEORY

Below we summarize the proposal of Chen and Kaplan [1]. Consider 2 species of nonrelativistic

fermions, $\psi \equiv \begin{pmatrix} \psi_1 \\ \psi_2 \end{pmatrix}$, with a local 4-fermion interaction with finite chemical potential μ

$$\mathcal{L} = \bar{\psi}_x (\psi_x - e^\mu \psi_{x-\hat{e}_0}) - \frac{1}{2M} \bar{\psi}_x (\nabla^2 \psi)_x + \frac{C_0}{2} (\bar{\psi}_x e^\mu \psi_{x-\hat{e}_0})^2. \quad (1)$$

The 2 species can be envisioned as spin degrees-of-freedom, although in the atomic gases they are two hyperfine states.

In order to perform simulations we must make the Lagrangian quadratic in the fermion field by introducing an auxiliary real scalar field ϕ so that

$$\frac{C_0}{2} (\bar{\psi}_x e^\mu \psi_{x-\hat{e}_0})^2 \rightarrow \frac{1}{2} m^2 \phi_x^2 - \phi_x \bar{\psi}_x e^\mu \psi_{x-\hat{e}_0} \quad (2)$$

where we identify m^2 with $-C_0^{-1}$.

Now one introduces a source for the fermion pairing, a necessary step to study spontaneous symmetry breaking in a finite volume

$$\mathcal{L}[J] = \mathcal{L} + \frac{1}{2} \left(J \psi^T \sigma_2 \psi + J^* \bar{\psi} \sigma_2 \bar{\psi}^T \right). \quad (3)$$

Combining ψ and $\bar{\psi}$ into a 4-component Majorana fermion, $\Psi^T \equiv (\psi^T, \bar{\psi}(i\sigma_2)^T)$, the action can be written in terms of $4VN_t \times 4VN_t$ matrices as $\Psi^T \mathcal{A} \Psi$, where

$$\mathcal{A} \equiv \begin{pmatrix} -iJ & K^\dagger \\ K & -iJ^* \end{pmatrix} \begin{pmatrix} i\sigma_2 & 0 \\ 0 & i\sigma_2 \end{pmatrix}. \quad (4)$$

$K = K[\phi]$ represents the fermion kinetic energy and interaction terms, and it is the only

$2VN_t \times 2VN_t$ block to contain nontrivial space-time terms. The only coupling between species is through the $i\sigma_2$ blocks. The partition function can be seen to have a real, nonnegative integrand:

$$\begin{aligned} \mathcal{Z}[J] &= \int [d\phi][d\Psi] \exp\left(-\frac{1}{2}m^2\phi^2 - \frac{1}{2}\Psi^T \mathcal{A}\Psi\right) \\ &= \int [d\phi] \det\left[|J|^2 + \tilde{K}^\dagger \tilde{K}\right] e^{-m^2\phi^2/2} \quad (5) \end{aligned}$$

where $K \equiv \tilde{K} \otimes I_2$ defines the $VN_t \times VN_t$ matrix \tilde{K} . We can perform Monte Carlo simulations of (5) using Hybrid Monte Carlo.

3. PARAMETER SPACE

We have 3 lattice parameters, m^2 , μ , and M which control the physical quantities a , n , and T (ignoring finite V and setting N_t fixed). $1/a$ is matched to the ratio m^2/M at $T, \mu = 0$ by requiring 2-body scattering amplitude to match the leading term in the effective range expansion as in pionless nuclear effective field theory [3]. n is mainly controlled by tuning μ . In a nonrelativistic theory, the fermion mass can be scaled away by rescaling the time coordinate, so it can be interpreted as the anisotropy. For fixed N_t , increasing M shortens the temporal length, increasing T .

4. CONTINUUM LIMIT

Some parts of parameter space do not describe the physics of a dilute Fermi gas. In a continuum theory with a local attractive interaction, the vacuum is unstable. Since we are not interested in physics at short distances, we simply have to regulate the theory. In our case, the lattice automatically provides a cutoff (see Fig. 1).

The continuum limit is defined as the limit where physical correlation lengths $a, n^{-1/3}$, become infinitely large compared to the lattice spacing b . This limit can only be taken using simulations where one finds $nb^3 \ll 1$ at the same time that $1/a$ is tuned to 0.

Figure 2 shows $n \ll 1$ (in lattice units) at $1/a = 0$ (and $\mu = 0$) which is necessary and sufficient for a continuum limit to exist. n is nonzero there due to finite V, N_t and J effects. As $1/a \rightarrow 1$, a rise in n is observed. It would




Figure 1. Mean field illustration of the instability at strong coupling. The solid line corresponds to a coupling which is small enough in lattice units for the ground state to be the physical one, the dashed line corresponds to a coupling so strong that the ground state favors a fermion at every site even at $\mu = 0$.

be satisfying to be able to increase $1/a$ enough so that we observe the artificial first-order transition to the filled-lattice state, but the HMC algorithm is becoming less efficient in this direction. For larger values of $1/a$ than plotted, 6000 HMC trajectories were not enough to equilibrate the lattice. The breakdown of the HMC algorithm is known from simulations of the Hubbard model. Indeed the physics in the uninteresting (from our perspective) region of parameter space is exactly that of the 3D attractive Hubbard model. Having established the existence of a continuum limit in the interesting part of parameter space, searching for the uninteresting region is not a priority.

5. CRITICAL TEMPERATURE

At low temperatures we expect condensation of fermion pairs, breaking the global $U(1)$ symmetry of fermion number conservation. The order




Figure 2. Number density vs. inverse scattering length (both in lattice units) computed on $6^3 \times 12$ lattices with $J = 0.05$ and $\mu = 0$. Since $n \ll 1$ in lattice units, we can conclude the ground state is the physical one and the continuum limit exists.

parameter is

$$\Sigma \equiv -\frac{1}{2} \langle \psi^T \sigma_2 \psi + \bar{\psi} \sigma_2 \bar{\psi}^T \rangle. \quad (6)$$

Figure 3 shows Σ as a function of J for several values of M . Extrapolating to $J = 0$ linearly indicates the vanishing of the order parameter as M increases. The validity of the linear extrapolations is under study, since we expect finite volume errors to increase as J decreases [4].

6. CONCLUSIONS

The results of this exploratory study show this lattice theory can describe the physics of dilute Fermi gases. These simulations show a continuum limit exists: it is defined by $b n^{1/3}, b/a \rightarrow 0$ with fixed $an^{1/3}$. Study of the critical temperature has begun. The first point on the critical line has been located in the space of lattice parameters (m^2, M) . The critical line can then be mapped to the physical space $(1/a, T)$. For precision results, one will have to repeat these steps for increasing




Figure 3. Order parameter vs. pairing source J (both in lattice units) computed on $8^3 \times 16$ lattices with $\mu = 0.4$ and $m^2 = 0.1456$. Increasing M corresponds to increasing T ; linearly extrapolating the order parameter to $J = 0$ indicates U(1) restoration close to $M = 1.4$.

volumes and decreasing lattice spacings. However, even with finite size and spacing effects, understanding the behavior of T_c as a function of $an^{1/3}$ will be an important accomplishment.

ACKNOWLEDGMENTS

I thank J.-W. Chen, Ph. de Forcrand, D.B. Kaplan, and S. Sharpe for helpful conversations. Work was supported by DOE grant DE-FG02-00ER41132.

REFERENCES

1. J.-W. Chen and D.B. Kaplan, Phys. Rev. Lett. **92**, 257002 (2004).
2. K.M. O'Hara *et al.*, Phys. Rev. A **66**, 041401(R) (2002).
3. D.B. Kaplan, M. Savage, and M. Wise, Nucl. Phys. **B534**, 329 (1998).
4. J.-W. Chen and M. Wingate, in preparation.