

Doubly Heavy Baryons and Quark-Diquark Symmetry in Quenched and Partially Quenched Chiral Perturbation Theory

Thomas Mehen^{1,2}, and Brian C. Tiburzi^{1,y}

¹Department of Physics, Duke University, Durham, NC 27708, USA

²Jefferson Laboratory, 12000 Jefferson Ave., Newport News, VA 23606, USA

(Dated: November 30, 2021)

Abstract

We extend the chiral Lagrangian with heavy quark-diquark symmetry to quenched and partially quenched theories. These theories are used to derive formulae for the chiral extrapolation of masses and hyperfine splittings of double heavy baryons in lattice QCD simulations. A quark-diquark symmetry prediction for the hyperfine splittings of heavy mesons and doubly heavy baryons is rather insensitive to chiral corrections in both quenched and partially quenched QCD. Extrapolation formulae for the doubly heavy baryon electromagnetic transition moments are also determined for the partially quenched theory.

PACS numbers: 12.38.Gc, 12.39.Fe, 14.20.Lq, 12.39.Hg

arXiv:hep-lat/0607023v2 21 Jul 2006

mehen@phy.duke.edu

^ybctiburzi@phy.duke.edu

I. INTRODUCTION

Two heavy antiquarks in a doubly heavy antibaryon feel an attractive force when they are in the 3 of color $SU(3)$ and therefore are expected to form a compact diquark whose size, $r_{QQ} \sim (m_Q v)^{-1}$, is much smaller than λ_{QCD}^{-1} . Here m_Q is the heavy quark mass and v is the typical velocity of the antiquarks in the diquark. To the light degrees of freedom in the doubly heavy antibaryon, this diquark is indistinguishable from a heavy quark in the $m_Q \rightarrow \infty$ limit. Therefore, the properties of ground state doubly heavy baryons should be related to properties of singly heavy mesons. Savage and Wise [1] first observed that quark-diquark symmetry is a consequence of the heavy quark limit and used Heavy Quark Effective Theory (HQET) [2] to derive a relation between hyperfine splittings of doubly heavy baryons and singly heavy mesons.¹ The correct framework for analyzing the properties of hadrons with two or more heavy quarks is Non-Relativistic QCD (NRQCD) [5, 6, 7] and recently the heavy quark-diquark symmetry prediction was rederived in both the potential NRQCD [8] and velocity NRQCD [9] formalisms. An extension of heavy hadron chiral perturbation theory (HHPT) [10, 11, 12] that incorporates heavy quark-diquark symmetry was developed in Ref. [13] and used to derive a novel quark-diquark symmetry prediction for the electromagnetic decays of the spin-3/2 members of the ground state doubly heavy baryon doublets, as well as study $O(1/m_Q)$ and chiral corrections to the electromagnetic decays and doubly heavy baryon masses. Some quark-diquark symmetry predictions for the exotic heavy tetraquarks as well as a critical discussion of the applicability of heavy quark-diquark symmetry to charm hadrons appeared in Ref. [14].

Much of the recent theoretical work has been stimulated by the SELEX collaboration reports of candidates for doubly charm baryons [15, 16, 17]. While the masses and hyperfine splittings of the observed states are consistent with theoretical calculations in the quark model [4] as well as quenched lattice QCD, other aspects of the data are poorly understood at the present time. For example, isospin splittings, weak decay rates, and production yields are in disagreement with theoretical expectations. Some excited states which would be expected to decay electromagnetically are instead observed via their weak decay. For a review of the experimental situation and discussion of theoretical interpretations of the SELEX results, see Refs. [13, 14, 18]. Recently, the BABAR [19] and BELLE [20] experiments searched for doubly charmed baryons in e^+e^- collisions but failed to observe the SELEX states. More experimental data and improved theoretical understanding of doubly charm baryons are necessary to clarify the situation.

Lattice gauge theory studies of the properties of doubly heavy baryons with dynamical quarks would be quite useful. First principles calculation of the spectroscopy of doubly heavy baryons could be directly compared with experiment to test our understanding of QCD and aid in the interpretation of candidate states. It would also be of interest to have first principle calculations of electromagnetic and weak decay matrix elements. Finally, the lattice could help determine the suitability of quark-diquark symmetry for charm and bottom hadrons. In these cases, it is possible that symmetry breaking corrections could be large enough to spoil symmetry predictions, but no systematic study of symmetry breaking corrections in NRQCD is yet available. A theoretical testing ground in which the heavy quark mass can be dialed at will could determine whether this symmetry will turn out to be useful for charm and bottom. Spectroscopic studies of doubly charm and bottom baryons

¹ The quark model has also been used to derive the relation [3, 4].

in quenched lattice QCD appeared in Refs. [3, 21, 22]. The heavy quark-diquark symmetry prediction for the hyperfine splitting holds at the 20% level, which is reasonable given the sources of error. In QCD, corrections are expected to be $O(v^2)$ and/or $O(m_{\text{QCD}}=m_Q)$. In existing lattice calculations there are also uncontrolled errors due to quenching, which, as we will see below, could be significant. Hopefully, the near future will bring studies of doubly heavy baryons in simulations with dynamical quarks.

As a step towards such studies, it is clearly useful to develop extensions of the heavy quark-diquark symmetric version of HQPT [13] that include lattice artifacts such as quenching and partial quenching. These theories provide the formulae needed for chiral extrapolations in lattice calculations. We will focus on chiral corrections to the doubly heavy baryon masses, especially hyperfine splittings and the quark-diquark symmetry prediction. We also apply the formalism to the electromagnetic decays of the spin-3/2 members of the ground state doubly heavy baryon doublets.

The organization of our paper is as follows. In section II, we review the basics of quenched chiral perturbation theory (QPT) and partially quenched chiral perturbation theory (PQPT) in the pseudoscalar meson sector. We then write down versions of PQPT and QPT that incorporate heavy quark-diquark symmetry. In section III, we calculate the one-loop chiral corrections to the masses of doubly heavy baryons in PQPT and QPT. These expressions can be used to perform the chiral extrapolation of lattice data on the spectrum of doubly heavy baryons provided one is in the range of applicability of the chiral effective theory. In section IV, we calculate to one loop the dipole and quadrupole transition moments in PQPT. For these transitions, the lattice can be used to test the predictions for the electromagnetic decay widths of the spin-3/2 members of the ground state doubly heavy baryon doublets [13]. Finally, we conclude with a brief summary in section V. Some useful formulae are collected in the Appendix.

II. PQPT AND QPT FOR DOUBLY HEAVY BARYONS

The formalism for constructing effective theories for quenched and partially quenched QCD is now well established. We use the technique of graded Lie algebras that was formulated in Refs. [23, 24, 25, 26, 27, 28, 29, 30]. Equivalently one could use the replica method [31]. We begin by briefly reviewing the pseudoscalar meson sector of QPT and PQPT, and then write down the quenched and partially quenched chiral Lagrangians incorporating heavy quark-diquark symmetry. These theories are an extension of QPT and PQPT for singly heavy mesons [32].

A. Pseudoscalar Mesons

In the pseudoscalar meson sector, we can treat PQPT and QPT as different limits of the same theory. This theory is described by the Lagrangian

$$L = \frac{f^2}{8} \text{str} (\partial_\mu \chi^\dagger \partial^\mu \chi) + \frac{1}{4} \text{str} m_q \chi^\dagger \chi + m_q \chi^\dagger \chi + \text{str} \chi^\dagger \chi \chi^\dagger \chi; \quad (1)$$

where the field χ is defined by

$$\chi = \exp \frac{2i}{f} \phi = \phi; \quad (2)$$

and the meson fields appear in the $U(6\beta)$ matrix,

$$= \begin{matrix} M & Y \\ & \tilde{M} \end{matrix} : \quad (3)$$

The quantities \tilde{m}_q and m_0 are non-vanishing in the chiral limit. The M and \tilde{M} matrices contain bosonic mesons (with quantum numbers of $q\bar{q}$ pairs and $q\bar{q}$ pairs, respectively, where q is a valence or sea quark and \bar{q} is a ghost quark), while the \tilde{M} and Y matrices contain fermionic mesons (with quantum numbers of $q\bar{q}$ pairs and $q\bar{q}$ pairs, respectively). The upper 3×3 block of the matrix M contains the familiar pions, kaons, and eta, while the remaining components consist of mesons formed from one or two sea quarks. The operation $\text{str}()$ in Eq. (1) is a supertrace over flavor indices, i.e., $\text{str}(A) = \sum_a A_{aa}$, where $a = ()^{1+}$, and a are the grading factors of the Lie algebra.

The quark mass matrix appearing above is given by

$$m_q = \text{diag}(m_u; m_d; m_s; m_j; m_l; m_r; m_u; m_d; m_s) : \quad (4)$$

We will work in the isospin limit of both the valence and sea sectors, where $m_d = m_u$ and $m_l = m_j$. Expanding the Lagrangian in Eq. (1) to lowest order in the fields, one finds that mesons with quark content $q\bar{q}^0$ are canonically normalized and their masses are given by

$$m_{q\bar{q}^0}^2 = \frac{1}{f^2} (m_q + m_{q^0}) : \quad (5)$$

The propagators of the off-diagonal mesons have the usual Klein-Gordon form.

On the flavor diagonal, the situation is more complicated and differs dramatically between the quenched and partially quenched theories. The flavor singlet field that appears above is defined to be $\phi_0 = \text{str}(\phi) = \frac{1}{6}$. Using this, the leading-order a_b propagator is [29]

$$G_{ab}(q^2) = \frac{i a_{ab}}{q^2 - m_{aa}^2 + i} - \frac{i}{3} \frac{q^2 - m_{jj}^2 (q^2 - m_{rr}^2) (q^2 - m_0^2)}{(q^2 - m_{aa}^2 + i) (q^2 - m_{bb}^2 + i) D(q^2)} : \quad (6)$$

The function $D(q^2)$ is defined as

$$D(q^2) = (q^2 - m_{jj}^2)(q^2 - m_{rr}^2) + (q^2 - m_0^2)(q^2 - m_x^2);$$

and the mass m_x is given by $m_x^2 = \frac{1}{3} m_{jj}^2 + 2m_{rr}^2$.

Because the fermionic determinant in quenched QCD is set to a constant, the sea quark masses are effectively infinite. The mesons containing sea quarks decouple from the theory, leaving a $U(3\beta)$ Goldstone manifold. In this limit, the flavor neutral propagator in Eq. (6) becomes

$$G_{ab}^Q(q^2) = \frac{i a_{ab}}{q^2 - m_{aa}^2 + i} - \frac{i}{3} \frac{q^2 - m_0^2}{(q^2 - m_{aa}^2 + i) (q^2 - m_{bb}^2 + i)}; \quad (7)$$

where we have appended a superscript Q to denote that this propagator is for the quenched theory. Notice the flavor singlet meson does not decouple as the theory still depends on the parameters \tilde{m}_q and m_0 . Numerical data suggest that \tilde{m}_q is small (for example, a recent study finds $\tilde{m}_q = 0.03 \pm 0.03$ [33]), and we neglect it below; m_0 will be treated in the standard quenched power counting [34].

In the partially quenched theory, sea quark contributions are retained and partially quenched QCD (PQQCD) has a strong axial anomaly. Because of this, the mass of the singlet field ϕ_0 can be taken to be on the order of the chiral symmetry breaking scale, and the ϕ_0 integrated out of the theory. The Goldstone manifold is reduced to $SU(6\beta)$ but the resulting flavor neutral two-point function, however, still deviates from the familiar form in PT. In PQQCD this propagator is given by the $\phi_0 \rightarrow 1$ limit of Eq. (6), namely

$$G_{ab}^{PQ}(\not{q}^2) = \frac{i_{ab}}{\not{q}^2 - m_{aa}^2 + i} - \frac{i}{3} \frac{\not{q}^2 - m_{jj}^2 (\not{q}^2 - m_{rr}^2)}{(\not{q}^2 - m_{aa}^2 + i) (\not{q}^2 - m_{bb}^2 + i) (\not{q}^2 - m_X^2 + i)} : \quad (8)$$

The PQ PT flavor neutral propagator can be conveniently rewritten as

$$G_{ab}^{PQ} = P_a \delta_{ab} + P_{ab}(P_a; P_b; P_X); \quad (9)$$

where

$$P_a = \frac{i}{\not{q}^2 - m_{aa}^2 + i}; \quad P_b = \frac{i}{\not{q}^2 - m_{bb}^2 + i}; \quad P_X = \frac{i}{\not{q}^2 - m_X^2 + i};$$

$$P_{ab}(A; B; C) = \frac{1}{3} \frac{m_{aa}^2 - m_{jj}^2 (m_{aa}^2 - m_{rr}^2)}{(m_{aa}^2 - m_{bb}^2) (m_{aa}^2 - m_X^2)} A + \frac{m_{bb}^2 - m_{jj}^2 (m_{bb}^2 - m_{rr}^2)}{(m_{bb}^2 - m_{aa}^2) (m_{bb}^2 - m_X^2)} B$$

$$+ \frac{m_X^2 - m_{jj}^2 (m_X^2 - m_{rr}^2)}{(m_X^2 - m_{aa}^2) (m_X^2 - m_{bb}^2)} C : \quad (10)$$

B. Doubly heavy baryons

As discussed earlier, in the heavy quark limit the properties of singly heavy mesons and doubly heavy baryons are related by a $U(5)$ quark-diquark symmetry [1]. Recently a chiral Lagrangian incorporating this $U(5)$ symmetry has been derived [13]. It is written in terms of a superfield H given by

$$H_{a; i} = H_{a; i} + T_{a; i}; \quad (11)$$

where a is an anti-fundamental flavor index, and i, j are Pauli spinor indices, while i is a vector spin index. The super-spin index $i = 1\{5$ and is related to i and j via

$$i = (j^1 + j^2) + (i + 2)(j^3 + j^4 + j^5); \quad (12)$$

The field $H_{a; i}$ is the heavy meson field of HH PT [10, 11, 12] in the rest frame of the heavy meson:

$$H_{a; i} = P_a + P_a : \quad (13)$$

H_a contains the vector (P) and pseudoscalar (P) field components. The field $T_{a; i}$ is the doubly heavy baryon field,

$$T_{a; i} = \frac{P}{2} T_{a; i} + \frac{1}{3} P_{a; i}^i; \quad (14)$$

that contains the spin-1=2 (T) and spin-3=2 (P) field components. The spin-3=2 field satisfies a non-relativistic Rarita-Schwinger constraint, $T_{a; i}^i = 0$.

In terms of the super-eld H , the Lagrangian for the quenched (Q) theory is

$$L^Q = H^\dagger (H \mathbb{D}_0) g^Q H^\dagger H A + g^{QQ} H^\dagger H \text{str}(A) + \frac{H}{4} H^\dagger H + g^Q H^\dagger H M + g^{QQ} H^\dagger H \text{str}(M); \quad (15)$$

while that for the partially quenched theory (PQ) is

$$L^{PQ} = H^\dagger (H \mathbb{D}_0) g^{PQ} H^\dagger H A + \frac{H^{PQ}}{4} H^\dagger H + g^{PQ} H^\dagger H M + g^{QPQ} H^\dagger H \text{str}(M); \quad (16)$$

Both Lagrangians include the leading quark-diquark symmetry breaking term. The bracket $()$ notation denotes the contraction of flavor, spinor and super-spin indices of eld bilinears. For a generic super-spin matrix X , flavor matrix Y , and spin matrix Z , $()$ denotes

$$H^\dagger X H Y Z = \begin{matrix} X & X & X \\ a & b & c \end{matrix} H_a^\dagger Y_{ab} X_b H_c Z_c; \quad (17)$$

In Eqs. (15) and (16), A is the spatial part of the axial-vector pion eld, $A = \frac{i}{2} \epsilon^{ijk} \sigma^j \tau^k$, and M is the mass operator defined by $M = \frac{1}{2} m_q + y m_q Y$. The U(5) super-spin matrix is given by

$$M = \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & T_{jk} & & \\ & & & & \\ & & & & 0 \end{pmatrix}; \quad (18)$$

where $(T^i)_{jk} = i \epsilon_{ijk}$.

Notice that there are separate Lagrangians for quenched and partially quenched QCD. The low-energy constants of the two theories are different, e.g. $g^{PQ} \neq g^Q$. Notice also that in QPT the flavor singlet eld cannot be integrated out. Consequently $\text{str}(A) \neq 0$ and the axial hairpin coupling g^{QQ} remains. Additionally in QPT, $\text{str}(M) = 0$ to leading order, and so we can effectively set $g^{QPQ} = 0$ for our calculations. There is no sector of QPT that contains PT, so in general quenched observables are unrelated to QCD observables. The sea sector of PQ QCD is QCD, and hence PQ PT contains the low-energy constants of PT. Restricting the flavor indices to the sea sector in Eq. (16), we recover the chiral Lagrangian of Ref. [13], so g^{PQ} , $\frac{H^{PQ}}{4}$, g^{PQ} and g^{QPQ} are identical to the low-energy constants g , $\frac{H}{4}$, γ , and β of PT, respectively. At this order there are no additional operators in the partially quenched Lagrangian. Carrying out the super-spin sums in Eqs. (15) and (16), we produce separate terms for heavy mesons and doubly heavy baryons but with the same low-energy constants. The heavy meson sector of the PQ PT and Q PT Lagrangians above is that of Ref. [32].

III. MASSES

In this section, we use the quenched and partially quenched chiral Lagrangians that incorporate quark-diquark symmetry to calculate chiral corrections to the masses of doubly heavy baryons. The results can be used for the chiral extrapolation of lattice data. PQ PT

is considered in subsection IIIA and Q PT is discussed in subsection IIIB. It is important to keep in mind that while PQ PT actually contains PT, there is no such connection between Q PT and PT.

Near the chiral limit, the masses of doubly heavy baryons generically have the form

$$\begin{aligned} M &= M_0 + \frac{1}{2} \Delta_H + M^{(1)} + M^{(3=2)} + \dots \\ M &= M_0 + \frac{1}{4} \Delta_H + M^{(1)} + M^{(3=2)} + \dots; \end{aligned}$$

where M_0 is the mass of the doubly heavy baryons in the chiral and heavy quark limit and Δ_H is the hyperfine mass splitting of the heavy mesons in the chiral limit.² The hyperfine splitting of the Σ and Λ baryons is $\frac{3}{4} \Delta_H$ in the chiral limit due to quark-diquark symmetry. The term $M^{(1)}$ is proportional to m_q and comes from tree-level counterterms. The term $M^{(3=2)}$ comes from the one-loop diagrams and depends on m , the Goldstone boson mass, and Δ , the mass splitting between the external and virtual doubly heavy baryons. Goldstone boson masses are $\propto \sqrt{m_q}$, while Δ receives contributions from SU(3) breaking counterterms, which are $\propto m_q$, and/or hyperfine splittings, which scale as $1/m_Q$. In our power counting we take $m \ll Q$ in which case $M^{(3=2)} \propto Q^3$. In the chiral and heavy quark limits, Δ can be neglected relative to m and then $M^{(3=2)} \propto m_q^{3=2}$. The ellipsis denotes higher-order terms in the Q expansion.

A. PQ PT

At linear order in the quark mass, the contributions to the masses of doubly heavy baryons at tree level come from the mass operators in Eq. (16):

$$M_a^{(1)} = M_a^{(1)} = (m_q)_a \text{str}(m_q); \quad (19)$$

where $\text{str}(m_q) = 2m_j + m_r$ in the isospin limit, and the subscript a denotes the light-quark anti-fundamental index of the doubly heavy baryon. Combining these results with the mass splitting in the chiral limit, we then define the baryon mass splittings

$$\Delta_{ba} = M_b - M_a = M_b - M_a = [(m_q)_a - (m_q)_b]; \quad (20)$$

$$\Delta_{ba} = M_b - M_a = \frac{3}{4} \Delta_H + [(m_q)_a - (m_q)_b]; \quad (21)$$

At one-loop order, the doubly heavy baryons receive mass contributions in PQ PT from the sunset diagrams shown in Fig. 1. The vertices in these diagrams are generated from

² Strictly speaking Δ_H is not the same parameter appearing in the Lagrangian above, Eq. (16), which in this footnote we denote by Δ_H^0 . This parameter, which transforms as a chiral singlet, is corrected by a string of operators that differ only by scaling with powers of $\Delta_H^0 = \Delta_H$. Thus we subsume this dependence as $\Delta_H = \Delta_H(\Delta_H^0)$ including also contributions to Δ_H arising from loop graphs. In this way we obtain Δ_H as the chiral limit value. The same is true of all other low-energy constants in Eq. (16). We similarly drop their polynomial dependence on $\Delta_H^0 = \Delta_H$, and absorb contributions from loop graphs to work with the values of the low-energy constants in the chiral limit. For more details see, e.g. Ref [35].




FIG. 1: One-loop graphs which give contributions to the masses of the doubly heavy baryons in PQPT. The single and double lines correspond to baryons or baryons, respectively, while the dashed lines correspond to mesons. The filled squares denote the non-singlet axial coupling given in Eq. (16), while crosses denote the hairpin interaction.

the pion-baryon interaction terms in Eq. (16), while the hairpin interaction corresponds to the disconnected part of the flavor neutral propagator in Eq. (8). Calculating these PQPT loop diagrams, we find

$$M_a^{(3=2)} = \frac{g^2}{(4f)^2} \sum_{b=i,j,l,r} X \left[\frac{5}{9} K(\bar{b}a; m_{ba};) + \frac{4}{9} K(\bar{a}b; m_{ba};) \right] + \frac{g^2}{(4f)^2} \left[\frac{5}{9} K(0; m_{aa}; m_{aa};) + \frac{4}{9} K(\bar{a}a; m_{aa}; m_{aa};) \right]; \quad (22)$$

for the spin-3/2 doubly heavy baryons, \bar{a} , and

$$M_a^{(3=2)} = \frac{g^2}{(4f)^2} \sum_{b=i,j,l,r} X \left[\frac{1}{9} K(\bar{b}a; m_{ba};) + \frac{8}{9} K(\bar{b}a; m_{ba};) \right] + \frac{g^2}{(4f)^2} \left[\frac{1}{9} K(0; m_{aa}; m_{aa};) + \frac{8}{9} K(\bar{a}a; m_{aa}; m_{aa};) \right]; \quad (23)$$

for the spin-1/2 doubly heavy baryons, \bar{a} . The non-analytic function $K(\bar{; m};)$ is

$$K(\bar{; m};) = 2(\bar{^2} - m^2) m R \frac{1}{m} \left[\log \frac{m^2}{2} + m^2 \log \frac{m^2}{2} \right]; \quad (24)$$

where $R(x)$ is defined by

$$R(x) = \frac{1}{x^2 - 1} \log \frac{x + \sqrt{x^2 - 1 + i}}{x - \sqrt{x^2 - 1 + i}}; \quad (25)$$

while the function $K(\bar{; m_{aa}; m_{bb};)$ which arises from hairpins is given by

$$K(\bar{; m_{aa}; m_{bb};) = P_{ab} K(\bar{; m_{aa};); K(\bar{; m_{bb};); K(\bar{; m_x};); \quad (26)$$

with P_{ab} appearing in Eq. (10), and a limit is understood when $b = a$.




FIG. 2: $\Delta\chi_{HF}$ as a function of $m_{\pi\text{ sea}}$ for different values of $m_{\pi\text{ val}}$. The width of the bands is the result of varying μ between 500 MeV and 1500 MeV.

To study the chiral non-analytic behavior of doubly heavy baryon mass splittings in PQ PT, we need values for the low-energy constants. We choose $g = 0.6$ [36], $m_H = 140$ MeV, and $f = 130$ MeV. The SU(3) splitting of the ground state D mesons is ≈ 100 MeV. Assuming the same SU(3) splitting for the doubly charm baryons we obtain $\mu = (337 \text{ MeV})^3$. We shall take the strange sea quark to be degenerate with its valence counterpart, $m_r = m_s$, and fix the value at the physical strange quark mass. This leaves three parameters that can be varied, the valence pion mass, $m_{\pi\text{ val}} = m_{uu}$, the sea pion mass, $m_{\pi\text{ sea}} = m_{jj}$, and the renormalization scale, μ . While the μ dependence in the above expressions is exactly canceled by μ -dependent counterterms at this order, we have not included these counterterms, nor do we have a reliable way to estimate them. Hence we vary μ to effectively obtain a range for the unknown counterterm contributions.

In Fig. 2, we assess the non-analytic chiral corrections to the hyperfine splitting of non-strange and strange baryons in PQ PT. To see the effect of partial quenching on this observable, we plot the difference between the chiral non-analytic contribution to the hyperfine splitting in PQ PT and the corresponding non-analytic correction in PT:

$$\Delta\chi_{HF} = M^{(3=2)}_{PQ\text{ PT}} - M^{(3=2)}_{PT} ; \quad (27)$$

as a function of $m_{\pi\text{ sea}}$ for a few different values of $m_{\pi\text{ val}}$. For each value of $m_{\pi\text{ val}}$, we let $m_{\pi\text{ sea}}$ range from $m_{\pi\text{ val}}$ up to the mass of eta-strange, $m_s = m_{ss} = m_{rr} \approx 700$ MeV. Thus the smallest value of $m_{\pi\text{ sea}}$ corresponds to an unquenched theory at $m = m_{\pi\text{ val}}$, while the largest value of $m_{\pi\text{ sea}}$ corresponds to a completely degenerate sea at the strange quark mass. The bands correspond to varying μ from 500 MeV to 1500 MeV, which is chosen to be the same

for both PQ PT and PT. For the PT hyperfine splitting, we use the physical Goldstone boson masses, so by construction $M_{HF} = 0$ when $m_{sea} = m_{val} = m_\pi$. As demonstrated by Fig. 2, chiral non-analytic corrections to the hyperfine splitting are sizable, and increase with increasing m_{sea} . Additionally corrections are enhanced as m_{val} increases. While we do not trust the chiral expansion at large pion masses, we have kept these in the plot to show that chiral corrections can become large and tend to push the hyperfine splitting down by tens of MeV.

We can use our results to check the chiral corrections to the mass splitting relation predicted by heavy quark-diquark symmetry [1, 8, 9]

$$M_{DQ} - M_{PQ} = \frac{3}{4} (M_P - M_{\bar{P}}) : \quad (28)$$

In Fig. 3, we plot the difference between the nonanalytic chiral corrections to this relation in PQ PT and PT, defined by

$$\Delta_{DQ} = M_{DQ}^{(3=2)} - M_{PQ}^{(3=2)} - \frac{3}{4} (M_P^{(3=2)} - M_{\bar{P}}^{(3=2)})_{PQ\ PT} - M_{DQ}^{(3=2)} + M_{PQ}^{(3=2)} - \frac{3}{4} (M_P^{(3=2)} - M_{\bar{P}}^{(3=2)})_{PT} ; \quad (29)$$

in the non-strange sector. Here $M_P^{(3=2)}$ and $M_{\bar{P}}^{(3=2)}$ are the chiral contributions to the masses of the vector and pseudoscalar heavy mesons, respectively [32]. As in Fig. 2, we vary m_{val} from 500 MeV to 1500 MeV for three different values of m_{sea} . Fig. 3 shows that the chiral corrections to the heavy quark-diquark symmetry prediction are remarkably small compared to chiral corrections to the hyperfine splittings and much less sensitive to m_{sea} . It was demonstrated in Ref. [13] that chiral corrections to the quark-diquark symmetry prediction for the hyperfine splittings are small and insensitive to m_{sea} . Interestingly, this conclusion is not changed as the pion mass is increased, nor is it altered by partial quenching. While the hyperfine splittings of heavy mesons and doubly heavy baryons receive considerable chiral corrections for larger pion masses, the quark-diquark symmetry relation fortuitously does not. Lattice calculations of this quantity will not suffer large uncertainty due to chiral extrapolation.

B. Q PT

In the quenched theory we can similarly calculate the masses of doubly heavy baryons to one-loop order. The tree level contribution from SU(3) breaking counterterms is

$$M_a^{(1)} = M_b^{(1)} = M_H^Q (m_q)_a : \quad (30)$$

The tree-level baryon mass splittings are

$$M_{ba} = M_b - M_a = M_b - M_a = M_H^Q [(m_q)_a - (m_q)_b]; \quad (31)$$

$$M_{ba} = M_b - M_a = \frac{3}{4} M_H^Q + M_H^Q [(m_q)_a - (m_q)_b]; \quad (32)$$

The differences in low-energy constants notwithstanding, the one-loop mass contributions in Q PT are considerably altered from those in PQ PT and PT. The sunset diagrams in




FIG. 3: χ_{QDQ} as a function of m_{sea} for different values of m_{val} . The width of the bands is the result of varying a between 500 MeV and 1500 MeV.




FIG. 4: Additional bop graphs which contribute to the masses of the doubly heavy baryons in QPT. The single and double lines correspond to baryons or baryons, respectively, while the dashed lines correspond to mesons. The filled squares denote the non-singlet axial coupling, and the cross denotes the axial hairpin interaction given in Eq. (15).

Fig. 1 again contribute, however, it is clear that only hairpin contributions survive quenching. There are additional one-bop diagrams present in the quenched theory that have no counterparts in PQPT or PT. These diagrams are depicted in Fig. 4 and involve the axial hairpin interaction in Eq. (15). Combining the hairpin diagrams and axial hairpin diagrams, we arrive at the one-loop quenched mass contributions

$$\begin{aligned}
M_a^{(3=2)} = & \frac{2g^0 g^{00}}{(4f)^2} \frac{5}{9} K(0; m_{aa};) + \frac{4}{9} K(\quad_{aa}; m_{aa};) \\
& + \frac{(g^0)^2}{3(4f)^2} \frac{2}{\partial m_{aa}^2} \frac{5}{9} K(0; m_{aa};) + \frac{4}{9} K(\quad_{aa}; m_{aa};) ; \quad (33)
\end{aligned}$$

for the spin-3=2 doubly heavy baryons, and

$$\begin{aligned}
M_a^{(3=2)} = & \frac{2g^0 g^{00}}{(4f)^2} \frac{1}{9} K(0; m_{aa};) + \frac{8}{9} K(\quad_{aa}; m_{aa};) \\
& + \frac{(g^0)^2}{3(4f)^2} \frac{2}{\partial m_{aa}^2} \frac{1}{9} K(0; m_{aa};) + \frac{8}{9} K(\quad_{aa}; m_{aa};) ; \quad (34)
\end{aligned}$$

for the spin-1=2 doubly heavy baryons.




FIG. 5: The quenched chiral correction to the hyper-ne splitting, $M_{\Xi}^{(3/2)} - M_{\Xi}^{(3=2)}$, as a function of the quenched pion mass. The χ PT band shows the corresponding range of chiral corrections to the hyper-ne splitting in QCD.

With the quenched chiral non-analytic corrections in hand, we can investigate their impact on the spectrum of doubly heavy baryons as we did for the partially quenched theory. To determine these corrections, we must estimate the parameters in the quenched chiral Lagrangian. Little is known about these parameters. We adopt a central value of $\Lambda_0 = 700$ MeV from the analysis of Ref. [33]. The remaining parameters are taken to be the same values as in χ PT, though there is no justification for this. Finally we use two values, $g^Q = 0.6$, for the parameter that has no analogue in the unquenched theory. In Fig. 5, we plot the quenched chiral correction to the hyper-ne splitting, $M_{\Xi}^{(3/2)} - M_{\Xi}^{(3=2)}$, as a function of the pion mass. The band plotted corresponds to a variation of Λ_0 from 500 MeV to 1500 MeV. As in the partially quenched theory, we see that the mass splitting receives sizable chiral corrections that increase with the pion mass. The trend is the same as in the partially quenched theory provided g^Q and $g^{\bar{Q}}$ have the same sign: quenched lattice calculations at larger pion masses than that in nature will underestimate the hyper-ne splitting. When g^Q and $g^{\bar{Q}}$ have opposite signs, it is possible for the splitting to be overestimated. Lastly we can test how well heavy quark-diquark symmetry holds up in the presence of quenched chiral corrections. In Fig. 6, we plot the quenched chiral correction to the quark-diquark symmetry relation for the hyper-ne splittings,

$$\Delta_{\text{QDQ}}^Q = M_{\Xi}^{(3=2)} - M_{\Xi}^{(3/2)} - \frac{3}{4} (M_{\text{P}}^{(3=2)} - M_{\text{P}}^{(3/2)}); \quad (35)$$

as a function of the pion mass, again varying Λ_0 from 500 to 1500 MeV. Surprisingly, the quenched chiral corrections do not sizably alter the quark-diquark symmetry prediction. The corrections are only a few MeV. The dependence of the result is minimal and in fact one can show analytically that the dependence vanishes for this observable if $g^{\bar{Q}} = 0$.

IV. ELECTROMAGNETIC TRANSITIONS IN PQPT

As a final application we determine the electromagnetic transition moments of the doubly heavy baryons in PQPT and χ PT. To include electromagnetism into the theory, we merely gauge a background U(1) field. In doing so, recall that the light-quark electric charge matrix




FIG . 6: The quenched chiral correction to the heavy quark-diquark symmetry breaking parameter, Δ_{QDQ}^Q , as a function of the quenched pion mass. The χ PT band shows the corresponding range of chiral corrections to quark-diquark symmetry in QCD.



FIG . 7: One-loop graphs contributing to the doubly heavy baryon electromagnetic transitions in PQPT. The single lines correspond to baryons, double lines to baryons. The dashed lines correspond to mesons, while the wiggly lines correspond to photons. The coupling of Goldstone mesons to baryons is the non-singlet axial coupling given in Eq. (16).

Q is not uniquely defined in PQCD [37]. The only constraint one must impose is that the charge matrix Q has vanishing supertrace. Following Refs. [38, 39], we use

$$Q = \text{diag}(q_1; q_1; q_2; q_2; q_3; q_3; q_4; q_4; q_5); \quad (36)$$

along with the condition $q_2 + q_3 + q_4 = 0$. QCD is recovered in the limit of degenerate valence and sea quarks only for the particular choice: $q_1 = q_2 = \frac{2}{3}$, and $q_3 = q_4 = q_5 = \frac{1}{3}$. Letting the charges be arbitrary, however, enables us to track the flow of charge in loop diagrams.

The relevant electromagnetic couplings in the chiral Lagrangian with quark-diquark symmetry are contained in the terms [13]

$$L = \frac{e}{2} H^Y H^Q B + \frac{Q^0 e}{2m_Q} H^Y \sim B H + \frac{E^2 e}{2m_Q} H^Y \sim^i H^Q \sim^j r_{fi} E_{jg}; \quad (37)$$

where Q^0 is the charge of the heavy quark and the 5×5 matrices \sim are

$$\sim = \begin{pmatrix} 0 & & & & \\ & 0 & & & \\ & & 0 & & \\ & & & 0 & \\ & & & & 0 \end{pmatrix}; \quad (38)$$

The first term in Eq. (37) gives the leading contribution to the decay rate in the $1/m_Q$ expansion. The contribution of this term at tree level is similar to the light quark magnetic moment contribution to the transition in the non-relativistic quark model, i.e. the quark model reproduces the tree-level result of HH PT upon the identification $1/m_q \rightarrow 1/m_Q$. The second term in Eq. (37) is the contribution from the magnetic moment of the heavy quark which is $O(1/m_Q)$. The third term in Eq. (37) is also $O(1/m_Q)$ suppressed. The notation f_{ijg} denotes symmetrization and tracelessness in the indices ij . In the heavy meson sector, this term contributes to the electric quadrupole moment of the P but not to the $P \rightarrow P'$ transition. It does give a contribution to the decay $\Gamma(P \rightarrow P' \gamma)$, which however is suppressed by $1/m_Q^2$ rather than $1/m_Q$ because it does not interfere with the leading-order contribution. Explicitly the formula for the decay width is given by

$$\Gamma(P \rightarrow P' \gamma) = \frac{4}{9} \left(\sum_{a,b} Q_a Q_b \left[\frac{Q_a^2}{2m_Q} + \frac{3}{4} E_2^2 \frac{E_{2Q_a}}{m_Q} + E_{2a} \right] \frac{M}{M} E^3 \right); \quad (39)$$

where $\sum_{a,b} Q_a Q_b$ and E_{2a} represent chiral corrections to the tree-level magnetic dipole and electric quadrupole transition moments, respectively. The loop diagrams shown in Fig. 7 give the leading non-analytic chiral corrections.

The Lagrangian in Eq. (37) is easily extended to PQ PT. Using the definition of the superfield H in Eq. (11), we can expand the Lagrangian in Eq. (37) to find the doubly heavy baryon part and thereby determine the contribution to the transition moments at tree level. As this contribution only involves valence quarks, the result is the same as in PT [13]. The coefficients $\sum_{a,b} Q_a Q_b$ and E_2 have the same value as in PT which can be demonstrated by matching to PQ PT in the sea sector. In the heavy quark limit, the operator proportional to E_2 can be neglected. When $O(1/m_Q)$ hyperfine splittings are included in the loop diagrams, there are non-analytic corrections to both $\sum_{a,b} Q_a Q_b$ and E_2 . At one-loop order in PQ PT, we must calculate the diagrams shown in Fig. 7. Explicitly we find

$$\sum_{a,b} Q_a Q_b = \frac{g^2}{24 f^2} \sum_{b=j,l,r} \left(Q_b Q_a \int_0^1 dx \sum_{i=1}^Z \left(\gamma_1(x) \log \frac{m_{ab}^2}{2} \right) \frac{1(x)}{m_{ab}} \right) + 5 \sum_{i=2}^Z \left(\gamma_2(x) \log \frac{m_{ab}^2}{2} \right) \frac{2(x)}{m_{ab}}; \quad (40)$$

$$E_{2a} = \frac{g^2}{24 f^2} \sum_{b=j,l,r} \left(Q_b Q_a \int_0^1 dx x (1-x) \left(\frac{1(x) m_{ab}}{[1(x)]^2 m_{ab}^2} R \frac{1(x)}{m_{ab}} \right) \right) + \sum_{i=2}^Z \left(\frac{2(x) m_{ab}}{[2(x)]^2 m_{ab}^2} R \frac{2(x)}{m_{ab}} \right); \quad (41)$$

In expressing the transition moments, we have made the following definitions

$$\gamma_1(x) = \sum_{ab} + x \sum_{aa}; \quad (42)$$

$$\gamma_2(x) = \sum_{ab} + x \sum_{aa}; \quad (43)$$

and used the function $R(x)$ defined in Eq. (25). The PT result can be deduced by replacing the sea charges and sea masses with the values of their valence counterparts.




FIG. 8: Additional loop graphs for the doubly heavy baryon electromagnetic transitions in QCD. The single lines correspond to Λ baryons, double lines to Σ baryons. The dashed lines correspond to mesons, while the wiggly lines correspond to photons. The filled circles denote the non-singlet axial coupling given in Eq. (15), the photon coupling is proportional to the magnetic couplings in Eq. (37), and crosses denote the quenched hairpin interaction.

Ref. [13] considered the loop corrections in PT in the approximation where $H_p = 0$ and $\mu = 0$. This corresponds to taking the heavy quark limit and keeping only $O(p^2/m_q)$ nonanalytic chiral corrections. In this limit the result can be inferred from the calculation of Ref. [40, 41] for heavy mesons:

$$a = \frac{g^2}{4f^2} \sum_b (Q_b - Q_a) m_{ab} \quad E_{2a} = 0; \quad (44)$$

For PT with physical parameters, using Eq. (40) instead of Eq. (44) has little effect on the transition magnetic moment, $Q_a - Q^0 = (2m_q) + a$. The transition magnetic moment changes by -14% (-1%) for $\mu = 500\text{MeV}$ (1500MeV) for $a = 1$ and by -1% (+3%) for $\mu = 500\text{MeV}$ (1500MeV) for $a = 2$. We conclude that keeping only the $O(p^2/m_q)$ nonanalytic chiral correction is an adequate approximation for estimating the electromagnetic decay width of the Λ in QCD. Away from the physical value of the quark mass, chiral corrections become increasingly important and hence formulae in Eqs. (40) and (41) should be considered for lattice extrapolations.

The x -integrals can be evaluated analytically as shown in the Appendix. In actual lattice calculations of transition moments, the current insertion method is limited by the discrete lattice momenta available. The matrix elements contributing to the electromagnetic decay are zero for vanishing photon three-momentum q . If we wish to extract these matrix elements from a lattice simulation, it is necessary to know the momentum transfer dependence of the corresponding form factors in order to perform a model independent zero-momentum extrapolation. Formulae for the multipole transition form factors can be recovered from our expressions above by merely using the replacement

$$m \rightarrow \frac{q^2}{m^2 - x(1-x)q^2}; \quad (45)$$

where q^2 is the virtuality of the photon.

While we do not wish to encourage quenched calculations of the transition moments, we shall comment briefly on the oddities of the quenched chiral calculation at one-loop

order. None of the diagrams in Fig. 7 survive quenching. There is, however, quark mass dependence at this order, but it arises from the hairpin interactions depicted in Fig. 8. These diagrams are enhanced over their PQ-PT and PT counterparts due to the size of the singlet parameter χ_0^2 . This was first observed for baryon magnetic moments in Ref. [42], and applied to baryonic transition moments in Ref. [43]. Thus in the chiral limit, the Q-PT transition moments have chiral singularities

$$\begin{aligned} \chi_{M1}^Q &= \chi_{M1}^{(0)Q} + \frac{Q^0}{m_Q} + \chi_{M1}^2 \log m_q + \dots \\ \chi_{E2}^Q &= \chi_{E2}^{(0)Q} + \chi_{E2}^2 \log m_q + \dots; \end{aligned}$$

which should be contrasted with the chiral limit behavior in both PQ-PT and PT

$$\begin{aligned} \chi_{M1} &= \chi_{M1}^{(0)} + \chi_{M1}^{(1=2)} \frac{P}{m_q} + \dots \\ \chi_{E2} &= \chi_{E2}^{(0)} + \chi_{E2}^{(1)} m_q \log m_q + \dots; \end{aligned}$$

which are non-singular.

The inclusion of dynamical quarks thus makes a dramatic difference in the chiral behavior of these observables. There are two types of dynamical contributions: those from the gauge configurations and those from the current operator self-contraction. The latter contributions are notoriously difficult to determine from the lattice. The effective theory can remedy this situation for the electromagnetic current in three-flavor simulations. Because we require $\text{str}Q = 0$, the choice $q_{\bar{1}} = q_{\bar{2}} = q_{\bar{3}} = 0$ is not excluded, i.e. one can ignore the self-contraction of the current. In this case, one is not determining the physical transition moments, but rather the physical parameters $\chi_{M1}^{(0)}$ and $\chi_{E2}^{(0)}$ in the effective theory. These in turn can be used to make physical predictions.

V. SUMMARY

In this paper, we have extended the chiral Lagrangian with heavy quark-diquark symmetry to quenched and partially quenched theories. This extension allows for the derivation of extrapolation formulae necessary to connect lattice QCD data to the chiral limit. These formulae are easily modified to include the effects of finite volume, see Ref. [44]. Furthermore light-quark discretization effects can be incorporated, but depend on the particular fermion discretization employed in the lattice calculation.

As an example, we determined the quark mass dependence of doubly heavy baryons in both quenched and partially quenched chiral perturbation theory. We demonstrated that the hyperfine splitting between Σ and Λ baryons is sensitive to chiral non-analytic corrections both in the quenched and partially quenched theories. Thus careful chiral extrapolation is required to connect lattice data to the doubly heavy baryon spectrum in the chiral regime. Surprisingly, we found that the heavy quark-diquark symmetry relation is rather insensitive to chiral corrections. Despite the possible enhancement by factors of χ_0^2 multiplying quenched chiral logarithms, this conclusion is true in the quenched theory.

We also investigated the chiral corrections to the electromagnetic transition moments. The width of the Σ baryons should be dominated by the decay $\Sigma \rightarrow \Lambda \gamma$. The lattice can be used to predict the electromagnetic decay widths and our results should be helpful for

chiral extrapolations in these calculations. Since the doubly heavy baryon hyperfine splitting is expected to be ~ 100 MeV, lattice pion masses will not allow on-shell intermediate states as one nears the chiral regime. This is in contrast to the $\Sigma \rightarrow N$ transitions where the pion mass must be $m_\pi \lesssim 300$ MeV to avoid the multi-particle continuum. Furthermore, lattice data can test heavy quark-diquark symmetry relations between the $\Sigma \rightarrow N$ and $P \rightarrow P$ decays.

Investigation of heavy mesons and doubly heavy baryons on the lattice provides a way to explore the implications of heavy quark-diquark symmetry. By varying the light quark mass, one will be able to see how well the predictions hold up against chiral corrections. Additionally by varying the heavy quark mass, one will be able to see in what regime this symmetry of the strong interaction emerges.

Acknowledgments

B.C.T. would like to thank the Institute for Nuclear Theory at the University of Washington for hospitality and partial support during the course of this project. This work was supported by the U.S. Department of Energy under Grants No. DE-FG 02-05ER 41368 (T.M. and B.C.T.), DE-FG 02-05ER 41376, and DE-AC 05-84ER 40150 (T.M.).

APPENDIX

Here we give some formulae which are useful for evaluating the x -integrals in Eqs. (40) and (41). A useful observation is that

$$\frac{dR(x)}{dx} = \frac{x}{x^2 - 1} R(x) \quad (46)$$

Using this it is straightforward to verify the following anti-derivatives:

$$\begin{aligned} \int dx R(x) &= \frac{1}{8} \frac{1}{x^2 - 1} R^2(x) + \frac{x}{2} R(x) + \frac{x^2}{2}; \\ \int dx \frac{1}{x^2 - 1} R(x) &= \frac{1}{4} \frac{1}{x^2 - 1} R^2(x); \\ \int dx \frac{x}{x^2 - 1} R(x) &= R(x) + 2x; \\ \int dx \frac{x^3}{x^2 - 1} R(x) &= \frac{x^2 + 2}{3} R(x) + \frac{2x^3 + 12x}{9}; \end{aligned} \quad (47)$$

All integrals in Eqs.(40) and (41) can be evaluated using these identities. For example, for one of the integrals appearing in \mathcal{A} , we find

$$\begin{aligned} \int_0^1 dx \frac{1}{x^2 - 1} \log \frac{m_{ab}^2}{2} &= m_{ab} R \frac{1}{m_{ab}} \\ &= \frac{m_{ab}^2}{m_{aa}} \int \frac{m_{aa} - m_{ab}}{m_{ab}} du \log \frac{m_{ab}^2}{2} R(u) \\ &= \frac{m_{ab}^2}{m_{aa}} G \left(\frac{m_{aa} - m_{ab}}{m_{ab}}; \frac{m_{ab}}{m_{ab}} \right) G \left(\frac{m_{ab}}{m_{ab}}; \frac{m_{ab}}{m_{ab}} \right); \end{aligned}$$

where

$$G(x; y) = x^2 \log(y) \frac{x^2}{2} \frac{1}{8} \log^2 \frac{x}{x + \sqrt{x^2 - 1 + i}} \frac{1}{\sqrt{x^2 - 1 + i}} - \frac{\sqrt{x^2 - 1 + i}}{2} \log \frac{x}{x + \sqrt{x^2 - 1 + i}} \frac{1}{\sqrt{x^2 - 1 + i}} :$$

-
- [1] M . J . Savage and M . B . W ise, Phys. Lett. B 248, 177 (1990).
- [2] A . V . M anohar and M . B . W ise, Cambridge Monogr. Part. Phys. Nucl. Phys. Cosmol. 10, 1 (2000).
- [3] R . Lewis, N . M athur, and R . M . W olshyn, Phys. Rev. D 64, 094509 (2001), hep-ph/0107037.
- [4] D . Ebert, R . N . Faustov, V . O . G alkin, and A . P . M artynenko, Phys. Rev. D 66, 014008 (2002), hep-ph/0201217.
- [5] G . T . Bodwin, E . Braaten, and G . P . Lepage, Phys. Rev. D 51, 1125 (1995), hep-ph/9407339.
- [6] M . E . Luke, A . V . M anohar, and I . Z . Rothstein, Phys. Rev. D 61, 074025 (2000), hep-ph/9910209.
- [7] N . Brambilla, A . Pineda, J . Soto, and A . Vairo, Nucl. Phys. B 566, 275 (2000), hep-ph/9907240.
- [8] N . Brambilla, A . Vairo, and T . Rosch, Phys. Rev. D 72, 034021 (2005), hep-ph/0506065.
- [9] S . Fleming and T . Mehen, Phys. Rev. D 73, 034502 (2006), hep-ph/0509313.
- [10] M . B . W ise, Phys. Rev. D 45, 2188 (1992).
- [11] G . Burdman and J . F . Donoghue, Phys. Lett. B 280, 287 (1992).
- [12] T . M . Yan et al., Phys. Rev. D 46, 1148 (1992).
- [13] J . Hu and T . Mehen, Phys. Rev. D 73, 054003 (2006), hep-ph/0511321.
- [14] T . D . Cohen and P . M . Hohler (2006), hep-ph/0606084.
- [15] M . A . M oinester et al. (SELEX), Czech. J. Phys. 53, B201 (2003), hep-ex/0212029.
- [16] M . Mattson et al. (SELEX), Phys. Rev. Lett. 89, 112001 (2002), hep-ex/0208014.
- [17] A . Ocherashvili et al. (SELEX), Phys. Lett. B 628, 18 (2005), hep-ex/0406033.
- [18] V . V . Kiselev and A . K . Likhoded (2002), hep-ph/0208231.
- [19] B . Aubert et al. (BABAR) (2006), hep-ex/0605075.
- [20] R . Chistov et al. (BELLE) (2006), hep-ex/0606051.
- [21] N . M athur, R . Lewis, and R . M . W olshyn, Phys. Rev. D 66, 014502 (2002), hep-ph/0203253.
- [22] J . M . Flynn, F . Mescia, and A . S . B . Tariq (UKQCD), JHEP 07, 066 (2003), hep-lat/0307025.
- [23] A . Morel, J. Phys. (France) 48, 1111 (1987).
- [24] S . R . Sharpe, Phys. Rev. D 46, 3146 (1992), hep-lat/9205020.
- [25] C . W . Bernard and M . F . L . Golterman, Phys. Rev. D 46, 853 (1992), hep-lat/9204007.
- [26] C . W . Bernard and M . F . L . Golterman, Phys. Rev. D 49, 486 (1994), hep-lat/9306005.
- [27] S . R . Sharpe, Phys. Rev. D 56, 7052 (1997), hep-lat/9707018.
- [28] M . F . L . Golterman and K . C . Leung, Phys. Rev. D 57, 5703 (1998), hep-lat/9711033.
- [29] S . R . Sharpe and N . Shoreh, Phys. Rev. D 62, 094503 (2000), hep-lat/0006017.
- [30] S . R . Sharpe and N . Shoreh, Phys. Rev. D 64, 114510 (2001), hep-lat/0108003.
- [31] P . H . Damgaard and K . Splittor, Phys. Rev. D 62, 054509 (2000), hep-lat/0003017.
- [32] S . R . Sharpe and Y . Zhang, Phys. Rev. D 53, 5125 (1996), hep-lat/9510037.
- [33] W . A . Bardeen, E . Eichten, and H . Thacker, Phys. Rev. D 69, 054502 (2004), hep-lat/0307023.

- [34] S. R. Sharpe, *Phys. Rev. D* 69, 034504 (2004), [hep-lat/0308009](#).
- [35] B. C. Tiburzi, *Phys. Rev. D* 71, 034501 (2005), [hep-lat/0410033](#).
- [36] S. Fajfer and J. Kamenik (2006), [hep-ph/0606278](#).
- [37] M. Golterman and E. Pallante, *JHEP* 10, 037 (2001), [hep-lat/0108010](#).
- [38] B. C. Tiburzi, *Phys. Rev. D* 71, 054504 (2005), [hep-lat/0412025](#).
- [39] W. Detmold and C. J. D. Lin, *Phys. Rev. D* 71, 054510 (2005), [hep-lat/0501007](#).
- [40] J. F. Amundson et al., *Phys. Lett. B* 296, 415 (1992), [hep-ph/9209241](#).
- [41] M. J. Savage, *Phys. Rev. D* 65, 034014 (2002), [hep-ph/0109190](#).
- [42] M. J. Savage, *Nucl. Phys. A* 700, 359 (2002), [nucl-th/0107038](#).
- [43] D. Amdt and B. C. Tiburzi, *Phys. Rev. D* 69, 014501 (2004), [hep-lat/0309013](#).
- [44] D. Amdt and C. J. D. Lin, *Phys. Rev. D* 70, 014503 (2004), [hep-lat/0403012](#).