# Toward P recision M easurem ent of B $_{\rm K}~$ with Q uenched K ogut-Susskind Q uarks

JLQCD Collaboration

S.Aok<sup>a</sup>, M.Fukugita<sup>b</sup>, S.Hashim oto<sup>c</sup>, Y.Iwasak<sup>a;d</sup>, K.Kanaya<sup>a;d</sup>, Y.Kuramash<sup>c</sup>, H.Mino<sup>e</sup>, M.Okawa<sup>c</sup>, A.Ukawa<sup>a</sup>, T.Yoshie<sup>a;d</sup>

<sup>a</sup> Institute of Physics, University of Tsukuba, Tsukuba, Ibaraki 305, Japan

<sup>b</sup>Yukawa Institute for Theoretical Physics, Kyoto University, Kyoto 606, Japan

 $^{
m c}$ N ational Laboratory for H igh Energy Physics (KEK), T sukuba, Ibaraki 305, Japan

<sup>d</sup>Center for Computational Physics, University of T sukuba, T sukuba, Ibaraki 305, Japan

<sup>e</sup>Faculty of Engineering, Yam anashi University, Kofu 400, Japan

W e present a status report of our ongoing e ort toward a precision determ ination of the kaon B parameter with the K ogut-Susskind quark action in quenched QCD.Results for  $B_K$  so far accumulated at = 5:85, 5.93, 6.0 and 6.2 corresponding to a <sup>1</sup> = 1:3 2:6 GeV do not exhibit the theoretically expected O (a<sup>2</sup>) behavior, but are apparently m ore consistent with an O (a) behavior.

### 1. Introduction

The kaon B parameter is one of the crucial hadron weak matrix elements for constraining the CP violation phase of the Cabibbo-Kobayashi-Maskawamatrix from experiments. A pioneering study was carried out by Gupta, Kilcup, Patel and Sharpe (GKPS) [1 {3], which laid a foundation for a lattice QCD determ ination of this quantity with the Kogut-Susskind guark action. A num ber of problem s, however, are posed to warrant viability of the lattice result in the continuum . W hile som e aspects have been exam ined previously [4,5], further studies are needed to establish the continuum value of  $\boldsymbol{B}_K$  . The statistics of GKPS are only 15 ( = 6.0) to 24 ( = 6.4) congurations: as a consequence the error of  $B_{K}$  at = 6:4 is so large that an O  $(a^2)$  dependence of BK on the lattice spacing inferred from a theoretical analysis[3] can not be ascertained. Furtherm ore the lattice size of  $32^3$  48 em ployed at = 62 and 6.4 m ay be too sm all in the tem poraldirection. We have therefore decided to carry

out a new set of simulations with improved statistics, incorporating advance in methods of analyses. Here we present a status report of this program which is being carried out on VPP 500/80 newly installed at KEK.

2. Sim ulation m ethod and run param eters

We basically employ the formalism developed by GKPS for calculating the B parameter. Following improvements are included in our simulations, however.

On each con guration we calculate  $2^3 = 8$ quark propagators corresponding to wall sources with a unit source placed at one of the corners of each spatial cube. This allows us to construct a wall source which creates only the pseudoscalar meson in the Nambu-Goldstone channel[6]. In contrast the even-odd source employed in previous studies[1{4] also creates vector mesons that contaminate signals. The sources are placed at the two edges of the lattice, and the Dirichlet boundary condition is employed in the temporal direction. G auge con gurations are xed to the Landau gauge.

presented by S.Aoki

Tabl	e1	
Run	param	eters

	5.85	5.93	6.0	62
L <sup>3</sup> T	16 <sup>3</sup> 32	20 <sup>3</sup> 40	24 <sup>3</sup> 40	32 <sup>3</sup> 64
# conf.	60	50	50	40
a <sup>1</sup>	1.34 (3)	1.58 (4)	1.88 (5)	2.62 (9)
m <sub>q</sub> a	0.01-0.04	0.01-0.04	0.01-0.03	0.005-0.02
m₃a=2	0.0197	0.0156	0.0125	0.0089
t	10-20	12–26	14-24	20-42

Four-quark operators written in terms of K ogut-Susskind quark elds are extended over a  $2^4$  hypercube. W e employ both gauge-invariant and non-invariant operators[4], which dier by an insertion of appropriate gauge link factors for the form er. The one-loop result for the lattice renormalization factors[7,8] is used to convert lattice values of B<sub>K</sub> to those in the continuum, renormalized with the naive dimensional regularization (NDR) scheme.

A source of uncertainty in the conversion is the choice of the gauge coupling constant used for the renormalization factors. We take the meaneld improved  $\overline{M}$  S coupling constant at the cuto scale = =a[9]. Correspondingly we improve the operators[10] through the replacement  $! \stackrel{P}{u_0}$  and U  $! u_0^{-1}U$ , where is the K ogut-Susskind quark eld, U the link variable and  $u_0 = P^{1=4}$  with P the average plaquette. Modi cations of renormalization factors due to this replacement [7,8] are also taken into account.

We note that we have ignored the mixing of non-N am bu-G oldstone four-quark operators in the present analysis. Studies of this mixing effect, which is believed to vanish in the continuum limit, are left for future work.

P aram eters of runs we have so far carried out are listed in Table 1. G auge con gurations are generated with the 5-hit heatbath algorithm at 2000(6:0) or 5000(=6:2) sweep intervals. The physical scale of lattice spacing is determ ined from the meson mass in the VT channel. Values of are chosen so that the lattice spacing is roughly equally spaced, and the lattice size is increased for weaker coupling to keep the physical lattice size approximately constant at La 2:4 2:5 fm. A sim ilar consideration is



Figure 1. C on parison of lattice value of  $B_K$  for gauge non-invariant operator at = 6.0 and 62. D at a for = 6.2 are shifted downward by 0.05 to avoid overlap.

m ade for the choice of the tting interval in extracting  $B_K$  from kaon G reen's functions. Q uark m ass is taken in steps of 0.01 for 6.0 and 0.005 at = 6.2. W hile we have evaluated  $B_K$  for non-degenerate strange and down quark m asses, we shall restrict ourselves to the degenerate case in the present analysis. The strange quark m ass yielding the observed ratio  $m_K = m = 0.648$  for this case is also given in Table 1. W e estimate errors by a single elimination jackknife procedure.

## 3. Com parison with previous work

In Fig.1 lattice values of  $B_K$  for the gauge noninvariant operator at = 6.0 and 6.2 are plotted. At = 6.2 our results are consistent with those of G K P S [2,3] at the level of one standard deviation. On the other hand, our values at = 6.0are system atically low er than those of G K P S and Ishizuka et al. (IFM O SU) [4] obtained on the sam e size of lattice.

In order to understand the discrepancy, we note that GKPS and IFM OSU respectively used 15 and 10 con gurations as compared to our 50, and that both groups employed the even-odd wall source, while we used the  $2^3$  source. For a com – parison under the same condition, we divided our ensemble into 5 sets of 10 con gurations and recalculated  $B_K$  for each set. In addition we made a new calculation for 3 sets of 10 con gurations using the even-odd wall source. The eight values



Figure 2. Lattice value of  $B_K$  at = 6.0 calculated for 10 con gurations with  $2^3$  (circles) and even-odd (diam onds) source as compared with previous results obtained with even-odd source (triangles). Solid lines are values for the full ensemble with errors indicated by dotted lines.

of  $B_K$  thus obtained are plotted in Fig. 2 and are compared with those of IFM OSU and GKPS. Solid lines represent values obtained for the full ensemble for each type of source.

W enote that, while results for the fullensem ble are consistent between the two types of sources, both errors and uctuations of central values are signi cantly reduced with the  $2^3$  source, dem onstrating the advantage of this method. Since values of GKPS and IFMOSU are consistent with ours for the even-odd source within one to two standard deviations, we conclude that previous results have overestim ated the value of  $B_K$  at = 6:0 due to insu cient statistics.

#### 4. Quark m ass dependence

Extraction of the physical value of B<sub>K</sub> requires an interpolation of results to the quark m ass corresponding to the physical kaon. We nd a suggestive result that the quark m ass dependence of our data is well described by the form predicted by chiral perturbation theory, which we employ for the above purpose. For the case of degenerate strange and down quark treated in the present analysis, the form ula is given by [11]

$$B_{K} = B [1 \quad 3c(m_{q}a) \log(m_{q}a) + b(m_{q}a)];$$
 (1)

where  $c = = (4 \text{ f})^2$  with  $= (m_K a)^2 = m_q a$  and b is an unknown constant. A representative ex-



Figure 3. M ean-eld improved  $B_K$  (NDR, 2G eV) as a function of  $m_q a$  at = 62 on a  $32^3$  64 lattice. Solid lines are ts to (1) and dashed line corresponds to physical value of  $m_K = m$ .

ample of the t taking B; c and b as parameters is plotted in Fig. 3 together with the quark m ass corresponding to  $m_{\rm K} = m = 0.648$  (dashed line). Several uncertainties exist, however, to take the agreement as evidence for the presence of chiral logarithm in our data. The pion decay constant f estimated from the tted values of c and the spectrum results for the slope turned out to have a large value of 150 300 M eV with 10 20% error as compared to the experimental value of 93 M eV. A lso a quadratic form without logarithm ts our data equally well.

## 5. Results for $B_K$

We plot our results for  $B_K$  renormalized at = 2 GeV in the NDR scheme as a function of m a in Fig. 4 both for gauge invariant (led circles) and non-invariant (open circles) operators. We observe that the data do not exhibit the theoretically predicted O ( $a^2$ ) dependence[3], and are apparently more consistent with a linear behavior, at least over the range m a = 0.57 0.29 (or 1=a = 1.3 2.6 GeV from m) corresponding to = 5.85 6.2. If we assume a linear form, we obtain  $B_K$  (a) = 0.4825(83) + 0.489(17)m a and 0.5097(83) + 0.464(17)m a for the gauge invariant and non-invariant operators.

We have varied the analysis procedure to see to what extent the values of  $B_K$  are modiled: (i) In-stead of  $m_K = m$  we examined the choice m = m



Figure 4. M ean-eld improved  $B_K$  (NDR, 2G eV) as a function of m a, together with linear ts (solid lines).

for estimating the strange quark mass. This only resulted in a common downward shift of the four values of B<sub>K</sub> by about 0:02, and hence does not change a linear dependence on the lattice spacing. (ii) W e also replaced them ean-eld in proved  $\overline{MS}$  coupling constant  $q_{\overline{MS}}$  (=a) in the renormalization factors by the running coupling constant at = 2 GeV evaluated by the two-loop renorm al-

ization group evolution starting from  $q_{MS}$  (=a). This increases the value of  $B_K$  by about 0.03-0.05 without, how ever, changing the linear behavior.

Let us comment on the fact that our results for  $B_K$  obtained with the two types of operators do not agree with each other and that the discrepancy does not become smaller toward the continuum limit. Possible sources of the discrepancy are an operator dependence of nite lattice spacing e ects, which disappears in the continuum limit, and 0 (g<sup>4</sup>) corrections in the renormalization factors which are not included in our analysis. Naively one would expect the magnitude of  $0 (q^4)$  corrections to also dim in ish tow ard the continuum limit. This is not the case, how ever, for  $B_K$  renorm alized at a xed scale = 2 GeV, since the gauge coupling constant g() goveming these corrections is independent of the lattice spacing. We have checked that increasing the scale leads to a smaller discrepancy. Thus the di erence of B  $_{\rm K}$  for the two types of operators extrapolated to the continuum lim it provides an estim ate of the magnitude of  $0 (g^4)$  corrections.

Finally we rem ark that an O ( $a^2$ ) dependence is not completely nuled out from our data if there exists an appropriate O ( $a^3$ ) or O ( $a^4$ ) term which just brings the curve to behave as if it is linear over a range of lattice spacing including that explored in our simulation. If this rather exceptional case happens, the value of B<sub>K</sub> extrapolated to the continuum lim it a = 0 would increase by about 0.05.

#### 6. Sum m ary

In this report we have described our e ort toward in proving previous work on the kaon B param eter with the K ogut-susskind quark action in quenched QCD. The results obtained at four values of show that  $B_K$  falls linearly with the lattice spacing over the range 1=a=1:3 2:6 GeV. If the theoretical prediction of an O (a<sup>2</sup>) dependence is valid, this m eans that such a behavior will become apparent only for sm aller values of lattice spacing. W e plan to carry out a simulation at = 6:4 6:5 em ploying the lattice size of  $40^3$  96.

## REFERENCES

- G.Kikup et al, Phys. Rev. Lett. 64 (1990) 25.
- S.Sharpe et al, Nucl. Phys. B (Proc. Suppl.) 26 (1992) 197.
- S. Shampe, Nucl. Phys. B (Proc. Suppl.)34 (1994) 403.
- N.Ishizuka et al, Phys.Rev.Lett. 71 (1993)
   24.
- W .Lee and M .K lom fass, Nucl. Phys.B (Proc. Suppl.)42 (1995) 418.
- N. Ishizuka et al, Nucl. Phys. B 411 (1994) 875.
- N. Ishizuka and Y. Shizawa, Phys. Rev. D 49 (1994) 3519.
- S. Sharpe and A. Patel, Nucl. Phys. B417 (1994) 307.
- A. ElKahdra et al, Phys. Rev. Lett. 69 (1992) 729.
- 10.G.P.Lepage and P.M ackenzie, Phys. Rev. D 48 (1993) 2250.
- 11.S.Shampe, Phys.Rev.D 46 (1992) 3146.