Undoubled Chiral Fermions on a Lattice

She-Sheng Xue [a]

[a] INFN - Milan Section, Via Celoria 16, Milan, Italy

arXiv:hep-lat/9607047v1 21 Jul 1996

We analyze the dynamics of an $SU_L(2)\otimes U_R(1)$ chiral theory on the lattice with a strong multifermion coupling. It is shown that no spontaneous symmetry breaking occurs; the "spectator" fermion ψ_R is a free mode; doublers are decoupled as massive Dirac fermions consistently with the chiral symmetries. In 1+1 dimension, we show that the right-handed three-fermion state disappears at the threshold and an undoubled left-handed chiral fermion remains in the continuum limit.

Since the demonstration of "no-go" theorem [1] of Nielsen and Ninomiya in 1981 the problem of chiral fermion "doubling" and "vector-like" phenomenon on a lattice still exists if one insists on preserving chiral symmetry. One of the ideas to get around this "no-go" theorem was proposed with a multifermion coupling model [2]. However, it was pointed out [4] that such a model of multifermion couplings fails to give chiral fermions in the continuum limit.

We have been studying multifermion couplings on the lattice for years [5] and we believe that such models still have a chance to work [6]. Let us consider the following fermion action of the $SU_L(2)\otimes U_R(1)$ chiral symmetries on the lattice with a multifermion coupling.

$$\begin{aligned} S &= S_k + g_2 \bar{\psi}^i_L(x)\cdot\Delta\psi_R\Delta\bar{\psi}_R\cdot\psi^i_L(x), \qquad (1)\\ S_k &= \frac{1}{2a}\Big(\bar{\psi}^i_L\partial\psi^i_L + \bar{\psi}_R\partial\psi_R\Big) \end{aligned}$$

where "a" is the lattice spacing; ψ^i_L $(i=1,2)$ is an $SU_L(2)$ doublet, ψ_R is an $SU_L(2)$ singlet and both are two-component Weyl fermions. The ψ_R is treated as a "spectator" fermion. The $\Delta(x)$ is a second order differential operator on the lattice. The multifermion coupling g_2 is a dimension-10 operator relevant only for doublers $p=\tilde{p}+\pi_A$, but irrelevant for the chiral fermions $p=\tilde{p}\sim 0$ of the ψ^i_L and ψ_R. Action (1) preserves the global chiral symmetry $SU_L(2)\otimes U_R(1)$ and the $\psi^i_L(x)$ can be gauged to have the exact local $SU_L(2)$ chiral gauge symmetry. In addition, action (1) possesses a ψ_R-shift-symmetry[7],

$$\psi_R(x)\to\psi_R(x)+\epsilon. \qquad (2)$$

The $U_L(1)$ global symmetry relating to the conservation of the fermion number of the $\psi^i_L(x)$ is explicit in eq.(1).

To seek a possible segment $g_2 \gg 1$, where an undoubled $SU_L(2)$-chiral fermion exists in the continuum limit, we are bound to demonstrate the following properties of the theory in this segment:

1. the normal mode of ψ_R is a free mode and decoupled;
2. no spontaneous chiral symmetry breaking occurs ;
3. all doublers $p=\tilde{p}+\pi_A$ are bound to be massive Dirac fermions and decoupled consistently with the $SU_L(2)\otimes U_R(1)$ chiral symmetry;
4. an undoubled chiral fermion $p=\tilde{p}$ of ψ^i_L exists in the low-energy spectrum.

To prove the first and second points, we use the Ward identity stemming from the ψ_R-shift-symmetry (2). Considering the generating functional $W(\eta)$ of the theory,

$$\begin{aligned} W(\eta) &= -\ell n Z(\eta), \qquad (3)\\ Z(\eta) &= \int [d\psi^i_L d\psi_R]\exp\Big(-S+\bar{\psi}\eta\Big), \end{aligned}$$

we define the effective action $\Gamma(\psi'^i_L,\psi'_R)$ as the Legendre transform of $W(\eta)$. Making the parameter ϵ in eq.(2) to be space-time dependent, and varying the generating functional (3) according to the transformation rule (2) for arbitrary $\epsilon(x)\neq 0$,

we arrive at the Ward identity corresponding to the ψ_R-shift-symmetry of the action (1):

$$\frac{1}{2a}\partial\psi'_R + g_2\langle\Delta\big(\bar\psi^i_L\cdot\Delta\psi_R\psi^i_L\big)\rangle - \frac{\delta\Gamma}{\delta\bar\psi'_R} = 0. \qquad (4)$$

From this Ward identity, one can obtain all one-particle irreducible vertices $\Gamma^{(n)}_R$ containing at least one external ψ_R. Taking functional derivatives of eq.(4) with respect to appropriate "primed" fields and then putting external sources $\eta = 0$, one can derive:

$$\int_x e^{-ipx}\frac{\delta^{(2)}\Gamma}{\delta\psi'_R(x)\delta\bar\psi'_R(0)} = \frac{i}{a}\gamma_\mu \sin p^\mu, \qquad (5)$$

$$\frac{1}{2}\Sigma^i(p) \sim w(p), \qquad (6)$$

where $w(p)$ is the Fourier transform of $\frac{1}{2}\Delta(x)$, which is the Wilson factor [8]. One has

$$w(p) = \sum_\mu (1-\cos p_\mu), \;\; \Sigma^i(0) = 0. \qquad (7)$$

Eq.(5) indicates an absence of the wave-function renormalization of the ψ_R. Eqs.(6,7) show the vanishing of the NJL symmetry breaking [3] for $p = 0$.

To adopt the technique of strong-coupling expansion in powers of $\frac{1}{g_2}$ and we make a rescaling of the fermion fields

$$\psi^i_L \to (g_2)^{\frac{1}{4}}\psi^i_L; \;\; \psi_R \to (g_2)^{\frac{1}{4}}\psi_R. \qquad (8)$$

For the strong coupling $g_2 \to \infty$, the kinetic terms S_k can be dropped and we compute the integral of $e^{-S(x)}$ in this limit:

$$Z = 2^{4N}\left(\det\Delta^2(x)\right)^4, \qquad (9)$$

where the determinant is taken only over the lattice-space-time and "N" is the number of lattice sites. For the non-zero eigenvalues of the operator $\Delta^2(x)$, eq. (9) shows an existence of the sensible strong-coupling limit. However, as for the zero eigenvalue of the operator $\Delta(x)$, this strong-coupling limit is not analytic and the strong-coupling expansion in powers of $\frac{1}{g_2}$ breaks down.

To show the second point concerning the NJL symmetry breaking, we calculate the two-point functions:

$$S^j_{RL}(x) \equiv \langle\psi_R(0), \bar\psi^j_L(x)\rangle, \qquad (10)$$

$$S^j_{MR}(x) \equiv \langle\psi_R(0), [\bar\psi^j_L\cdot\psi_R]\bar\psi_R(x)\rangle. \qquad (11)$$

At non-trivial leading order $O(\frac{1}{g_2})$, we get the recursion relations:

$$S^j_{RL}(x) = \frac{\sum^\dagger_\mu S^j_{MR}(x+\mu)\gamma_\mu}{8a^3 g_2\Delta^2(x)}, \qquad (12)$$

$$S^j_{MR}(x) = \frac{\sum^\dagger_\mu S^j_{RL}(x+\mu)\gamma_\mu}{2a g_2\Delta^2(x)}. \qquad (13)$$

These recursion relations are not valid where the operator $\Delta(x)$ has zero eigenvalues. For $p \neq 0$ and $\Delta(p) = 2w(p) \neq 0$, the Fourier transform of these recursion relations leads to the solution ($p \neq 0$),

$$S^j_{RL}(p) \equiv \Sigma^j(p) = 0, \quad S^j_{MR}(p) = 0. \qquad (14)$$

Together with eq.(7), we prove the second point.

We turn to the third point that concerns decoupling of doublers. On this extreme strong coupling ($g_2 \gg 1$) symmetric segment, the ψ^i_L and ψ_R in (1) are bound up to form the three-fermion states[2,6]:

$$\Psi^i_R = \frac{(\bar\psi_R\cdot\psi^i_L)\psi_R}{2a}; \;\; \Psi^n_L = \frac{(\bar\psi^i_L\cdot\psi_R)\psi^i_L}{2a}, \qquad (15)$$

which carry the appropriate quantum numbers of the chiral group that accommodates ψ^i_L and ψ_R. These three-fermion states are Weyl fermions and respectively pair up with the $\bar\psi_R$ and $\bar\psi^i_L$ to be massive, neutral Ψ_n and charged Ψ^i_c Dirac fermions,

$$\Psi^i_c = (\psi^i_L, \Psi^i_R), \quad \Psi_n = (\Psi^n_L, \psi_R). \qquad (16)$$

To show this phenomenon, we compute the following two-point functions,

$$S^{ij}_{LL}(x) = \langle\psi^i_L(0)\bar\psi^j_L(x)\rangle,$$
$$S^{ij}_{ML}(x) = \langle\psi^i_L(0)\bar\Psi^j_R(x)\rangle,$$
$$S^{ij}_{MM}(x) = \langle\Psi^i_R(0)\bar\Psi^j_R(x)\rangle. \qquad (17)$$

In the lowest non-trivial order $O(\frac{1}{g_2})$, we obtain the following recursion relations

$$S^{ij}_{LL}(x) = \frac{\sum^\dagger_\mu S^{ij}_{ML}(x+\mu)\gamma_\mu}{8a^3 g_2\Delta^2(x)}, \qquad (18)$$

$$S^{ij}_{MM}(x) = \frac{\sum^{\dagger}_{\mu}\gamma_\mu\gamma_\circ S^{ij\dagger}_{ML}(x+\mu)\gamma_\circ}{8a^3g_2\Delta^2(x)}, \quad (19)$$

$$S^{ij}_{ML}(x) = \frac{\delta(x)\delta_{ij}}{4ag_2\Delta^2(x)} + \frac{\sum^{\dagger}_{\mu}S^{ij}_{LL}(x+\mu)\gamma_\mu}{8a^2g_2\Delta^2(x)}. \quad (20)$$

Eq.(20) suggestes that the states coupling to operators $\psi^i_L(x)$ and $\Psi^i_R(x)$ are mixed, producing a massive four-component Dirac fermion. To find the masses, we make the Fourier transform of these recursion equations for $p \neq 0$ and $\Delta^2(p) = 4w^2(p) \neq 0$ and obtain the inverse propagators of the charged Dirac fermions,

$$S^{-1}_c(p) = \frac{i}{a}\sum_\mu \gamma_\mu \sin p^\mu + 8ag_2w^2(p). \quad (21)$$

This shows that all doublers $p = \tilde{p}+\pi_A$ are decoupled as massive Dirac fermions consistently with the $SU_L(2)\otimes U_R(1)$ chiral symmetries.

We are left with the last point that undoubled chiral fermions ($p = \tilde{p} \sim 0$) of ψ^i_L exist in this segment. We repeat the same calculations and discussions in 1+1 dimensions, where the time direction is continuous and space is discrete. We obtain the dispersion relation corresponding to the Dirac fermion (21) for $P \neq 0$,

$$E(P) = \pm\sqrt{\sin^2 P + (8a^2gw^2(P))^2}, \quad (22)$$

which is analytically continuous to low momentum states $P \to 0$, unless it hits the threshold. For a given total momentum P, we consider the system that contains three free chiral fermions, i.e. right-movers $\bar{\psi}_R, \psi_R$ with momenta p_1 and p_2 and a left-mover ψ^i_L with momentum p_3 ($|p_i| \ll \frac{\pi}{2}, i = 1,2,3$),

$$P = p_1 + p_2 + p_3 \ll \frac{\pi}{2}. \quad (23)$$

Since NJL spontaneous symmetry breaking does not occur when $|p_i| \to 0$, the total energy of such a system is given by

$$E = E_1(p_1) + E_2(p_2) + E_3(p_3)$$
$$E_i = \pm p_i, \quad (24)$$

where all negative-energy states have been filled. The lowest energy (the threshold) of such a system and corresponding configuration are

$$E_t(P) = 3\sin P, \quad (25)$$
$$p_3 = -P, p_1 = p_2 = P,$$

The three-fermion state is stable if there is a gap δ between the threshold (25) and the energy (22) of the three-fermion state, i.e.

$$\delta = E_t(P) - E(P) > 0. \quad (26)$$

The three-fermion state disappears, when the gap δ goes to zero,

$$\delta = E_t(P) - E(P) = 0. \quad (27)$$

Substituting eqs.(22) and (25) into eq.(27), we obtain in the continuum limit $P \to 0$, the gap

$$\delta(P) \to 0, \quad (28)$$

where the three-fermion-state spectrum dissolves into free chiral fermion spectra. As a result, an undoubled chiral fermion exists in the continuum limit.

I thanks Profs. G. Preparata, H. B. Nielsen and E. Eichten for discussions on the last point.

REFERENCES

[1] H.B. Nielsen and M. Ninomiya, *Nucl. Phys.* **B185** (1981) 20; *ibid.* **B193** (1981) 173; *Phys. Lett.* **B105** (1981) 219.

[2] E. Eichten and J. Preskill, *Nucl. Phys.* **B268** (1986) 179.

[3] Y. Nambu and G. Jona-Lasinio, *Phys. Rev.* **122** (1961) 345.

[4] M.F.L. Golterman, D.N. Petcher and E. Rivas, *Nucl. Phys.* **B395** (1993) 597.

[5] G. Preparata and S.-S. Xue, *Phys. Lett.* **B264** (1991) 35; *ibid.* **B335** (1994) 192; *ibid.* **B377** (1996)124; *Nucl. Phys.* **B26** (Proc. Suppl.) (1992) 501; *ibid.* **B30** (1993) 647; **B47** (1996) 583.

[6] S.-S. Xue, hep-lat/9605003, to appear in *Phys. Lett. B* and hep-lat/9605005.

[7] M.F.L. Golterman, D.N. Petcher, *Phys. Lett.* **B225** (1989) 159.

[8] K. Wilson, in *New phenomena in subnuclear physics* (Erice, 1975), ed. A. Zichichi (Plenum, New York, 1977).