Singularities of the Partition Function for the Ising M odel C oupled to 2d Q uantum G ravity

J.Ambj rn^y, K.N.Anagnostopoulos^y and U.M agnea^z

^YT he Niels Bohr Institute ^ZNORDITA Blegdam svej 17, DK-2100 Copenhagen , Denmark

A bstract

We study the zeros in the complex plane of the partition function for the Ising model coupled to 2d quantum gravity for complex magnetic eld and real temperature, and for complex tem perature and realmagnetic eld, respectively. We compute the zeros by using the exact solution coming from a two matrix model and by M onte C arlo simulations of Ising spins on dynamical triangulations. We present evidence that the zeros form simple one-dimensional curves in the complex plane, and that the critical behaviour of the system is governed by the scaling of the distribution of the singularities near the critical point. Despite the small size of the system s studied, we can obtain a reasonable estimate of the (known) critical exponents.

e-m ail addresses: ^y am bjorn, konstant, ^z um agnea@ nbidk

1 Introduction

The understanding of the phase transition of a statistical mechanical model has long been connected to the study of its partition function zeros. This line of research was pioneered by Yang and Lee [1], and has subsequently been pursued by many authors (for recent work see β] and references therein). In the therm odynam ic lim it, the zeros of the partition function of such a system in the complex coupling space accumulate in nitely close to the critical coupling on the real axis. This de nes, in the in nite volum e lim it, disconnected regions in the complex plane, with di erent analytic structure for the therm odynam ic functions which describe the phases of the system .

The critical behaviour in the neighbourhood of a continuous phase transition can be extracted from the behaviour of the density of zeros near its singular points [2, 4, 5]. In the best-understood m odels, these points occur at the ends of lines on which the zeros accumulate in the in nite volum e lim it. Such singular points, even when they occur at non {physical values of the coupling, can be considered as ordinary critical points with distinctive critical exponents [7] (see also [8]). For a physical phase transition, these singularities coincide with the (real) critical coupling of the system.

Substantial progress, also from a practical point of view, has been made by applying renormalization group arguments to the motion of zeros, thereby deriving nite size scaling relations by which the critical exponents can be computed [6]. This nite-size scaling technique applied to complex zeros has been proven powerful in numerical computations of critical exponents.

A special class of statistical system s are spin system s de ned on random surfaces. In addition to being interesting system s per se, they have received special attention for their relation to conform alm atter system s coupled to two dimensional quantum gravity. At criticality, the spin system s are described by conform all editheories (CFT) coupled to gravity. The critical exponents of such conform all theories have been found by K nizhnik, Polyakov and Zam obdchikov (KPZ) to be simply related to the critical exponents of the same CFT de ned in at space [9]. In particular, the critical Ising m odel coupled to 2d gravity can be recast in the form of a c = 1=2 conform all editheory coupled to 2d gravity. In addition, the equivalence of the Ising m odel on a random surface to a herm itean two m atrix m odel, discovered and studied by K azakov and B oulatov [10, 11], renders it exactly solvable. The critical exponents were rst com puted this way, outside of the CFT context. The system is found to have a third order phase transition from a high tem perature disordered phase to a low tem perature ordered phase. The coupling between m atter and the geom etry of the surface is found to change the critical exponents associated to geom etry only at the phase transition.

The location of the partition function zeros of the Ising model in the complex fugacity plane, the so called Lee{Yang zeros, in the presence of gravity has been studied by Staudacher [12]. The som ewhat surprising discovery that the Lee{Yang zeros lie on the unit circle in the complex fugacity plane for each lattice size N will be con med below for slightly bigger lattices than those considered in [12]. As discussed by Staudacher, the Lee{Yang theorem [1] guarantees that the roots of the partition function of the Ising model on a xed lattice Z at (G_N) for a large class of not too pathological lattices G_N will lie on the unit circle, but it is not a priori expected that the roots of the partition function Z_N , obtained by summing the Z at (G_N)'s over all dynamical

random lattices G_N ,

$$Z_{N} = \begin{array}{c} X \\ Z_{M} = \begin{array}{c} X \\ G_{M} \end{array} \qquad (1.1)$$

will also be located on the unit circle. The summation in (1.1) corresponds to an integration over the additional quantum degree of freedom introduced by a uctuating metric. The Lee{ Yang theorem for this case remains unproven. A priori the zeros can be located on di erent curves or even on sets with more complicated topology, or on two-dimensional regions. The latter cases have been observed for the complex temperature zeros of 2d Ising and Potts models on hierarchical lattices [27], for anisotropic lattices [28], and interestingly, also for a regular lattice made of di erent polygons [29]. (See also ref. [30], where a very interesting, albeit som ewhat inconclusive attempt at studying the partition function zeros of Ising spin glasses is made, and ref. [31], where the zeros for aperiodic system s were studied.)

In this paper we use the same method as in [12] in order to compute the exact partition function for system s with up to 14 squares. We compute the Lee{Yang zeros of the partition function and con m that they are located on the unit circle of the complex fugacity plane. We also compute the zeros of the partition function for complex temperature (Fisher zeros) and nd that they move on curves in the complex temperature plane as the lattice size is varied. This indicates that in the in nite volume limit, the zeros accumulate to form dense sets on these curves. We test the scaling relations derived by Itzykson, Pearson and Zuber in the context of regular lattices [6] and extract from there some (combinations of) critical exponents. These scaling relations are based on the assumption that the spin (spin correlation length diverges at the critical point. Despite the sm all size of the systems studied, we nd reasonable agreement with the critical exponents given in [10, 11]. This result is expected from the non{trivial fact that the matter correlation length diverges at the phase transition [13] (see [15] for an example where this is not true). This way we provide strong evidence that the phase transition is governed by the singularities of the distribution densities at H = 0 and $_{c}$ of the Lee(Y and A risher zeros respectively as is the case for the Ising model on a xed lattice. The pattern of zeros we observe con m s the observation in [14] that no antiferrom agnetic transition occurs for the Ising m odel on a square lattice when the spins are placed on the faces of the lattice. U sing duality, we can study the pattern of the Fisher zeros of the dualm odel where the spins are placed on the ~. This indicates vertices. We observe that the set of Fisher zeros maps onto itself under ~! the existence of an antiferrom agnetic transition whose critical exponents are identical to the ones of the ferrom agnetic transition . This con rm s the results reported in [14].

One of the m ethods for computing complex zeros is provided by M onte C arb simulations and histogram m ing techniques. These techniques have been applied successfully to system s de ned in at space [16, 17]. They have also been tried on several system s coupled to gravity (random surfaces coupled to extrinsic curvature, Potts m odels, 4d quantum gravity [22]), but it proved di cult to observe the partition function complex zeros. Here, we apply the m ethod to Ising spins on dynam ical triangulations [18], and we are able to observe the zeros lying closest to the real axis for a range of lattice sizes w ith up to 256 vertices. In general, it was easier to nd the Lee{Y ang zeros than the F isher zeros. In the latter case, we were able to locate only the rst

W e thank Desm ond Johnston for pointing this out to us.

zero for any given lattice size. A fler determ ining the position of the zeros as a function of lattice size, we test their scaling and the critical exponents com puted are in reasonable agreem ent with their exactly known values.

In section 2 we de ne our model and review the expected scaling behaviour of the zeros on the assumption of a diverging matter correlation length. In section 3 we describe in detail the methods we use in order to compute the zeros, and in section 4 we test their nite size scaling behaviour. Section 5 contains som e concluding remarks.

2 Themodel

W e will study one Ising model on a dynamical square or triangular lattice. In the form er case the spins are located on the N faces (squares) of the lattice, in the latter on the N $_v$ vertices. The partition function for a xed lattice G $_N$ is given by

$$Z_{at}(G_{N}; ; H) = \begin{cases} X & P & P \\ e & {}^{hi;ji & i & j^{+}H} & {}^{i & i}; \\ fg & fg \end{cases}$$
 (2.2)

and the partition function Z (N; ;H) for the model coupled to quantum gravity is obtained by summing Z $_{at}$ (G $_{N}$; ;H) over all lattices G $_{N}$ with spherical topology and N faces

$$Z(N; ; H) = \sum_{fG_N g}^{X} Z_{at}(G_N; ; H):$$
 (2.3)

In eq. (2.2) $\int_{hi;ji}^{P} denotes a sum over the neighbouring pairs of faces (in case of a square lattice) or vertices (in case of a triangular lattice) of <math>G_N$. $\int_{fg} denotes the sum pover all the possible spin con gurations, is the inverse temperature and H the magnetic eld. <math>\int_{G_N g} denotes the discrete analogue of sum m ing over all possible metrics up to di eom orphism s on a spherical surface of xed volume, since <math>G_N$ de nes a metric on the discretized surface if we de ne each link to have the sam e xed length. The class of lattices we consider include degenerate ones which have double links and vertices of order one.

The system (2.3) undergoes a third order phase transition whose universal properties are independent of the m icroscopic details of the lattice [10, 11]. For the square lattice, the critical inverse temperature is given by $_{c} = \ln 2$ 0:69314718, and for the triangular lattice, $_{c} = \frac{1}{2}\ln \frac{13+\frac{p}{7}}{14-\frac{p}{7}}$ 0:16030370 [14]. The critical exponents are given by [11]

$$=$$
 1=2 (2.4)

= 2 (2.5)

$$d_{\rm H} = 3;$$
 (2.7)

which are dened in the usual way by the behaviour of the magnetization M j $_{\rm c}$ j, M H $^{1=}$ and the magnetic susceptibility M j $_{\rm c}$ j in the critical region. is the spin (spin correlation length exponent j $_{\rm c}$ j, and d_H denotes the Hausdor dimension of space.

For a given lattice G_N , the partition function (2.2) can be written (up to a multiplicative constant) as a polynom ial

$$Z_{at}(G_{N}; ; H) = \sum_{m=0}^{X^{N}} C_{mn} y^{m} c^{n}; \qquad (2.8)$$

where $c = e^2$, $y = e^{2H}$, N is the number of faces in G_N (when we put the spins on the faces) and N₁ the number of links. y is called the fugacity. In the following we will also use the notation $u = e^4$ and K = u u_c .

Eq. (2.8) is a polynom ial in two variables, and its analytic structure is entirely determ ined by its zeros in the complex plane. If we x the temperature , Z _{at} (G_N; ;H) is a polynom ial in y and its zeros were shown by Y ang and Lee [1] to lie on the unit circle in the complex fugacity plane. We expect the theorem to be true for a large class of (not too pathological) lattices G_N. Then one can write the free energy of the system as

$$F_{at}(G_{N}; ; y) = \ln (y - y_{k}()); \qquad (2.9)$$

where a factor has been absorbed into F $_{at}$ (G $_{N}$; ;y) and an additive constant has been suppressed (y e^{2H}). Here y_k () are the zeros of Z $_{at}$ (G $_{N}$; ;H) in the y plane, called the Lee {Y ang zeros. In the therm odynamic limit these zeros form dense sets on lines, which are Stokes lines for F $_{at}$ (G $_{1}$; ;y). All the relevant inform ation is then encoded in the density of zeros $_{YL}$ (;) (H = i). The free energy per spin in the limit of in nite volume N ! 1 is then given by

$$F_{at}(G_{1}; ;y) = d_{YL}(;) \ln(y e^{2i}):$$
 (2.10)

Here 2 denotes the polar angle on the unit circle of the partition function zero in the complex y-plane. This notation is motivated by the Lee{Y ang theorem. The singularities of the partition function are expected to occur at the ends of the lines on which the partition function zeros condense. In the case of (2.2), in the high temperature phase the zeros form a gap $_0 > 0$ such that $_{YL}(;) = 0$ for j j < $_0$. The points $y_0 = e^{2i_0}$ are the Lee{Yang edge singularities and they can be regarded as conventional critical points [7]. The (only) characteristic critical exponent is

$$Y_{L}(;) (0); (2.11)$$

which implies that M $(_0)$. In [26] it was shown that = 1=6 in two dimensions.

As we approach the critical temperature from above, the gap closes ($_0$! 0) and the zeros pinch the real axis at y = 1. This signals the onset of the phase transition, since the partition function has di erent analytic behaviour in the two disconnected parts of the y-plane. The information on the universal scaling of the partition function is given by the scaling of $_{\rm Y\,L}$ (;) near = 0. At = $_{\rm c}$, $_0$ = 0 and if we de ne M H $^{1-}$ we obtain

$$Y_{L}(c;)^{1=}:$$
 (2.12)

O ther critical exponents can be derived from scaling relations discussed in [6] and later in this section. For a nite system the zeros are never on the positive real axis since C_{mn} is positive and c > 0 for real in eq. (2.8). When the system is critical, i.e. when the correlation length L (L is the linear size of the system), the zeros approach the real axis in nitely close as L ! 1 and one can apply nite size scaling in order to extract critical exponents.

If we x y in eq. (2.8), we obtain a polynom ial of order N₁ in c. Its zeros in the c-plane are the so-called Fisher zeros [2]. For many systems the Fisher zeros are located on curves C, but this is not necessarily true in general. For the 2d (zero-eld) Ising model on a square lattice, these curves C are two intersecting circles (both in the c-plane and in the tanh () plane). One of the curves intersects the real positive axis at the physical critical point. Sim ilarly to the case of Lee{Yang zeros, the Fisher zeros condense on these two curves in the therm odynam ic lim it and the free energy per spin is given, in this lim it, by their density $_{\rm F}$ (; H):

$$F_{at}(G_{1}; ; H) = \int_{C}^{C} d_{F}^{0}(; H) \ln(c()) e^{2^{0}}; (2.13)$$

where 0 is a complex number. Near the critical tem perature and at H = 0,

$$_{\rm F}$$
 (;H = 0) j $_{\rm c}$ j⁺¹; (2.14)

which im plies that F at (G1; ; H = 0) j cj⁺². Therefore is the speci c heat exponent. The possibility of extracting scaling exponents from the study of com plex zeros of the partition function relies on their scaling behaviour under the renorm alization group [6]. Applying nite size scaling is a convenient m ethod often used in num erical simulations. Sim ple scaling argum ents [6] give the position of the jth Lee{Yang zero as

$$H_{j}^{2}N^{2} = (d_{H}) = f_{j}(K N^{1} (d_{H})) :$$
(2.15)

where we have substituted N $^{1=d_{H}}$ for the linear size L of the system (N is the volum e and d_{H} is the Hausdor dimension). f_{j} is an analytic function and the Lee(Y ang theorem in plies that $f_{j}(0) < 0$. If we invert the above relation we obtain the positions of the F isher zeros. Hence we can deduce that the jth Lee(Y ang zero will scale as

$$H_{i} N = (d_{H});$$
 (2.16)

and the jth Fisher zero as

$$K_{i} = N^{1=(d_{H})}$$
: (2.17)

The gap of the Lee{Y and edge singularity will scale as

$$H_0^2$$
 CK²; (2.18)

where C > 0 and H_0 i₀. Conversely, in the scaling region the dependence of the jth F isher zero on the value of the (real) m agnetic eld will be

$$K_{j} e^{\frac{i}{2}} \frac{H}{P} = \frac{1}{C}$$
 (2.19)

Eq. (2.19) implies that the trajectories of the motion of the Fisher zeros in the K plane with varying real nonzero H will form an angle

$$=\frac{1}{2}$$
 (2.20)

with the real K axis. Using eq. (2.15) a stronger scaling relation can be derived:

$$H_{j}^{2} (N = j)^{2 = (+1)} = F K (N = j)^{1 = (d_{H})};$$
 (2.21)

where F is a universal analytic scaling function such that F (0) < 0, H _j is the jth Lee{Yang zero. Similarly

$$K_{j} = \frac{j}{N} F^{1}(0); \qquad (2.22)$$

where K_{i} is the jth Fisher zero.

In [13], we provided evidence that there exists a divergent correlation length associated to matter elds coupled to gravity in the range of central charge 0 < c < 1. Based on these results, we expected that the observed zeros, for large enough lattices, m ight show a behaviour com patible with the scaling hypothesis. It is the main purpose of this work to provide evidence that this is indeed the case.

3 Computation of Complex Zeros

Our computation of Lee{Yang zeros (real , complex H) and Fisher zeros (real H, complex) of the partition function Z (N; ; H) was done using exact results from m atrix m odels and M onte C arlo multihistogram m ing techniques. We describe these m ethods in this section.

3.1 Exact determ ination of partition functions

The exact solution of the Ising model on a square dynam ical lattice came from solving the planar lim it of the two matrix model de ned in [10, 11]. It was noticed that in this lim it the free energy of the model

$$F(n;g; ;H) = \ln d^{n^{2}} d^{n^{2}}$$

where $_{1;2}$ are n n herm itean matrices, equals the grand canonical partition function Z (g; ;H) for the Ising model coupled to gravity:

$$\lim_{n! = 1} \frac{1}{n^2} F(n;g; ;H) = Z(g; ;H) \qquad \overset{X^{1}}{\underset{N=1}{\overset{P}{\longrightarrow}}} e^{N} Z(N; ;H); \qquad (3.24)$$

where $c = 4gc = (1 c^2)^2$. The solution for Z (g; ;H) is given by [10, 11]

$$Z (g; ;H) = \frac{1}{2} \ln \frac{z(g)}{g} + \frac{1}{2g^2} \int_{0}^{Z (g)} \frac{dz^0}{z^0} g(z^0)^2 - \frac{1}{g} \int_{0}^{Z (g)} \frac{dz^0}{z^0} g(z^0); \qquad (3.25)$$

where

$$g(z) = \frac{1}{9}c^{2}z^{3} + \frac{1}{3}z \frac{1}{(1-z)^{2}} c^{2} + \frac{2z(\cosh H - 1)}{(1-z^{2})^{2}} :$$
(3.26)

U sing eq. (325) and eq. (326) we can expand Z (g; ;H) in powers of g and read o the coe cients Z (N; ;H) (corresponding to lattice size with N squares) in eq. (324).

We have done this using the symbolic manipulation program M athematica for N 14. We obtain

$$Z(N; ; H) = C^{N} y \frac{\frac{N}{2}}{2} D_{mn} y^{m} C^{n}; \qquad (3.27)$$

$$m = 0 n = 0$$

where [:] denotes integer part. The same calculation for N 6 was done in [12] and our results are in complete agreement. Then the roots of Z (N; ;H) for either xed or xed H were computed num erically. Note that, apart from a trivial multiplicative factor, the partition function is a polynom ial and has a nite number of zeros in the complex c and y planes.

As already discussed above, and in [12] for N 6, the Lee{Yang zeros can be seen, som ew hat surprisingly, to lie on the unit circle in the com plex y-plane. No Lee{Yang theorem has so far been proven for the partition function of the dynam ical lattice. Fig. 1 shows the results for lattice sizes 8 N 14, at the (bulk) critical temperature. This gure clearly shows that the zero closest to the real positive axis approaches the point y = 1 as we increase the lattice size, indicating a vanishing gap at $_{\rm c}$ as N ! 1. We will study this approach quantitatively using nite size scaling in the next section.

In Fig. 2 we show the motion of Lee{Yang zeros with varying for the N = 14 lattice as we approach $_{\rm c}$ from the hot phase. The zeros close up towards the point y = 1 (H = 0) on the real axis, rejected vanishing of the gap in $_{\rm YL}$ (;) for $!_{\rm c}$ in the in nite volume limit.

We note from Fig. 3 and Fig. 4 that also the Fisher zeros form curves in the complex plane. In Fig. 3 (a) we display the approach of the Fisher zeros for H = 0 to the (physical and unphysical) critical points on the realaxis, c = 1=4 for increasing lattice size. There are also Fisher zeros on the in aginary axis, which ow to c = i1. In Fig. 3 (b) we show the same curves m apped onto the cplane, where the tilde refers to the dual spin m odel. The usual duality relation is given by $e = \tanh()$. We note that our model is not self(dual, so the critical point $c_c = 1=4$ is di erent from the dual critical point $e_c = 3=5$. Fig. 3 (b) suggests the existence of an antiferrom agnetic phase transition for the Ising m odel on a square lattice with spins placed on the vertices [14]. It happens at $e_c^{(af)} = 5=3$. This corresponds to the unphysical value of c = 1=4, re exting the fact that no antiferrom agnetic transition occurs for the Ising m odel coupled to 2d gravity on a square lattice with spins placed on the faces. We notice that the zeros in the equal are m apped exactly onto each other under the transform ation e ! 1=e. This in plies that the critical exponents of the antiferrom agnetic transition are identical to the ones of the farm agnetic transition, as was noted in [14].

Fig. 4(a) shows the ow of F isher zeros with varying magnetic eld H for a xed lattice size N = 14. The zeros on the arcs ow away from the imaginary axis as H increases, while the zeros on the imaginary axis again appear to move towards c = i1 (note that these points are mapped onto c = (1;0) in Fig. 4(c)). For H = 0 the zeros should pinch the real axis like in Fig. 3, but for $H \notin 0$ they seem to avoid the real axis. This is compatible with the absence of a phase transition in the presence of a symmetry breaking eld. In Fig. 4(c) we show the same ow in the c-plane.

3.2 Multihistogramming of Monte Carlo data

For system s with more than a few spins, the study of complex zeros must rely on numerical methods. Monte C arlo methods are very powerful in studying a system using the fullham iltonian. They can be used to study singularities of observables in the complex plane as was done in the pioneering work ref. [16]. With the development of (single) histogram ming techniques the partition function could be calculated for a continuous region in the coupling space and its zeros determined in the complex plane [17]. U sing multihistogram ming [20], where M onte C arlo data taken at di erent couplings are combined optim ally, a more accurate determination of the partition function is possible over a wide range of couplings. This method for computing complex zeros was rst used in [21]. Here we provide a brief description of the method, in order to describe the procedure we followed. For details see [32].

The M onte C arlo updating of the triangulations was performed by the so (called ip algorithm and the spins were updated using a standard cluster algorithm. One updating sweep of the lattice consisted of approximately N_1 accepted ips where N_1 is the number of links of the triangulated surface. A firer a sweep of the lattice we updated the spin system. All this is by now standard and we refer to [18, 19] for details. We use the high quality random number generator RANLUX [34] whose excellent statistical properties are due to its close relation to the Kolm ogorov K (system originally proposed by Savvidy et al. [33].

The lattice sizes that we simulated ranged from 32 to 256 vertices. In order to minimize nite size e ects, we took the spins to be on the vertices of the triangulations and included degenerate triangulations with double links and vertices of order one [19]. First we made a rough map of the partition function. Since the partition function is invariant under ! + ik = 2 and H ! H + ik = 2 it is not necessary to calculate $j_{2N} \cdot j_{1}^{2}$ for Im () > =2 or Im (H) > =2. In our calculation we took H to be purely in aginary. Then we scanned the region where we expected to see the rst zeros, and ultimately we took a denser grid of measurements around the points where the partition function touched zero within errors. For the largest lattice, 55 million sw eeps were performed at each of 11 values k around the location of the zeros, in order to get a su ciently accurate determination of the partition function in this region. For each k we computed energy and magnetization histogram sh^k (E; M), where E is the total energy and M the (absolute value of the) magnetization, from which one can obtain an approximate estimate of the density of states: <math>X

$$(E;M) = {X \atop k} w_{k}(E) _{k}(E;M) = {X \atop k} w_{k}(E)h^{k}(E;M)e^{kE} : (3.28)$$

The coe cients w $_{\rm k}$ (E) are the appropriate weights for an optim all determination of (E;M)

$$w_{k} (E; M) = \frac{1}{\frac{n^{1}}{n^{k}} + \frac{k}{1}} \exp((k + F_{1})) + F_{1}};$$
(3.29)

where $n^{k;l}$ is the number of measurements at $_{k;l}$, and $_{k;l}$ the respective autocorrelation times. F₁ is the free energy at $_{l}$, determined self{consistently from

$$e^{F_{1}} Z(_{1}) = \frac{X \qquad h^{k}(E;M)}{\sum_{E \in M, j^{k}} \frac{n^{j}}{n^{k}} \sum_{j=1}^{k} \exp((_{1} _{j})E + F_{j})} :$$
(3.30)

Then the partition function Z () and any observable hO (E; M) i for real is given by:

$$Z() = {}^{X} (E;M) e^{E}$$
 (3.31)

hO (E; M) i =
$$\frac{1}{Z()} X O (E; M) (E; M) e^{E}$$
: (3.32)

Lee{Yang zeros are computed from the minim a of

$$\frac{Z(;H)}{Z()} = h\cos(\text{Im}(H)M)i \text{ ihsin}(\text{Im}(H)M)i \qquad (3.33)$$

for real and imaginary H and Z () $\,$ Z (;0). The Fisher zeros were computed from the minima of

$$\frac{Z()}{Z(Re)} = h\cos(\operatorname{Im}()E)i \quad \text{ihsin}(\operatorname{Im}()E)i \quad (3.34)$$

for complex values of A = 0. The errors in the partition function and in the position of the complex zeros were computed by a standard binning procedure.

Table 1 and Table 2 contain our results for the zeros. We rst checked that multihistogram – ming was working quite well with our data by looking at multihistogram s of the speci c heat and susceptibility. It was relatively easy to observe the rst few Lee{Yang zeros. For F isher zeros the computation was harder and in a similar way we could only clearly observe the zero closest to the real axis. D elicate cancellations in $\frac{1}{2}$ $\frac{2}{3}$ between contributions from the two terms in eq. (3.34) make the zeros located further away from the real axis disappear in the statistical noise. This sets limitations to the size of the surface on which we were able to observe F isher zeros.

N _v			Im (H)	Im (H)
64	0.1718	0.0005	0.203	0.003
96	0.1666	0.0004	0.1484	0.0002
	0.2114	0.0004	0,2454	0.0004
128	0.1643	0.0014	0.1182	0.0005
	0.2037	0.0012	0.195	0.005
	0,2386	0.0018	0,260	0.003
256	0.1596	0.0010	0.069	0.003
	0.1850	0.0017	0.112	0.004
	0,2069	0.0009	0.148	0.005
	0,2262	0.0009	0.185	0.001
	0.2497	0.0034	0.175	0.005

Table 1. Læ{Yang zeros observed using multihistogram ming.

N _v	Re()	Re()	Im ()	Im ()
32	0.1428	0.0016	0.1773	0.0005
64	0.1481	0.0049	0.1348	0.0022
96	0.1533	0.0021	0.1121	0.0034
128	0.1556	0.0300	0.1030	0.0050
256	0.1527	0.0035	0.0788	0.0047

Table 2. Fisher zeros observed using multihistogram ming.

4 Veri cation of scaling relations

In this section we discuss the extent to which the scaling relations eqs. (2.15{2.22) hold for the Ising model on a dynamical lattice. There is no a priori reason to expect these relations to hold. As discussed earlier, for the xed lattice, their validity is due to the divergence of the spin {spin correlation length in the critical region. For the Ising model on a dynamical lattice, we expect the correlation length associated to geometry uctuations to diverge [25, 23, 24], but the same need not necessarily be true for the spin {spin correlation length even at a continuous transition [15]. Recently how ever, num erical evidence indicated that this is indeed the case, and that the system behaves as an ordinary statistical system near a third order phase transition [13].

Num erical sinulations [23, 24, 13] further indicate that the Hausdor dimension of the system is very close to 4. From this value we can estimate the linear size L N^{1=d_H} of the system s that we are studying and nd that it is indeed quite small. In spite of substantial nite size e ects, our results will provide evidence that the scaling relations are indeed satis ed by the motion of the complex zeros with varying lattice size, couplings, or the order of the zero. We can, in general, observe reasonable agreement with the known critical exponents (we will refer to these as the

KPZ exponents). In the cases where a deviation from the expected value is observed, we can observe an asymptotic approach to this value with increasing lattice size.

4.1 Zeros of the exact partition functions for sm all square lattices

At K = 0 and large enough N eq. (2.16) in plies

$$\ln j_{\rm H_j j=} - \frac{1}{d_{\rm H_j}} \ln (N_{\rm J}) + C_{\rm j}:$$
 (4.35)

The slope on a log(log plot of H j vs. N is expected to take the value

$$-\frac{1}{d_{\rm H}} = 5 = 6 = 0$$
:8333:::: (4.36)

for all j. Fig. 5 and Table 3 show our results for the stathaet Lee{Yang zeros. The extracted exponent combination is in reasonable agreement with the KPZ exponents, especially for the state. The errors reported in Table 2 are not true statistical errors (which are meaningless in this case). They are computed from the standard formula giving least squares linear terrors and in this case they are simply a measure of the systematic deviation of the points from a straight line. We will follow this practice for our ts throughout this section.

j	Slope		
1	-0.871	0.002	
2	-0.935	0.002	
3	-0.951	0.002	

Table 3. Predictions for the combination of critical exponents $\frac{1}{d_H}$ from scaling of the rst three Lee{Yang zeros of exactly known partition functions Z_N .

Eq. (2.22) im plies

$$h_{j}K_{j}j = h_{j}u_{c}j = \frac{1}{d_{H}}h_{N} + C :$$
 (4.37)

Similarly, since χ_{jj}/j_{j} cj+ 0 (j j c²), we would expect that

$$\ln j_{j} c_{j} = \frac{1}{d_{H}} \ln \frac{j}{N} + C; \qquad (4.38)$$

where the slope should be given by

$$\frac{1}{d_{\rm H}} = \frac{1}{3}$$
; (4.39)

and the constant C should be independent of j. Table 4 shows the results of ts to eq. (4.38), and similar ts with N replaced by N_v, the number of vertices of the lattice. The corresponding ts using the scaling variable K did not yield straight lines in a log(log plot, as can be seen from Fig. 6. In this gure, we have plotted K_1 j (lower curve) resp. j₁ _____c j (upper curve) vs. N on a logarithm ic scale. The di erence between the scaling behaviour of the two di erent scaling variables (which are expected to be identical for very large system s), shows that we are not deep in the scaling region and that nite size e ects are important for the system sizes that we consider. The same must be said about the di erence between the slopes and intercepts C of Table 4. From eq. (4.38) it is expected that the slopes and C should take the same value for all j, so that the data points would fall on one and the same curve. The di erence between the results obtained by using N and N_v should asymptotically decrease as the lattice size goes to in nity. (In the table, n denotes the number of points included in the t.)

Quantity tted		j	Slope	С	n
lnj	cjvs. ln(j=N)	1	0.327 0.001	0.295 0.002	8
		2	0.377 0.002	0.490 0.005	4
lnj	cjvs.ln(j=Nv)	1	0.383 0.002	0.492 0.006	6
		2	0.438 0.004	0.667 0.009	4

Table 4. Slopes and intercepts from the ts to eq. (4.38). The theoretical value for the slope is $\frac{1}{d_{H}} = 1=3$. The dependence of the slope and of C on j indicates nite size e ects, as does the di erence between N and N_v. n is the number of degrees of freedom in the ts.

The scaling relation eq. (2.19) predicts the angle that the trajectories of the zeros will form with respect to the Re(K) axis as H is varied for large L. In our case the angle is expected to be

$$=\frac{1}{2}=36$$
: (4.40)

This prediction is valid for large system sizes L and small magnetic eld. In Fig. 7(a) we show the trajectories of the rst Fisher zero for N = 4; 6; 8; :::; 14 and magnetic elds in the interval 0.2 < H < 0.2. As expected from eq. (2.19), H = 0 corresponds to the \tuming point" closest to the Re(K) axis. Fig. 7(b) and Table 5 show the corresponding values of tan , calculated from ts in the small H range 0.03 < H < 0.02. A though does not reach its in nite volum e value, Fig. 7(b) shows the approach to it with increasing lattice size. The breakdown of scaling for very strong H can be observed in Fig. 7(c) where a trajectory for H jbetween zero and 2.6 are shown.

Ν	tan	tan	
6	6.489	0.162	
8	3,295	0.036	
10	2.308	0.001	
12	1.792	0.006	
14	1.450	0.002	
1	0.727		

Table 5. Measured (N = 6; 8; :::; 14) and expected (eq. (4.40)) slopes of trajectories of zeros in the complex K plane as the magnetic eld H is varied, as a function of the size of the lattice.

We also checked the validity of the scaling relation given by eq. (2.21). Taking K real, we plotted H_j(N=j)⁵⁼⁶ vs. ($_{c}$)(N=j)¹⁼³. The scaling function F is shown in Fig. 8. All zeros of order j > 1 for lattice sizes (10 N 14) are included in the same graph. We observe that there exists a range of in the hot phase where the data points lie approximately on a univeral curve de ning the function F.

By varying the value of the exponent =(+1) = 5=6 of the scaling variable = j=N on the y-axis away from its KPZ value, we observed a broadening of the curve and in this way we obtained a (som ewhat subjective) determination of the exponent combinations

(the equality of these follow from the general exponent equalities and was used in the derivation of the scaling relation (2.21)). This is in excellent agreem ent with the KPZ value of 5=6.

4.2 Zeros determ ined from multihistogramming

In this section we describe the scaling of the zeros determ ined numerically from M onte C arlo data. The lattice sizes investigated here range from 32 to 256 vertices (60{508 triangles}). We were able to determ ine the rst ve Lee{Yang zeros for the biggest lattice size, while for the sm aller ones at m ost three zeros were visible. For the F isher zeros, only the zero closest to the real axis was observed for any lattice size.

The Lee{Yang zeros we observed with multihistogram m ing are listed in Table 1. The least-squares ts to eq. (4.35) give for the rst and second Lee{Yang zero,

$$d_{\rm H} = 0.773 \quad 0.013 \quad (j = 1)$$
 (4.42)

$$d_{\rm H} = 0.788 \quad 0.033 \quad (j = 2):$$
 (4.43)

All data points with j 2 shown in Table 1 were included in the ts. U sing N $_{\rm v}$ instead of N for the volume of the system we obtain:

$$d_{\rm H} = 0.787 \quad 0.013 \quad (j = 1)$$
 (4.44)

$$-\frac{1}{d_H} = 0.800 \quad 0.034 \quad (j = 2);$$
 (4.45)

which gives a measure of the presence of nite size e ects. The results are in quite good agreement with the expected value of = d = 5=6. From the exactly known value $d_{\rm H} = 3$ we obtain

$$= 2:36 \quad 0:04 \quad (j = 1)$$
 (4.46)

$$= 2:40 \quad 0:10 \quad (j = 2) :$$
 (4.47)

The exact value of this combination of exponents is 2.5.

$$\frac{1}{d_{\rm H}} = 0.350 \quad 0.067: \tag{4.48}$$

Quantity tted		$1=(d_{\rm H})$	
ln∱ jvs.ln(N)		0.392	0.016
lnjK jvs. ln (N _v)		0.407	0.017
lnj	_c jvs. ln(N)	0.386	0.014
lnj	_c jvs. ln (N _v)	0.401	0.014

Table 6. Finite size scaling for the rst Fisher zero.

5 Conclusion

We have computed the positions of the singularities of the partition function for the two{ dimensional Ising model coupled to gravity in the complex plane. Both Lee{Yang and Fisher zeros were studied.

We veried the result in [12] that the Lee{Yang zeros are located on the unit circle of the complex fugacity plane. This presents us with the challenge of proving a corresponding Lee{ Yang theorem for the case when a uctuating metric contributes an additional quantum degree of freedom. We also observed that the Fisher zeros form one dimensional curves in the complex tem perature plane.

Given the small size of the system swe studied, we obtained reasonable agreement with scaling laws derived from ordinary renormalization group arguments. Although one cannot use our results to accurately determine the scaling exponents, the extracted exponent combinations show a reasonable agreement with their exactly known values [11]. This, together with the fact that a diverging matter correlation length exists in the system [13], gives us con dence to conclude that the critical behaviour of the system is given by the scaling of the distribution of the complex zeros of the partition function.

References

C.N.Yang and T.D.L@Phys.Rev.87 (1952) 404;
 T.D.L@ and C.N.Yang Phys.Rev.87 (1952) 410

- [2] M.Fisher, in Lectures in Theoretical Physics VIIC (University of Colorado Press, Boulder, 1965)
- [3] V.M atveev and R.Shrock, Phys. Rev. E 53 (1996) 254;
 V.M atveev and R.Shrock, Phys. Lett. A 215 (1996) 271
- [4] R.Abe, Prog. Theor. Phys. 37 (1967) 1070;
 R.Abe, Prog. Theor. Phys. 38 (1967) 72;
 R.Abe, Prog. Theor. Phys. 38 (1967) 568
- [5] S.Ono, Y.Karaki, M. Suzukiand C.Kawabata, J.Phys. Soc. Japan 25 (1968) 54
- [6] C. Itzykson, R.B. Pearson and J.B. Zuber, Nucl. Phys. B 220 FS8] (1983) 415
- [7] M.E.Fisher Phys. Rev. Lett. 40 (1978)1610;
 P.J.Kortm an and R.B.Gri ths, Phys. Rev. Lett. 27 (1971) 1439
- [8] P.H.Dam gaard and U.Heller, Nucl. Phys. B 410 (1993) 494
- [9] V.Knizhnik, A.Polyakov and A.Zamolodchikov, Mod. Phys. Lett. A 3 (1988) 819
- [10] V.A.Kazakov, Phys. Lett. A 119 (1986) 140
- [11] D.V.Boulatov and V.A.Kazakov, Phys. Lett. B 186 (1987) 379
- [12] M . Staudacher, Nucl. Phys. B 336 (1990) 349
- [13] J. Ambjm, K. N. Anagnostopoulos, U. Magnea and G. Thorleifsson, Phys. Lett. B 388 (1996) 713;
 J. Ambjm and K. N. Anagnostopoulos, \Quantum Geometry of 2d Gravity Coupled to Unitary Matter", NBIHE-96-69 (hep{lat/9701006)
- [14] D.A.Johnston, Phys.Lett. B 314 (1993) 69;
 C.F.Baillie and D.A.Johnston, Phys.Lett. B 357 (1995) 287
- [15] M.G.Harris and J.Ambj m, NuclPhys.B474 (1996) 575
- [16] M. Falcioni, E. Marinari, M. L. Paciello, G. Parisi and B. Taglienti, Phys. Lett. B 102 (1981) 270
- [17] M.Falcioni, E.Marinari, M.L.Paciello, G.Parisi and B.Taglienti, Phys.Lett.B108 (1982)
 331;
 E.Marinari Nucl. Phys. B 235 [FS11] (1984) 123
- [18] J. Jurkiew icz, A. Krzywicki, B. Petersson and B. Soderberg, PhysLett. B 213 (1988) 511;
 C.F. Baillie and D.A. Johnston, PhysLett. B 286 (1992) 44;
 S. Catterall, J. Kogut and R. Renken, PhysLett. B 292 (1992) 277;
 J. Ambjm, B. Durhuus, T. Jonsson and G. Thorleifsson, Nucl. Phys. B 398 (1993) 568

- [19] J.Ambjm, G.Thorleifsson and M.Wexler, NuclPhys. B 439 (1995) 187
- [20] A.M. Ferrenberg and R.H. Swendsen Phys. Rev. Lett. 61 (1988) 2635
- [21] N.A.A lves, B.A.Berg and R.V illanova Phys.Rev.B 41 (1990) 383; Phys.Rev.B 43 (1991) 5846
- [22] K.N.Anagnostopoulos, M.J.Bowick and E.Marinari, unpublished.
- [23] S.Catterall, G.Thorleifsson, M.Bowick and V.John, Phys. Lett. B 354 (1995) 58
- [24] J.Ambj m, J.Jurkiew icz and Y.W atabiki, NuclPhys. B 454 (1995) 313
- [25] J.Ambjm and Y.W atabiki, NuclPhys.B 445 (1995) 129
- [26] D.Dhar Phys. Rev. Lett. 51 (1983) 853;
 J.L.Cardy Phys. Rev. Lett. 54 (1985) 1354
- [27] B.Derrida, L.De Seze and C. Itzykson J. Stat. Phys. 33 (1983) 559
- [28] W .van Saarlos and D .K urtze J.Phys. A 17 (1984) 1301;
 J.Stephenson and R.Couzens Physica 129A (1984) 201
- [29] V.Matveev and R.Shrock, J.Phys. A 28 (1995) 5235
- [30] P.H.Dam gaard and J.Lacki, Int. J.M od. Phys. C 6 (1995) 819
- [31] M. Baake, U. Grimm and C. Pisani, J. Stat. Phys. 78 (1995) 285
- [32] S.Huang, K.J.M.Moriarty, E.Myers and J.Potvin Z.Phys.C 50 (1991) 221
- [33] G.K.Savvidy and N.G.Ter-A nutyunyan Savvidy, EPI-865-16-86 (1986); J.Com put. Phys. 97 (1991) 566;
 N.Z.Akopov, G.K.Savvidy and N.G.Ter-A nutyunyan Savvidy, J.Com put. Phys. 97 (1991) 573
- [34] M. Luscher, Comput. Phys. Commun. 79 (1994) 100;
 F. James, Comput. Phys. Commun. 79 (1994) 111; Erratum 97 (1996) 357



Figure 1: Lee{Yang zeros in the complex fugacity ($y = e^{2H}$) plane for the Ising model on square dynam ical lattices of varying size N = 8(+), 9(), 10(), 11(2), 12**\square**), 13(), 14() at the bulk critical tem perature.



Figure 2: Lee{Yang zeros in the complex fugacity plane for a square lattice with N = 14 at the critical point ($c_c = e^{2} = 0.25(+)$) and in the hot phase (c = 0.35(), 0.45(), 0.55(2), 0.65()).



Figure 3: (a) Fisher zeros in the complex c plane ($c = e^2$) for the Ising model with spins on the faces of square dynam ical lattices of varying size (N = 8; 10; 12; 14) at H = 0. The zeros m ove on arcs and on the imaginary axis. The zeros ow towards if on the Im (c) axis, and towards the critical points c = (1=4;0) and c = (1=4;0) on the real axis as the therm odynam ic lim it is approached. The physical critical point is c = 1=4. (b) The trajectories in (a) show n in the complex tanh () plane. Recalling the duality relation $e = \tanh()$, this corresponds to Fig. 3(a) for the model with spins on the vertices (with e related to the dual inverse temperature \sim by $e = e^2$). The zeros approach the points (1;0), (3=5;0) and (5=3;0) as N ! 1. The point e = 3=5 corresponds to the ferrom agnetic and the point e = 5=3 to the antiferrom agnetic transition of the model. The zeros are mapped onto each other under e ! 1=e:



Figure 4: (a) Fisher zeros in the complex c plane for a square dynamical lattice of size N = 14 and at varying magnetic eld ($y = e^{2H} = 1.0$; 1:6; 2:5). For zero magnetic eld, the zeros with Re(c) > 0 end in the vicinity of the physical critical point c = (1=4;0). (b) The rst quadrant in (a) show n magni ed for several values of the fugacity y = 1.0; 1:6; 2:5; 5:0. (c) The trajectories in (a) show n in the complex c-plane for zero and nonzero magnetic eld. The fugacity takes the values y = 1:0; 1:3; 1:6; 1:9; 2:5 and N = 14.



Figure 5: Scaling of the rst three Lee{Y ang zeros for the Ising m odelon square dynam ical lattices of size N (j labels the zero). The slopes are expected to be given by the exponent combination $=(q_1) = 5=6$ (d_H is the Hausdor dimension). The tted slopes are given in Table 3.



Figure 6: Illustration of the scaling relation $jK j / N^{1=(d_H)}$ for the rst Fisher zero at H = 0. Due to nite size e ects, the curve for jK j is not a straight line. On the other hand, j cj scales well and yields a slope of 0:327 0:01. The expected slope is 1=3.



Figure 7: (a) Trajectories of the rstFisher zero in the complex K plane for nonzero m agnetic eld 0.2 < Im(H) < 0.2. The trajectories are for small square lattices of size N = 4; 6; 8; 10; 12; 14 in order from left to right. The angle that the trajectories form with the Re(K) axis is expected to approach = = (2) = 36 as N ! 1 . (b) tan versus N for small lattices. The dashed line corresponds to the expected asymptotic value of tan for large N . (c) B reakdown of scaling for strong m agnetic eld (up to H j 2:6). The angle changes drastically for strong H .



Figure 8: Check of the scaling relation for the jth Lee{Yang zero H $_j^2$ $^2 = (+1) = F$ (K $^{1=(d_H)}$), where = j=N and F is a universal function. All values of j > 1 for lattice sizes in the range 10 N 14 have been plotted on the same graph. The points are expected to fall on the same universal curve for large j.



Figure 9: The rst (j = 1) and second (j = 2) Lee{Yang zero observed using multihistogram ming on dynamical triangular lattices in the size range 64 N_v 256 (124 N 508). The slopes of the lines should be given by = (q_1) = 5=6.



Figure 10: The str Fisher zero observed using multihistogramming on dynamical triangular lattices in the size range 32 N_v 256 (60 N 508). The slopes of the lines are expected to be $1=(d_{\rm H}) = 1=3$.