E ects of CP-V iolation in Neutralino Scattering and Annihilation

Paolo Gondolo^a and Katherine Freese^{a;b}

^aM ax-P lanck-Institut fur P hysik, F ohringer R ing 6, 80805 M unchen, G erm any (gondolo@ m ppm u m pg de)

^bPhysics Department, University of Michigan, Ann Arbor, MI 48109, USA (ktfræsse@umichedu)

W e show that in som e regions of supersymm etric parameter space, CP violating e ects that m ix the CP-even and CP-odd Higgs bosons can enhance the neutralino annihilation rate, and hence the indirect detection rate of neutralino dark m atter, by factors of 10^6 . The same CP violating e ects can reduce the neutralino scattering rate o nucleons, and hence the direct detection rate of neutralino dark m atter, by factors of 10^{-7} . W e study the dependence of these e ects on the phase of the trilinear coupling A, and nd cases in the region being probed by dark m atter searches which are experimentally excluded when CP is conserved but are allowed when CP is violated.

The neutralino elastic scattering cross section (in pb) is plotted in g.1 as a function of neutralino m ass (in GeV) for 10⁶ values in SUSY param eter space. The upper panel is for the case of CP violation via Im (A) 6 0 while the lower panel is for the case of no CP violation. In the upper panel, it is the maxim ally enhanced cross section (as a function of arg(A)) that is plotted. The dark points refer to those values of param eter space which have the maximum value of the cross section for nonzero Im (A) and which are experim entally excluded at zero Im (A). The grey region refer to those values of parameter space which are enhanced when CP violation is included and which are allowed also at zero Im (A). The light grey empty squares refer to those values of param eter space which have no enhancem ent when CP violation is included. The solid lines indicate the current experim ental bounds placed by DAMA and CDMS; the dashed lines indicate the future reach of the CDMS (Soudan), GENIUS,



and CRESST proposals.

In g. 2 we show the enhancement and suppression factors of the elastic scattering cross section for the case of CP violating arg (A). The plot shows the ratios $R_{max} = {}^{max} = max[(0); ()] > 1$ and $R_{min} = {}^{min} = min[(0); ()]$ as a function of the values A of the phase of A where them aximum /m inim um occur H ere ${}^{max} ({}^{min})$ is the enhanced (suppressed) scattering cross section and the superscript max (min) indicates the maximal enhancement (suppression) as one goes through the phase of A. The denominator of the ratio $R_{max} (R_{min})$ chooses the larger (smaller) value of the scattering cross section without CP violation, i.e., for phase = 0 or phase = .

Fig. 3 shows the enhancement and suppression factors of the neutralino annihilation cross section times relative velocity v (at v = 0). The ratios $R_{m \, ax}^{ann}$ and $R_{m \, in}^{ann}$ are dened similarly to $R_{m \, ax}$ and



 $R_{\,\text{m in}}$ but with $\,\,v\,$ replacing the scattering cross section $\,$.

Finally, in gs. 4 and 5 we show two exam ples of the behavior of the scattering and annihilation cross sections with the phase of A. The four panels from top to bottom display the following: the scattering cross section p in pb, the annihilation cross section v in cm^3/s , the branching ratio BR (b! s) 104, and the lightest Higgs boson mass m_h, in GeV as a function of the phase A of A. CP conserving phases are A = 0; while all other values are CP violating. In the third and fourth panels we hatch the regions currently ruled out by accelerator experiments. In all four panels we denote the part of the curves that is experim entally allowed by thickened solid lines, and the part that is experimentally ruled out by thinner solid lines. In this gure, the possible phases are bound by the limit on the b! s branching ratio. In the allow ed regions, the scattering cross section at CPviolating phases is suppressed, while the annihilation cross section is enhanced. The latter takes its maximum allowed value when the b! s lim it is reached. In the case plotted in g. 4, both CP conserving cases are experim entally excluded while som e CP violating cases are allowed. The scattering cross section is of the order of 10^{-6} pb, and lies in the region being probed by direct detection experiments. The annihilation cross section peaks at A = 3 = 4; notice that this value is not the point of maxim alCP violation A = = 2.



A detailed presentation and references can be found in G ondolo and Freese, hep-ph/9908390.