Net-baryon scaling near the QCD critical point

N.G. Antoniou¹, F.K. Diakonos and A.S. Kapoyannis Department of Physics, University of Athens, GR-15771 Athens, Greece

A bstract

The net-baryon density at m idrapidity is proposed as an order parameter in the search for the QCD critical point in heavy ion collisions. As a function of the initial energy and the total number of participants, this quantity obeys a scaling law, dictated by the critical exponents of the appropriate universality class. The corresponding scaling variable speci es the proximity of a given experiment to the critical point. W ithin this fram ework, measurements at the SPS are discussed and predictions for RHIC and LHC are extracted.

¹em ailmantonio@ cc.uoa.gr

1 Introduction

A remarkable property of the QCD phase diagram (T) is the existence of an endpoint along the critical line of the st order quark-hadron phase transition [1]. It de nes a critical point of second order, belonging to the universality class of a 3d Ising system and located on a line of nonzero baryonic density = c. The QCD critical point is associated with the chiral phase transition in the sense that it is the rem nant of a tricritical point corresponding to the chiral lim it $m_u = m_d = 0$ [2]. In other words, the existence of a second-order critical point, at nonzero baryonic density, is a fundam ental property of realQCD with small but nonzero quark masses ($m_u; m_d \in 0$). The QCD critical point communicates with the hadronic world through the uctuations of a scalar eld (-eld) which carries the quantum numbers of an isoscalar (-m eson) as the manifestation of a quark condensate, hqqi, in them al environment. At the critical temperature (and in nite volume) the isoscalar has zero m ass, in order to provide the in nite wavelength m ode required by the divergence of the correlation length. It remains, therefore, stable in the therm al environm ent of a nuclear collision as long as the tem perature stays close to the critical value. At the freeze-out stage the mass of the $-\operatorname{eld} \mathfrak{m}$ ay reach the two-pion threshold (m $\stackrel{\scriptstyle >}{}$ 2m) and become accessible to observation [2, 3]. In the elective theory of the QCD critical point, the classical - eld is a natural order parameter, the uctuations of which obey scaling laws dictated by the critical exponents of the 3d Ising $\frac{2}{3}$). In a baryonic environment, howsystem (0; ±_; 5; ever, the chiral condensate is expected to have, at $T = T_c$, a strong dependence on the net-baryon density, driving the - eld close to zero for c : $-\frac{c}{c}$ hqqi₀ + 0 [(c)²] where is a dimensionless constant hqqi of the order of unity [4]. This dependence suggests a new order parameter, m =c, associated with the critical properties of the baryonic uid created in a quark-hadron phase transition. In fact, approaching the critical point in the phase diagram, both the -eld uctuations and the uctuations of the order parameter m (x) obey the same scaling laws (hqui m (x)).

In this work we exploit the scaling properties of the order parameter m(x), properly adjusted to measurable quantities, in heavy ion collisions. For this purpose we consider net-baryon production in collisions of heavy nuclei with total number of participants A_t and initial energy corresponding to a total size in rapidity y = L. The paper is organized as follows: In Section 2 we discuss in detail the scaling properties of the net-baryon uid

near the critical point. In Section 3 we present shortly phenom enological consequences of the presence of critical uctuations in the baryonic sector. Finally in Section 4 we present our conclusions concerning the possibility to approach and observe the QCD critical point in heavy-ion colliders.

2 Net-baryon scaling

The created baryons in the process of quark-hadron phase transition occupy a cylindrical volum e with transverse radius $R_2 = A_2^{1=3}$ and longitudinal size L (in rapidity). The parameter A₂ species the elective number of participants, contributing to the transverse geometry of the collision, and it $\frac{A_{t}}{2}$, valid both for central (A₂ = A_{m in}) and non-central is assumed $A_{?}$ collisions. Projecting out the net-baryon system onto the longitudinal direction we end up with a 1d liquid con ned in a nite rapidity region of size L with local density (y) = $\frac{n_{b}(y)}{R_{2}^{2} f}$, directly related to the measurable net-baryon density in rapidity $n_b(y) = \frac{dN_b}{dy}$. Putting $R_2 = R_0 A_2^{1=3}$, we introduce a characteristic volum e $V_o = R_o^2 f$ in terms of the freeze-out time 8 fm [5] and the nuclear-size scale R which contains also scale f 6 any growth e ects near the critical point (R $_{\circ}$ 12 fm).). Using V 1 as a scale for baryonic, freeze-out densities, the order parameter m (y) of the 1d baryonic liquid is written:

$$m(y) = A_{2}^{2=3}n_{b}(y)$$
 _c ; 0 y L (1)

In what follows, our basic assumption is that the decon-ned phase of quark matter, in heavy-ion collisions, approaches the QCD critical point in local thermal equilibrium. In the framework of inside-outside mechanism, this process is in plan ented by considering the isothermal space-time hyperbolas $t^2 = x_k^2 = t^2$ and employing the corresponding rapidity variable as the appropriate longitudinal coordinate. In this description, the assumed local equilibrium in the new geometry $(\mathbf{x}_k; \mathbf{x}_2; t)$ is translated as global equilibrium in the new geometry $(\mathbf{y}; \mathbf{x}_2; t)$ is translated as global and one may consistently impose, near criticality, static scaling laws on the order parameter m (y). In particular, the expected singularity at $T = T_c$ is incorporated in a general expansion of the form :

$$m(y) = t[F_{\circ}(y=L) + tF_{1}(y=L) + :::]$$
 (2)

where t $\frac{T_c - T}{T_c}$ and is the appropriate critical exponent $(\frac{1}{3})$. The leading term F_o (y=L) in the expansion (2) has a universal power-law behaviour near the walls (y = 0;L), shared by all uids belonging to the same universality class and con ned in a nite, one-dimensional region [6, 7]. At m idrapidity (y $\frac{L}{2}$), eq.(2) gives the deviation of the measurable, bulk density of net baryons, from the critical value $_c$, as we approach the critical point, not along the critical isochore (= $_c$) but along the freeze-out line:

$$A_{?}^{2=3}n_{b} = c + t_{f} [F_{o} + t_{f}F_{1} + :::]$$
(3)

where $t_f = \frac{T_c T_f}{T_c}$, $n_b = n_b$ (L=2), $F_i = F_i$ (1=2) and T_f is the freeze-out tem perature. Integrating now eq.(2) in the interval 0 y L we obtain at $T = T_f$:

$$A_{?}^{2=3}A_{t}L^{1} = _{c} + t_{f} (I_{o} + t_{f}I_{1} + :::)$$
 (4)

where $I_i = {R_1 \atop 0} F_i()d$. Introducing the variable $z_c = A_2^{2=3}A_tL^{-1}$ we nd from eqs.(3) and (4) a scaling law for the net-baryon density at m idrapidity:

$$A_{2}^{2=3}n_{b} = (z_{c}; c) ; z_{c} c$$
 (5)

where the scaling function $(z_c; c)$ has the property $(z_c = c; c) = c$. In the crossover regime $z_c < c$, where critical uctuations disappear, the local density at m idrapidity is, to a good approximation, $n_b = A_t L^{-1}$ suggesting a continuous extension of the scaling law (5) in this region with $(z_c; c) = z_c$ $(z_c < c)$. It is of interest to note that although the scaling function $(z_c; c)$ is continuous at the critical point $(z_c = c)$, the rst derivative is expected to be discontinuous at this point in accordance with the nature of the phase transition (critical point of second order).

In order to study in detail the critical behaviour of the baryonic uid in terms of the new variables n_b , z_c a further investigation of the structure of the scaling function (z_c ; $_c$) for z_c $_c$ is necessary. For this purpose we approximate the nearby part of the freeze-out line, close to the critical point (t_f 1), by truncating the series (2) keeping only the next to the leading term F_1 (y=L). As a result, the equations (3) and (4) are written correspondingly:

$$A_{2}^{2=3}n_{b}$$
 $_{c} + t_{f}(F_{o} + t_{f}F_{1})$ (6-a)

$$z_{c} \quad c \quad t_{f} (I_{o} + t_{f} I_{1})$$
 (6-b)

Neglecting terms of order O (t_f^2) in the power expansion of the quantity (I_o + $t_f I_1$)¹⁼ and combining eqs.(6) we nally obtain:

$$A_{2}^{2=3}n_{b} = c + \frac{F_{o}}{F_{1}}[f(z_{c}; c)] + C[f(z_{c}; c)]^{+1}$$
(7-a)

$$A_{2}^{2=3}n_{b} = _{c} + F_{o}t_{f} (1 + t_{f} \frac{C I_{o}^{1+-}}{F_{o}})$$
 (7-b)

$$f(z_c; _c) = \frac{1}{G} (1 + \frac{q}{1 + 2G} (z_c _c)^{1=}); z_c _c (7-c)$$

where $C = \frac{F_1}{I_0^{1+\frac{1}{L}}}$, $G = \frac{2I_1}{I_0^{1+\frac{1}{L}}}$. The component F_0 (y=L) which dominates the order parameter m (y) in $\frac{L}{2}$; L 1), far from the walls (at the points y = 0;L) due to the approximate translational invariance of the nite system in this region. On the other hand, approaching the walls, F_o (y=L) describes the density correlation with the endpoints and obeys a universal power law. In sum mary:

$$F_{o}()$$
 const: $(\frac{1}{2})$ (8-a)

$$F_{o}() = (0) (8-b)$$

 $F_{\circ}()$ (1) = (1) (8-c)

The solution which ful is these requirements is:

$$F_{\circ}() = g[(1)]^{=}(0 1)$$
 (9-a)

$$\frac{F_{o}}{I_{o}} = \frac{4^{=}}{B(1 - i)} ; B(n;m) = \frac{(m)(n)}{(m + n)}$$
(9-b)

Inserting the constant $\frac{F_{\circ}}{I_{\circ}}$ into the eq.(7-b), the nal form of the scaling function $(z_c; c)$ is obtained:

$$(z_{c}; c) = c + \frac{4^{=}}{B(1 - ; 1 -)} [f(z_{c}; c)] + C [f(z_{c}; c)]^{+1}; z_{c} c$$
(10-a)

$$(z_{c}; c) = z_{c}; z_{c} < c$$
 (10-b)

where $f(z_c; c)$ is given by eq.(7-c). It is straightforward to show that the discontinuity of the rst derivative of $(z_c; c)$ at $z_c = c$ is a nonzero universal

constant: disc $\frac{d}{dz_c}$ 1 $\frac{2}{c}$ (- $\frac{1}{2}$) as expected from the characteristic properties of a second-order phase transition. Combining eqs.(5-10) one m ay propose a fram ework for the treatm ent of certain phenom enological aspects of the QCD critical point. The scaling law (5) combined with eq.(10) involves three nonuniversal parameters: (a) the critical density $_{\rm c}$ and (b) the constants C; G which give a measure of the nonleading e ects, allowing to accom odate in the scaling function (10) processes not very close to the critical point. We have used measurements at the SPS in order to x these parameters on the basis of eqs.(5) and (10). More specially, in a series of experiments (Pb+Pb, S+Au, S+Ag, S+S) with central and noncentral (Pb+Pb) collisions at the SPS [8, 9] net baryons have been measured at m idrapidity whereas the scaling variable $z_c = A_2^{2=3}A_tL^{-1}$, associated with these experiments, covers a su ciently wide range of values (1 $\mathbf{Z}_{\mathbf{C}}$ 2) allowing for a best t solution. The outcome of the t is consistent with the 0 and the equations (7) are simplied as follows: choiœ G

$$A_{2}^{2=3}n_{b} = c + \frac{2}{-}(z_{c} c) + C(z_{c} c)^{4}$$
 (11-a)

$$A_{2}^{2=3}n_{b} = c + \frac{2I_{o}}{2}t_{f}^{1=3}(1 + t_{f} - \frac{CI_{o}^{1=3}}{2})$$
 (11-b)

$$A_{2}^{2=3}n_{b}(y) = _{c} + \frac{(z_{c} _{c})L}{y(L _{y})} + O[(z_{c} _{c})^{4}]$$
 (11-c)

In eqs.(11) we have used the approximate values of the critical exponents $\frac{1}{3}$, - $\frac{1}{2}$ in the 3d Ising universality class [10]. We have also added eq.(11-c) which gives the universal behaviour of the net-baryon density $n_b(y)$, in the vicinity of the critical point (z_c) с 1). The tted values of the parameters in eq.(11-b) are $_{c} = 0.81$, C = 0.68 and the overall behaviour of the solution is shown in Fig.1. In turns out that the critical density is rather 0:17 fm ³: $_{\rm c}$ $\frac{\circ}{5}$ small, compared to the normal nuclear density 12 fm, f = 6 fm), suggesting that the critical tem perature remains (R_o close to the value T_c 140 M eV obtained in studies of QCD on the lattice at zero chem ical potential [11]. The di erence $d_c = \dot{z}_c$ _cjin Fig. 1 is a m easure of the proxim ity of a given experiment $(A_+;L)$ to the critical point. W e observe that the central Pb+Pb collisions at the SPS $(d_c$ 1:1) drive the system into the most distant freeze-out area, from the critical point, as com pared to other processes at the sam e energy. In fact, the most suitable experiments to bring quark matter close to the critical point at the SPS are:

S+S (d_c) 0:18), S+S₁ (d_c) 0.20) and C+C (d. 0:06)), central collisions. One may even reach the critical point (d_c) 0) at the SPS with medium size nuclei, either in noncentral collisions: $S + S_i S_i + S_i$ (A_t 29) or at lower 130 A_{C}^{GeV}). Also, the experiments at RHIC, with energies: C+C (P 0:27) collisions, com e close to the critical point and m ay centralAu+Au (d_c 0) either with lighter nuclei or with noncentral even reach it exactly (d_c 165). Finally at the LHC $\left(\overline{s} \right)$ 5:5 TeV), Pb+Pb Au+Au collisions (A₊ collisions are expected to drive quark m atter into the crossover area ($z_c < c_r$ 0:13) where no critical uctuations occur. At lower energies however d_s 1:4 TeV), Pb+Pb central collisions may reach the critical point even (s at the LHC. Obviously, in order to have a clear answer on the proximity of a given experiment to the critical point, precision measurements of the net-baryon density at m idrapidity, in a wide range of energies (between SPS and RHIC), are needed, both for central and noncentral collisions. With such measurements, the scaling law (5) may be tested in its full extent and a sharp determination of the critical density can be achieved through eq.(7a).

The freeze-out line $(n_b \text{ versus } T_f)$ given by eq.(11b) is shown in Fig. 2. We have used the freeze-out tem peratures for the processes [12]: Pb+Pb 115 M eV) and S+S ($T_{\rm f}$ 145 M eV) at the SPS, in order to de-(T_f term ine the critical tem perature, T_c 145 M eV, and x the parameter 1:86. Again, we notice that the relatively sm all value of the freeze-out L tem perature in Pb+Pb collisions (at the SPS) is associated with the fact that in this process the system of quark matter is not driven close to the critical point. On the contrary the freeze-out tem perature in S+S collisions remains practically equal to the critical value, due to the proximity of this process to the critical point, at the freeze-out stage (Fig. 1). Finally, in Fig. 3, the rapidity distribution of net baryons in Au+Au collisions at RH IC 11;z 1:1) is shown, relying upon the fact that in this process the **(**L distance from the critical point is small (d_c 0:27) and eq.(11-c) is valid.

3 Baryonic critical uctuations

O noe the phase of quark m atter has reached the critical point in a particular class of experiments, as discussed in the previous section, strong critical uctuations are expected to form interm ittency patterns both in the pion and net-baryon sector. As already mentioned in the introduction, the origin of these uctuations can be traced in the presence, at $T = T_c$, of a zero m ass

eld with a classical pro le (-eld) which, under the assumption of a phase transition in local therm al equilibrium, is described by an elective action in 3 d, the projection of which onto rapidity space is written as follows [3]:

$$_{c} = \frac{R_{2}^{2}}{C_{A}} \sum_{y}^{Z} dy \frac{4}{2} \frac{1}{2} \frac{\theta}{\theta y} + 2C_{A}^{2} \frac{4}{c} + \frac{15}{2} ; \quad C_{A} = \frac{-c}{c}; \quad c = T_{c}^{1}$$
(12)

Equation (12) gives the free energy of the - eld within a cluster of size y in rapidity and R₂ in transverse space. The critical uctuations generated by (12) in the pion sector have been studied extensively in our previous work [3], therefore, in what follows, we are going to discuss the uctuations induced by the - eld in the net-baryon sector, noting that a direct measurement of these uctuations may become feasible in current and future heavy-ion experiments. For this purpose we introduce in eq.(12) the order parameter m (y) through the follow ing equations:

(y)
$$F_{c}^{2}m(y)$$
; $F_{-}\frac{hqqi_{o}}{3}$; $hqqi_{o}$ 3 fm³
 $_{c}$ $g_{y}^{2}dy^{4}\frac{1}{2}\frac{e^{m}}{e^{y}} + g_{2}jmj^{+15}$; $m(y) = {}_{c}^{3}m(y)$ (13)

where: $g_1 = F^2 \frac{R_2}{C_A c_c^2}$, $g_2 = 2C_A^2 F^4$. The partition function $Z = {}^R D$ [f]e c^{ff}] for each cluster is saturated by instanton-like con gurations [13] which for $y c_c$ lead to self-sim ilar structures, characterized by a pair-correlation function of the form :

hm (y)m (0)i
$$\frac{5}{6} \frac{(1=3)}{(1=6)} = \frac{R_2^2 C_A}{C_c^2} F^1 y^{\frac{1}{+1}}$$
 (14)

The size, in rapidity, of these fractal clusters is $c = \frac{R_2^2}{16 \frac{1}{c} C_A^2}^2$ according to the geom etrical description of the critical system s [13]. Integrating eq.(14) we not the uctuation h n_bi of the net-baryon multiplicity with respect to the critical occupation number within each cluster, as follow s:

h n_bi F⁻¹
$$\frac{R_{2}^{2}C_{A}}{2_{c}^{2}} \frac{2^{2=3}}{(1=3)} = \frac{5=6}{c}$$
 (15)

The dimensionless parameter F is of the order 10^2 and the size $_{\rm c}$, on general grounds $(R_2 \leq 2_c)$ is of the order of one $(c \leq 1)$. This is in agreement with the fact that the rapidity separation of two causally correlated space-time events is logarithm ically bounded, $y = \ln 1 + \frac{-c}{c}$, leading, practically, to a direct correlation length of the order of one (y $^{<}$ 1). However, in reality, the global baryonic system develops uctuations at all scales in rapidity since the direct correlation (O ($_{\rm c}$)) propagates along the entire system through the cooperation of m any self-sim ilar clusters of relatively sm all size ($_{\rm c}$ 0:35 and h n_bi 140). We have quantied this mechanism in a Monte-Carlo sim ulation for the conditions of the experiments at RHIC (in RHIC the size of 11) in order to generate baryons with critical uctuations. the system is L The distribution of the order parameter in (y) j describing the uctuations of the \critical" baryons in the rapidity space for a typical event as well as the corresponding interm ittency analysis in terms of factorial moments are presented in Fig. 4. The interm ittency exponent of the second m om ent F_2 0:18 which is very close to the theoretically in rapidity is found to be s_2 expected value $(\frac{1}{6})$ of a monofractal 1 d set with fractal dimension $\frac{5}{6}$. The quantitative details of this new class of phenom ena, concerning critical uctuations in the net-baryon sector, and in particular the interm ittency pattern of the observable net-baryon density, go beyond the purpose of this Letter and will be presented elsewhere [14].

4 Concluding rem arks

In conclusion, we have shown that the baryonic sector in an experiment with heavy ions, possess valuable information regarding the proximity and observation of the second order QCD critical point. Complementary suggestions for the signi cance of net-baryon uctuations in order to trace the critical line of rst order in the phase diagram are described in [16]. In particular the measurements of net-baryon spectra in rapidity provide a valuable set of observables in heavy ion experiments, in connection with the phenomenology of the QCD critical point. The trend of the existing data at the SPS suggests the presence of a critical point of second order in the phase diagram (Figs. 1,2) specified by the critical values of temperature and density: $T_{\rm c}$ 145 MeV, $_{\rm c}$ $-\frac{\circ}{5}$. A scaling law for the net-baryon density at midrapidity $n_{\rm b}$, as a function of the initial energy and the number of participants, has been established in the neighbourhood of the critical point. The scaling function

incorporates the indices of the universality class (critical exponents) whereas the scaling variable gives a measure of the proximity of a given experiment to the critical point. On the basis of this investigation, the experiments at RHIC are very likely to reach the QCD critical point and as a rst sign of this new phenomenon we have predicted a rather unconventional prole of net-baryon density in rapidity (in Au+Au collisions), associated with the presence of a critical point, nearby (Fig. 3). Finally we have indicated the presence of strong intermittency elects in the rapidity spectrum of the net baryons which can in principle be used to reveal experimentally the QCD critical point and con monits universality class (Fig. 4).

References

- [1] F.W ilczek, hep-ph/0003183.
- M. Stephanov, K. Rajagopaland E. Shuryak, Phys. Rev. Lett. 81, 4816 (1998); J. Berges, D.-U. Jungnickel and C. W etterich, Phys. Rev. D 59 (1999) 034010; Eur. Phys. J. C13 (2000) 323.
- [3] N.G. Antoniou, Y.F. Contoyiannis and F.K. Diakonos, Nucl. Phys. A 661, 399c (1999).
- [4] R.Brockm ann and W.Weise, Phys.Lett.B 367, 40 (1996).
- [5] B.Muller, Nucl. Phys. A 661, 272c (1999).
- [6] A.Maciolek, R.Evans and C.R.W ilding, Phys. Rev. E 60, 7105 (1999).
- [7] M E. Fisher and P.G. de Gennes, C R. Seances A cad. Sci. B 287, 207 (1978).
- [8] The NA35 Collaboration, Eur. Phys. J. C 2, 634 (1998); D. Rohrich
 (NA35 Collaboration), Nucl. Phys. A 566, 35c (1994); F. Sikler (NA49
 Collaboration), Nucl. Phys. A 661, 45c (1999).
- [9] G E. Cooper (NA 49 Collaboration), Nucl. Phys. A 661, 362c (1999).
- [10] S.K.Ma, Modern Theory of Critical Phenomena, (Benjamin/Cummings, Reading, MA, 1976).

- [11] E. Laerm ann, Nucl. Phys. A 610, 1c (1996).
- [12] V.Heinz, J.Phys.G 25, 263 (1999); R.Stock, Nucl. Phys. A 661, 282c (1999).
- [13] N.G. Antoniou, Y.F. Contoyiannis, F.K. Diakonos and C.G. Papadopoulos, Phys. Rev. Lett. 81, 4289 (1998); N.G. Antoniou, Y.F. Contoyiannis and F.K. Diakonos, Phys. Rev. 62E, 3125 (2000).
- [14] N.G. Antoniou, F.K. Diakonos and A.S. Kapoyannis (in preparation).
- [15] A.Bialas and R.Peschanski, Nucl. Phys. B 273, 703 (1986); B 308 857 (1988).
- [16] S.Gavin, nucl-th/9908070.

Figure Captions

FIG.1. The scaling law (10) is illustrated together with m easurements at the SPS. The critical point and the corresponding break in the slope of $(z_c; _c)$ are also shown.

FIG.2. The freeze-out line, n_b as a function of t_f , is illustrated together with m easurem ents at the SPS (S+S, Pb+Pb). The extracted freeze-out points for S+Au and S+Ag are also shown in this diagram.

FIG.3. The net-baryon prole in rapidity for Au + Au collisions at RHIC energies is shown, as predicted by eq.(11-c).

FIG.4. (a) The distribution of jn (y) j in rapidity for a MC-generated event. (b) The rst three factorialm on ents for the event shown in (a) in a log-log plot. A linear t determining the slope s_2 (0:18) of the second moment is also shown.







Fig. 1



Fig. 2



Fig. 3