

In uence of perturbations on the electron wave function inside the nucleus

M .Yu. Kuchiev y and V .V .Flambaum z

School of Physics, University of New South Wales, Sydney 2052, Australia

A b s t r a c t. A variation of the valence electron wave function inside a nucleus induced by a perturbative potential is expressed in term s of the potential m o m e n t a . A s an application we consider Q E D vacuum polarization corrections due to the Uehling and W ichm ann-K roll potentials to the weak interaction m a t r i x e l e m e n t s .

P A C S n u m b e r s : 32.80.Y s, 11.30.E r, 31.30.J v

Short title: In uence of perturbations on the electron wave function inside the nucleus

January 28, 2022

y E-m ail: kuchiev@newt.phys.unsw.edu.au

z E-m ail: flambaum@newt.phys.unsw.edu.au

1. Introduction

Precise low-energy experiments on parity nonconservation (PNC) in atoms provide a test of the standard model of elementary particle interactions. In a recent PNC experiment with cesium by Wood et al [1] the PNC E1 transition amplitude between the 6s and 7s states has been determined with an unprecedented accuracy of 0.3 %. At this level of accuracy a small perturbative potential (such as the Breit interaction, QED vacuum polarization) may influence the result. The PNC weak interaction matrix elements are determined by the electron wave function inside a nucleus. Having this, as well as some other possible applications, in mind we present in this paper a simple analytical approach which allows one to calculate corrections to the electron wave function near the origin due to a perturbative potential. As an application we consider corrections to weak matrix elements produced by the QED vacuum polarization (Uehling potential [2], Wichmann-Kroll potential [3]) and a QED correction to the electron potential. Relativistic units $c = \hbar = 1$; $e^2 = 1/137.04$; $a_0 = m^{-1}$ (a_0 is the Bohr radius) are used below everywhere, if not specified otherwise..

2. General formalism

Our goal is to describe a variation of an electron wave function inside the nucleus due to some local potential $V(r)$ considered as a perturbation. Let us presume that the electron motion in an atom is described with the help of a single-electron spherically symmetrical Hamiltonian H , which is, generally speaking, relativistic, i.e. of the Dirac type. The wave function for the n -th energy level $\psi_{njl}(r)$ which is the Dirac 4-spinor characterized by the total momentum j , orbital momentum l and projection of the total momentum m , the latter index is suppressed, satisfies the Schroedinger equation

$$E_{njl} \psi_{njl}(r) = H \psi_{njl}(r); \quad (1)$$

with appropriate boundary conditions at $r = 0$ and 1 . The spherically symmetrical potential $V(r)$ considered as a perturbation results in a variation of the wave function $\psi_{njl}(r)$ that we are looking for. This variation can obviously be presented as

$$\psi_{njl}(r) = \int \tilde{G}_{jl}(r; r^0; E_n) V(r^0) \psi_{njl}(r^0) d^3 r^0; \quad (2)$$

where $\tilde{G}_{jl}(r; r^0; E_n)$ is the corresponding Green function for the operator H , that can be expanded as a series over a full set of solutions of the Schroedinger equation (1) for the j, l wave

$$\tilde{G}_{jl}(r; r^0; E_n) = \sum_{n^0 \neq n} \frac{\psi_{n^0jl}(r) \psi_{n^0jl}^+(r^0)}{E_n - E_{n^0}}; \quad (3)$$

This Green function is necessarily a 4×4 matrix in the Dirac indices that are suppressed. Remember that the n -th electron level is excluded from summation in (3). To keep trace of this fact a tilde hat is used for $\tilde{G}_{j1}(r; r^0; E_n)$ thus distinguishing it from the Green function $G_{j1}(r; r^0; E)$ defined for arbitrary, non-specified energy E . Using orthogonality and a completeness of a full set of the wave functions $\psi_{nj1}(r)$ that satisfy the Schroedinger equation (1), i.e. the fact that

$$\int_V \psi_{nj1}^+(r) \psi_{nj1}(r) d^3r = \delta_{nn^0}; \quad (4)$$

$$\sum_{n,j1} \psi_{nj1}^+(r) \psi_{nj1}(r^0) = \delta(r - r^0); \quad (5)$$

one finds from (3) that $\tilde{G}_{j1}(r; r^0; E_n)$ satisfies the following non-homogeneous equation

$$(E_n - H) \tilde{G}_{j1}(r; r^0; E_n) = \delta(r - r^0) \sum_{n,j1} \psi_{nj1}(r) \psi_{nj1}^+(r^0); \quad (6)$$

as well as an integral condition

$$\int_V \tilde{G}_{j1}(r; r^0; E_n) \psi_{nj1}(r^0) d^3r = 0; \quad (7)$$

The last term in the right-hand side of (6) as well as condition (7) arise due to the same, mentioned above reason, namely that the n -th energy level is excluded from summation in (3).

Our goal is to describe the wave function variation inside the nucleus. To this end it is sufficient to find its variation only at one point inside, for example at the nuclear center which we consider as an origin. Having done that one recovers the wave function everywhere inside (and in close vicinity of) the nucleus simply by scaling its non-perturbed value to comply with the variation found at the chosen point. This statement follows from the fact that the potentials considered in this paper are supposed to produce negligible effects inside the nucleus. They contribute only due to their existence in a nuclear exterior region. This means that equations governing the electron wave function inside the nucleus remain intact by the perturbation. As a result the perturbation can only scale the wave function inside. We will use this fact looking for the wave function specially at the origin, i.e. hunting for $\psi_{nj1}(0)$.

In order to find the wave function at the origin one needs to find the Green function $G(0; r^0; E_n)$. This task can be conveniently fulfilled using the following method. Consider an energy E as an arbitrary parameter assuming only that E is located in some vicinity of a chosen atomic energy level E_n . Let us call $\psi_{j1}(r; E)$ a solution of the Schroedinger equation

$$(E - H) \psi_{j1}(r; E) = 0; \quad (8)$$

Obviously for an arbitrary energy E this solution cannot satisfy proper boundary conditions both at infinity and at the origin. However, we can always consider the

proper condition at one of the two points. We may assume therefore that $j_l(r; E)$ decreases at infinity

$$j_l(r; E) \rightarrow 0; \quad r \rightarrow \infty : \quad (9)$$

Solutions of this type are necessarily singular at the origin, if the energy parameter E does not coincide with some physical energy level. It is convenient, nevertheless, to introduce the following normalization integral

$$\int_0^\infty j_l(r; E) j_l(r; E_n) dr = 1 : \quad (10)$$

which converges well because the mentioned singularity at the origin is compensated for by a zero-type behaviour of the regular solution.

The function $j_l(r; E)$ possesses several important for us properties. Firstly, at the point $E = E_n$ the boundary condition at the origin can obviously be satisfied. Combining this statement with equations (8), (9) and (10) we conclude that for $E = E_n$ the function $j_l(r; E)$ coincides with the wave function for the n -th energy level

$$j_l(r; E_n) = j_l(r) : \quad (11)$$

To reveal another useful property of $j_l(r; E)$, let us differentiate the Schroedinger equation (1) over the energy at the point $E = E_n$

$$(E_n - H) \frac{\partial j_l(r; E_n)}{\partial E} = j_l(r; E_n); \quad r > 0 : \quad (12)$$

Keeping in mind that $j_l(r; E)$ is irregular at the origin we have to suspect that the right-hand side of (12) may include the delta-term $\delta(r)$, or its derivatives. This suspicion, justified below, prompts to remember that $r > 0$ when (12) is taken literally.

To proceed let us differentiate the normalization condition (10) over E at the point $E = E_n$

$$\int_0^\infty j_l(r) \frac{\partial j_l(r; E_n)}{\partial E} dr = 0 : \quad (13)$$

Compare now a set of equations (12) and (13) with (6) taken at $r = 0$; $r^0 > 0$ and (7). Their obvious identity indicates that the Green function at the origin $G_{j_l}(0; r; E_n)$ can be presented as

$$G_{j_l}(0; r; E_n) = j_l(0) \frac{\partial j_l(r; E_n)}{\partial E} : \quad (14)$$

We can now clarify behaviour of the right-hand side of equation (12) at $r = 0$. From (14) and (6) we deduce that our expectation was correct, at $r = 0$ the right-hand side of (12) should indeed be modified to include an additional delta-term $\delta(r)$ (though for our purposes it suffices to consider this equation only at $r > 0$).

We find that $\psi_{jl}(r; E)$ is a very convenient object. It allows one to describe simultaneously a set of wave functions $\psi_{jl}(r)$ (11) as well as the Green function $G(0; r; E_n)$ (14). It follows from (2), (14) that the variation of the wave function due to perturbation $V(r)$ can be expressed as

$$\psi_{jl}(0) = \psi_{jl}(0) + \frac{\partial \psi_{jl}^+(r; E_n)}{\partial E} V(r) \psi_{jl}(r) d^3r : \quad (15)$$

This convenient presentation is the main result of this Section. Its applications are discussed below.

3. Perturbation at small distances

Let us consider perturbative potentials $V(r)$ located at small separations from the nucleus. We will assume, however, that a potential considered gives a significant contribution mainly outside the atomic nucleus, i.e. assume that a region of distances where the potential is important satisfies

$$r_N \ll r \ll a_0 = Z^{-1} ; \quad (16)$$

where r_N is the nuclear radius. A wave function variation in the nuclear interior can be described by a scaling factor, which we find below.

Conventional presentation for the Dirac four-spinor for spherically symmetrical potentials reads

$$\psi_{jl}(r) = \frac{1}{r} \begin{pmatrix} f_n(r) \chi_{jl}(n) \\ ig_n(r) \chi_{j\bar{l}}(n) \end{pmatrix} : \quad (17)$$

Here $f_n(r)$ and $g_n(r)$ are the large and small radial components of the spinor, the indices jl for them are suppressed. They are normalized as

$$\int_0^\infty (f_n^2(r) + g_n^2(r)) dr = 1 ; \quad (18)$$

$\chi_{jl}(n)$ and $\chi_{j\bar{l}}(n) = (-1)^n \chi_{jlm}(n)$ are spherical spinors, and $l + \bar{l} = 2j$. The Dirac equation (1) for a motion in the potential $U(r)$ in this notation takes the familiar form

$$f_n^0(r) + \frac{1}{r} f_m(r) = (m + E_n - U(r)) g_n(r) \quad (19)$$

$$g_n^0(r) + \frac{1}{r} g_m(r) = (m - E_n - U(r)) f_n(r) ;$$

where $\epsilon = l(l+1) - j(j+1) = 1 - 4 = -(j+1/2)^2$. For small separations of an electron from the atomic nucleus (16) the potential $U(r)$ can be approximated by the pure Coulomb potential created by the nuclear charge Z

$$U(r) = -\frac{Ze^2}{r} : \quad (20)$$

Additional simplification for this region comes from the fact that the energy of a valence electron is low and therefore for small separations (16) one can safely assume that $E \approx m$. With these simplifications the Dirac equation (19) reads

$$\begin{aligned} f_n^0(r) + \frac{1}{r} f_n(r) &= -2m + \frac{Ze^2}{r} g_n(r) \\ g_n^0(r) + \frac{1}{r} g_n(r) &= \frac{Ze^2}{r} f_n(r) : \end{aligned} \quad (21)$$

There are two sets of solutions for these equations. One of them is regular at the origin. From (21) one finds that it behaves as

$$\begin{aligned} f_+(r) &= a_+ r ; \\ g_+(r) &= b_+ r ; \quad b_+ = \frac{+}{Z} : \end{aligned} \quad (22)$$

Here $\gamma = (\frac{1}{2} - (Z\alpha)^2)^{1/2}$. Clearly, this regular solution can be used to describe the electron wave function for small distances (16): $f_n(r) = f_+(r)$; $g_n(r) = g_+(r)$. The subscript + is used to distinguish this solution from the singular one. The latter, that will be called $f_-(r)$; $g_-(r)$, behaves as

$$\begin{aligned} f_-(r) &= a_- r^{-\gamma} ; \\ g_-(r) &= b_- r^{-\gamma} ; \quad b_- = \frac{+}{Z} : \end{aligned} \quad (23)$$

This solution is used below for the description of the Green function. The explicit form for both sets of solutions found from (21) reads,

$$f_-(r) = a_- \frac{(\frac{1}{2} - \gamma + 1)}{(8Z\alpha m)} J_{2-\gamma}(x) - \frac{1}{2(\frac{1}{2} - \gamma)} x J_{2-\gamma+1}(x) \quad (24)$$

$$g_-(r) = a_- \frac{+}{Z} + \frac{(\frac{1}{2} - \gamma + 1)}{(8Z\alpha m)} J_{2-\gamma}(x) : \quad (25)$$

Here $J_\nu(x)$ is the Bessel function, $x = \frac{p}{8Z\alpha m} r$. Numerical coefficients in (24), (25) are chosen to satisfy (22), (23). The coefficient a_+ for the regular solution that represents the electron wave function should be found from the normalization condition for this function. A proper normalization of the coefficient a_- is discussed in detail below, see (31).

Following the approach of Section 2 we need to replace E_n in (19) by an arbitrary value E , assuming that the corresponding solution $\psi_{j1}(r; E)$ with components $f(r; E)$; $g(r; E)$ behaves regularly at infinity as specified in (9). Further, we need to consider derivatives over the energy $\partial f(r; E) = \partial E f_E(r; E)$ and $\partial g(r; E) = \partial E g_E(r; E)$. Our task is to find these functions at small r . Since $f(r; E)$; $g(r; E)$ are singular at the origin, we have to expect that $f_E(r; E_n)$; $g_E(r; E_n)$ are singular as well. Bearing this in mind we can neglect the regular, and therefore small, right-hand side of (12) when $r \rightarrow 0$. We deduce from this that for small r the functions

$f_E(r; E_n); g_E(r; E_n)$ satisfy the homogeneous Dirac equation (19) behaving singularly in the vicinity of $r = 0$. The notation $f(r); g(r)$ introduced above specifies exactly this solution of the Dirac equation. It follows from (23) that for small separations the following asymptotic conditions hold

$$\begin{aligned} f_E(r; E_n) &= f(r) = a r^{-1} ; \\ g_E(r; E_n) &= g(r) = b r^0 : \end{aligned} \quad (26)$$

We need to continue this line of argumentation and find corrections of the order of $m r$ to the right-hand sides of (26). Observe firstly that according to (22) the right-hand side of non-homogeneous equation (12) is small enough for short distances to produce no corrections of the order of $m r$. Therefore the main correction to (26) arises from the homogeneous Dirac equation (21) when the mass term in the right-hand side of this equation is taken into account. Technically the easiest way to recover the correction due to the mass term is through an expansion of the explicit solutions (24), (25) in powers of $m r$. We will present the result below, in (32), where a similar correction for the regular solution (22) is also included.

To proceed we need to rewrite (15) in terms of large and small components of the Dirac spinor. Relations (22) ensure that variations of both components due to perturbation are proportional at the origin. This means that if we define the component ratios at the origin as a limit $f_+(0) = f_+(0) \quad [f(r) = f_+(r)]_{r \rightarrow 0}$ and $g_+(0) = g_+(0) \quad [g(r) = g_+(r)]_{r \rightarrow 0}$ (the limit is necessary since one of the components may turn zero at the origin) then

$$f_+(0) = f_+(0) = g_+(0) = g_+(0) : \quad (27)$$

Using this fact we derive from (15)

$$\frac{f_+(0)}{f_+(0)} = \frac{g_+(0)}{g_+(0)} = \int_0^{Z^{-1}} [f_n(r) f_E(r; E_n) + g_n(r) g_E(r; E_n)] V(r) dr \quad (28)$$

$$= \int_0^{Z^{-1}} [f_+(r) f(r) + g_+(r) g(r)] V(r) dr : \quad (29)$$

The last identity here arises because, as explained above, for small separations satisfying (16) the wave function $f_n(r); g_n(r)$ and derivatives over energy $f_E(r; E_n); g_E(r; E_n)$ can be replaced by the plus and minus solutions $f(r); g(r)$ respectively. The magnitude of these latter solutions is governed by coefficients a in asymptotic formulae (26). We need therefore to find the product $a_+ a_-$. This can be achieved using the following transformation. Multiply the non-homogeneous equation (12) by $\psi_{njl}^+(r)$ and integrate over the full 3D space from which the interior of a sphere S_n of radius n around the origin is excluded. Consider $n > 0$ as a small parameter which is to be put to zero at the end of the calculations, $n \rightarrow 0$. The normalization condition for $\psi_n(r)$ ensures that this procedure gives 1 in the right-hand side of (12). Integrating the Dirac Hamiltonian H

in the left-hand side of this equation by parts one observes that only the surface term sitting on the sphere S'' survives

$$\begin{aligned}
 \int_{r=0}^Z \psi_{nj1}^+(r; E) (E - H) \frac{\partial \psi_{nj1}(r; E)}{\partial E} d^3 r &= \int_{S''} \psi_{nj1}^+(r) \left(i \frac{\partial}{\partial r} \psi_{nj1}(r; E_n) \right) dS \\
 &= f_n'' \frac{\partial g''(E_n)}{\partial E} - g_n'' \frac{\partial f''(E_n)}{\partial E} = (f_+'' g_-'' - g_+'' f_-'') \\
 &= (a_+ b_- - b_+ a_-) :
 \end{aligned} \tag{30}$$

Here we use representation of the spinors $\psi_{nj1}(r)$ and $\partial \psi_{nj1}(r; E) = \partial E$ in terms of their large and small components, compare (17), as well as the fact that in unity $r = 1$ gives no contribution to the surface term since $\psi_{nj1}(r)$ and $\partial \psi_{nj1}(r; E) = \partial E$ are regular there. For small radius $r = ''$ one can express $\psi_{nj1}(r)$ and $\partial \psi_{nj1}(r; E) = \partial E$ in terms of $f(r); g(r)$, as was explained above, and use asymptotic formulae (22), (23), expressing thus the surface term via the coefficients $a_{\pm}; b_{\pm}$ in the last identity in (30). We find from these transformations that $a_+ b_- - b_+ a_- = 1$, or, remembering expressions of $b_{\pm}$ in terms of $a_{\pm}$ in (22), (23), find the product in question

$$a_+ a_- = \frac{1}{2} \frac{Z}{2} : \tag{31}$$

We possess now all ingredients necessary to derive the final result. Take equation (29). Substitute in its right-hand side expressions (24) and (25) for $f(r); g(r)$ that are supplemented by condition (31) on the coefficients $a_{\pm}$. After that expand the resulting integrand that arises from the right-hand side of (29) in powers of $m r$. This expansion is both justified and necessary in view of the following reasons. The expansion is allowed because the perturbation $V(r)$ is located in the region of small separations (16). A typical radius where the potential is located is a parameter for this expansion. Necessity for this expansion is twofold. Firstly, the approach developed neglects the screening of the Coulomb field by an electron cloud, which is a good approximation only in the close vicinity of the nucleus. Secondly, the procedure described neglects the regular solution in the right-hand side of the non-homogeneous Dirac equation, which is justified only for small distances where this solution is small.

Analytical calculations described above are straightforward. The final result reads

$$\frac{f_+(0)}{f_-(0)} = \frac{g_+(0)}{g_-(0)} = \frac{m}{h^2} \int_0^Z V(r) (a + kr) dr : \tag{32}$$

Here a is a parameter with length dimension while k is a dimensionless coefficient

$$a = \frac{Z}{2} \frac{h}{m c} ; \tag{33}$$

$$k = \frac{2(2-1)}{(4-2-1)} : \tag{34}$$

Relations (32),(33) are presented in absolute units, to make them more accessible for different applications.

Simple formula (32) is one of the most important results of this paper. It solves the main problem formulated in this section presenting a variation of the wave function at the origin in very transparent terms, as a linear combination of the zeroth and first moments of the perturbative potential.

Note that for the short-range potentials relative corrections to the energy and wave function are quite different. Indeed, we can approximate the energy variation in this case using asymptotic relations (22) as

$$E_n = \ln |j| \int_0^{\infty} V(r) r^2 dr : \quad (35)$$

For an arbitrary perturbation $V(r)$ the integral in the right-hand side of this identity may deviate significantly from the integral in (32). Therefore for short-range perturbations the energy variation, generally speaking, cannot serve as estimate what happens with the wave function.

An interesting comparison can be made with the nonrelativistic limit of (32) that reads

$$\frac{\psi_{n1}(0)}{\psi_{n1}(0)} = \frac{m}{h^2} \frac{2}{2l+1} \int_0^{\infty} V(r) r dr : \quad (36)$$

Deriving this identity we use the fact that according to (33) the parameter a turns zero in the limit $Z \rightarrow 0$, while from (34) one derives $k \rightarrow 2/(2l+1)$. There is a simple short cut derivation that leads to (36) and can be used for verification of this result. Assume that in the nonrelativistic limit the electron motion is dominated by the kinetic term, which is true for short separations. Derive from this an approximation

$$G_1(r; r^0; E_n) \approx G_1^{(0)}(r; r^0; E_n) \approx \frac{2m}{h^2} \frac{1}{2l+1} \frac{r_{<}^l}{r_{>}^{l+1}} ; \quad (37)$$

where $G_1^{(0)}(r; r^0; E_n)$ is the Green function for the free motion in the l -th partial wave and the last identity takes into account the fact that the binding energy is negligible for short distances. Remembering also that for nonrelativistic motion the wave function behaves as $\psi_1(r) \propto r^l$; $r \rightarrow 0$ one immediately derives (36) directly from (2) thus verifying relativistic equation (32) that we used above. To comply with absolute units used in (32) we use the same units in nonrelativistic formulae (36),(37).

The nonrelativistic result (36) shows that the parameter that governs variation of the wave function is $\int_0^{\infty} V(r) r dr$. This is almost an obvious result valid for a variety of quantum mechanical problems [4]. The relativistic result (32) shows that there exists another parameter $\int_0^{\infty} V(r) dr$. It is suppressed compared with the nonrelativistic parameter only by a factor Z which is not small for heavy atoms. This suppression can be well compensated for if the potential considered increases at small separations which

makes $\int^R V(r)dr$ larger than $\int^R V(r)rdr$. In this case the found relativistic parameter becomes more important than the nonrelativistic one. A perturbation due to the QED vacuum polarization discussed in Section 4 presents an example important for applications.

We can apply the results obtained above for a specific interesting case. Consider the parity-violating weak interaction of an atomic electron with the nucleus that mixes $s_{1=2}$ and $p_{1=2}$ states of an outer electron. The matrix element for this mixing $\langle p_{1=2} | \hat{W} | s_{1=2} \rangle$ is saturated inside the nucleus. Therefore the variation of the matrix element for the weak interaction can be found simply by adding variations of $s_{1=2}$ and $p_{1=2}$ states given in (43)

$$\frac{\langle p_{1=2} | \hat{W} | s_{1=2} \rangle}{\langle p_{1=2} | \hat{W} | p_{1=2} \rangle} = \frac{2m}{h^2} \int_0^{Z_1} V(r) \left(a + \frac{2}{3} kr \right) dr : \quad (38)$$

Deriving this result we take into account that essential parameters for $s_{1=2}$ and $p_{1=2}$ states are $s = 1$; $p = 1$; $s = p = (1 - (Z\alpha)^2)^{1/2}$ and assumed that $k = 6 = (4 - 1)$.

4. Vacuum polarization

Let us apply (32) to a specific case when perturbation originates from polarization of the QED vacuum caused by the Coulomb field of the nucleus. In the lowest, second order of the QED perturbation theory this polarization is described by the Uehling potential [2] $V_{VP}(r)$

$$V_{VP}(r) = \frac{2}{3} \frac{Ze^2}{r} \int_0^{Z_1} \exp(-2mr) Y(\rho) d\rho ; \quad (39)$$

$$Y(\rho) = 1 + \frac{1}{2\rho^2} \frac{1}{1 + \frac{\rho^2}{2}} : \quad (40)$$

The Uehling potential (39) is singular at the origin

$$V_{VP}(r) = \frac{2}{3} \frac{Ze^2}{r} \left(\ln \frac{1}{mr} - C - \frac{5}{6} \right) ; \quad mr \ll 1 : \quad (41)$$

Here $C = 0.577$ is the Euler constant. A $\ln mr$ function in (41) describes conventional scaling of the QED coupling constant e^2 that manifests itself for short distances. This scaling factor has an interesting consequence for the problem at hand. Being introduced in (32) it results in the $\ln^2 mr$ divergence of the integral $\int^R V_{VP}(r)dr$ at small r . This divergence is eliminated by the finite nuclear size. As a result we find an estimate for the variation of the weak matrix element (38)

$$\frac{\langle p_{1=2} | \hat{W} | s_{1=2} \rangle}{\langle p_{1=2} | \hat{W} | p_{1=2} \rangle} = \frac{2m}{h^2} \int_0^{Z_1} V_{VP}(r)dr = \frac{2}{3} (Z\alpha)^2 \ln^2(mr_N) : \quad (42)$$

This shows that there exists the $\ln^2 m r_N$ enhancement in the problem, as was firstly discovered by Milstein and Sushkov [5] using other methods. In our approach this result is linked with the relativistic parameter $\alpha = \int_0^R V(r) dr$ (33) introduced in Section 3. For heavy atoms the $\ln^2 m r_N$ enhancement compensates for the additional suppressing factor Z in the relativistic parameter α (33) thus making this parameter dominant.

In order to present more accurate results describing the influence of the vacuum polarization on the wave function let us substitute (39) in (32). Integrating over r we find

$$\begin{aligned} \frac{f_+(0)}{g_+(0)} &= \frac{g_+(0)}{g_+(0)} = \frac{2Z}{3} \int_0^1 \int_0^1 \alpha m E_1(m r_N) + \frac{k}{2} Y(\alpha) d\alpha \\ &= \frac{Z^2}{16} \frac{3}{4} \frac{(2-1)}{1} + Z \int_0^1 \frac{2}{3} \int_0^1 E_1(m r_N) Y(\alpha) d\alpha ; \end{aligned} \quad (43)$$

where $Y(\alpha)$ is defined in (40) and we used (33), (34) to present α, k explicitly. As is evident from (42) it is essential to take the finite size r_N of the nucleus into account. We follow in (43) the simplest way cutting the divergent integral in (43) at $r = r_N$. The symbol $E_1(x)$ in (43) represents the known integral-exponent function

$$E_1(x) = \int_0^1 \exp(-xt) \frac{dt}{t} : \quad (44)$$

Formula (43) solves the problem formulated above, giving a simple transparent presentation for variation of the atomic electron wave function due to vacuum polarization. Similarly we can find contribution of the QED vacuum polarization to the parity-violating weak interaction. Substituting (39) into (38) and making transformations similar to the ones used in (43) we find

$$\frac{\langle p_{1=2} | \hat{W} | p_{1=2} \rangle}{\langle p_{1=2} | \hat{W} | p_{1=2} \rangle} = \frac{Z^2}{4} \frac{3}{4} \frac{1}{4} \frac{1}{1} + Z \int_0^1 \frac{4}{3} \int_0^1 E_1(m r_N) Y(\alpha) d\alpha : \quad (45)$$

Equation (45) presents the weak interaction matrix element for an arbitrary atom in a transparent analytical form without fitting parameters. Numerical results are easily obtained by a straightforward one-dimensional integration in (45). One only needs to specify the nuclear size that can be taken as $r_N = 1.2 \cdot 10^{-13} A^{1/3}$ cm where A is the atomic number, see [6]. For the most interesting case of the ^{13}C s formula (45) gives correction produced by the Uehling potential 0.47%.

Alternatively the right-hand side of (45) can be calculated using an expansion in powers of $m r_N \ll 1$ that reads $-f \frac{3}{4(4^{3/2}-1)} Z^2 + \frac{2}{3} (Z^2)^2 [(\ln \frac{2}{m r_N} - C - \frac{5}{6})^2 + 0.759] g + O(m r_N)$, where $C = 0.577$. This expansion brings (45) to a form that is close, but not identical to the one derived in [5]. We will not pursue an origin for this discrepancy since calculations in the cited paper were fulfilled up to a constant that was eventually used as a fitting parameter.

Compare this result with other results obtained recently. Johnson, Bednyakov and So in Ref. [8] calculated correction due to the Uehling potential for the parity-nonconservation in the 6s-7s amplitude in ^{133}Cs . It proves to be large 0.4%, which agrees with qualitative expectations expressed by Sushkov in [9] previously. The result of [8] includes, along with variation of the weak matrix element, variations of the dipole matrix element and the corresponding energy denominator that, combined together, describe a satisfactory mixing measured experimentally. Ref. [7] of Dzuba, Flambaum and Ginges confirms this result and supplies more details providing separate variations for all three quantities mentioned above. It was found that variations of the dipole matrix element and the energy denominator, being not small, compensate each other almost completely. Thus the variation of the weak matrix element proves to be 0.4%. Numerical calculations in [5] were restricted by the logarithmic accuracy that was improved by using a constant as a fitting parameter to obtain 0.4% in line with [8].

We are interested in heavy atoms where the parameter Z is not small, therefore the lowest order polarization potential (Uehling potential) may be not sufficient. The higher order polarization potential (Weichmann-Kroll potential) was obtained in [3]. To calculate the correction to the weak matrix element with the logarithmic accuracy it is enough to know this potential at small distances [10]:

$$V_{WK}(r) = 0.092 \frac{2}{3} (Z\alpha)^2 \frac{Ze^2}{r} \ln r \quad (46)$$

The calculation with the logarithmic accuracy gives the following ratio of the Weichmann-Kroll correction to the Uehling correction for the weak matrix element (see (38)):

$$\frac{W_{WK}}{W_U} = 0.184 \frac{(Z\alpha)^2}{\ln 1/(m r_N)} \quad (47)$$

For ^{133}Cs this ratio is about -0.007. This confirms the statement of [5] that the higher order corrections to the polarization potential are not important (this may probably be explained by high momenta of the electron-positron pair in the polarization loop).

5. Large separations, non-relativistic case

Let us apply formula (15) for the case when the perturbative potential $V(r)$ is located in a region of distances r that satisfy the following conditions

$$a_0 = Z\alpha \ll r \ll a : \quad (48)$$

An example of an application here may be the calculation of QED corrections to the weak matrix element which originate from the atomic electron potential (see below).

Two simplifications are possible here. Firstly, the motion can be described by non-relativistic equations, and, secondly, the semiclassical approach is valid here [4]. We

can therefore assume that the Dirac spinors $\chi_{j1}(r;E)$; $\chi_{nj1}(r)$ can be expressed in terms of the single-component nonrelativistic wave functions. The angular components of the nonrelativistic wave functions will be called $\psi_1(r;E)$ and $\psi_{n1}(r)$ respectively. Applying conventional semiclassical methods [4] in the classically allowed region, which includes all distances specified in (48), we can write

$$\chi_1(r;E) = \frac{1}{r} \frac{2\psi_1(E)}{v(r)} \sin \int_r^Z p(r^0) dr^0 ; \quad (49)$$

where $p(r)$ and $v(r)$ are a classical momentum and velocity, $p(r) = mv(r)$, and $\psi_1(E)$ is a classical frequency

$$\psi_1(E) = \frac{2}{T(E)} ; \quad T(E) = \int \frac{dr}{v(r)} ; \quad (50)$$

Generally speaking, the velocity $v(r)$ depends on the energy E , making the period of the classical motion $T(E)$ and the frequency $\psi_1(E)$ energy dependent as well. Recall, however, that we are interested in the behaviour of an outer electron whose binding energy is much lower than an atomic potential when r satisfies (48). This fact makes the velocity and momentum in the integrand in (49) almost independent on energy E in the vicinity of the n -th energy level. In contrast, $\psi_1(E)$ exhibits rapid variation with energy because the integral for the period $T(E)$ in (50) is saturated at large distances, where velocity sharply depends on energy. Taking this into account we deduce from (49) that in the region of interest (48) the following equality holds

$$\frac{\partial \chi_1(r;E)}{\partial E} = \frac{1}{2\psi_1(E)} \frac{d\psi_1(E)}{dE} \chi_1(r;E) ; \quad (51)$$

This shows that in the region (48) the derivative of the wave function $\partial \chi_1(r;E)/\partial E$ can be described by a simple scaling factor $(1=2\psi_1(E))(d\psi_1(E)/dE)$. This statement remains true for the Dirac spinor $\partial \chi_{j1}(r;E)/\partial E$ as well because in the considered region (48) the spinor is proportional to the nonrelativistic wave function (51). Using now the fact that for shorter separations $r < a_0=Z$ the perturbation is assumed insignificant, we conclude that description of the perturbation by the scaling factor remains valid all the way down to the nucleus. This means that for all distances inside an atomic core $0 < r < a$ the derivative of the Dirac spinor over energy $\partial \chi_{j1}(r;E)/\partial E$ remains proportional to the spinor itself $\chi_{j1}(r;E)$ with a scaling coefficient identical to the one in the right-hand side of (13). Using this result in (15) we find

$$\chi_{nj1}(r) = \frac{\psi_1^0(E_n)}{2\psi_1(E_n)} \chi_{nj1}(r) E_n \quad (52)$$

In what follows we will not need an explicit form for relations expressing the spinor via the nonrelativistic wave function. A simple fact of their linear dependence will be sufficient.

Here $\dot{\psi}^{(0)}(E)$ is a shortcut notation for a derivative of $\psi(E)$ over E , while E_n is an energy variation due to the potential $V(r)$

$$E_n = \int \psi_{n1}^* V(r) \psi_{n1} d^3r : \quad (53)$$

We conclude from (52) that the relative variation of the wave function at the origin is proportional to the variation of the energy level, being independent on any specific features of the potential. The coefficient $\dot{\psi}^{(0)}(E_n) = (2\dot{\psi}(E_n))$ in this formula is expressed in terms of the classical frequency for the electron motion. It is very simple for calculations, but can be simplified even further, if one needs only an estimation. Remember again that large separations from the atom $r > a_0$ give large contribution to the classical period $T(E)$. For these distances an atomic field can be approximated by the Coulomb potential $\phi = e^2/r$ created by a singly charged atomic residue. This fact allows one to approximate the frequency by conventional formula of Newtonian celestial mechanics for the Kepler problem which for the potential $\phi = e^2/r$ read $\dot{\psi}(E) = (\pi e^2) (2B_n^3/m)^{1/2}$, where $B_n = m E_n$ is the electron binding energy. Substituting this in (52) one finds

$$\psi_{n1}(0) \approx \frac{3}{4} \frac{B_n}{B_n} \psi_{n1}(0) ; \quad (54)$$

where $B_n = m E_n$ is the variation of the binding energy. Thus the behaviour of the wave function can be described in terms of the binding energy only. There is, of course, a short-cut way to derive this result. The wave function of an outer electron at small r is known to depend on the binding energy according to $\psi_{n1}(r) = \text{const} \cdot r^{\frac{3}{2}}$, where n is an effective radial quantum number defined by the binding energy $B_n = m e^4 = (2n^2)$. Taking variation of this relation and assuming a weak influence of the perturbation on constant immediately reproduces (54).

The numerical simulation performed shows that for the $6s_{1/2}$ state in cesium atom the Kepler approximation (54) ensures an accuracy of $\sim 20\%$ for all perturbative potentials $V(r) = \text{const} \cdot \exp(-pr)$ with $1/a_0 \leq p \leq 50/a_0$, while an accuracy of slightly more sophisticated formula (52) is even higher, of the order of 2% .

We verified in this Section that the approach based on (15) gives sensible results (52), (54) for the region (48). Now we can use this approach to estimate the influence of QED radiative corrections to electron-electron interaction on the weak matrix element. In the non-relativistic limit the Uehling potential can be replaced by a zero-range potential (proportional to the δ -function). A larger correction comes from the self-energy operator $\Sigma(r; r^0; E)$ which also reduces to δ -function in the non-relativistic limit,

It is amusing to observe that the coefficient in the right-hand side of (54) originates directly from the Kepler law that relates cubes of periods with squares of separations.

An atomic potential was approximated by some local potential which reasonably reproduces the valence electron wave function both inside and outside the atomic core.

ie. $\langle r; r^0; E \rangle = \langle r | \frac{1}{2} r^2 U(r) \rangle$. A semiclassical formula for $U(r)$ was obtained by Flambaum and Zelevinsky [11]

$$U(r) = \frac{Z^2}{3m^2} \ln \frac{m}{\frac{1}{2} U_p(r) - E} r^2 U(r): \quad (55)$$

Here $U(r)$ is the atomic potential, while $U_p(r)$ is the atomic potential with a correction that takes into account the centrifugal potential which influence the p-wave electron in the intermediate state (see details in [11]). As usual, this semiclassical expression is not valid near the turning points where $U_p(r) = E$. However, a very weak logarithmic singularity does not produce any practical limitations on the applicability of (55). For the electrostatic potential $r^2 U(r) = \frac{1}{4} \rho(r)$ where $\rho(r)$ is the electric charge density. The main contribution to the Lamb shift of the energy level produced by $U(r)$ is given by the nuclear charge (this contribution was calculated in [7]). The contribution of the electron charge density can be calculated using the Thomas-Fermi approximation. The main contribution here comes from the interval $a_0 = Z^{-1} < r < a_0 = Z^{-1/3}$. A simple estimate shows that the squared semiclassical electron wave function at $r = a_0 = Z^{-1/3}$ is Z times smaller than near the origin. Therefore, the electron-electron contribution to the Lamb shift of s-wave electron is Z times smaller than the electron-nucleus contribution. According to [7] the Lamb shift of the s-levels in ^{133}Cs is 0.1% . Then using equation (54) we obtain that the correction to the weak matrix element produced by the electron density contribution to the $U(r)$ is 0.001% .

6. Summary and conclusions

Equation (15) provides a convenient framework to calculate a variation of the wave function of an atomic electron inside the atomic nucleus that arises due to a perturbative potential in an atom. Applied to the region of large distances ($a_0 = Z^{-1} < r < a_0$) it results in equation (52) and its simplified version (54). There is a reason that makes them interesting for applications for the atomic parity nonconservation. The most difficult and cumbersome part of theoretical investigation in the latter problem present many-electron correlations [12, 13, 14, 15, 7]. The correlations take place exactly in the region of large distances discussed in Section 5. We can deduce from (52), (54) that an accuracy of calculations of the atomic spectrum provides a direct test for an accuracy of the weak matrix element calculation. There are, of course, other problems which require knowledge of an electron wave function in the vicinity of a nucleus such as hyperfine interaction, field isotopic shift, and time invariance violation.

In the region of small separations $r < a_0 = Z^{-1}$ our main result (32) gives simple transparent presentation for the influence of perturbation on the wave function. We deduce from it that for potentials singular at the origin a dimensionless parameter $m \int_0^R V(r) dr$ (33) measures the strength of the potential.

Formula (32) was applied to find a variation of the weak electron-nucleus matrix element due to the vacuum polarization (Uehling potential). The result is 0.47% for ^{133}Cs atom, which agrees with the results reported recently in [8, 5, 7]. The contributions of the Wichmann-Kroll potential and QED corrections to the electron-electron interaction were found to be very small.

Acknowledgments

We are grateful to M Kozlov, A Milstein and O Sushkov for useful discussions. This work was supported by the Australian Research Council.

- [1] Wood C S, Bennett S C, Cho D, Masterson B P, Roberts J L, Tanner C E, and Wieman C E 1997, *Science* 275 1759
- [2] Uehling E A 1935 *Phys.Rev.* 48 55
- [3] Wichmann E H and Kroll N M 1956 *Phys.Rev.* 101 843
- [4] Landau L D and Lifshitz E M 1977 *Quantum mechanics : non-relativistic theory* (Oxford, New York, Pergamon Press)
- [5] Milstein A I, Sushkov O P 2001 hep-ph/0109257
- [6] Berestetskii V B, Lifshitz E M, and Pitaevskii L P, *Relativistic quantum theory* (Oxford, New York, Pergamon Press, 1982).
- [7] Dzubov V A, Flambaum V V andGINGES J S 2001 hep-ph/0111019
- [8] Johnson W R, Bednyakov I and Song G 2001 *Phys. Rev. Lett.* 87 233001
- [9] Sushkov O P 2001 *Phys. Rev. A*, 63 042504
- [10] Milstein A I and Strakhovenko V M 1983 *Sov.ZhETF* 84 1247
- [11] Flambaum V V and Zelevinsky V G 1999 *Phys. Rev. Lett.* 83 3108
- [12] Dzubov V A, Flambaum V V and Sushkov O P 1989 *Phys. Lett. A* 141 147
- [13] Blundell S A, Johnson W R and Sapirstein J 1990 *Phys. Rev. Lett.* 65 1411
- [14] Blundell S A, Sapirstein J and Johnson W R 1992 *Phys. Rev. D* 45 1602
- [15] Kozlov M G, Porsev S G and Tupitsyn I I 2001 *Phys. Rev. Lett.* 86, 3260