

Charge Neutrality Effects on 2-flavor Color Superconductivity

M ei Huang¹, Pengfei Zhuang¹, Wei Qin Chao²

¹ Physics Department, Tsinghua University, Beijing 100084, China

huangmei@mails.tsinghua.edu.cn

zhuangpf@mails.tsinghua.edu.cn

² CCAST, Beijing 100080, China

Institute of High Energy Physics,

Chinese Academy of Sciences, Beijing 100039, China

chaowq@hp.ccast.ac.cn

The effect of electric and color charge neutrality on 2-flavor color superconductivity has been investigated. It has been found that the effect of the charge neutrality on a 2-flavor quark system is very different from that on a 3-flavor system. The BCS diquark pair in 2-flavor color superconducting phase has been largely reduced by the large Fermi momentum of electron, the diquark condensate first increases with the increase of quark's chemical potential, then decreases rapidly and finally disappears at about $\mu = 535 \text{ MeV}$, at which the thermodynamic potential equals to that in the neutral normal 2-flavor quark matter.

I. INTRODUCTION

Because of the existence of an attractive interaction in the anti-triplet quark-quark channel in QCD, the cold and dense quark matter has been believed to favor the formation of diquark condensate and being in the superconducting phase [1]– [4].

The QCD phase structure at high baryon density is determined by how many kinds of quarks can participate in the pairing. If we have an ideal system with two massless quarks u and d , and the three colored u quarks and three colored d quarks have the same Fermi surface, the system will be in 2-flavor color superconducting phase (2SC); and if we have an ideal system with three massless quarks u , d and s , and all the nine quarks have the same Fermi surface, then all the quarks will participate in pairing and form a color-flavor-locking phase (CFL) [5].

In the real world, the strange quark mass m_s should be considered, which will reduce the strange quark's Fermi surface. If m_s is large enough, there will be no pairing between us and ds , and a 2SC + s phase will be favored [6].

Also, in reality, the system should be electric and color neutral, the existence of electrons in the system will shift the Fermi surface of the two pairing quarks, and the system can be in BCS phase or crystalline phase [7] or normal quark matter, depending on how large the difference in the chemical potentials of the two pairing quarks is.

In a recent paper [8], it has been argued that the 2SC + s phase would not appear if the electric and color charge neutrality condition is added, which is also agreed with the results in [9] by using SU(3) NJL model and taking into account the self-consistent effective quark mass $m_f(\mu)$, where f refers to $u; d; s$. It is found in [9] that the charge neutrality has a large effect on the s quark mass, i.e., $m_s(\mu)$ begins to decrease at a smaller $\mu \sim 400$ MeV than the case if no charge neutrality is considered like in [10] [11], where $m_s(\mu)$ begins to decrease at about $\mu \sim 550$ MeV.

In this paper, complementary to [9], we investigate the effect of charge neutrality on 2-flavor color superconductivity assuming that there is no strange quark involved in the

chemical potential regime $e < 550 \text{ MeV}$.

This paper is organized as following: in Sec. II, we extend our method in [12] to derive the thermodynamic potential when charge neutrality is considered; the gap equations and charge neutrality conditions will be derived in Sec. III, and in Sec. IV, we will give our numerical results; the conclusions and discussions will be given at the end.

II. THERMODYNAMIC POTENTIAL

A. The Lagrangian

Assuming that the strange quark does not appear in the system, we use the SU(2) Nambu-Jona-Lasinio model, and only consider scalar, pseudoscalar mesons and scalar diquark. The Lagrangian density has the form as

$$L = \bar{q}(i \not{\partial} - m_0)q + G_S [(q\bar{q})^2 + (q_1 \sim q)^2] + G_D [(i\bar{q}^c \tau^b \gamma_5 q)(i\bar{q}^c \tau^b \gamma_5 q^c)]; \quad (1)$$

where $q^c = C q^T$, $\bar{q}^c = \bar{q}^T C$ are charge-conjugate spinors, $C = i\gamma^2 \gamma^0$ is the charge conjugation matrix (the superscript T denotes the transposition operation), the quark field $q = (q_i)$ with $i = 1, 2$ and $\alpha = 1, 2, 3$ is a flavor doublet and color triplet, as well as a four-component Dirac spinor, $\tau^a = (\tau^1; \tau^2; \tau^3)$ are Pauli matrices in the flavor space, and $(\tau^a)^{ik} = -(\tau^a)^{ki}$, $(\tau^a)^{bc} = -(\tau^a)^{cb}$ are antisymmetric tensors in the flavor and color spaces respectively. G_S and G_D are independent effective coupling constants in the scalar quark-antiquark and scalar diquark channel.

After bosonization, one can obtain the linearized version of the model (1)

$$\Gamma = \bar{q}(i \not{\partial} - m)q - \frac{1}{2} \bar{q}^c \tau^b \gamma_5 q^c - \frac{1}{2} \bar{q}^c \tau^b \gamma_5 q^c \quad (2)$$

$$\frac{1}{4G_S} - \frac{1}{4G_D};$$

where we have assumed that there will be no pion condensate and introduced the constituent quark mass

$$m = m_0 + \dots \quad (3)$$

From general considerations, there should be eight scalar diquark condensates [13] [14]. In the case of the NJL type model, the diquark condensates related to momentum vanish, and there is only one 0^+ diquark gap with Dirac structure γ_5 for massless quark, and another 0^+ diquark condensate with Dirac structure $\gamma_0 \gamma_5$ at nonzero quark mass. In this paper, we assume that the contribution of the diquark condensate with $\gamma_0 \gamma_5$ is small, and only consider the diquark condensate with γ_5 . The diquark condensate with Dirac structure $\gamma_5 \gamma_0$ has been recently discussed in [15].

We can choose the diquark condensates in the third color direction, i.e., only the first two colors participate in the condensate, while the third one does not.

The model is non-renormalizable, and a momentum cut-off should be introduced. The parameters G_S and μ in the chiral limit $m_0 = 0$ are fixed as

$$G_S = 5.0163G \text{ eV}^{-2}; \quad \mu = 0.6533G \text{ eV} \quad (4)$$

The corresponding effective mass $m = 0.314G \text{ eV}$, and we will choose $G_D = 3=4G_S$ in our numerical calculations.

B. Partition function and thermodynamic potential

The partition function of the grand canonical ensemble can be evaluated by using standard method [16] [17],

$$Z = N^0 \int [dq] [d\bar{q}] \exp \int_0^Z d^3x (\bar{\psi} + q_0 q) g; \quad (5)$$

where $\beta = 1/T$ is the inverse of temperature T , and μ is the chemical potential. When electric and color charge neutrality is considered, the chemical potential is a diagonal 6×6 matrix in flavor and color space, and can be expressed as

$$= \text{diag}(\mu_1; \mu_2; \mu_3; \mu_4; \mu_5; \mu_6); \quad (6)$$

where $q_1; q_2; q_3$ are for the three colored u quarks, and $q_4; q_5; q_6$ are for the three colored d quarks.

Like in [8] [9], the chemical potential for each color and flavor quark is specified by its electric and color charges $\mu_i = \mu_e Q_e + \mu_3 T_3 + \mu_8 T_8$, where $Q_e; T_3$ and T_8 are generators of $U(1)_Q; U(1)_3$ and $U(1)_8$. Because the diquark condenses in the third color direction, and the first two colored quarks are degenerate, we can assume $\mu_3 = 0$. For the same flavor, the difference of chemical potentials between the first two colored quarks and the third colored quark is induced by μ_8 , and for the same color, the difference of chemical potentials between u and d is induced by μ_e .

The explicit expressions for each color and flavor quark's chemical potential are:

$$\begin{aligned} \mu_1 = \mu_2 &= \frac{2}{3} \mu_e + \frac{1}{3} \mu_8; \\ \mu_4 = \mu_5 &= \frac{1}{3} \mu_e + \frac{1}{3} \mu_8; \\ \mu_3 &= \frac{2}{3} \mu_e + \frac{2}{3} \mu_8; \\ \mu_6 &= \frac{1}{3} \mu_e + \frac{2}{3} \mu_8. \end{aligned} \quad (7)$$

For the convenience of calculations, we define the mean chemical potential for the pairing quarks $q_1; q_5$, and $q_2; q_4$

$$\mu = \frac{\mu_1 + \mu_5}{2} = \frac{\mu_2 + \mu_4}{2} = \frac{1}{6} \mu_e + \frac{1}{3} \mu_8 \quad (8)$$

and the difference of the chemical potential

$$\mu_e = \frac{\mu_5 - \mu_1}{2} = \frac{\mu_4 - \mu_2}{2} = \mu_e = 2 \mu_8 \quad (9)$$

Because the third colored u and d, i.e., the 3rd and 6th quarks do not participate in the diquark condensate, the partition function can be written as a product of three parts,

$$Z = Z_{\text{const}} Z_{36} Z_{15;24} \quad (10)$$

The constant part is

$$Z_{\text{const}} = N^0 \exp \int_0^Z d^3 \mathbf{x} \left[\frac{2}{4G_S} + \frac{2}{4G_D} g \right] \quad (11)$$

Z_{36} part is for the unpairing quarks q_3 (u_3) and q_6 (d_3), and $Z_{15;24}$ part is for the quarks participating in pairing, q_1 (u_1) paired with q_5 (d_2), and q_2 (u_2) paired with q_4 (d_1). In the following two subsections, we will derive the contributions of Z_{36} and $Z_{15;24}$.

1. Calculation of Z_{36}

Introducing the 8-spinors for q_3 and q_6 ,

$$Z_{36} = (q_3; q_6; q_3^C; q_6^C); \quad (12)$$

we can express Z_{36} as

$$\begin{aligned} Z_{36} &= \int_0^Z d^3 \mathbf{x} \exp \left[\frac{1}{2} \int_{\mathbf{n}, \mathbf{p}} \frac{[G_0^{-1}]_{36}}{T} \right] Z_{36} \\ &= \text{Det}^{1=2} ([G_0^{-1}]_{36}); \end{aligned} \quad (13)$$

where the determinant operation Det is to be carried out over the Dirac, color, flavor and the momentum-frequency space, and $[G_0^{-1}]_{36}$ has the form of

$$[G_0^{-1}]_{36} = \begin{pmatrix} 0 & & & & 1 \\ \vdots & [G_0^+]_3^{-1} & 0 & 0 & 0 \\ \vdots & 0 & [G_0^+]_6^{-1} & 0 & 0 \\ \vdots & 0 & 0 & [G_0]_3^{-1} & 0 \\ \vdots & 0 & 0 & 0 & [G_0]_6^{-1} \end{pmatrix}^A; \quad (14)$$

with

$$[G_0]_i^{-1} = \delta_{ij} - g_i - m; \quad (15)$$

Here we have used $\delta = p$ and $g_i = i \cdot 0$.

For the two quarks not participating in the diquark condensate, from Eq. (13), we can have

$$G_{0,15;24}^{-1} = \begin{pmatrix} 0 & & & & 1 \\ G_{0,1}^{-1} & 0 & 0 & 0 & \\ 0 & G_{0,2}^{-1} & 0 & 0 & \\ 0 & 0 & G_{0,4}^{-1} & 0 & \\ 0 & 0 & 0 & G_{0,5}^{-1} & 0 \end{pmatrix}; \quad (22)$$

and the matrix form for is

$$= i \begin{pmatrix} 0 & & & & 1 \\ 0 & 0 & 0 & 1 & \\ 0 & 0 & 1 & 0 & \\ 0 & 1 & 0 & 0 & \\ 1 & 0 & 0 & 0 & \end{pmatrix}; \quad + = 0 ()^y 0: \quad (23)$$

From Eq. (20), we have

$$\ln Z_{15;24} = \frac{1}{2} \ln \text{Det} (G_{15;24}^{-1}): \quad (24)$$

For a 2 × 2 matrix with elements A; B; C and D, we have the identity

$$\text{Det} \begin{pmatrix} A & B \\ C & D \end{pmatrix} = \text{Det} (CB + CAC^{-1}D) = \text{Det} (BC + BDB^{-1}A): \quad (25)$$

Replacing A; B; C and D with the corresponding elements of $G_{15;24}^{-1}$, we have

$$\begin{aligned} \text{Det} (G_{15;24}^{-1}) &= \text{Det} D_1 = \text{Det} (+ + + G_{0,15;24}^{-1} []^{-1} G_{0,15;24}^{-1}) \\ &= \text{Det} D_2 = \text{Det} (+ + G_{0,15;24}^{-1} []^{-1} G_{0,15;24}^{-1}): \end{aligned} \quad (26)$$

Using the massive energy projectors in [12] for each flavor and color quark, we can work out the diagonal matrix D_1 and D_2 as

$$\begin{aligned} (D_1)_{11} &= (D_1)_{22} = [(p_0 +)^2 [E -]^2] + [(p_0 +)^2 [E +]^2] + \\ (D_1)_{33} &= (D_1)_{44} = [(p_0)^2 [E -]^2] + [(p_0)^2 [E +]^2] + \\ (D_2)_{11} &= (D_2)_{22} = [(p_0)^2 [E +]^2] + [(p_0)^2 [E -]^2] + \\ (D_2)_{33} &= (D_2)_{44} = [(p_0 +)^2 [E +]^2] + [(p_0 +)^2 [E -]^2] +; \end{aligned} \quad (27)$$

where $E = \frac{q}{(E^+)^2 + \dots}$.

With the above equations, Eq. (24) can be expressed as

$$\ln Z_{15;24} = 2 \sum_n \sum_p \left[\ln [(p_0^2 (E^+ + \dots))^2] + \ln [(p_0^2 (E^+ + \dots))^2] + \ln [(p_0^2 (E^+ + \dots))^2] + \ln [(p_0^2 (E^+ + \dots))^2] \right]; \quad (28)$$

C. The thermodynamic potential

Using the helpful relation

$$\begin{aligned} \ln Z_f &= \sum_n \ln [(p_0^2 + E_p^2)] \\ &= [E_p + 2T \ln(1 + e^{-E_p})]; \end{aligned} \quad (29)$$

we can evaluate the thermodynamic potential of the quark system

$$\begin{aligned} \Omega_q &= \frac{m^2}{4G_S} + \frac{\dots}{4G_D} \int \frac{d^3p}{(2\pi)^3} [2E + T \ln(1 + \exp(-E_3^+)) \\ &\quad + T \ln(1 + \exp(-E_3^-)) + T \ln(1 + \exp(-E_6^+)) + T \ln(1 + \exp(-E_6^-)) \\ &\quad + 2E^+ + 2E^- + 2T \ln(1 + \exp(-E^+)) + 2T \ln(1 + \exp(-E^-)) \\ &\quad + 2T \ln(1 + \exp(-E_+)) + 2T \ln(1 + \exp(-E_-))] \end{aligned} \quad (30)$$

with $E = E$.

For the total thermodynamic potential, we should include the contribution from the electron gas, Ω_e . Assuming the electron's mass is zero, we have

$$\Omega_e = \frac{4}{12} \frac{e}{2}; \quad (31)$$

The total thermodynamic potential of the system is

$$\Omega = \Omega_q + \Omega_e; \quad (32)$$

III. GAP EQUATIONS AND CHARGE NEUTRALITY CONDITION

From the thermodynamic potential Eq.(32), we can derive the gap equations of the order parameters m and \bar{m} for the chiral and color superconducting phase transitions.

A. Gap equation for quark mass

The gap equation for quark mass can be derived by using

$$\frac{\partial \Omega}{\partial m} = 0; \quad (33)$$

The explicit expression for the above equation is

$$\begin{aligned} m \left[4G_S \int \frac{d^3p}{(2\pi)^3} \frac{1}{E} \left[2 \frac{E}{E} (1 - f(E_+) - f(E_-)) \right. \right. \\ \left. \left. + 2 \frac{E^+}{E^+} (1 - f(E^+) - f(E^+)) \right. \right. \\ \left. \left. + (2 - f(E_3^+) - f(E_3) - f(E_6^+) - f(E_6)) \right] = 0; \quad (34) \end{aligned}$$

with the Fermi distribution function $f(x) = 1/(\exp(x/T) + 1)$. $m = 0$ corresponds to the chiral symmetric phase, $m \neq 0$ corresponds to the chiral symmetry breaking phase.

B. Gap equation for diquark condensate

Using

$$\frac{\partial \Omega}{\partial \bar{m}} = 0; \quad (35)$$

we can derive the gap equation for diquark condensate

$$\begin{aligned} \left[4G_D \int \frac{d^3p}{(2\pi)^3} \frac{1}{E} \left(1 - f(E_+) - f(E_-) \right) \right. \\ \left. + 2 \frac{1}{E^+} (1 - f(E^+) - f(E^+)) \right] = 0; \quad (36) \end{aligned}$$

C . C olor charge neutrality

The color charge neutrality condition is to choose q_{8c} such that the system has zero net charge T_8 , i.e.,

$$T_8 = \frac{Q}{Q_{8c}} = 0: \quad (37)$$

Evaluating the above equation, we have the color charge neutrality condition as

$$\begin{aligned} & \sum^Z \frac{d^3p}{(2\pi)^3} \left[\frac{E}{E} (1 - f(E_+) - f(E_-)) \right. \\ & \quad + \frac{E^+}{E^+} (1 - f(E^+_{+}) - f(E^+_{-})) \\ & \quad \left. + (f(E^+_3) - f(E^-_3)) + (f(E^+_6) - f(E^-_6)) \right] = 0: \end{aligned} \quad (38)$$

D . E lectric charge neutrality

Similarly, the electric charge neutrality condition is to choose q_e such that the system has zero net electric charge Q_e , i.e.,

$$Q_e = \frac{Q}{Q_e} = 0: \quad (39)$$

From the above equation, we obtain

$$\begin{aligned} & \sum^Z \frac{d^3p}{(2\pi)^3} \left[2 \frac{E}{E} (1 - f(E_+) - f(E_-)) \right. \\ & \quad + 2 \frac{E^+}{E^+} (1 - f(E^+_{+}) - f(E^+_{-})) \\ & \quad + 6 [f(E_{+}) + f(E^+_{+}) - f(E^+_{-}) - f(E_{-})] \\ & \quad \left. + 4 (f(E^+_3) - f(E^-_3)) - 2 (f(E^+_6) - f(E^-_6)) \right] + \frac{3}{2} = 0: \end{aligned} \quad (40)$$

IV . N U M E R I C A L R E S U L T S

In the numerical results, we investigate the following four different cases: 1), no color superconducting phase, no charge neutrality condition; 2), no color superconducting phase,

electric charge neutrality condition added, the electron's chemical potential in this phase will be characterized by μ_e^0 ; 3), color superconducting phase exists, no charge neutrality condition, the diquark condensate in this phase will be characterized by Δ_0 ; 4), color superconducting phase exists, electric and color charge neutrality condition added, the electron's chemical potential and diquark condensate in this phase will be characterized by μ_e and Δ_0 respectively.

Fig. 1 is for the chiral phase transition in the case of no electric neutrality considered, which is familiar to all of us, the quark possesses a large constituent mass in the low baryon density regime due to chiral symmetry breaking, and restores chiral symmetry at high baryon density.

Fig. 2 is the chiral phase diagram when the electric charge neutrality is considered, the solid squares and the solid circles are for quark mass m and electron's chemical potential μ_e^0 respectively. It is found that, in the chiral symmetric phase, there is a large Fermi surface for electrons, and the electron's chemical potential μ_e^0 increases linearly with the increase of quark's chemical potential μ .

Fig. 3 is the phase diagram for color superconductivity without charge neutrality constraints, the quark mass m (squares) and diquark condensate Δ_0 (circles) are plotted as functions of quark's chemical potential μ . It is found that in the color superconducting phase, the magnitude of the diquark condensate Δ_0 is about 100 MeV by using the parameter $G_D = 3=4G_S$. Δ_0 first increases with increasing μ , and this tendency stops at about $\mu = 530$ MeV, which is related to the momentum cut-off Λ .

Fig. 4 is the phase diagram for color superconductivity when electric and color charge neutrality condition is considered, the quark mass m (squares), diquark condensate Δ_0 (triangles) and the electron's chemical potential μ_e (circles) are plotted as functions of quark's chemical potential μ . It is found that when chiral symmetry restores, the color superconducting phase appears. In the color superconducting phase, it is found that both the diquark condensate Δ_0 and the electron's chemical potential μ_e first increase with increasing μ , and reaches their maximum at about $\mu = 475$ MeV, then decrease rapidly with increasing μ .

, and the diquark condensate disappears at about $\mu = 535\text{M eV}$.

In Fig. 5, we show the chemical potential μ_{8c} as a function of μ , which ensures the system having zero net color charge T_8 . It is found that μ_{8c} can be negative and positive, but its value is very small, only about several M eV. It means that for the same flavor quark u or d, the difference of Fermi momenta between the third colored quark and the first two colored quarks can be neglected.

In order to explicitly see what happens to a color superconducting phase when charge neutrality condition is added, we compare the electron's chemical potential in the neutral normal quark matter μ_e^0 (light circles) and in the neutral superconducting phase μ_e (solid circles) in Fig. 6, and compare the diquark condensate Δ_0 (light squares) and Δ (solid circles) in the electric charged and neutral superconducting phases in Fig. 7.

From Fig. 6, we find that the electron has a larger Fermi surface in the neutral superconducting phase than that in the neutral normal quark matter. This is because in the superconducting phase, the isospin is mainly carried by unpaired quarks only, and μ_e has to be larger in order to get the same isospin density. μ_e is equal to μ_e^0 at about $\mu = 535\text{M eV}$, at which the diquark condensate disappears. We separate μ_e into two parts as $\mu_e = \mu_e^0 + \delta\mu_e$, where $\delta\mu_e$ reflects the effect induced by diquark condensate, comparing μ_e (light squares) and μ_e^0 (solid squares) in Fig. 6, we can see that $\delta\mu_e$ increases when μ increases, and decreases when μ decreases, and when $\mu = 0$, $\delta\mu_e = 0$, too.

From Fig. 7, we can see that the diquark condensate Δ in the neutral superconducting phase has a much smaller value than Δ_0 in the charged superconducting phase, it means that the difference of the Fermi surfaces of the two pairing quarks reduces the magnitude of the diquark condensate Δ .

From Fig. 6 and Fig. 7, we can see a distinguished characteristic in the neutral superconducting phase, i.e., both the electron's chemical potential μ_e and the magnitude of the diquark condensate Δ first increase with increasing μ , then at about $\mu = 475\text{M eV}$, decrease with increasing μ . Now we try to understand this phenomena.

The magnitude of the diquark condensate is not only affected by μ , but also by $\mu_e = \mu_e^0 + \delta\mu_e$,

which describes the difference of Fermi surface between the two paired quarks and reduces the diquark condensate as seen in Fig. 7. The diquark condensate first increases with increasing μ , which is the result of the increasing density of states. At about $\mu = 475\text{M eV}$, where $\mu_d = \mu + 1=3 \mu_e$ is about 530M eV , the diquark condensate starts to be affected by the momentum cut-off. Like that in the charged color superconducting phase, the diquark condensate stops increasing with μ , so does μ_e . However, when $\mu > 475\text{M eV}$, μ_e^0 keeps increasing with μ , which reduces the diquark condensate largely. This is why we see μ_e decreases with increasing μ . Finally at a certain chemical potential $\mu = 535\text{M eV}$, where μ_d is about 580M eV , the diquark condensate disappears. Because μ_e decreases with the decrease of μ , the total μ_e decreases in the chemical potential regime $\mu > 475\text{M eV}$.

At last, in Fig.8, we show the thermodynamic potentials in the four different cases as functions of chemical potential μ , the light squares and circles are for the cases with and without charge neutrality and without diquark condensate, the dark circles and squares are for the cases with and without charge neutrality and with diquark condensate. It can be seen that in the chiral symmetric phase, the charged superconducting phase has the lowest thermodynamic potential, and the charged normal quark matter has the second lowest. The neutral superconducting phase has a little bit lower than that of the neutral quark matter in the chemical potential region $\mu < 535\text{M eV}$, and at $\mu = 535\text{M eV}$, the two coincide with each other. Therefore, in the chiral symmetric phase, the stable state of the neutral system is the color superconducting phase for $\mu < 535\text{M eV}$. At $\mu = 535\text{M eV}$, the stable phase is the normal neutral quark matter.

V. CONCLUSIONS

We investigated the effect of charge neutrality on a two flavor quark system. It has been found that the BCS diquark pair has been largely reduced by the large Fermi momentum of electron, the diquark condensate first increases with the increase of quark's chemical potential, then decreases rapidly and disappears at about $\mu = 535\text{M eV}$, at which the

thermodynamic potential equals to that in the neutral normal quark matter. In the chemical potential region $330\text{ MeV} < \mu < 535\text{ MeV}$, the stable neutral system is in the color superconducting phase if no strange quark involved.

As we mentioned in the introduction, we did not consider the strange quark in the system, which should be considered and has been investigated self-consistently in [9]. Comparing our results with their results about the diquark gap, the electron's chemical potential μ_e and the chemical potential μ_{sc} in 2SC, it can be found that the main difference lies in the chemical potential region $\mu > 400\text{ MeV}$. In [9], the diquark condensate in the neutral superconducting phase is largely reduced only in a small chemical potential region $370\text{ MeV} < \mu < 400\text{ MeV}$. μ_e first increases with μ then begins to decrease at about $\mu = 400\text{ MeV}$. As for μ_{sc} , the tendency also becomes different from our results at about $\mu = 400\text{ MeV}$. The reason lies in that, in [9], the strange quark involves in the system when $\mu > 400\text{ MeV}$. This shows that the effect of the charge neutrality condition on three- and two-flavor quark system is quite different.

A C K N O W L E D G E M E N T S

One of the authors (M.H.) thanks valuable discussions with Dr. T. Schafer. This work was supported in part by China Postdoctoral Science Foundation, the NSFC under Grant Nos. 10105005, 10135030 and 19925519, and the Major State Basic Research Development Program under Contract No. G2000077407.

-
- [1] D Bailin and A Love, *Phys. Rep.* 107, 325 (1984).
- [2] M Alford, K Rajagopal and F Wilczek, *Phys.Lett.B* 422,247 (1998).
- [3] R Rapp, T Schafer, E V Shuryak and M Velkovsky, *Phys.Rev.Lett.*81,53 (1998).
- [4] K Rajagopal and F Wilczek, hep-ph/0011333.
- [5] M Alford, K Rajagopal, F Wilczek, *NuclPhys.B* 537 (1999) 443.
- [6] M Alford, J. Berges, K Rajagopal, *NuclPhys.B* 538 (1999) 219.
- [7] M Alford, J. A Bowers, K Rajagopal, *Phys.Rev.D* 63 (2001) 074016; J. A Bowers, J. Kundu, K Rajagopal, E. Shuster, *Phys.Rev.D* 64 (2001) 014024; A. K. Leibovich, K Rajagopal, E. Shuster, *Phys.Rev.D* 64 (2001), 094005; J. Kundu, K Rajagopal, *Phys.Rev.D* 65 (2002), 094022; J. A Bowers, K Rajagopal, hep-ph/0204079.
- [8] M Alford, K Rajagopal, hep-ph/0204001.
- [9] A W Steiner, S Reddy, M Prakash, hep-ph/0205201.
- [10] F Gastineau, R Nebauer, J. Aichelin, *Phys. Rev. C* 65 (2002) 045204.
- [11] M Buballa, M Oertel, *NuclPhys. A* 703 (2002) 770.
- [12] M Huang, P Zhuang and W Q Chao, *Phys. Rev.D* 65 (2002) 076012.J
- [13] R D Pisarski, D H Rischke, *Phys.Rev.D* 60 (1999) 094013.
- [14] R D Pisarski, D H Rischke, *Phys.Rev.Lett.*83 (1999) 37.
- [15] T Fugleberg, hep-ph/0206033.
- [16] J Kapusta, "Finite-temperature Field Theory", Cambridge University Press, 1989.
- [17] M L Bellac, "Thermal Field Theory", Cambridge University Press, 1996.




FIG .1. The quark mass m as a function of chemical potential μ in the case of no color superconducting phase and no charge neutrality condition.




FIG .2. The quark mass m (squares) and the electron's chemical potential μ_e^0 (circles) as functions of chemical potential μ in the case of neutral normal quark matter.




FIG . 3. The quark mass m (squares) and the diquark condensate Δ_0 (circles) as functions of chemical potential μ in the case of no charge neutrality on color superconducting phase.




FIG .4. The quark mass m (squares), the diquark condensate Δ (triangles) and the electron's chemical potential μ_e (circles) as functions of chemical potential μ in the case of considering charge neutrality on color superconducting phase.




FIG .5. The μ_{8c} as a function of chemical potential μ in the case of considering charge neutrality on color superconducting phase.




FIG .6. The electron's chemical potential in the neutral normal quark matter μ_e^0 (light circles) and in the neutral superconducting phase μ_e (solid circles) as functions of chemical potential μ , and $\mu_e = \mu_e^0$ (light squares) comparing with Δ (solid squares) as functions of μ .




FIG .7. The diquark condensate in the electric charged Δ_0 (light squares) and in the neutral (solid circles) superconducting phase as a function of chemical potential μ .




FIG .8. The thermodynamic potentials as functions of chemical potential μ , the light squares and circles are for the cases with and without charge neutrality and without diquark condensate, the dark circles and squares are for the cases with and without charge neutrality and with diquark condensate.