

Impact of Tau Polarization for the determination of high $\tan\beta$ and $A^\pm$

E. Boos^{a,b}, G. Moortgat-Pick^{b,c}, H. J. Martyn^d, M. Sachwitz^e,
A. Vologdin^a

^a Skobeltsyn Institute of Nuclear Physics, Moscow State University, 119992 Moscow, Russia

^b DESY, Deutsches Elektronen-Synchrotron, D-22603 Hamburg, Germany

^c II. Institute for Theoret. Physics, University of Hamburg, D-22761 Hamburg, Germany

^d Rheinisch-Westfälische Technische Hochschule, D-52074 Aachen, Germany

^e DESY, Deutsches Elektronen-Synchrotron, D-15738 Zeuthen, Germany

Abstract

In order to determine the fundamental MSSM parameters $M_1, M_2,$ and $\tan\beta$ the tau polarization from $\tilde{\tau}$ decays can be explored as a 'bridge' between the gaugino/higgsino and the stau sector in particular in the high $\tan\beta$ range. Even in the case of high $\tan\beta$ an accuracy of $(\tan\beta) \sim 5\%$ with an simultaneous determination of $A^\pm$ is possible without assuming a specific SUSY breaking scheme.

1 Introduction

The minimal Supersymmetric extension of the Standard Model (MSSM) is one of the most promising extensions of the Standard Model (SM). However, SUSY has to be broken and in the unconstrained version of the MSSM a parameterization of all possible soft SUSY breaking terms leads to 105 new parameters in addition to the ones of the SM.

A linear collider (LC) is, due to its clear signatures, the most promising tool for revealing the underlying structure of the physics beyond the SM as e.g. the precise determination of these parameters.

Strategies to determine the fundamental parameters in the chargino/neutralino sector have already been worked out in [1, 2] and references therein. However, for $\tan\beta > 10$, the gaugino/higgsino sector is weak dependent on $\tan\beta$ so that its determination becomes rather inaccurate. We therefore concentrate in this paper on the production of the SUSY partners $\tilde{\nu}_{1,2}$ and their decays into $\tilde{\nu}_1^0$ and using the polarization of the $\tilde{\nu}_1^0$'s for the accurate determination of high $\tan\beta$. Since the polarization involves simultaneously

*Talks given at the 10th International Conference on Supersymmetry and Unification of Fundamental Interactions, SUSY 02, DESY, Hamburg 2002

Speaker at SUSY 02, boos@theory.sinp.msu.ru

Speaker at SUSY 02, gudrid@maildesy.de

mixing parameters from both neutralino and stau sectors it plays the role of a ‘bridge’ between these two sectors.

The importance of the τ polarization has already been pointed out in [3, 4]. We derive the compact formula for the polarization including the general mixing in the neutralino sector and show in which regions of MSSM parameter space the polarization will be a suitable observable. Contrary to the former studies we assume no specific character of the LSP $\tilde{\chi}_1^0$ or specific GUT relations between the underlying gaugino parameters. We show with an numerical example that the polarization on τ ’s is well suited for a rather accurate determination even of high $\tan\beta$ as well as for a simultaneous determination of A_0 .

2 The Stau Sector

2.1 Masses and Mixing

Since the τ lepton has the largest Yukawa coupling of the three lepton families the weak eigenstates $\tilde{\tau}_{L,R}$ mix to the mass eigenstates $\tilde{\tau}_{1,2}$, where the mass matrix is given by:

$$M_{\tilde{\tau}}^2 = \begin{pmatrix} M_L^2 + m^2 + D_L & m(A \tan\beta) \\ m(A \tan\beta) & M_E^2 + m^2 + D_R \end{pmatrix} = \begin{pmatrix} m_{LL}^2 & m_{LR}^2 \\ m_{LR}^2 & m_{RR}^2 \end{pmatrix} \quad (1)$$

with the D-terms $D_L = (\frac{1}{2} + \sin^2\theta_W)m_Z^2$ and $D_R = -\sin^2\theta_W m_Z^2$, the trilinear slepton-Higgs $\tilde{\chi}_R^0\tilde{\chi}_L^0 H_1$ coupling A , the higgsino mass parameter μ , the ratio of the Higgs expectation values $\tan\beta = v_2/v_1$ and the SU(2) doublet (singlet) mass parameters M_L (M_E). The mass parameters m_{LL}^2, m_{RR}^2 have to be positive for $\tan\beta > 1$, whereas the sign of the off-diagonal terms m_{LR}^2 depends on A and μ .

The mass eigenvalues are given by

$$m_{\tilde{\tau}_{1,2}}^2 = \frac{1}{2} [m_{LL}^2 + m_{RR}^2 \mp \sqrt{(m_{LL}^2 - m_{RR}^2)^2 + 4(m_{LR}^2)^2}] \quad (2)$$

and a large mass difference may be quite natural. In many SUSY scenarios $\tilde{\tau}_1$ is similar light as light charginos/neutralinos. A future high L/LC will be well suited to measure the masses with an high accuracy of e.g. about $(m_{\tilde{\tau}_1}) \sim 0.6 \text{ GeV}$ [5].

The stau mixing angle $\tilde{\alpha}$, $[0; \pi/2]$ is given by:

$$\tan(2\tilde{\alpha}) = \frac{2m_{LR}^2}{m_{RR}^2 - m_{LL}^2} = : \quad (3)$$

so that the mass parameters $m_{LL}^2, m_{RR}^2, m_{LR}^2$ can also be expressed via the measurable observables $m_{\tilde{\tau}_{1,2}}^2$:

$$m_{LL}^2 = \frac{m_{\tilde{\tau}_1}^2 + m_{\tilde{\tau}_2}^2}{2} - \frac{m_{\tilde{\tau}_2}^2 - m_{\tilde{\tau}_1}^2}{2} \cos(2\tilde{\alpha}); \quad (4)$$

$$m_{RR}^2 = \frac{m_{\tilde{\tau}_1}^2 + m_{\tilde{\tau}_2}^2}{2} + \frac{m_{\tilde{\tau}_2}^2 - m_{\tilde{\tau}_1}^2}{2} \cos(2\tilde{\alpha}); \quad (5)$$

$$m_{LR}^2 = \frac{1}{2} (m_{\tilde{\tau}_1}^2 - m_{\tilde{\tau}_2}^2) \sin(2\tilde{\alpha}); \quad (6)$$

One sees from (4) and (5) that one can distinguish the two cases

$$1. \text{ if } m_{LL}^2 > m_{RR}^2 \text{ then } \cos(2\tilde{\alpha}) = -\frac{1}{1+\tilde{\alpha}^2} < 0$$

$$2. \text{ if } m_{LL}^2 < m_{RR}^2 \text{ then } \cos(2\tilde{\alpha}) = +\frac{1}{1+\tilde{\alpha}^2} > 0$$

The cross section for $\tilde{\chi}_1\tilde{\chi}_1$ production depends on the mixing angle $\cos 2\tilde{\alpha}$ ([6] and references therein) and $\cos \tilde{\alpha}$ can be accurately determined, with a two-fold ambiguity, via the cross section for $\tilde{\chi}_1$ production with polarized beams or via a polarization asymmetry $A_{Pol} = (\sigma_L - \sigma_R)/(\sigma_L + \sigma_R)$.

In the following we discuss the polarization from $\tilde{\chi}_1$ decays $P_{\tilde{\chi}_1!}$. The tau polarization P_τ can be derived from the energy distribution of the decay products of the tau lepton.

2.2 polarization from $\tilde{\chi}_i$ decays

In this section we study the polarization P_τ from the decays

$$\tilde{\chi}_1! \tilde{\chi}_i^0; \text{ and } \tilde{\chi}_2! \tilde{\chi}_i^0; \quad i=1, \dots, 4 \quad (7)$$

taking into account a general neutralino mixing in the MSSM [7].

The tau polarization for (7) is given by (using the narrow width approximation) [3]:

$$P_{\tilde{\chi}_1!} = \frac{(a_{1i}^R)^2 - (a_{1i}^L)^2}{(a_{1i}^R)^2 + (a_{1i}^L)^2}; \text{ and } P_{\tilde{\chi}_2!} = \frac{(a_{2i}^R)^2 - (a_{2i}^L)^2}{(a_{2i}^R)^2 + (a_{2i}^L)^2}; \quad (8)$$

with

$$a_{1i}^{L,R} = \cos \tilde{\alpha} a_{Li}^{L,R} + \sin \tilde{\alpha} a_{Ri}^{L,R}; \text{ and } a_{2i}^{L,R} = \sin \tilde{\alpha} a_{Li}^{L,R} + \cos \tilde{\alpha} a_{Ri}^{L,R}; \quad (9)$$

where the coefficients a_{ij}^k are defined by the Lagrangian

$$\mathcal{L} = \sum_{i=1,2} \tilde{\chi}_i (P_L a_{ij}^R + P_R a_{ij}^L) \tilde{\chi}_j^0; \quad (10)$$

in the neutralino basis $(\tilde{B}, \tilde{W}_3, \tilde{H}_1, \tilde{H}_2)$:

$$a_{Lj}^R = -\frac{g}{2m_W \cos \tilde{\alpha}} N_{j3}; \quad a_{Rj}^L = a_{Lj}^R \quad (11)$$

$$a_{Lj}^L = +\frac{g}{2} [N_{j2} + N_{j1} \tan \tilde{\alpha}]; \quad a_{Rj}^R = -\frac{2g}{2} N_{j1} \tan \tilde{\alpha}; \quad (12)$$

Taking into account a general mixing in the neutralino sector we derive the polarization:

$$P_{\tilde{\chi}_1!} = \frac{(4 - x_W^2) - (4 + x_W^2 - 2y_h^2) \cos 2\tilde{\alpha} + 2(2 + x_W)y_h \sin 2\tilde{\alpha}}{(4 + x_W^2 + 2y_h^2) - (4 - x_W^2) \cos 2\tilde{\alpha} + 2(2 - x_W)y_h \sin 2\tilde{\alpha}}; \quad (13)$$

And for the case $m_{LL}^2 > m_{RR}^2$ the formula takes the form :

$$= \frac{(4 - x_W^2) - \frac{1}{1+\tilde{\alpha}^2} + (4 + x_W^2 - 2y_h^2) - 2(2 + x_W)y_h}{(4 + x_W^2 + 2y_h^2) - \frac{1}{1+\tilde{\alpha}^2} + (4 - x_W^2) - 2(2 - x_W)y_h}; \quad (14)$$

where the mixing angle $\tilde{\alpha}$ i.e. $\tilde{\alpha}$ is given by (3) and $P_{\tilde{\gamma}2}$ can be obtained from eq. (13) by changing the sign of $\cos 2\tilde{\alpha}$ and $\sin 2\tilde{\alpha}$.

The coefficient x_W contains the complete contribution from the gaugino components, y_h is a combination of the factors of the Yukawa coupling and the complete contribution from the higgsino components x_h .

$$x_W = \frac{\tan \beta N_{11} + N_{12}}{\tan \beta N_{11}} \quad (15)$$

$$y_h = \frac{\frac{1}{\cos \beta} \frac{m}{m_W} N_{13}}{\tan \beta N_{11}} = : \frac{1}{\cos \beta} \frac{m}{m_W} x_h : \quad (16)$$

One sees from eq. (13) that the transformation between the two cases $m_{LL}^2 > m_{RR}^2$, whose hierarchy is motivated by the MSSM as far as no additional D terms have to be included, and $m_{LL}^2 < m_{RR}^2$ lead only to an exchange of $P_{\tilde{\gamma}1}$ & $P_{\tilde{\gamma}2}$.

2.2.1 $\tan \beta$ dependence of the polarization

With the coefficients x_W and x_h in eqn. (13) (14) the complete $\tan \beta$ dependence from the neutralino sector has been separated and the coefficient y_h shows the interplay between the Yukawa coupling and the $\tilde{\chi}_i^0$ higgsino admixture. All components of (15), (16) are given explicitly as function of the fundamental MSSM parameters in [2] and in an approximation, which is valid for high values of $\tan \beta$ in [7].

One can summarize that the dependence on $\tan \beta$ of $P_{\tilde{\gamma}1,2}$ is given in a three-fold way:

1. by the $\tan \beta$ dependence of the mixing angle $\tilde{\alpha}$ (3) and the off-diagonal term in (1);
2. by the coefficients x_W and x_h , which corresponds to the $\tan \beta$ dependence of the neutralino sector,
3. by the coefficient y_h which corresponds to the $\sim$ Yukawa coupling $\frac{1}{\cos \beta} \frac{m}{m_W}$.

One sees clearly from (16) that for a given mixing angle $\tilde{\alpha}$ or only the coefficient y_h contains a strong dependence on $\tan \beta$ of $P_{\tilde{\gamma}1}$ from the Yukawa coupling. Obviously a sufficient large higgsino admixture part x_h has to be for that.

The $\tan \beta$ dependence of x_W and x_h is weak since the neutralino mass eigenvalues $m_{\tilde{\chi}_i^0}$ as well as the components of the eigenvectors N_{ij} become in the high $\tan \beta$ approximation $f(1 + \text{const} \frac{1}{\tan \beta})$. Therefore it is a good approximation to take all neutralino mixing contributions in the high $\tan \beta$ approximation.

$\tilde{\chi}_1^0$	$P_{\tilde{\chi}_1^!}$	$P_{\tilde{\chi}_1^!}^{-1}$	$P_{\tilde{\chi}_1^!}^{+0}$	$P_{\tilde{\chi}_2^!}$	$P_{\tilde{\chi}_2^!}^{-1}$	$P_{\tilde{\chi}_2^!}^{+0}$
a) N_{11}	$\frac{p}{5} \frac{1+2+5}{1+2+3}$	$\frac{3}{5}$	+ 1	$\frac{p}{5} \frac{1+2+5}{1+2+3}$	$\frac{3}{5}$	1
b) $C_{11}N_{11}$ + $C_{12}N_{12}$	$\frac{(4+x_W^2)}{(4+x_W^2)} \frac{p}{1+2+(4+x_W^2)}$	$\frac{4+x_W^2}{4+x_W^2}$	+ 1	$\frac{(4+x_W^2)}{(4+x_W^2)} \frac{p}{1+2+(4+x_W^2)}$	$\frac{4+x_W^2}{4+x_W^2}$	1
c) N_{13}	$\frac{p}{1+2}$	0	1	$\frac{p+1}{1+2}$	0	+ 1

Table 1: polarization for extrem a for the mixing angle and speci c neutralino mixing.

2.3 Limiting cases: extrem a for the mixing angle

Assuming no constraints for the mass parameters one can vary the mixing angle $\tilde{\alpha}$ from 0 to π , i.e. from $0 \leq \tilde{\alpha} \leq \pi$, and we get for these extrem a the following expressions:

$$P_{\tilde{\chi}_1^!}^{-1} = \frac{4+x_W^2+2(2+x_W)y_h}{4+x_W^2+2(2+y_h-x_W)y_h} \quad (17)$$

$$P_{\tilde{\chi}_1^!}^{+0} = \frac{4+y_h^2}{4+y_h^2} \quad (18)$$

$$P_{\tilde{\chi}_2^!}^{-1} = \frac{4+x_W^2+2(2+x_W)y_h}{4+x_W^2+2(2+y_h-x_W)y_h} \quad (19)$$

$$P_{\tilde{\chi}_2^!}^{+0} = \frac{x_W^2+y_h^2}{x_W^2+y_h^2} \quad (20)$$

The formulae for the tau polarization even simplifies for speci c neutralino mixing cases. We give the corresponding expressions in Table 1 for the processes $\tilde{\chi}_1^! \rightarrow \tilde{\chi}_1^0$ and $\tilde{\chi}_2^! \rightarrow \tilde{\chi}_1^0$. The conventions are chosen so that in a no mixing case, i.e. $\tilde{\chi}_1^! \rightarrow \tilde{\chi}_R < \tilde{\chi}_2^! \rightarrow \tilde{\chi}_L$, and when only a pure U(1) gauge coupling interacts between $\tilde{\chi}^0$, the polarization $P_{\tilde{\chi}_1^! \rightarrow \tilde{\chi}_R} = +1$ and $P_{\tilde{\chi}_2^! \rightarrow \tilde{\chi}_L} = -1$. Studying the polarization of the antiparticle $\tilde{\chi}^+$ one has to take into account that the SUSY partner of $\tilde{\chi}_{L,R}^+$ is $\tilde{\chi}_{R,L}^+$, so that $P_{\tilde{\chi}_R^! \rightarrow \tilde{\chi}_L^+} = -1$ and $P_{\tilde{\chi}_L^! \rightarrow \tilde{\chi}_R^+} = +1$. We show in Fig. 1 representative examples for illustration.

The corresponding plot for case a), Table 1, is given in Fig. 1a. The asymptotic limit $\frac{3}{5}$ for $\tilde{\alpha} \rightarrow \pi$ can clearly be seen.

The corresponding plot for case b), Table 1, is given in Fig. 1b. The asymptotic limit depends now on the gaugino parameters and since the wino fraction x_W^2 is less than 1 the polarization $P_{\tilde{\chi}_1^!}$ gets closer to 1 and will always be higher than the asymptotic value $\frac{3}{5}$ of the pure bino case a).

In the pure higgsino case c), Table 1, the no mixing case shows a helicity flipping behaviour, $P_{\tilde{\chi}_1^!} = -1$ and $P_{\tilde{\chi}_2^!} = +1$, which is the typical feature of the Yukawa couplings.

To give a feeling even for the neutralino mixing effects we plot in Fig. 2a the polarizations $P_{\tilde{\chi}_1^!}$, $P_{\tilde{\chi}_2^!}$ for the mixed case $x_W = 0.8$ and $y_h = 0.6$. The polarizations even interchange for a speci c mixing angle.




Figure 1: The dependence of the tau polarizations $P_{\tilde{\tau}_1}$ and $P_{\tilde{\tau}_2}$ from the mixing angles ξ for a pure bino case (left) with the neutralino mixing variables $x_W = 1, y_h = 0$ and for a pure gaugino case (right) with $x_W = 0.5, y_h = 0$.

So if the fraction of the higgsino mixing part is small compared to the gaugino mixing one expects $P_{\tilde{\tau}_1}$ close to unity and $P_{\tilde{\tau}_2}$ variable in a large range, eqn. (17)–(20). This is the case for all the SPS scenarios [8] as shown in the Table 2.

In order to demonstrate the $\tan \beta$ dependence coming from the interplay between the Yukawa coupling and the higgsino admixture of the LSP we show in Fig. 2b the polarizations $P_{\tilde{\tau}_1}$ and $P_{\tilde{\tau}_2}$ for the case with a rather large higgsino admixture and set $\mu = 0$, i.e. $\tilde{\chi}_1^0 = \tilde{\chi}_R$ and $\tilde{\chi}_2^0 = \tilde{\chi}_L$.

Parameter Point	$\tan \beta$	Polarization		slopes	
		$P_{\tilde{\tau}_1}$	$P_{\tilde{\tau}_2}$	$d(P_{\tilde{\tau}_1})/d(\tan \beta)$	$d(P_{\tilde{\tau}_2})/d(\tan \beta)$
SPS 1a	10	98.1 %	-50 %	0.3%	5.0%
SPS 1b	30	97.0 %	-40 %	0.1%	1.6%
SPS 3	10	99.2 %	-80 %	0.1%	2.4%
SPS 4	50	99.6 %	-62 %	+0.1%	2.0%
SPS 5	5	97.8 %	-60 %	0.6%	7.0%
SPS 6	10	99.0 %	-65 %	0.2%	4.0%

Table 2: Tau polarization and $\tan \beta$ slopes for the SPS scenarios [8]. The program ISAJET 7.58 [9] has been used for the parameter evaluation. The SPS 2 focuspoint point scenario does not have a stable slope.

In order to get an impression in which ranges of the parameter space the tau polarizations from $\tilde{\tau}_{1,2}$ decays may have large variations, we show in Fig. 3 the polarizations $P_{\tilde{\tau}_{1,2}}$ as functions of M_2 and μ . The other relevant parameters have been chosen to $\tan \beta = 40, M_L = 300 \text{ GeV}, M_E = 150 \text{ GeV}, A = 254.2 \text{ GeV}$. Both plots show that one could get high values for the polarization for a large region of the M_2 parameter space. In particular in the higgsino-like region with $\mu < M_2$ the polarization $P_{\tilde{\tau}_1}$ is very variable.




Figure 2: The dependence of the tau polarizations $P_{\tilde{\tau}_1}$ and $P_{\tilde{\tau}_2}$ from the mixing angles (left) for a mixed case with the neutralino mixing variables $x_w = 0.8$, $y_h = 0.6$ and as function of $\tan \beta$ (right) where the mixing angle is chosen to be small $\beta \rightarrow 0$. In this case $\tilde{\gamma}_1 \rightarrow \tilde{\chi}_R, \tilde{\gamma}_2 \rightarrow \tilde{\chi}_L$.

3 M SSM parameter determination for high $\tan \beta$

3.1 Parameter determination in the $\tilde{\gamma}_{1,2}$ sector

We assume that one can measure at a high LHC the masses with rather high accuracy of about % level via mass threshold scans.

We study the light system $e^+e^- \rightarrow \tilde{\gamma}_1\tilde{\gamma}_1$ and determine the mixing angle $\cos \tilde{\alpha}$ via polarized cross sections $(\sigma_{LR}^+ - \sigma_{RL}^+) / (\sigma_{LR}^+ + \sigma_{RL}^+) = \cos(2\tilde{\alpha})$ or via the asymmetry $A_{pol} = (\sigma_{LR} - \sigma_{RL}) / (\sigma_{LR} + \sigma_{RL})$. If one wants to derive $\tilde{\alpha}$ unambiguously one would also need $(\sigma_{e^+e^- \rightarrow \tilde{\gamma}_1\tilde{\gamma}_2} - \sigma_{e^+e^- \rightarrow \tilde{\gamma}_2\tilde{\gamma}_1}) / (\sigma_{e^+e^- \rightarrow \tilde{\gamma}_1\tilde{\gamma}_1} + \sigma_{e^+e^- \rightarrow \tilde{\gamma}_2\tilde{\gamma}_2}) = \sin(2\tilde{\alpha})$, which is more tricky because of the difficult reconstruction of the $\tilde{\gamma}_1, \tilde{\gamma}_2$ decay products from the $\tilde{\gamma}_2$ decays.

We determine the mixing angle via measuring the production $e^+e^- \rightarrow \tilde{\gamma}_1\tilde{\gamma}_1^+$ in the configuration (RL), $P(e^-) = +80\%$, $P(e^+) = -60\%$, since in this case the worse background from W^+W^- is strongly suppressed and in contrary the signal is enhanced. From the rates R_{RL} we could derive the mixing angle rather accurately, $\cos \tilde{\alpha} = 0.15 \pm 0.01$, Fig.4, if we measure $R_{RL}(\tilde{\gamma}_1\tilde{\gamma}_1^+) = 112 \text{ fb}$ at $\sqrt{s} = 500 \text{ GeV}$ and assume that a statistical error of $\pm 1\%$ was taken into account. In Table 3 we list the corresponding cross sections (σ_{ij}) for $\sqrt{s} = 500 \text{ GeV}$ and $\sqrt{s} = 800 \text{ GeV}$.

In principle one can alternatively also derive the mixing angle via measuring the polarization asymmetry A_{pol} of the production process. Taking rates with left polarized electrons leads, however, to an enhancement of the strong W^+W^- background and a tricky analysis would be needed to get the wanted experimental information.

Generically the cross section is a quadratic polynomial in $\cos(2\tilde{\alpha})$. With a suitably high degree of beam polarization, however, the quadratic terms are suppressed and only one solution survives.




Figure 3: Contourplots in the M_2 - μ plane for $P_{\tilde{\tau}_1}$ and $P_{\tilde{\tau}_2}$.




Figure 4: Mixing angle $\cos \tilde{\theta} = 0.15 \pm 0.01$ via measurement of the polarized cross section $(e^+ e^- \rightarrow \tilde{\tau}_1 \tilde{\tau}_1)$ at $\sqrt{s} = 500$ GeV and $P(e^-) = +80\%$, $P(e^+) = -60\%$ and on the assumption that 1 standard deviation as statistical error was taken into account.

3.2 Measurement of the polarization

We have computed the cross sections and distributions for the final state particle processes with all spin correlations taken into account:

$$e^+e^- \rightarrow \tilde{\gamma}_1^+ \tilde{\gamma}_1^- \text{ with } \tilde{\gamma}_1 \rightarrow_{L,R} \tilde{\gamma}_1^0 \text{ and } \tilde{\gamma}_1 \rightarrow_{L,R} \tau^\pm \quad (21)$$

In order to measure the polarization of the $\tilde{\gamma}$'s one has to study their decays into leptons or mesons and to kinematically the energy distributions of the decay products [3]. The decays of polarized $\tilde{\gamma}$ to π - and η -mesons are implemented into ComPHEP and are cross checked with TAUOLA [10].

The calculations are performed by means of the ComPHEP V41 [11] with implemented MSSM and mSUGRA models. In order to estimate an accuracy of polarization measurements we have generated a number of unweighted events which correspond to the production cross section with the corresponding decay branching fractions and the overall registration efficiency of signal events:

$$N = (e^+e^- \rightarrow \tilde{\gamma}_1^+ \tilde{\gamma}_1^-) \text{Br}(\tilde{\gamma}_1 \rightarrow \tilde{\gamma}_1^0) \text{Br}(\tilde{\gamma}_1 \rightarrow \tau^\pm) L_e \quad (22)$$

Based on the results of MC analysis given in [3] we assumed 30% for the efficiency and we assume 500 fb^{-1} for a luminosity. So in total about 3300 unweighted events have been simulated.

The pion energy distribution is shown in the Fig. 5 left. The distribution is described by the formula presented in a slightly different form as that one given in [3].

$$\begin{aligned} & \frac{1}{dy} (e^+e^- \rightarrow \tilde{\gamma}_1^+ \tilde{\gamma}_1^- \rightarrow \tilde{\gamma}_1^0 \tilde{\gamma}_1^0 \rightarrow \pi^+ \pi^-) \\ &= \frac{1}{x_{\text{max}} - x_{\text{min}}} \left((1 - P_{\tilde{\gamma}_1}) \log \frac{x_{\text{max}}}{x_{\text{min}}} + 2P_{\tilde{\gamma}_1} y \left(\frac{1}{x_{\text{min}}} - \frac{1}{x_{\text{max}}} \right); 0 < y < x_{\text{min}} \right) \\ & \quad (1 - P_{\tilde{\gamma}_1}) \log \frac{x_{\text{max}}}{y} + 2P_{\tilde{\gamma}_1} \left(1 - \frac{y}{x_{\text{max}}} \right); x_{\text{min}} < y \end{aligned} \quad (23)$$

where $y = \frac{2E_{\pi}^{\text{CMS}}}{\sqrt{s}}$, $x_{\text{min}} = \frac{2E_{\pi}^{\text{min}}}{\sqrt{s}}$, $x_{\text{max}} = \frac{2E_{\pi}^{\text{max}}}{\sqrt{s}}$, $E_{\pi}^{\text{max,min}} = \frac{E_{\tilde{\gamma}} - p_{\tilde{\gamma}}}{2}$, $\tilde{\gamma} = 1 - \frac{4M_{\tilde{\gamma}}^2}{s}$,
 $E_{\tilde{\gamma}} = \frac{M_{\tilde{\gamma}}^2 + M_{\tilde{\gamma}_1^0}^2 + M_{\pi}^2}{2M_{\tilde{\gamma}_1}}$, $p_{\tilde{\gamma}} = \frac{p_{\tilde{\gamma}_1^0}^2 + M_{\pi}^2}{E_{\tilde{\gamma}_1^0}^2 - M_{\pi}^2}$.

The fit of the energy distribution by the above formula (23) gives for the tau polarization $P_{\tilde{\gamma}_1}$ about 57% $\pm$ 3%, compared to the theoretical value of polarization for the reference point of 56%.

(P(e^-); P(e^+)))	$\sqrt{s} = 500 \text{ GeV}$		$\sqrt{s} = 800 \text{ GeV}$	
	($e e^+ \rightarrow \tilde{\gamma}_1 \tilde{\gamma}_1$)	($e e^+ \rightarrow \tilde{\gamma}_1 \tilde{\gamma}_1$)	($e e^+ \rightarrow \tilde{\gamma}_1 \tilde{\gamma}_2$)	($e e^+ \rightarrow \tilde{\gamma}_2 \tilde{\gamma}_2$)
unpolarized	48.6 fb	29.7 fb	0.2 fb	12.0 fb
(-0.8; 0)	25.6 fb	15.9 fb	0.3 fb	18.3 fb
(+0.8; 0)	71.6 fb	43.5 fb	0.2 fb	5.7 fb
(-0.8; 0.6)	31.6 fb	19.8 fb	0.4 fb	28.8 fb
(+0.8; -0.6)	112.1 fb	68.1 fb	0.3 fb	6.7 fb

Table 3: Cross sections for the reference scenario with polarized beams. The pairs $\tilde{\gamma}_1 \tilde{\gamma}_2$, although kinematically accessible at $\sqrt{s} = 500 \text{ GeV}$, lead to rates less than 0.1 fb.




Figure 5: Left: The pion energy histogram for the unweighted events and the fit by the distribution formula (23). The total event number is 3300 events which corresponds to an integrated luminosity of $L = 500 \text{ fb}^{-1}$. For beam polarization we choose $P_{e^-} = +80\%$, $P_{e^+} = -60\%$. Right: The determination of $\tan\beta = 40 \pm 2$ as function of the measured polarization $P_{\tilde{\tau}_1 \rightarrow \tau} = 57 \pm 3\%$.

The presented example is meant as an illustration only. Measurements of the lepton decay mode together with its subsequent decays may help to get even a better accuracy.

3.3 Determination of $\tan\beta$ and the A parameter

In the last section we have shown that $P_{\tilde{\tau}_1 \rightarrow \tau}$ could be measured within about 5% accuracy. We derive the value of $\tan\beta$ via inversion of (13) for the calculated values of x_W and x_h and the measured mixing angle $\cos\tilde{\alpha}$, see Fig. 5 right, and we determine in our scenario the high $\tan\beta$ with an error of about 5%:

$$\tan\beta = 40 \pm 2 \quad (24)$$

In the case that also the heavier mass of $\tilde{\tau}_2$ can be determined in the experiment, we can even determine the parameter A without assuming anything about the SUSY breaking scheme and GUT relations:

$$A = \frac{1}{m} \left(\frac{1}{2} (m_{\tilde{\tau}_1}^2 - m_{\tilde{\tau}_2}^2) \sin(2\tilde{\alpha}) + m \tan\beta \right) \quad (25)$$

With an error of about 5% in $\tan\beta$ and of about 5% in $m_{\tilde{\tau}_2}$ one could derive A with an accuracy about 8%. For the application of this method on the $\tilde{b}$ and $\tilde{t}$ sector and a detailed simulation study see [7].

4 Conclusions

Tau polarization from stau decays might be an important variable for the determination of the fundamental MSSM parameters. It serves as a 'bridge' between chargino/neutralino and stau sectors.

The stau mixing angles can be precisely determined via polarized rates and in case that the decay neutralino has a suitable higgsino admixture, the study of $P_{\tilde{\chi}_1^0}$ leads to an accurate determination of $\tan\beta$ even in the case of high $\tan\beta$. For these procedure it is enough to measure only the light system: $m_{\tilde{\chi}_1}$ and $(e^+e^- \rightarrow \tilde{\chi}_1\tilde{\chi}_1)$. In case that also $m_{\tilde{\chi}_2}$ can be measured, even the parameter A_0 can be derived without an assumption about the underlying SUSY breaking scheme.

For a given example we explored $P_{\tilde{\chi}_1^0}$ and showed that one can determine simultaneously A_0 and $\tan\beta$, e.g. in the case of high $\tan\beta$, within 5% accuracy.

A systematic simulation of the process as well as the application of this method on the $\tilde{b}$ and $\tilde{t}$ sector will be presented in a following paper [7].

The work of E.B. and A.V. was partly supported by the INTAS 00-0679, CERN-INTAS 99-377, and RFBR 01-02-16710 grants. E.B. thanks the Humboldt Foundation for the Bessel Research Award and DESY for the kind hospitality.

References

- [1] J.L.Feng, M.E.Peskin, H.Murayama, X.Tata, Phys.Rev.D 52 (1995) 1418;
T.Tsukamoto, K.Fujii, H.Murayama, M.Yamaguchi, Y.Okada, Phys.Rev.D 51 (1995) 3153.
- [2] S.Y.Choi, J.Kalinowski, G.Moortgat-Pick and P.M.Zerwas, Eur.Phys.J.C 22 (2001) 563 [hep-ph/0108117];
S.Y.Choi, J.Kalinowski, G.Moortgat-Pick and P.M.Zerwas, Eur.Phys.J.C 23 (2002) 769 [hep-ph/0202039];
G.Moortgat-Pick, [hep-ph/0211039], to appear in the Proceedings of the 10th International Conference on Supersymmetry and Unification of Fundamental Interactions, SUSY02, DESY, Hamburg 2002.
- [3] M.M.Nojiri, Phys.Rev.D 51 (1995) 6281 [hep-ph/9412374];
M.M.Nojiri, K.Fujii and T.Tsukamoto, Phys.Rev.D 54 (1996) 6756 [hep-ph/9606370].
- [4] M.Guchait and D.P.Roy, Phys.Lett.B 535 (2002) 243 [hep-ph/0205015].
- [5] J.A.Aguilar-Saavedra et al., ECFA/DESY LC Physics Working Group Collaboration, [hep-ph/0106315].
- [6] A.Bartl, H.Eberl, S.Kraml, W.Majerotto and W.Porod, Z.Phys.C 73 (1997) 469 [hep-ph/9603410]; A.Bartl, H.Eberl, S.Kraml, W.Majerotto, W.Porod and A.Sopczak, Z.Phys.C 76 (1997) 549 [hep-ph/9701336]; S.Kraml, PhD Thesis, HEPHY Vienna, [hep-ph/9903257]; A.Bartl, H.Eberl, S.Kraml, W.Majerotto and W.Porod, Eur.Phys.J.direct C 2 (2000) 6 [hep-ph/0002115]; A.Bartl, K.Hidaka, T.Kemreiter and W.Porod, [hep-ph/0207186].
- [7] E.Boos, H.U.Martyn, G.Moortgat-Pick, M.Sachwitz, A.Sherstnev, P.M.Zerwas, hep-ph/0303110.
- [8] N.Ghodbane and H.U.Martyn, in Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics [hep-ph/0201233]; B.C.Allanach et al., in Proc. of the APS/DPF/DPB Summer Study on the Future of Particle Physics, Eur.Phys.J.C 25 (2002) 113 [hep-ph/0202233].

- [9] H Baer, F Paige, S Protopopescu and X Tata, hep-ph/0001086.
- [10] S Jadach, Z Was, R Decker, and JH Kuehn, Comp Phys Comm . 76 (1993) 361.
- [11] A Pukhov, E Boos, M Dubinin, V Ilyin, D Kovalenko, A Kryukov, V Savrin, S Shichanin and A Semenov, Report INP-M SU 98-41/542, [hep-ph/9908288]; A Semenov, Comp Phys Comm . 115 (1998) 124 and [hep-ph/0205020].