

Single Squark Production at TeV Scale p and e Colliders

Z.Z. Aydın, N.K. Aragoz and A.J. Yılmaz

Ankara University, Faculty of Engineering
Department of Engineering Physics
06100 Tandoğan, Ankara – Turkey

We calculate the total cross sections for squark productions at future p and e colliders in R_p violating MSSM. The λ_{ijk}^0 couplings offer several ways of producing single squarks in photon colliders. The results are discussed and compared with those of the existing colliders.

PACS number(s) : 14.80.Ly, 12.60.Jv.

I. INTRODUCTION

Supersymmetric particles can be produced singly in the R -parity violating version of the minimal supersymmetric standard model (R_p - MSSM). Although the production cross section is suppressed by R_p coupling constants the relevant processes have higher kinematical reaches compared with the case of pair (or associated) productions of superparticles through the R_p -conserving mechanisms. Also R_p violation leads to a totally different phenomenology and the detailed studies for the specific colliders (LEP2, HERA, TEVATRON) have already appeared in the literature [1].

On the other hand in recent years in addition to the existing colliders the possibilities of the realization of e , and p colliders have been proposed and discussed in detail [2]. Colliding the beam of high energy photons produced by Compton backscattering of laser photons on linac electrons with the beam of a proton ring is the idea leading to TeV scale high luminosity p colliders. Also e option has been included in the conceptual design of the linear colliders such as TESLA. Physics program of the photon colliders are studied in [3]. Also search for SUSY in polarized p collisions have been investigated in [4].

In searching the superpartners squarks (except the third generation ones) might be too heavy to be produced at HERA, LEP or TEVATRON, but sleptons are generally expected to be lighter than squarks so single slepton production would be interesting. Also pair production of sleptons via R_p conserving mechanisms might be closed kinematically. The s -channel slepton resonance production via R_p interactions in e^+e^- collisions through $e^+e^- \rightarrow \tilde{l}^+ \tilde{l}^-$, and in pp collisions through $pp \rightarrow \tilde{l}^+ \tilde{l}^-$, $pp \rightarrow \tilde{l}^+ \tilde{l}^+$ have been examined in the literature [5]. In e^+p collisions at HERA squark resonance productions via $e^+d_R^* \rightarrow u_L^j$ ($u^j = u; c; t$), $e^+u_L^j \rightarrow d_R^*$, ($d^* = d; s; b$) have been investigated in [6]. Although HERA, LEP, FERMILAB and LHC should be sufficient to check the low energy SUSY however experiments at all possible types of colliding beams would be useful to explore the new physics around the TeV scale, and hence with their additional polarization facilities future e and p colliders might play complementary roles to the existing accelerators.

In this letter we mainly focus on the single squark productions via R_p - MSSM interactions at photon-proton and photon-electron collisions as an alternative to the above-mentioned s -channel resonance production.

II. SINGLE SQUARK PRODUCTION WITH AN ESCAPING NEUTRINO AT GAMMA-PROTON COLLIDERS

R_p violating superpotential terms can be written in four-component Dirac notation as

¹Abant İzzet Baysal University, Physics Department, Bolu – Turkey

$$\begin{aligned}
L_{R_p} = & \sum_{i,j,k} \left[\tilde{e}_{iL} e_{kR} e_{jL} + e_{jL} e_{kR} \tilde{e}_{iL} + e_{kR}^2 (\tilde{e}_{iL})^c e_{jL} + \tilde{e}_{jL} e_{kR} e_{iL} \right. \\
& e_{iL} e_{kR} \tilde{e}_{jL} + e_{kR}^2 (\tilde{e}_{jL})^c e_{iL} + \sum_{i,j,k} \left[\tilde{e}_{iL} d_{kR} d_{jL} + \tilde{d}_{jL} d_{kR} \tilde{e}_{iL} \right. \\
& + d_{kR}^2 (\tilde{e}_{iL})^c d_{jL} + e_{iL} d_{kR} u_{jL} + \tilde{u}_{jL} d_{kR} e_{iL} + d_{kR}^2 (\tilde{e}_{iL})^c u_{jL} + \sum_{i,j,k} \left[\right. \\
& \left. \left. \tilde{u}_{iR} d_{kR} \tilde{d}_{jR} + \tilde{d}_{jR} e_{kR} u_{iR} + d_{kR}^2 (\tilde{u}_{iR})^c d_{jR} + h.c. \right] \right] \quad (1)
\end{aligned}$$

where i, j, k are the generation indices. In photon-proton collisions a single slepton or a single squark production is possible. Single productions of charged sleptons (stau) and sneutrinos have been considered in a previous letter [7]. Here let us examine the single squark production associated with an escaping neutrino $p \rightarrow \tilde{d}_R^j L X$. We note that $p \rightarrow \tilde{d}_L^j L X$ is also possible from the second 0 term in Eq.(1). One of the relevant subprocesses, $d \rightarrow \tilde{d}_R^j L$, proceeds via the s-channel d-quark, t-channel squark exchanges. The invariant amplitude in four-component Dirac notation (which could be written equally in two-component Weyl language) is

$$M = e_d g_e \epsilon^\mu(k) u(p^0) \bar{Q}_\mu u(p) \quad (2)$$

$$Q = \frac{1}{2} (1 - \gamma_5) \left[\frac{\not{k} + \not{p}}{\hat{s} - m_d^2} + \frac{(\not{p} - \not{p}^0 + k^0)}{\hat{t} - m_d^2} \right] \quad (3)$$

where $g_e = \frac{P}{4}$, $\epsilon^\mu(k)$ is the photon polarisation, q_d is the charge factor and k, p, k^0 and p^0 are the four momenta of the photon, d-quark in the proton, squark and escaping neutrino respectively. The differential cross-section for the subprocess is given by

$$\frac{d\hat{\sigma}}{d\hat{t}} = \frac{1}{16 \hat{s}^2} M^2 \quad (4)$$

and to obtain the total cross section for the subprocess $d \rightarrow \tilde{d}_R^j L$ one should perform integration over $\hat{t}$. The total cross-section for the main process $p \rightarrow \tilde{d}_R^j L X$ is obtained after the integration of $\hat{\sigma}$ over the quark and photon distributions. For this purpose we make the following change of variables: first expressing $\hat{s}$ as $\hat{s} = x_1 x_2 s$ where $\hat{s} = s_q$, $s = s_{ep}$, $x_1 = E_q/E_e$, $x_2 = E_q/E_p$ and furthermore calling $x_1 = x_1 x_2$, $x_2 = x$ then one obtains $dx_1 dx_2 = dx/dx$. The limiting values are $x_{1,max} = 0.83$ in order to get rid of the background effects in the Compton backscattering, particularly e^+e^- pair production in the collision of the laser with the high energy photon in the conversion region, $x_{1,min} = 0$, $x_{2,max} = 1$, $x_{2,min} = 0.83$, $\hat{s}_{min} = m_d^2$. Then we can write the total cross-section as :

$$\begin{aligned}
& \int_{0.83}^1 \int_{m_d^2/s}^1 d\hat{s} \int_{0.83}^1 \frac{dx}{x} f_q(x) f(\gamma) \quad (5)
\end{aligned}$$

where $f_q(x)$ is the distribution of down-quarks inside the proton [8]

$$f_q(x) = 0.67 x^{-0.6} (1 - x^{1.5})^{4.5} \quad (6)$$

and $f(\gamma)$ is the energy spectrum of the high energy realphotons (Ginzburg et.al. in ref.[2])

$$f(\gamma) = \frac{1}{D(\gamma)} \left[1 - \gamma + \frac{1}{1 - \gamma} - \frac{4\gamma}{(1 - \gamma)^2} + \frac{4\gamma^2}{(1 - \gamma)^3} \right] \quad (7)$$

with $\gamma = \frac{E}{E_e}$; $D(\gamma) = 4.8$, and $\gamma_{max} = 0.83$. Q^2 independent proton structure function used above is satisfactory for the present analysis; use of the Q^2 dependent quark distributions $f(x; Q^2)$ does not alter the results significantly.

The current upper bounds on the relevant R_p coupling constants are $g_{311}^0 = g_{312}^0 = g_{313}^0 = 0.11 \frac{m_{\tilde{g}}}{100 \text{ GeV}}$ [12], consequently we may look for the productions of $\tilde{d}_R$, $\tilde{s}_R$ or $\tilde{b}_R$. The results of the numerical integration for $p \rightarrow \tilde{q}_R L X$, where $\tilde{q}_R$ being either one of the above three down-type squarks is plotted in Fig 1.

As can be seen from the figure the total cross-section for a $\tilde{d}$ mass of 400 GeV is about 0.1 pb. Hence around one thousand events per running year can be seen at HERA+LC up to $\tilde{d}$ masses of 400 GeV. For comparison we note

that for the resonant production of squarks of masses up to 200 GeV at the HERA the total cross-section is 0.1 - 1 pb (see E Perez et al. in ref.[6]). On the other hand, if one uses the Weizsacker-Williams approximations for the quasi-real photon distribution at the HERA machine a similar $p \rightarrow \tilde{q}_R^i \gamma_L^k X$ process is possible but with an almost hundred times smaller cross-section since WW-spectrum is much softer than the real γ -spectrum. Clearly for the HERA machine resonant production of the particles in R-parity violating MSSM is the dominant process.

Signature: In models with R-parity violation the LSP is unstable, which leads to signatures which differ strongly from the characteristic missing energy signals in usual MSSM. In our case the produced squark ($\tilde{q}_R$ or $\tilde{b}$) will decay either by direct $\tilde{R}_p$ couplings into matter fermions $\tilde{q}_R \rightarrow u$ (or d) through the same coupling $\tilde{g}_{ijk}^0$ leading to the signals 1 lepton + 1 jet + $\cancel{E}_T$ or 1 jet + $\cancel{E}_T$ (however there are large SM backgrounds); or by cascading through MSSM to the LSP which in turn decays via $\tilde{R}_p$: $\tilde{q}_R \rightarrow \tilde{q}_R^0 \rightarrow \tilde{q}_1^0 \rightarrow u$ or d leading to the signals 2 leptons + 1 jet + $\cancel{E}_T$ or 1 lepton + 3 jets + $\cancel{E}_T$. Within MSSM it is forbidden for $\tilde{q}_R$ to decay into charginos. These LSP decays depend on couplings $\tilde{g}^0$ but also on the supersymmetry parameters M_2 , and $\tan \beta$ (see E Perez et al. ibid). In the special cases involving only the operators $L_i Q_j D_j$ the LSP can also dominantly decay via the radiative process $\tilde{q}_1^0 \rightarrow \gamma + [9]$. On the other hand the above $\tilde{R}_p$ decays of the produced squark have the following partial decay widths :

$$\Gamma(\tilde{q}_R \rightarrow u + q_d) = \frac{m_q (\tilde{g}_{313}^0)^2}{16} \quad (8)$$

$$\Gamma(\tilde{q}_R \rightarrow q_u) = \frac{m_q (\tilde{g}_{313}^0)^2}{16} \quad (9)$$

thus they are equal to each other.

III. SINGLE SQUARK PRODUCTION ASSOCIATING TAU LEPTON AT GAMMA-PROTON COLLIDERS

$\tilde{R}_p$ Yukawa couplings $\tilde{g}_{ijk}^0$ offer also the opportunity to produce a single squark with an associating lepton in gamma-proton collisions. The relevant subprocess $u \rightarrow \tilde{q}_R^i e_L^{+k}$, proceeds via the s-channel u-quark, t-channel squark and u-channel lepton exchanges. The invariant amplitude is given as in Equ.(2), with Q as,

$$Q = \frac{1}{2} (1 - \gamma_5) e_u \frac{\cancel{k} + \cancel{p}}{\hat{s} - m_u^2} + e_d \frac{(\cancel{p} - \cancel{p}^0 + k^0)}{\hat{t} - m_{\tilde{q}}^2} + e \frac{\cancel{p}^0 \cancel{k}}{\hat{u} - m^2} \quad (10)$$

$\tilde{R}_p$ coupling constants $\tilde{g}_{311}^0 = \tilde{g}_{312}^0 = \tilde{g}_{313}^0 = 0.11 \frac{m_{\tilde{q}}}{100 \text{ GeV}}$ favor the productions of $\tilde{q}_R, \tilde{s}_R$ or $\tilde{b}_R$ respectively, associating the τ -lepton. Since instead of a heavy sparticle, a τ -lepton is exchanged in the u-channel the cross section for squark production becomes considerably bigger than the case analysed in the previous section. The details of the calculation is similar and taking again the simplistic Q^2 independent distribution of up-quarks inside the proton as [8]

$$f_q(x) = 2.751 x^{0.412} (1 - x)^{2.69} \quad (11)$$

the results of the numerical integration for $p \rightarrow \tilde{q}_R^i \gamma_L^k X$ is plotted in Fig 2. Similarly d-quark in the proton gives rise to the production $d \rightarrow \tilde{q}_R^i \gamma_L^k X$, which contributes roughly half of the u-quark distribution, and leads to the same signature. Hence around 10^3 events per running year can be seen at HERA+LC up to $\tilde{q}_R$ masses of 300 GeV. For single squark production at LEP and hadron colliders see [1],[5] and [9-10].

Signature: Considering leptonic τ -decay via SM-W boson to $\tau \rightarrow \mu + W \rightarrow \mu + \gamma + \nu_\mu$ and also taking into account the squark decays listed above we hence point out that the signature might be 2 leptons + 1 jet + $\cancel{E}_T$ or 1 leptons + 1 jet + $\cancel{E}_T$ or 3 leptons + 2 jets + $\cancel{E}_T$ or 3 leptons + $\cancel{E}_T$.

IV. SINGLE SQUARK PRODUCTION AT ELECTRON-PHOTON COLLIDERS

The same $\tilde{R}_p$ Yukawa couplings $\tilde{g}_{ijk}^0$ (i.e. eleventh term in Equ.(1)) offer the opportunity to produce squarks in photon-electron collisions. One of the relevant subprocess $e \rightarrow \gamma_{jL} \tilde{q}_{kR}$, proceeds via electron (left-handed)

exchange in s-channel, squark exchange in the t-channel and quark (right-handed) exchange in u-channel. The invariant amplitude in two-component MSSM language is given as

$$M = g_e^0 (k + p) + (p^0) Q (p) \quad (12)$$

$$Q = e_d \frac{(k + p)}{s - m_e^2} + e_t \frac{(p - p^0 + k^0)}{t - m_{\tilde{q}}^2} + e_s \frac{(p^0 - k)}{u - m_q^2} \quad (13)$$

where $u = (p^0) (p)$ is the Weyl spinor for the right-handed quark (left-handed electron). Taking $\alpha_{132}^0 = 0.33$ (present upper bound) the total cross section for the particular process $e^+ e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$ may be written as

$$\sigma(s; e^+ e^-) = \int_{t_{min}}^{t_{max}} \frac{1}{16 s^2} |M|^2 dt \quad (14)$$

where

$$t_{max} = t_{min} = \left(\frac{m_{\tilde{t}}^2 - m_{\tilde{q}}^2}{2s} \right)^2 \quad \frac{p - s}{2} \quad \frac{(s + m_{\tilde{t}}^2 - m_{\tilde{q}}^2)^2}{4s} \quad m_{\tilde{t}}^2 \quad (15)$$

The total cross-section for $e^+ e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$ is given by

$$\sigma(s; e^+ e^-) = 2 \int_{m_{\tilde{\chi}_1^0}^2}^{0.83} f_{\gamma}(y) \sigma_{\gamma}(ys; m_{\tilde{\chi}_1^0}) dy \quad (16)$$

where f_{γ} is the distribution of high energy real photons at a given fraction $y = \frac{E}{E_e}$. The results of numerical integration for $e^+ e^- \rightarrow \tilde{\chi}_1^0 \tilde{\chi}_1^0$ is plotted in Fig.3 for a linear collider with center-of-mass energy of 500 GeV. As can be seen from the figure around ten thousands events per running year can be observed up to masses of 275 GeV. On the other hand using the Weizsacker-Williams approximation for the photon flux the same process can also be studied at the ordinary modes of LEP2 and NLC. The relevant calculations are completely similar to the above analysis, the only difference is in taking the upper limit of the integral as unity in the Equ.(16) and the insertion of the WW distribution, which is given by

$$f_{\gamma}(y) = \frac{1}{2} \frac{1 + (1 - y)^2}{y} \ln \frac{Q_{max}^2}{Q_{min}^2} - 2m_e^2 \left(\frac{1}{Q_{min}^2} - \frac{1}{Q_{max}^2} \right) \quad (17)$$

The results of the numerical integration for the total cross section at a center of mass energy of 500 GeV is shown in Fig.(4) as a function of the sneutrino mass. The total cross section is approximately a hundred times smaller than the value obtained at the e collider mode of NLC, and consequently the discovery mass reach for $\tilde{\chi}_1^0$ is less than 150 GeV.

Here we should note that the production of a single scalar leptoquark in $e^+ e^-$ collisions is a similar process and has been studied in some detail in a series of papers by Doncheski and Godfrey [11]. The total cross section is the same up to a trivial relabeling of the couplings. They have also taken into account the resolved photon contribution, which can surprisingly give values higher than the direct photon interactions. On the other hand the best place to produce squarks singly is an ep-collider (namely HERA) [1].

In this paper we have first investigated three different ways of producing single squarks (down-type) at TeV scale pp colliders considering R-parity violating LQD interactions. In one of them the particle associating the squark is a neutrino and in the latter case it is a tau-lepton. Then single production of squarks through again LQD interactions at e mode of the NLC colliders has been discussed. For comparison, it is shown that for $\tilde{\chi}_1^0$ production the e collider mode of a future NLC gives much higher values of the total cross section than the normal operation mode through soft WW photons. The production cross sections are functions of only sparticle mass and R_p coupling constants, and lead to detectable signals. Polarizations of the initial photon beam [8-9], which can be accomplished relatively easily, constitute additional advantages. Our results show that e and pp colliders can play complementary role in searching for supersymmetry in the future.

-
- [1] H E Haber and G L Kane, Phys.Rep.C 117, (1986)75
H P Nilles, Phys.Rep.C 110, (1984)1
Perspectives on Supersymmetry, ed by G L Kane, World Scientific (1998)
Jonathan L Feng et.al preprint hep-ph/0210390
D Choudhury and R Raychoudhuri, Phys.Lett.B 401 (1997)54
H Abramowicz and A Caldwell, DESY report 98-192.
R Barbier et.al preprint hep-ph/9810232
B Allanach et.al preprint, hep-ph/9906224
G Moreau, preprint hep-ph/0012156 (Ph.D thesis)
J C Romao, preprint hep-ph/9811454
Y Okada, preprint hep-ph/0304286
L3 Collaboration, preprint hep-ex/0011087
ALEPH Collaboration, preprint hep-ex/0011008, hep-ex/0210014
D0 Collaboration, preprint hep-ex/0207100
H1 Collaboration, DESY preprint 96-056.
- [2] I F Ginzburg et al, Nucl.Instrum. and Meth. 205 (1983)47
V I Telnov, Nucl.Instrum. and Meth. A 294 (1990)72
V Telnov, preprint hep-ph/0001029 (to be published in Nucl.Instrum. Meth. B)
I F Ginzburg, preprint hep-ph/0101029.
- [3] Z Z Aydin et al, Int.J.Mod.Phys.A 11 (1996)2019 and references therein.
J A Garfols and R Pascual, Phys.Lett.B 135 (1984)319
E Boos et al, Phys.Lett.B 273 (1991)273
J E Cieza Montalvo and O J P Eboli, Phys.Rev.D 47 (1993)837
M Nadeau and D London, Phys.Rev.D 47 (1993)3742.
G Jikia, Nucl.Phys.B 333, (1990)317.
- [4] A Kandemir and A U Y Imazer, Phys.Lett.B, 208 (1998)175
A Kandemir and A U Y Imazer, Nuovo Cim. A, 112 (1999)597
Z Aydin and O Y Imaz, Europhys.Lett. 50 (2000)22
- [5] J K alinowski, preprint hep-ph/9807312
P Richardson, preprint hep-ph/0101105 (Ph.D thesis).
- [6] J Butterworth and H Dreiner, Nucl.Phys.B 397 (1993)3
E Perez, Y Sirois and H Dreiner, preprint, hep-ph/9703444. published in the Proc. of Workshop on Future Physics at HERA, Sept.1996 (Ed. G. Ingelman, A De Roeck and R Kanner)
- [7] Z Z Aydin et al. Phys.Lett.b 515 (2001) 125-130
- [8] E Eichten et al, Rev.of Mod.Phys. 56, (1984)579
A D Martin and W J Stirling, Phys.Rev.D 50 (1994)6734
- [9] H Dreiner and G G Ross, Nucl.Phys.B 365 (1991)597.
- [10] T Plehn, preprint, hep-ph/9809319.
- [11] M Doncheski and S Godfrey, Phys.Rev.D 49, 6220 (1994)
M Doncheski and S Godfrey, Phys.Rev.D 51, 1040 (1994)
M Doncheski and S Godfrey, Phys.Lett.B 393, 355 (1997).
- [12] M Herz, preprint, hep-ph/0301079.

FIG .1. Production cross section of a down-squark ($\tilde{d}_R$, $\tilde{s}_R$ or $\tilde{b}_R$) as a function of its mass for HERA + LC $p\bar{p}$ collider via the process $p \rightarrow \tilde{d}_R^i + \bar{L} + X$.

FIG .2. Production cross section of a down-squark ($\tilde{d}_R$, $\tilde{s}_R$ or $\tilde{b}_R$) as a function of its mass for HERA + LC $p\bar{p}$ collider via the process $p \rightarrow \tilde{d}_R^i + L + X$.

FIG .3. Production cross section of a stop ($\tilde{t}_R$) as a function of its mass for e^+e^- mode of LC collider via the process $e^+e^- \rightarrow \tilde{t}_R + \bar{e}$.

FIG. 4. Production cross section of a stop (t_R) as a function of its mass for NLC ordinary mode with $W W$ -photons via the process e^+st .




Fig.1




Fig.2




Fig.3




Fig.4