

Resummed jet rates for e^+e^- annihilation into massive quarksFrank Krauss^y and Germán Rodrigo^zTheory Division, CERN
CH-1211 Geneva 23, Switzerland

(Dated: March 5, 2003)

Expressions for Sudakov form factors for heavy quarks are presented. They are used to construct resummed jet rates for up to four jets in e^+e^- annihilation. The coefficients of leading and next-to-leading logarithmic corrections, mandatory for a combination with higher order matrix elements, are evaluated up to second order in α_s .

PACS numbers: 12.38.Bx, 12.38.Cy, 13.66.Bc, 14.65.Fy

The formation of jets is the most prominent feature of perturbative QCD in e^+e^- annihilation into hadrons. Jets can be visualized as large portions of hadronic energy or, equivalently, as a set of hadrons confined to an angular region in the detector. In the past, this qualitative definition was replaced by quantitatively precise schemes to define and measure jets, such as the cone algorithms of the Weinberg (Sternan [1]) type or clustering algorithms, e.g. the Jade [2, 3] or the Durham scheme (k_T scheme) [4]. A refinement of the latter one is provided by the Cambridge algorithm [5]. Within the context of this paper, however, the Durham and the Cambridge algorithms are equivalent and they will be referred to as k_T algorithm. A clustering according to the relative transverse momenta has a number of properties that minimize the effect of hadronization corrections and allow an exponentiation of leading (LL) and next-to-leading logarithms (NLL) [4, 6] stemming from soft and collinear emission of secondary partons.

Equipped with a precise jet definition the determination of jet production cross sections and their intrinsic properties is one of the traditional tools to investigate the structure of the strong interaction and to deduce its fundamental parameters. In the past decade, precision measurements, especially in e^+e^- annihilation, have established both the gauge group structure underlying QCD [7, 8, 9, 10, 11] and the running of its coupling constant α_s over a wide range of scales [12]. In a similar way, also the quark masses should vary with the scale.

A typical strategy to determine the mass of, say, the bottom-quark at the centre-of-mass (c.m.) energy of the collider is to compare the ratio of three-jet production cross sections for heavy and light quarks [13, 14, 15, 16, 17]. At jet resolution scales below the mass of the quark, i.e. for gluons emitted by the quark with a relative transverse momentum k_T smaller than the mass, the collinear divergences are regularized by the quark mass. In this region mass effects are enhanced by logarithms $\ln(m_b/k_T)$, increasing the significance of the measurement. Indeed, this leads to a multiscale problem since in this kinematical region also large logarithms $\ln(\frac{p_{\perp}}{s} = k_T)$ appear such that both logarithms need to be resummed simultaneously. A solution to a somewhat similar two-scale problem, namely for the average subjet multiplicities in two- and three-jet events in e^+e^- annihilation was given in [35].

These large logarithms can be deduced by inspection of the corresponding splitting functions for massive quarks [18, 19, 20, 21, 22] and of the boundaries for their integration over the energy fractions of the outgoing particles. The resulting integrated splitting functions exhibit the LL and NLL behaviour, and resummation is achieved by means of Sudakov form factors. Jet rates can then be expressed, up to NLL accuracy, via the integrated splitting functions and the corresponding Sudakov form factors obtained from them. Following the work of Catani et al. [4], the resummation of such logarithms will be discussed in this paper for the case of heavy quark production. Furthermore, the corresponding LL and NLL coefficients will be calculated. These coefficients are mandatory for the combination with next-to-leading order calculations of the three-jet rate, involving heavy quarks in e^+e^- annihilation [13, 23, 24, 25, 26, 27, 28, 29]. In fact, they exhibit the correct logarithmic behaviour and thus provide a good estimate for the size of mass effects. Such a matching procedure was first defined for event shapes in [30], and later applications include the matching of fixed-order and resummed expressions for the four-jet rate in e^+e^- annihilation into massless quarks [31, 32]. A similar scheme for the matching of tree-level matrix elements with resummed expressions in the framework of Monte Carlo event generators for e^+e^- processes was suggested in [33] and extended to general collision types in [34]. Finally, mass effects are briefly highlighted for the case of two- and three-jet events both for bottom quarks at LEP 1 energies and for top quarks at a potential linear collider operating at c.m. energies of 500 GeV.

Work supported by the EC 5th Framework Programme under contract numbers HPMF-CT-2000-00989 and HPMF-CT-2002-01663

^yFrank.Krauss@cern.ch

^zGerman.Rodrigo@cern.ch

To obtain expressions for the splitting function of the process $a \rightarrow b(p_1) + c(p_2)$, which involves massive particles, we use the following Sudakov parameterization

$$\begin{aligned} p_1 &= z p \quad q + \frac{q^2 + p_1^2}{z} \frac{n}{2n \cdot p}; \\ p_2 &= (1-z)p + q_2 + \frac{q^2 + p_2^2}{1-z} \frac{n}{2n \cdot p}; \end{aligned} \quad (1)$$

where p and n are light-like vectors: $p^2 = n^2 = 0$, and q_2 is the space-like transverse momentum, $p \cdot q = n \cdot q = 0$, with $q^2 = -q_\perp^2 > 0$. Furthermore, $q = \eta j$. The quasi-collinear limit, as discussed in [19], is obtained by rescaling q and m through $q \rightarrow \bar{q}$ and $m \rightarrow \bar{m}$, respectively. Taking the limit $\bar{m} \rightarrow 0$, and keeping only terms of order $1 = \bar{m}^2$ in the squared matrix element we reproduce their result. It should be stressed, however, that this result is independent of the ratio $q = m$. In the limit discussed above, the squared amplitude at the tree-level fulfils a factorization formula, and it also contains naturally the two limits $q = m$ and $q = \bar{m}$ as particular cases.

For the splitting process

$$Q \rightarrow Q(p_1) + g(p_2); \quad (2)$$

Q being a heavy quark, $p_1^2 = m^2$, the squared matrix element factorizes as

$$\begin{aligned} \mathcal{M}(p_1; p_2; \dots) &\approx \mathcal{M}(p_1 + p_2; \dots) \frac{4}{p_1 \cdot p} P_{QQ}(z; q) \\ &\approx \mathcal{M}(p_1 + p_2; \dots) \frac{z(1-z)}{q^2 + (1-z)\bar{m}^2} P_{QQ}(z; q); \end{aligned} \quad (3)$$

where the unregularized spin-averaged splitting function in $D = 4 - 2\epsilon$ dimensions is given by

$$\begin{aligned} P_{QQ}(z; q) &= C_F \frac{1+z^2}{1-z} (1-z) \frac{m^2}{p_1 \cdot p} \\ &= C_F \frac{1+z^2}{1-z} (1-z) \frac{2z(1-z)m^2}{q^2 + (1-z)\bar{m}^2}; \end{aligned} \quad (4)$$

Analogously, for the $g \rightarrow Q(p_1) + Q(p_2)$ branching a similar factorization formula holds, where

$$\begin{aligned} P_{gQ}(z; q) &= T_R (1-z) \frac{2z(1-z)}{1} + \frac{2m^2}{(1-z)\bar{m}^2} \\ &= T_R (1-z) \frac{2z(1-z)}{1} + \frac{2z(1-z)m^2}{(1-z)(q^2 + m^2)}; \end{aligned} \quad (5)$$

As expected, these splitting functions match the massless splitting functions in the limit $m \rightarrow 0$ for q^2 fixed. Finally, for the $g \rightarrow g(p_1) + g(p_2)$ branching

$$P_{gg}(z) = C_A \frac{z}{1-z} + \frac{1}{z} + z(1-z); \quad (6)$$

In the above equations, $C_F = 4/3$ and $C_A = 3$ are the structure constants of $SU(3)$ in the fundamental and adjoint representation, respectively, and $T_R = 1/2$ is the normalization of its generators. For the purposes of this investigation, however, the splitting functions can be taken in $D = 4$ dimensions.

A counting for the number $n_f^{(i,h)}$ of active light or heavy fermions (quarks), respectively, P_{gQ} can be replaced by

$$P_{gf}(z; q) = n_f^{(l)} P_{gQ}(z; q)_{m=0} + \sum_{i \geq 2} n_f^{(h)} P_{gQ}(z; q; m_i); \quad (7)$$

where the sum runs over all $n_f^{(h)}$ flavours of heavy quarks. For massless particles the corresponding integrated splitting functions or branching probabilities yield

$$\int_{q=Q}^{1/z=Q} dz P_{QQ}(z) = 2C_F \ln \frac{Q}{q} \approx \frac{3}{4};$$

$$\begin{aligned}
g(Q; q; m=0) &= \int_{q=Q}^{1/z^q=Q} dz P_{gg}(z) = 2C_A \ln \frac{Q}{q} \frac{11}{12} ; \\
f(Q; q; m=0) &= \int_{q=Q}^{1/z^q=Q} dz P_{gf}(z) = \frac{2n_f T_R}{3} ;
\end{aligned} \tag{8}$$

The Sudakov form factors, which yield the probability for a parton experiencing no emission of a secondary parton between transverse momentum scales Q down to Q_0 , read

$$\begin{aligned}
Q(Q; Q_0) &= \exp \int_{Q_0}^Q \frac{dz}{z} \frac{d\alpha_s(z)}{\alpha_s(z)} P_{gg}(z; q) ; \\
g(Q; Q_0) &= \exp \int_{Q_0}^Q \frac{dz}{z} \frac{d\alpha_s(z)}{\alpha_s(z)} (g(Q; q) + f(Q; q)) ; \\
f(Q; Q_0) &= [Q(Q; Q_0)]^2 = g(Q; Q_0) ;
\end{aligned} \tag{9}$$

The fact that the above equations (8) and (9) are for massless particles only is reflected by the "propagator-like" structure $1/z^q$. In contrast, for massive particles the propagator term is given by $1/[q^2 + (1/z^q)m^2]$ for $Q \neq Q_0$, and by $1/[q^2 + m^2]$ for $Q = Q_0$ instead. In order to compensate for this we define the integrated splitting functions involving heavy quarks through

$$\begin{aligned}
Q(Q; q; m) &= \int_{q=Q}^{1/z^q=Q} dz \frac{q^2}{q^2 + (1/z^q)m^2} P_{gQ}(z; q) \\
&= Q(Q; q; m=0) + C_F \frac{1}{2} \frac{q}{m} \arctan \frac{m}{q} \frac{2m^2}{2m^2} \ln \frac{m^2 + q^2}{q^2} ; \\
f(Q; q; m) &= \int_{q=Q}^{1/z^q=Q} dz \frac{q^2}{q^2 + m^2} P_{gQ}(z; q) = T_R \frac{q^2}{q^2 + m^2} \left[1 - \frac{1}{3} \frac{q^2}{q^2 + m^2} \right] ;
\end{aligned} \tag{10}$$

Written in such a fashion, the Sudakov form factor for massive quarks is obtained by a mere replacement of the integrated splitting function according to Eq. (9). Similarly, for every heavy quark Q occurring in a splitting $g \rightarrow Q\bar{Q}$, the integrated splitting function f is modified correspondingly. In accordance with these replacements the running coupling constant α_s changes as well, since the number of active quarks changes when passing a heavy quark threshold.

The integrated splitting functions and the corresponding Sudakov form factors can be employed to calculate multi-jet rates in e^+e^- annihilation in the k_T schemes. In both schemes, the jet resolution parameter y_{ij} is given by

$$y_{ij} = \frac{2m \ln \frac{E_i^2 + E_j^2}{s} (1 - \cos \theta_{ij})}{s} ; \tag{11}$$

where $s = Q^2$ is the c.m. energy squared of the colliding electrons. For massive particles, it might be more suitable to replace the energies with the absolute values of the three-momenta in order to identify the relative transverse momentum properly, i.e.

$$y_{ij} = \frac{2m \ln \frac{p_i^2 + p_j^2}{s} (1 - \cos \theta_{ij})}{s} ; \tag{12}$$

However for our discussion this difference is subleading only.

Rates for two-, three-, and four-jet events can be expressed by the integrated splitting functions and the Sudakov form factors as

$$R_2 = [Q(Q; Q_0)]^2 ;$$

$$\begin{aligned}
R_3 &= 2 \left[\int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \right]^2 \int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \int_{Q_0}^q \frac{dq'}{q'} \frac{s(q')}{s(Q)} \int_{Q_0}^{q'} \frac{dq''}{q''} \frac{s(q'')}{s(Q)} ; \\
R_4 &= 2 \left[\int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \right]^2 \int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \int_{Q_0}^q \frac{dq'}{q'} \frac{s(q')}{s(Q)} \int_{Q_0}^{q'} \frac{dq''}{q''} \frac{s(q'')}{s(Q)} \int_{Q_0}^{q''} \frac{dq'''}{q'''} \frac{s(q''')}{s(Q)} ; \\
&\quad + \int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \int_{Q_0}^q \frac{dq'}{q'} \frac{s(q')}{s(Q)} \int_{Q_0}^{q'} \frac{dq''}{q''} \frac{s(q'')}{s(Q)} \int_{Q_0}^{q''} \frac{dq'''}{q'''} \frac{s(q''')}{s(Q)} \int_{Q_0}^{q'''} \frac{dq''''}{q''''} \frac{s(q''''')}{s(Q)} ; \\
&\quad + \int_{Q_0}^Q \frac{dq}{q} \frac{s(q)}{s(Q)} \int_{Q_0}^q \frac{dq'}{q'} \frac{s(q')}{s(Q)} \int_{Q_0}^{q'} \frac{dq''}{q''} \frac{s(q'')}{s(Q)} \int_{Q_0}^{q''} \frac{dq'''}{q'''} \frac{s(q''')}{s(Q)} \int_{Q_0}^{q'''} \frac{dq''''}{q''''} \frac{s(q''''')}{s(Q)} ;
\end{aligned} \tag{13}$$

where Q_0 now plays the role of the jet resolution scale in the k_T algorithm, $Q_0^2 = y_{\text{cut}} Q^2$, and Q is the c.m. energy of the colliding e^+e^- . Single-flavour jet rates in Eq. (13) are defined from the flavour of the primary vertex, i.e. events with gluon splitting into heavy quarks where the gluon has been emitted on primary light quarks are not included in the heavy jet rates but would be considered in the jet rates for light quarks.

In order to catch which kind of logarithmic corrections are resummed with these expressions it is illustrative to study the above formulae in the kinematical regime such that $Q \gg m \gg Q_0$. Expanding in powers of s , jet rates can be formally expressed as

$$R_n = c_{n2}^{(n)} + \sum_{k=n-2}^{\infty} \frac{s(Q)}{s(Q_0)^k} c_{k1}^{(n)} ; \tag{14}$$

where the coefficients $c_{k1}^{(n)}$ are polynomials of order 1 in $L_y = \ln(1-y_{\text{cut}})$ and $L_m = \ln(m^2=Q_0^2)$. At the given NLL accuracy it is sufficient to treat the running of s to first order (one-loop),

$$s(q) = \frac{s(Q)}{1 + \sum_{n=1}^{\infty} \frac{s(Q)}{s(Q_0)^n} \ln(q^2=Q^2)} ; \tag{15}$$

where the β -function β_n for n active quarks is given by

$$\beta_n = \frac{11 C_A - 2n}{12} ; \tag{16}$$

Alternatively, instead of using the transverse momentum q as the scale for s in the Sudakov form factors, the expression showing up in the propagator term s , i.e. $q^2 + (1-z)^2 m^2$ for the heavy quark splitting $Q \rightarrow Qg$, or $q^2 + m^2$ for the gluon splitting into two heavy quarks $g \rightarrow Q\bar{Q}$, can be chosen. However, this is a subleading effect.

The coefficients for the first order in s are given by

$$\begin{aligned}
c_{12}^{(2)} &= c_{12}^{(3)} = \frac{1}{2} C_F (L_y^2 - L_m^2) ; \\
c_{11}^{(2)} &= c_{11}^{(3)} = \frac{3}{2} C_F L_y + \frac{1}{2} C_F L_m ;
\end{aligned} \tag{17}$$

all coefficients for higher jet multiplicities being 0. For second order s with n active flavours at the high scale, the LL and NLL coefficients read

$$\begin{aligned}
c_{24}^{(2)} &= \frac{1}{8} C_F^2 L_y^2 L_m^2 ; \\
c_{24}^{(3)} &= \frac{1}{4} C_F^2 L_y^2 L_m^2 + \frac{1}{48} C_F C_A L_y^4 L_m^4 ; \\
c_{24}^{(4)} &= \frac{1}{8} C_F^2 L_y^2 L_m^2 + \frac{1}{48} C_F C_A L_y^4 L_m^4 ;
\end{aligned}$$




FIG. 1: Effect of a b-mass of 5 GeV on the single-, two- and three-jet rate at LEP1 energies as a function of the jet resolution parameter in the Durham scheme. In the left plot this effect is treated through the full inclusion of m masses into the splitting function, see Eq. (10), whereas in the plot on the right hand side this effect is modeled through the dead cone, see Eq. (19).

$$\begin{aligned}
C_{23}^{(2)} &= C_F^2 L_Y^2 L_m^2 \left(\frac{3}{4} L_Y - \frac{1}{4} L_m \right) - \frac{1}{3} n C_F L_Y^3 - \frac{3}{2} L_Y L_m^2 + \frac{1}{2} L_m^3 \\
&\quad - \frac{1}{3} (n - n - 1) C_F L_m^3 ; \\
C_{23}^{(3)} &= \frac{1}{2} C_F^2 L_Y^2 L_m^2 (3L_Y - L_m) + \frac{1}{2} n C_F L_Y L_Y^2 L_m^2 + \frac{1}{24} C_F C_A (3L_Y^3 - L_m^3) \\
&\quad + \frac{1}{6} (n - n - 1) C_F L_m L_Y^2 L_Y L_m + 2L_m^2 ; \\
C_{23}^{(4)} &= C_F^2 L_Y^2 L_m^2 \left(\frac{3}{4} L_Y - \frac{1}{4} L_m \right) - \frac{1}{6} n C_F L_Y^3 - \frac{1}{8} C_F C_A L_Y^3 - \frac{1}{3} L_m^3 \\
&\quad - \frac{1}{6} (n - n - 1) C_F L_Y L_m (L_Y - L_m) ;
\end{aligned} \tag{18}$$

Terms $(n - n - 1)$ in the NLL coefficients are due to the combined effect of the gluon splitting into massive quarks and of the running of α_s below the threshold of the heavy quarks, with a corresponding change in the number of active flavours. With our definition of jet rates with primary quarks the jet rates add up to one at NLL accuracy. This statement is obviously realized in the result above order by order in α_s .

The corresponding massless result [4] is obtained from Eqs. (17) and (18) by setting $L_m \rightarrow 0$. Notice that Eqs. (17) and (18) are valid only for $m \ll Q_0$ and therefore $m \rightarrow 0$ does not reproduce the correct limit, which has to be smooth as given by Eq.(13). Let us also mention that for $Q \approx m$ there is a strong cancellation of leading logarithms and therefore subleading effects become more pronounced.

An approximate way of including mass effects in massless calculations, that is sometimes used, is the "dead cone" [36] approximation. The dead cone relies on the observation that, at leading logarithmic order, there is no radiation of soft and collinear gluons off heavy quarks. This effect can be easily understood from the splitting function in Eq. (4). For $q = (1 - z)m$ the splitting function is not any more enhanced at $z \rightarrow 1$. This can be expressed via the modified integrated splitting function

$$d_{Q \rightarrow q}^{d.c.}(Q; q; m) = d_{Q \rightarrow q}(Q; q; m = 0) + 2C_F \ln \frac{Q}{m} (m - q) ; \tag{19}$$

To obtain this result the massless splitting function has been used, which is integrated with the additional constraint $z > 1 - q/m$. We also compare our results with this approximation.

The impact of mass effects can be highlighted by two examples, namely by the effect of the b-mass in e^+e^- annihilation at the Z-pole and by the effect of the t-mass at a potential linear collider operating at a cm. energy of 500 GeV. With $m_b = 5$ GeV, $M_Z = 91.2$ GeV, and $\alpha_s(M_Z) = 0.118$, the effect of the b-mass at the Z-pole on the two- and three-jet rates is depicted in Fig. 1. Clearly, using the full massive splitting function, the onset of mass effects in




FIG. 2: The effect of a top mass of 175 GeV on the single-, two- and three-jet rate as a function of the jet resolution parameter, at a potential linear collider operating at c.m. energies of 500 GeV. A gain, in the left plot this effect is treated through the full inclusion of masses into the splitting function, see Eq. (10), whereas in the plot on the right hand side this effect is modeled through the dead cone, see Eq. (9).

the jet rates is not abrupt as in the dead cone case and becomes visible much earlier. Already at the rather modest value of the jet resolution parameters of $y_{\text{cut}} = 0.004$, the two-jet rate, including mass effects, is enhanced by roughly 4% with respect to the massless case, whereas the three-jet rate is decreased by roughly 3.5%. For even smaller jet resolution parameters, the two-jet rate experiences an increasing enhancement, whereas the massive three-jet rate starts being larger than the massless one at values of the jet resolution parameters of the order of $y_{\text{cut}} = 0.001$. The curves have been obtained by numerical integration of Eq. (13). Furthermore, in order to obtain physical results the branching probabilities have been set to one whenever they exceed one or to zero whenever they become negative.

While in the case of bottom quarks at LEP1 energies the overall effect of the quark mass is at the few-percent level, this effect becomes tremendous for top quarks at a potential linear collider operating at 500 GeV; see Fig. 2. Owing to the size of mass effects, the difference between the LL treatment through the dead cone and the full NLL result becomes visible, reflecting the fact that $Q = m$ and the respective logarithms canceling each other. In other words, for this case a full next-order treatment is mandatory.

In this paper Sudakov form factors involving heavy quarks have been employed to estimate the size of their mass effects in jet rates in e^+e^- annihilation into hadrons. These effects are sizeable and therefore observable in the experimentally relevant region. In addition, the coefficients for the leading logarithms, both in the jet resolution parameter and in the quark mass, have been deduced up to second order in α_s . They are mandatory for the combination with next-order calculations of the two-, three- and four-jet rates, and resummed expressions as obtained by Sudakov form factors. The matching between next-order calculations and resummed results will be presented in a forthcoming article.

Acknowledgments

The authors want to acknowledge financial support from the EC 5th Framework Program under contract numbers HPMF-CT-2000-00989 (G.R.) and HPMF-CT-2002-01663 (F.K.). The authors are grateful for stimulating discussions with S. Catani, B. Webber and M. L. Mangano. Valuable comments from J. Fuster and K. Hamacher concerning the experimental situation are also acknowledged. Special thanks to S. Vasotto for carefully reading the manuscript. G.R. acknowledges also partial support from Generalitat Valenciana under grant CTID IB/2002/24 and MCyT under grants FPA-2001-3031 and BFM 2002-00568.

-
- [1] G. Sterman and S. Weinberg, Phys. Rev. Lett. 39 (1977) 1436.
 - [2] W. Bartel et al. [JADE Collaboration], Z. Phys. C 33 (1986) 23.
 - [3] S. Bethke et al. [JADE Collaboration], Phys. Lett. B 213 (1988) 235.

- [4] S. Catani, Y. L. Dokshitzer, M. Olsson, G. Tumock and B. R. Webber, Phys. Lett. B 269 (1991) 432.
- [5] Y. L. Dokshitzer, G. D. Leder, S. Moretti and B. R. Webber, JHEP 9708 (1997) 001 [arXiv:hep-ph/9707323].
- [6] S. Catani, CERN-TH-6281-91
- [7] P. Abreu et al. [DELPHI Collaboration], Phys. Lett. B 255 (1991) 466.
- [8] P. Abreu et al. [DELPHI Collaboration], Z. Phys. C 59 (1993) 357.
- [9] P. Abreu et al. [DELPHI Collaboration], Phys. Lett. B 414 (1997) 401.
- [10] G. Abbiendi et al. [OPAL Collaboration], Eur. Phys. J. C 20 (2001) 601 [arXiv:hep-ex/0101044].
- [11] A. Heister et al. [ALEPH Collaboration], CERN-EP/2002-029, submitted to Eur. Phys. J. C.
- [12] S. Bethke, J. Phys. G 26 (2000) R27 [arXiv:hep-ex/0004021].
- [13] G. Rodrigo, A. Santamaria and M. S. Bilenky, Phys. Rev. Lett. 79 (1997) 193 [arXiv:hep-ph/9703358].
- [14] P. Abreu et al. [DELPHI Collaboration], Phys. Lett. B 418 (1998) 430.
- [15] A. Brandenburg, P. N. Burrows, D. Muller, N. Oishi and P. Uwer, Phys. Lett. B 468 (1999) 168 [arXiv:hep-ph/9905495].
- [16] R. Barate et al. [ALEPH Collaboration], Eur. Phys. J. C 18 (2000) 1 [arXiv:hep-ex/0008013].
- [17] G. Abbiendi et al. [OPAL Collaboration], Eur. Phys. J. C 21 (2001) 411 [arXiv:hep-ex/0105046].
- [18] Y. L. Dokshitzer, V. A. Khoze and S. I. Troian, Phys. Rev. D 53 (1996) 89 [arXiv:hep-ph/9506425].
- [19] S. Catani, S. Dittmaier and Z. Trocsanyi, Phys. Lett. B 500 (2001) 149 [arXiv:hep-ph/0011222].
- [20] M. Cacciari and S. Catani, Nucl. Phys. B 617 (2001) 253 [arXiv:hep-ph/0107138].
- [21] S. Catani, S. Dittmaier, M. H. Seymour and Z. Trocsanyi, Nucl. Phys. B 627 (2002) 189 [arXiv:hep-ph/0201036].
- [22] M. Cacciari and E. Gardi, Nucl. Phys. B 664 (2003) 299 [arXiv:hep-ph/0301047].
- [23] G. Rodrigo, Nucl. Phys. Proc. Suppl. 54A (1997) 60 [arXiv:hep-ph/9609213].
- [24] M. S. Bilenky, S. Cabrera, J. Fuster, S. Marti, G. Rodrigo and A. Santamaria, Phys. Rev. D 60 (1999) 114006 [arXiv:hep-ph/9807489].
- [25] G. Rodrigo, M. S. Bilenky and A. Santamaria, Nucl. Phys. B 554 (1999) 257 [arXiv:hep-ph/9905276].
- [26] P. Nason and C. Oleari, Phys. Lett. B 407 (1997) 57 [arXiv:hep-ph/9705295].
- [27] P. Nason and C. Oleari, Nucl. Phys. B 521 (1998) 237 [arXiv:hep-ph/9709360].
- [28] W. Bemreuther, A. Brandenburg and P. Uwer, Phys. Rev. Lett. 79 (1997) 189 [arXiv:hep-ph/9703305].
- [29] A. Brandenburg and P. Uwer, Nucl. Phys. B 515 (1998) 279 [arXiv:hep-ph/9708350].
- [30] S. Catani, L. Trentadue, G. Tumock and B. R. Webber, Nucl. Phys. B 407 (1993) 3.
- [31] L. J. Dixon and A. Signer, Phys. Rev. D 56 (1997) 4031 [arXiv:hep-ph/9706285].
- [32] Z. Nagy and Z. Trocsanyi, Phys. Rev. D 59 (1999) 014020 [Erratum-ibid. D 62 (2000) 099902] [arXiv:hep-ph/9806317].
- [33] S. Catani, F. Krauss, R. Kuhn and B. R. Webber, JHEP 0111 (2001) 063 [arXiv:hep-ph/0109231].
- [34] F. Krauss, JHEP 0208 (2002) 015 [arXiv:hep-ph/0205283].
- [35] S. Catani, B. R. Webber, Y. L. Dokshitzer and F. Fiorani, Nucl. Phys. B 383 (1992) 419.
- [36] Y. L. Dokshitzer, V. A. Khoze and S. I. Troian, J. Phys. G 17 (1991) 1602.