Radiative muon (pion) pair production in high energy electron-positron annihilation process

A B. Arbuzov, E. Bartos, V.V. Bytev, and E A. Kuraev Joint Institute for Nuclear Research, 141980 Dubna, Russia (Dated: March 24, 2024)

Process of muon (pion) pair production with sm all invariant mass in the electron { positron high {energy annihilation, accompanied by emission of hard photon at large angles, is considered. We nd that the Dell {Yan picture for dimential cross section is valid in the charge {even experimental set {up. Radiative corrections both for electron block and for nal state block are taken into account.

I. IN TRODUCTION

Initial state radiation processes in high energy electron-positron annihilation processes with creation of a hadronic system provide the laboratory for studying the hadron states created by virtual photon. Special interest is paid to the case when the invariant mass of hadron system $p_{\overline{s_1}}$ is small compared with the center{of{mass total energy $p_{\overline{s}} = 2$ ". Here such interesting physical quantities as the form factor of pion and nucleon can be investigated. The process of radiative annihilation into muon and pion pairs, considered here, plays a crucial role as a norm alization one as well as the ones, providing an essential background process in studying the hadron creation. D escription of its di erential cross section with a rather high level of accuracy (better than 0.5%) is the goal of our paper.

W e specify the kinem atics of radiative m uon pair creation process

$$e(p) + e_{+}(p_{+})! (q) + {}_{+}(q_{+}) + (k_{1});$$
 (1)

Department of Theoretical Physics, Comenius University, 84248 Bratislava, Slovakia.

as follow s:

$$p^{2} = m^{2}; q^{2} = M^{2}; k_{1}^{2} = 0;$$

= 2k₁p; ⁰ = 2k₁q; s = (p + p₊)²; s₁ = (q + q₊)²;
t = (p q)²; t₁ = (p₊ q)²; u = (p q)²; u₁ = (p₊ q)²; (2)

where m and M are the electron and muon (pion) masses, respectively. Throughout the paper we will suppose

$$s t_1 t u_1 u = s_1 M^2 m^2$$
: (3)

Situation where s s_1 M 2 is allowed too.

W e will system atically om it the term s of order M 2 =s and m 2 =s₁ com pared with ones of order of magnitude of unity. In O () radiative corrections we drop also term s suppressed by the factor s₁=s. A kinem atical diagram of the process under consideration is drawn in Fig.1.



Fig. 1: K inem atical diagram for a radiative event with a sm all invariant m ass s_1 .

In this paper we will consider only the charge {even part of the di erential cross section, which can be measured in an experimental set {up blind to charges of the created particles. A detailed study of the charge {odd part of the radiative annihilation cross section in general kinematics will be presented elsewhere.

Our paper is organized as follows. The next Section is devoted to the Bom {level cross section. Radiative corrections to the nal and initial states are considered in Sect. III. In C onclusion we give some num erical illustration and discuss the resulting precision. Some details of calculations and useful form ulae are given in Appendix.

W ithin the Born approximation, the matrix element of the initial state emission process has the form :

$$M_{B} = \frac{(4)^{3=2}}{s_{1}} v(p_{+}) \qquad \frac{\hat{p} \qquad \hat{k}_{1} + m}{2p k_{1}} e^{\frac{\hat{p}}{2} + \hat{k}_{1} + m} \qquad u(p) J \qquad (4)$$

with

$$J = u(q) u(q_{+})$$
 (5)

for muon pair production, and

$$J = (q \quad q_1) F^{str}(s_1)$$
(6)

for the case of charged pions, F $^{str}(s_1)$ is the pion strong { interaction form factor.

The corresponding contributions to the cross section is

$$\frac{d}{d} = \frac{3}{8^{2} s s_{1}^{2}} R^{j}; \quad R^{j} = B \quad i^{(0j)}; \quad i^{(0)j} = X \quad J^{j} (J^{j})^{2}; \quad j = ;;$$

$$B = B_{g}g + B_{11} (p p) + B_{22} (p_{+} p_{+});$$

$$B_{g} = \frac{(s_{1} + \frac{1}{2})^{2} + (s_{1} + \frac{1}{2})^{2}}{+}; \quad B_{11} = \frac{4s_{1}}{+}; \quad B_{22} = \frac{4s_{1}}{+}; \quad (7)$$

where we used the short notation (qq) = qq, (pq) = pq + qp. For the case with the muon nalstate

$$i^{(0)} = 4 (q, q) \quad g \frac{s_1}{2}$$
 (8)

For the case of pions,

$$\dot{z}^{(0)} = f^{str}(s_1) f(q - q) (q - q) :$$
(9)

Here we put also the form ulae for B orn cross section in the case when only term s of order $(m^2=s)$ are om itted and hard photon are em itted on large angle (see [9]):

$$\frac{d}{d} \frac{B}{d} = \frac{3}{2^{2}s} \frac{jq_{+}j!}{2^{2}s} A; \qquad (10)$$

$$A = \frac{t^{2} + t_{1}^{2} + u^{2} + u_{1}^{2}}{ss_{1}} W_{even} + W_{odd}];$$

$$W_{even} = \frac{s}{t_{+}} + \frac{s_{1}}{t_{+}} + \frac{s_{1}}{t_{+}} + \frac{M^{2}}{s};$$

$$W_{odd} = \frac{t_{1}}{t_{+}} \frac{t_{-}}{t_{+}} + \frac{t_{-}}{t_{+}} + \frac{u_{-}}{t_{+}} + \frac{u$$

for m uon pair production, and for pion pair production

$$\frac{d_{B}}{d} = \frac{3}{4^{2}s} \frac{tu + t_{1}u_{1}}{ss_{1}} [W_{even} + W_{odd}]:$$
(11)

Functions W_{even} and W_{odd} give rise to the charge{even and charge{odd contributions to the cross section. In quantity W_{even} the term s=(+) is the most important in our kinem atic region. In what follows, only the even part will be taken into account.

In our kinem atics som e sim pli cation takes place:

$$B_{g} = 2\frac{1+c^{2}}{1-c^{2}}; \quad B_{11} = B_{22} = -\frac{16s_{1}}{s^{2}(1-c^{2})}: \quad (12)$$

with

$$c = \cos(\theta q) = \cos ; \qquad (13)$$

We will suppose that the emission angle of hard photon lies outside the narrow cones around the beam axis $_0 < _1 < _0$, with $_0 = 1$, $_0$ " M. For initial state radiative process (in paper [2] the scattering channel was considered). The phase volume of the nal particles is

$$d = \frac{d^{3}q}{"} \frac{d^{3}q_{+}}{"_{+}} \frac{d^{3}k_{1}}{!_{1}} {}^{4}(p_{+} + p \quad q \quad q \quad k_{1}):$$
(14)

For the case of sm all invariant m ass of created pair s_1 s it can be rewritten as (see g.1):

$$d = {}^{2} dx \ d\alpha ds_{1}; \tag{15}$$

with

$$x = \frac{"}{"}; x_{+} + x = 1; ! = \frac{1p}{2}; = \frac{s(1 - c)}{2};$$

$$t = \frac{sx(1 - c)}{2}; t_{1} = \frac{sx_{+}(1 + c)}{2}; u = \frac{sx_{+}(1 - c)}{2}; u_{1} = \frac{sx(1 + c)}{2}:$$

In the lim iting case of sm all invariant m ass of the created pair om itting term s of the order s_1 =s and M²=s, the Born level cross section takes a simple form :

$$d_{B}^{()}(p;p_{+};k_{1};q;q_{+}) = \frac{3(1+c^{2})}{ss_{1}(1-c^{2})} + x x_{+} dx dc ds_{1}; \quad (16)$$

$$d_{B}^{()}(p;p_{+};k_{1};q;q_{+}) = \frac{3(1+c^{2})}{ss_{1}(1-c^{2})} + x x_{+} dx dc ds_{1}; \quad (16)$$

$$\frac{1}{2}(1-c^{2}) + x x_{+} dx dc ds_{1}; \quad (16)$$

$$\frac{1}{2}(1-c^{2}) + x x_{+} dx dc ds_{1}; \quad (16)$$

Here is the velocity of a pair component in the center {of {m ass reference fram e of the pair.

Radiative corrections (RC) can be separated into three gauge{invariant parts. They can be taken into account by the form al replacement (see(7)):

$$\frac{R^{j}}{s_{1}^{2}} \cdot \frac{K J^{j}}{s_{1}^{2} J (s_{1}) f}$$

$$(17)$$

where (s_1) describes the vacuum polarization of the virtual photon (see Appendix); K is the initial state emission C ompton tensor with RC taken into account; J^j is the nal state current tensor with O () RC.

First we consider the explicit formulae for RC due to virtual, soft, and hard collinear nal state em ission. A sconcerning RC to the initial state for the charge (blind experimental set {up considered here, we will use the explicit expression for the C on pton tensor with heavy photon K calculated in the paper of one of us [2] for the scattering channel and apply the crossing transformation. Possible contribution due to emission of an additional real photon from the initial state will be taken into account too. In conclusion we give the explicit formulae for cross section, consider separate the kinematics of collinear emission, and estimate the contribution of higher orders of perturbation theory (PT).

A. Corrections to the nalstate

The third part is related with lowest order RC to muon and pion current

$$J = i^{(v)} + i^{(s)} + i^{(h)}$$
(18)

The virtual photon contribution $i^{(v)}$ takes into account the D irac and P auli form factors of m uon current

$$J^{(v)} = u(q) [F_1(s_1) + \frac{\hat{q}}{4M} F_2(s_1)]v(q_*); \quad q = q_* + q; \quad s_1 = q^2:$$
(19)

W e have

$$B \quad i^{(v)} = B_{g} \qquad j \quad j^{(v)} \quad j^{2} + B_{11} \qquad p \quad J^{(v)} \quad j^{2} + \qquad p_{+} \quad j_{+} \quad J^{(v)} \quad j^{2} : \qquad (20)$$

Here means sum over muon spin states and

$$X \qquad jJ^{(v)} f = - 8(s + 2M^{2})f^{()} \quad 12s f_{2}^{()};$$

$$X \qquad jJ^{(v)} p f = -s^{2}(1 - c)(x + x - f_{1}^{()} + \frac{1}{4}f_{2}^{()});$$
(21)

see Appendix for details. For the pion nal state we have

B
$$i^{(v)} = 2B i^{(0)} f^{QED};$$

B $i^{(0)} = F^{str}(s_1)f^{2}(4M^2 s_1)B_g + \frac{1}{8}s^2B_{11}(x_+ x_-)^2(1+c^2);$ (22)

The explicit expression for the electrom agnetic form factor of pion is given in Appendix. The soft photon correction to the nal state currents reads

$$i^{(s)} = - {}_{1^{0}2^{0}}i^{(0)}; \qquad i^{(s)} = - {}_{1^{0}2^{0}}i^{(0)};$$

$${}_{1^{0}2^{0}} = \frac{1}{4} \frac{Z}{4} \frac{d^{3}k}{!} \frac{q_{+}}{q_{+}k} \frac{q}{q_{k}k}^{2} = (\frac{1+\frac{2}{2}}{2}\ln\frac{1+\frac{1}{1}}{1} - 1)\ln\frac{\frac{\pi^{2}M^{2}}{r_{+}\pi^{2}}}{1} (23)$$

$$+ \frac{1+\frac{2}{2}}{g} \frac{1}{2}\ln^{2}\frac{1+\frac{1}{1}}{1} - \ln\frac{1+\frac{1}{2}}{h}\ln\frac{1+\frac{2}{4}}{4} - \frac{2}{6} - 2L\frac{1}{2} - \frac{1}{+1};$$

$$g = 2 \frac{Z^{1}}{1} \frac{dt}{1-\frac{2}{2}t^{2}}\ln 1 + \frac{1}{4} \frac{t}{4} \frac{(x_{+}-x_{-})^{2}}{x_{+}x} = \ln \frac{1+\frac{1}{1}}{h} \ln(1+z-z=^{2})$$

$$+ \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}}\ln 1 + \frac{Li_{2}}{1-\frac{1}{2}t^{2}} - \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} = \ln \frac{1+\frac{1}{1}}{h} \ln(1+z-z=^{2})$$

$$+ \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1-\frac{1}{2}}{2}t^{2}} - \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1+\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{2}t^{2}} + \frac{Li_{2}}{1+\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{1-\frac{1+\frac{1}{2}}{2}t^{2}} + \frac{Li_{2}}{2}t^{2}} + \frac{Li_$$

The contribution of an additional hard photon (with momentum k_2) emission by the muon block, provided $s_1 = (q_+ + q_- + k_2)^2 = s_1$ s can be found by the expression

$$B \quad i^{(h)} = \frac{Z}{4^{2}} \quad \frac{d^{3}k_{2}}{!_{2}} B \qquad J^{()}(J^{()}) \qquad ; \qquad (24)$$

with

X

$$jJ^{()}J^{2} = 4Q^{2}(s_{1} + 2k_{2}q + 2k_{2}q_{+} + 2M^{2}) = 8\frac{(k_{2}q_{-})^{2} + (k_{2}q_{+})^{2}}{(k_{2}q_{-})(k_{2}q_{+})};$$

 $Q = \frac{q}{q_{-}k_{2}} = \frac{q_{+}}{q_{+}k_{2}};$

and

$$X = jJ^{()}p^{-2}j = 8Q^{2}(q p)(q_{+}p) + 8(p k_{2}) Qq_{+}\frac{p q}{q_{+}k_{2}} = Qq\frac{p q_{+}}{q k_{2}} + 8(p k_{2}) \frac{p q}{q_{+}k_{2}} + 8(p Q)(p q_{+} p q) = 8\frac{(k_{2}p)^{2}}{(k_{2}q_{+})(k_{2}q)}; \quad (25)$$

For the case of charged pion pair production the radiative current tensor has the form

$$i^{(h_{1})} = \frac{2}{4^{2}} f^{str}(\mathbf{s}_{1}) f^{2} \frac{d^{3}k_{2}}{!_{2}} \frac{M^{2}}{!_{2}} (Q_{1}Q_{1}) + \frac{M^{2}}{!_{2}} (Q_{2}Q_{2}) \frac{q_{+}q}{!_{2}} (Q_{1};Q_{2}) + q \frac{1}{!_{2}} (Q_{1};q_{1}) + \frac{1}{!_{2}} (Q_{2};q_{+}) ; \qquad (26)$$

$$Q_{1} = q \quad q_{+} k_{2}; \quad Q_{2} = q \quad q_{1} \quad k_{2}; \quad 2 = 2k_{2}q :$$

O ne can check that the Bose symmetry and the gauge invariance condition is valid for the pionic current tensor. Namely it is invariant regarding the permutation of the pion momenta operation and turns to zero after conversion with 4-vector q.

The total sum of RC to the muon current does not depend on "=".

${\tt B}$. ${\tt C}\xspace$ or ${\tt C}\xspace$ or ${\tt C}\xspace$ of the initial state

Let us now consider RC to the Compton tensor with RC, which describe virtual corrections to the initial state. In our kinem atical region it will be convenient to rewrite the tensor explicitly extracting large logarithm s. W e will distinguish two kinds of large logarithm s:

$$l_{s} = \ln \frac{s}{m^{2}}; \quad l_{1} = \ln \frac{s}{s_{1}};$$
 (27)

W e rewrite the Compton tensor (17) in the form :

$$K = (1 + \frac{1}{2})B + \frac{1}{2}gg + \frac{1}{1}(pp) + \frac{1}{2}(p+p+) = \frac{1}{2}(pp+); \quad (28)$$
$$= 4\ln^{\frac{m}{2}}(l_{s} 1) \frac{1}{s} + 3l_{s} - 3\frac{1}{2} + \frac{4}{3} \frac{2}{2} - \frac{9}{2};$$

with $i = a_i l_i + b_i$ and

$$a_{11} = \frac{2s_1}{+} \frac{2b^2}{+} + \frac{4s}{a} + \frac{4(s^2 + b)}{a^2} \frac{b^2(2c)}{c^2} \frac{2s + \frac{1}{+}}{+}; \quad (29)$$

$$b_{11} = \frac{2}{+} s_1(1 + \frac{s^2}{+})G s_1(2 + \frac{b^2}{-2})G s_1(2 + \frac{b^2}{-2})G s_1(2 + \frac{b^2}{-2}) s_1(2 + \frac{b^2}{-2})S s_1(2 + \frac{$$

$$a_{12} = \frac{2s_1}{+} + \frac{4ss_1}{+} + \frac{8(++)s_1}{a^2} + \frac{4s}{a} + 4ss_1 + \frac{1}{c} + \frac{1}{b_+} + (2ss_1 + 4_+) + \frac{1}{c^2} + \frac{1}{b^2}; \qquad (31)$$

$$b_{12} = \frac{2}{+} \frac{2s_1}{\frac{2}{+}} (sc_{+})G_{+} + \frac{2s_1}{2} (sb_{+})G_{+}$$

$$+ s_1 \frac{2ss_1 + 4}{c^2} + \frac{4ss_1}{c} \ln \frac{s_{+}}{+} + s_1 \frac{2ss_1 + 4}{b^2} + \frac{4ss_1}{b_{+}} \ln \frac{s_{-}}{b_{+}}$$

$$+ \frac{8(s^2_{+})}{a} 2s \frac{s_{+}}{c} + \frac{s_1}{b_{+}} + 2s_1 + 10s; \qquad (32)$$

$$a_{g} = 2s \frac{s_{1}}{+} \frac{2}{a} + \frac{c}{+} \frac{3s}{b} 1 + \frac{b}{-} \frac{3s}{c} 1;$$
 (33)

$$b_{g} = \frac{1}{+} \left(\frac{ss_{1}}{+} + \frac{2sb + \frac{2}{+}}{+} \right) G \qquad \frac{1}{-} \left(\frac{ss_{1}}{+} + \frac{2sc + \frac{2}{+}}{+} \right) G_{+}
- \frac{c}{+} \frac{3s}{b} \qquad 1 \ln \frac{s}{-} \frac{b}{-} \frac{3s}{c} \qquad 1 \ln \frac{s}{+} + \frac{2s^{2} - \frac{2}{+} - \frac{2}{+}}{2 +}; \quad (34)$$

$$a = (+ +); \quad b = s_{1} + ; \quad c = s_{1} + +; ;$$

$$G = \ln^{2} \frac{-}{s} + \frac{2^{-2}}{3} \qquad 2L_{\frac{1}{2}} \qquad 1 \quad \frac{s_{1}}{s} \qquad 2L_{\frac{1}{2}} \qquad 1 + \frac{s_{1}}{-} ; ;$$

$$G_{+} = \ln^{2} \frac{+}{s} + \frac{2^{-2}}{3} \qquad 2L_{\frac{1}{2}} \qquad 1 \quad \frac{s_{1}}{s} \qquad 2L_{\frac{1}{2}} \qquad 1 + \frac{s_{1}}{-} ; ;$$

$$g_{+} = \ln^{2} \frac{+}{s} + \frac{2^{-2}}{3} \qquad 2L_{\frac{1}{2}} \qquad 1 \quad \frac{s_{1}}{s} \qquad 2L_{\frac{1}{2}} \qquad 1 + \frac{s_{1}}{-} ; ;$$

Infrared singularity (the presence of photon mass in) is compensated by taking into account soft photon emission from the initial particles:

$$d^{\text{soft}} = d_{0} - \frac{1}{12};$$

$$(35)$$

$$_{12} = \frac{1}{4} \frac{Z}{l_{1}} \frac{d^{3}k}{l_{1}} \frac{p_{+}}{p_{+}k} \frac{p}{p_{-}k} \frac{2}{l_{1}} = 2(l_{s} - 1) \ln \frac{m}{m} + \frac{1}{2}l_{s}^{2} - \frac{2}{3}$$

as a result the quantity in formulae (28) will be

$$! = (4 \ln \frac{"}{"} + 3) (l_s - 1) - 3l + \frac{2^{-2}}{3} - \frac{3}{2};$$
(36)

Cross section of two hard photon emission for the case when one of them is emitted collinearly to the incoming electron or positron can be obtained by means of the quasi{real electron method [7]:

$$\frac{d^{j}}{dx \ dc \ ds_{1}} = dW_{p} \ (k_{3}) \frac{d^{j} (p \ (1 \ x_{3}); p_{+}; k_{1}; q_{+}; q)}{dx \ dc \ ds_{1}} + dW_{p_{+}} \ (k_{3}) \frac{d^{j} (p \ ; p_{+} \ (1 \ x_{3}); k_{1}; q_{+}; q)}{dx \ dc \ ds_{1}};$$
(37)

with

$$dW_{p}(k_{3}) = -[(1 \quad x_{3} \quad \frac{x_{3}^{2}}{2}) \ln \frac{("_{0})^{2}}{m^{2}} \quad (1 \quad x_{3})]\frac{dx_{3}}{x_{3}}; \quad x_{3} = \frac{!_{3}}{"}; \quad x_{3} > \frac{"}{"}: \quad (38)$$

Here we suppose that the polar angle $_3$ between the directions of the additional collinear photon and the beam axis does not exceed some small value $_0$ 1, " $_0$ m.

The boosted di erential cross section $d\sim_B^j$ (p x; p+y; k_1; q+; q) with reduced m on enta of the incoming particles reads (compare with Eq. (16))

$$\frac{d\sim_{B} (p_{+} x_{2}; p x_{1}; k_{1}; q_{+}; q)}{dx \ dcds_{1}} = \frac{{}^{3} (1+2 2_{+}) (x_{1}^{2} (1 c_{1}^{2} + x_{2}^{2} (1+c)^{2})}{s_{1} sx_{1} x_{2} (1 c_{1}^{2}) (x_{1} + x_{2} + c(x_{2} x_{1}))};$$

$$\frac{d\sim_{B} (p_{+} x_{2}; p x_{1}; k_{1}; q_{+}; q)}{dx \ dcds_{1}} = \frac{{}^{3} ((1)) (x_{1}^{2} (1 c_{1}^{2} + x_{2}^{2} (1+c)^{2})}{s_{1} sx_{1}^{2} x_{2}^{2} (1 c_{1}^{2}) (x_{1} + x_{2} + c(x_{2} x_{1}))};$$

$$= \frac{x}{y_{2}}; \qquad y_{2} = \frac{2x_{1} x_{2}}{x_{1} + x_{2} + c(x_{2} x_{1})};$$
(39)

In a certain experim ental situation, an estim ate of the contribution of the additional hard photon em ission outside the narrow cones around the beam axes. It can be estimated by

$$\frac{d^{j}}{dx \ dc \ ds_{1}} = \frac{2}{4^{2}} \frac{d^{3}k_{3}}{!_{3}} \frac{\mathbf{u}^{2} + (\mathbf{u}^{2} \cdot !_{3})^{2}}{\mathbf{u}^{2} \cdot \mathbf{u}^{2}} \frac{1}{k_{3}p} \frac{d^{j}}{p} (p \ (1 \ x_{3}); p_{+}; k_{1}; q_{+}; q)}{dx \ dc \ ds_{1}} + \frac{1}{k_{3}p_{+}} \frac{d^{j}}{p} (p \ ; p_{+} \ (1 \ x_{3}); k_{1}; q_{+}; q)}{dx \ dc \ ds_{1}} ; \qquad x_{3} = \frac{!_{3}}{\mathbf{u}} : \qquad (40)$$

It is a simplied expression for the two {photon initial state emission cross section. Deviation for the case of large angles emission of our estimation from the exact quantity is small. It does not depend on s and slightly depends on $_0$. For $_0$ 10² we have

$$-\frac{\overset{R}{(d_{j,noncoll}} d_{j,noncollexact})}{\overset{d}{\underset{B}{}^{j}} < 10^{-1}:$$
(41)

By sum m ing all contributions for charge {even set {up we can put the cross section of radiative production in the form :

$$\frac{d^{-j}(p_{+};p_{-};k_{1};q_{+};q_{-})}{dx \ dc \ ds_{1}} = \frac{Z^{1} \ Z^{1}}{j!} \frac{dx_{1}dx_{2}}{(sx_{1}x_{2})^{\frac{1}{2}}} \frac{d^{-j}_{B}(p_{+},x_{2};p_{-}x_{1};k_{1};q_{+};q_{-})}{dx \ dc \ ds_{1}}$$
(42)

$$D(x_{1};l_{s})D(x_{2};l_{s}) \ 1 + -K^{-j} + \frac{Z^{1}}{2} \ dx \ \frac{1 + (1 - x)^{2}}{x} \ \ln \frac{2}{q} + x$$

$$\frac{d^{-j}_{B}(p((1 - x);p_{+};k_{1};q_{+};q_{-}))}{dx \ dc \ ds_{1}} + \frac{d^{-j}_{B}(p(p);p_{+}(1 - x);k_{1};q_{+};q_{-})}{dx \ dc \ ds_{1}} + \frac{d^{-j}_{-};noncoll}{dx \ dc \ ds_{1}};$$

$$D(x_{1};l_{s}) = (1 - x) + \frac{2}{2} P^{(1)}(x) \ (l_{s} - 1) + ::::;$$

$$P^{(1)}(x) = \frac{1 + x^{2}}{1 - x}; \quad j = ; :$$

The boosted cross sections d^{\sim} is de ned above in Eq. (39). The lower lim its of the integrals over $x_{1,2}$ depend on experimental conditions.

The structure function D include all dependence on the large logarithm l_s . The so{called K -factor reads

$$K^{j} = \frac{1}{R^{j}}B \qquad i^{(v)j} + i^{(s)j} + i^{(h)j} + \frac{2}{3} - \frac{3}{4} + R^{(j)}:$$
(43)

It includes the RC which connected with nal state interaction.

Q uantities R $^{(j)}$ include the "non-leading" contributions from the initial state radiation. Generally, they are rather cum bersom e expressions for the case s_1 s. For the case q M 2 s we obtain

$$R^{(1)} = \frac{1}{16(1+c^{2})} \frac{2\frac{1+c^{2}}{1-c}}{2} 32\frac{5+2c+c^{2}}{1-c}\ln^{2} \frac{2}{1+c} + 4\frac{5-c}{1+c}\ln \frac{2}{1+c} + \frac{2}{1+c} + \frac{2}{1+c} + \frac{2}{1+c} + \frac{2}{1+c} + 4x_{+}x_{-} 2\frac{1+c^{2}}{1-c} + 4 + (c!-c); \quad (44)$$

$$R^{(1)} = \frac{1-c^{2}}{16(1+c^{2})} 4\frac{5+2c+c^{2}}{1-c}\ln^{2} \frac{2}{1+c} + \frac{45-c}{1+c}\ln \frac{2}{1+c} + \frac{1+c^{2}}{1+c} + 2 + (c!-c); \quad (45)$$

Here we see the remarkable phenomena: the cancellation of terms containing $\ln(\frac{s}{s_1})$. In such a way only one kind of large logarithm $\ln(s=m^2)$. This fact is the sequence of renormalization group.

In paper [5] the RC to one {photon annihilation of ee to hadrons was calculated beyond the leading logarithm ic (LL) approximation. In paper [2] the heavy photon C ompton tensor was calculated for the scattering channel.

We had considered the charge-even contributions, which correspond to the charge-blind experimental set{up. The charge{odd contribution to the cross section under consideration appears starting from the Born level [see Eq. (10)]. In the kinematics discussed here, the odd part of the cross section is suppressed by the factor $s_1=s$ 1 with respect to the even one.

In papers [4] the inclusive event selection set-up was considered for radiative pion pair production process. In such a set-up the hard photon is emitted preferably in direction close to the beam axes, unlike the kinematics considered here.

We see that cross section in our quasi 2 ! 2 kinematics can be written down in the form of cross{section of D rell{Yan process. It is not a trivial fact as well as that two types of large logarithms are present in the problem. Possible background from peripherical process ee ! ee is negligible in our kinematics: it is suppressed by factor $-\frac{s_1}{s}$ and can be eliminated if the registration of the primary hard photon (see Eq. (1)) is required by experimental conditions for event selection.

A cknow ledgm ents

One of us (E A K.) is grateful to Slovak Institute for Physics, Bratislava for hospitality when the nalpart of this work was done. This work was supported by RFBR grant 01-02-17437 and INTAS grant 00366.

Appendix

The one { bop QED form factors of m uon and pion are

$$\begin{aligned} \operatorname{ReF}_{1}^{(1)}(\mathbf{s}_{1}) &= 1 + -f_{1}^{(1)}(\mathbf{s}_{1}); \quad \operatorname{ReF}_{2}^{(1)}(\mathbf{s}_{1}) &= -f_{2}^{(1)}(\mathbf{s}_{1}) \\ f_{1}^{(1)}(\mathbf{s}_{1}) &= \ln \frac{M}{-} \quad 1 \quad 1 \quad \frac{1 + \frac{2}{2}}{2} \mathbf{1} \quad + \frac{1 + \frac{2}{2}}{2} \quad \frac{1}{4}\mathbf{l}^{2} + 1 \ln \frac{1 + \frac{2}{2}}{2} + \frac{2}{3} \\ &+ 2\operatorname{Li}_{2} \quad \frac{1}{1 + } \quad \frac{1}{4}\mathbf{l}; \\ f_{2}^{(1)}(\mathbf{s}_{1}) &= \quad \frac{1 - \frac{2}{3}}{8}\mathbf{l}; \\ \operatorname{ReF}^{QED}(\mathbf{s}_{1}) &= \quad 1 + -f^{QED}(\mathbf{s}_{1}); \\ f^{QED}(\mathbf{s}_{1}) &= \quad \ln \frac{M}{-} \quad 1 \quad 1 \quad \frac{1 + \frac{2}{2}}{2}\mathbf{1} \quad + \frac{1 + \frac{2}{2}}{2} \quad \frac{1}{4}\mathbf{l}^{2} + 1 \ln \frac{1 + \frac{2}{2}}{2} + \frac{2}{3} \\ &+ 2\operatorname{Li}_{2} \quad \frac{1}{1 + } \quad ; \quad ^{2} = 1 \quad \frac{4\operatorname{M}^{2}}{\mathbf{s}_{1}}; \quad \mathbf{1} = \ln \frac{1 + \frac{1}{1} : \quad (A.1) \end{aligned}$$

W e put here the expressions for leptonic and hadronic [8] contributions into the vacuum polarization operator (s):

$$(s) = {}_{1}(s) + {}_{h}(s);$$

$${}_{1}(s) = {}_{-1}(s) + {}_{-2}(s) + {}_{-3}(s) + :::$$

$${}_{1}(s) = \frac{s}{4} PV \xrightarrow{Z^{1}} \frac{e^{+}e^{+} hadrons(s^{0})}{s^{0} s} ds^{0} i^{e^{+}e^{+} hadrons}(s) :$$

$$(A 2)$$

The rst order leptonic contribution is well known [1]:

$$f_{1}(s) = \frac{1}{3}L + \frac{5}{9} + f(x) + f(x) + i + \frac{1}{3} + (x) + (x)$$

where

$$f(x) = \begin{cases} 8 \\ < \frac{5}{9} & \frac{x}{3} + \frac{1}{6} (2 + x)^{p} \overline{1 - x} \ln \frac{1 + \frac{p}{1 - x}}{1 + \frac{p}{1 - x}} & \text{for } x \quad 1; \\ \vdots & \frac{5}{9} & \frac{x}{3} + \frac{1}{6} (2 + x)^{p} \overline{x - 1} \arctan \frac{p - 1}{1 - x + 1} & \text{for } x > 1; \\ (x) = \frac{1}{6} (2 + x)^{p} \overline{1 - x}; \quad x_{i} = \frac{4m^{2}}{s} : \end{cases}$$

In the second order it is enough to take only the logarithm ic term from the electron contribution

$$_{2}(s) = \frac{1}{4}(L \quad i) + (3) \quad \frac{5}{24}:$$
 (A.4)

- [1] V.V.Bytev, E.A.Kuraev and B.G.Shaikhatdenov, JINR preprint E2-2002-56.
- [2] E.A.Kuraev N.P.M erenkov and V.S.Fadin, Sov.J.Nucl. Phys. 45 (1987) 486.
- [3] A. I. Akhiezer and V. B. Berestetski, Quantum Electrodynamics, 1981.
- [4] V.A.Khoze et al, hep-ph/0202021; Eur. Phys. J.C 18 (2001) 481.
- [5] E.A.Kuraev, V.S.Fadin, Sov.J.Nucl.Phys.41 (1985) 466; O.Nikrosini, L.Trentadue, Phys.
 Lett. 196B (1987) 551.
- [6] H Anlaufet al.
- [7] P.Kessler, Nuovo C in ento 17 (1960) 809; V.N.Baier, V.S.Fadin and V.A.Khoze, NuclPhys.
 B 65 (1973) 381; V.N.Baier, V.S.Fadin and V.A.Khoze and E.A.Kuraev, PhysReports 78 (1981) 293.
- [8] S.Eidelm an and F. Jegerlehner, Z. Phys. C 67 (1995) 585.
- [9] F A. Berends and R. K leiss Nucl. Phys. B177 (1981) 237-262