

phase shifts and CP Violation in $\Lambda \rightarrow \Lambda^0 \pi^0$ Decay

C.C. Barros Jr.

Nuclear Theory and Elementary Particle Phenomenology Group
Instituto de Física, Universidade de São Paulo,
C.P. 66318, 05315-970, São Paulo, SP, Brazil
(Dated: March 24, 2024)

In the study of CP violation signals in $\Lambda \rightarrow \Lambda^0 \pi^0$ nonleptonic decays, the strong $J = 3/2$ P and D phase shifts for the Λ -state interactions are needed. These phases are calculated using an effective Lagrangian model, that considers π , η (1530), and the η' -term, in the intermediate states. The η' -term is calculated in terms of the scalar form factor of the baryon.

I. INTRODUCTION

In the search for physics beyond the standard model, the observation of CP violation could be a very useful tool. Many models show CP nonconservation effects, as, for example, the superweak model [1], the Kobayashi-Maskawa model [2], the penguin model [3] or the Weinberg-Higgs model [4].

Today, there are three systems where CP violation has been observed. The first one was the $K_L^0 \rightarrow \pi^0 \pi^0$ decay, where it was shown [5] that K_2^0 is not a pure eigenstate of CP, and then, the parameter ϵ is nonzero. Later, direct CP violation in $K \rightarrow \pi \pi$ decays has been observed. Very recently, CP violation has also been observed in the $B \rightarrow J/\psi K_S$ and other related modes (for a review, see [6]).

In 1957, Okubo [7] noted that CP violation could cause differences in the branching ratios of the $\Lambda \rightarrow \Lambda^0 \pi^0$ decays. Pais [8] extended this proposal also to $\Lambda \rightarrow \Lambda^0 \eta$ decays. In more recent works the CP violation signs were also investigated using nonleptonic hyperon decays [9]–[13] where decays were studied too. In [14] the study has been made in the framework of the standard model, and in [15], [16], new physics was considered. At the experimental level, there are experiments searching for CP violation in hyperon decays [17].

The nonleptonic decays have only been studied in [18] where the strong phases were estimated at leading order in heavy-baryon chiral perturbation theory. The aim of this paper is to calculate the strong phases using an effective Lagrangian model, without the heavy-baryon approximation, and their effects to the asymmetry parameters in the $\Lambda \rightarrow \Lambda^0 \pi^0$ decays.

The size of the CP violation depends on the Λ -state strong interaction between the produced particles. So, in order to perform the calculations, the strong phase shifts are needed. At the moment, the situation of the $\Lambda \rightarrow \Lambda^0 \pi^0$ and $\Lambda \rightarrow \Lambda^0 \eta$ decays is very comfortable, the Λ -state strong phase shift analysis is very well known experimentally [19]. At the theoretical level, this system is very well described, and, at least at low energies, the chiral perturbation theory is very precise [20], [21].

However, in the decays that produce hyperons, the situation is not so good, no experimental data is available to the Λ -state interactions. In fact, some information can be

obtained in the study of hyperonic atoms (see, for example, [22], where the coupling constant is estimated), but it is not enough to fully understand the interactions. Thus, to investigate the Λ -state interactions, the only way is to use a model.

As it was said, the chiral perturbation theory is very accurate when applied to the Λ -state interactions, so, we hope that it works in the Λ -system too. In [10]–[13], [23] that was done to the $\Lambda \rightarrow \Lambda^0 \pi^0$ decay, and the calculated phases were very small.

In this work, the $\Lambda \rightarrow \Lambda^0 \pi^0$ is studied, and the strong phase shifts for Λ -state interactions are calculated using the model presented in [24]. In [24], chiral lagrangians are used, describing processes with π , η and η' in the intermediate states. The η' -term is also included, but not only as a parametrization (as it was done in [24]), but relating it with the scalar form factor $\langle \pi | \bar{\psi} \psi | \pi \rangle$, based in the results of [25], [26].

This paper will show the following contents: In section II it will be shown the $\Lambda \rightarrow \Lambda^0 \pi^0$ decay and how to calculate the observables. In section III we will calculate the phase shifts in the Λ -state interactions. The results and conclusions are in section IV.

II. NONLEPTONIC DECAY

In the $\Lambda \rightarrow \Lambda^0 \pi^0$ ($J^P = \frac{3}{2}^+$) decays, the transitions are of the form

$$spin\ 3/2 \rightarrow spin\ 0 + spin\ 1/2; \quad (1)$$

and the contributing phases are the $J = \frac{3}{2}$ P (parity conserving) and D (parity violating) waves in the rest frame. The $S = 1$ nonleptonic decays are $\Lambda \rightarrow \Lambda^0 \pi^0$ and $\Lambda \rightarrow \Lambda^0 \eta$.

The experimental observables are the total rate Γ , and the asymmetry parameters, that can be written as

$$\alpha = 2 \operatorname{Re}(P/D) = (P^2/D^2 + D^2/P^2) \quad (2)$$

$$\beta = 2 \operatorname{Im}(P/D) = (P^2/D^2 - D^2/P^2) \quad (3)$$

$$\gamma = (P^2/D^2 - D^2/P^2) = (P^2/D^2 + D^2/P^2) \quad (4)$$

and obeys the relation

$$P^2 + D^2 + F^2 = 1 : \quad (5)$$

For antihyperons decays the expressions are

$$\overline{A} = 2 \operatorname{Re}(\overline{P} \overline{D}) = (\overline{P} f + \overline{D} f) \quad (6)$$

$$\overline{B} = 2 \operatorname{Im}(\overline{P} \overline{D}) = (\overline{P} f + \overline{D} f) : \quad (7)$$

The P and D amplitudes can be parametrized as

$$P = \sum_I a_{2I} P_{2I} e^{i(\frac{2I}{P} + \frac{2I}{P})} \quad (8)$$

$$D = \sum_I a_{2I} D_{2I} e^{i(\frac{2I}{D} + \frac{2I}{D})} ; \quad (9)$$

where I is the isospin state, $\frac{2I}{P}$ are the strong phase shifts and $\frac{2I}{D}$ are the weak CP violating phases. The respective CP conjugated amplitudes are

$$\overline{P} = \sum_I a_{2I} P_{2I} e^{i(\frac{2I}{P} - \frac{2I}{P})} \quad (10)$$

$$\overline{D} = \sum_I a_{2I} D_{2I} e^{i(\frac{2I}{D} - \frac{2I}{D})} ; \quad (11)$$

Using eq. (8), (9) for the Λ^0 decay we have

$$P(\Lambda^0) = \frac{r}{2} P_1 e^{i(\frac{1}{P} + \frac{1}{P})} + \frac{r}{3} P_3 e^{i(\frac{3}{P} + \frac{3}{P})} \quad (12)$$

$$D(\Lambda^0) = \frac{r}{2} D_1 e^{i(\frac{1}{D} + \frac{1}{D})} + \frac{r}{3} D_3 e^{i(\frac{3}{D} + \frac{3}{D})} \quad (13)$$

and for Σ^0

$$P(\Sigma^0) = \frac{r}{3} P_1 e^{i(\frac{1}{P} + \frac{1}{P})} + \frac{r}{3} P_3 e^{i(\frac{3}{P} + \frac{3}{P})} \quad (14)$$

$$D(\Sigma^0) = \frac{r}{3} D_1 e^{i(\frac{1}{D} + \frac{1}{D})} + \frac{r}{3} D_3 e^{i(\frac{3}{D} + \frac{3}{D})} : \quad (15)$$

In the limit of CP conservation, the CP asymmetry parameters

$$A = \frac{+}{-} \quad (16)$$

and

$$B = \frac{+}{-} \quad (17)$$

vanish, since $= -$ and $= -$. In hyperon decays, the $I=3/2$ amplitudes are much smaller than the $I=1/2$, then, in the first order in $I=3/2$ amplitudes,

$$\begin{aligned} A(\Lambda^0) &= \tan\left(\frac{1}{P} - \frac{1}{D}\right) \tan\left(\frac{1}{P} - \frac{1}{D}\right) + 1 + \\ &+ \frac{1}{P} \frac{P_3}{2 P_1} \frac{\cos\left(\frac{3}{P} - \frac{1}{D}\right) \cos\left(\frac{3}{P} - \frac{1}{D}\right)}{\cos\left(\frac{1}{P} - \frac{1}{D}\right) \cos\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &\frac{\sin\left(\frac{3}{P} - \frac{1}{D}\right) \sin\left(\frac{3}{P} - \frac{1}{D}\right)}{\sin\left(\frac{1}{P} - \frac{1}{D}\right) \sin\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &+ \frac{1}{P} \frac{D_3}{2 D_1} \frac{\cos\left(\frac{3}{P} - \frac{1}{D}\right) \cos\left(\frac{3}{P} - \frac{1}{D}\right)}{\cos\left(\frac{1}{P} - \frac{1}{D}\right) \cos\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &\frac{\sin\left(\frac{3}{P} - \frac{1}{D}\right) \sin\left(\frac{3}{P} - \frac{1}{D}\right)}{\sin\left(\frac{1}{P} - \frac{1}{D}\right) \sin\left(\frac{1}{P} - \frac{1}{D}\right)} \end{aligned} \quad (18)$$

$$\begin{aligned} B(\Lambda^0) &= \cot\left(\frac{1}{P} - \frac{1}{D}\right) \tan\left(\frac{1}{P} - \frac{1}{D}\right) + 1 + \\ &+ \frac{1}{P} \frac{P_3}{2 P_1} \frac{\sin\left(\frac{3}{P} - \frac{1}{D}\right) \cos\left(\frac{3}{P} - \frac{1}{D}\right)}{\sin\left(\frac{1}{P} - \frac{1}{D}\right) \cos\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &\frac{\cos\left(\frac{3}{P} - \frac{1}{D}\right) \sin\left(\frac{3}{P} - \frac{1}{D}\right)}{\cos\left(\frac{1}{P} - \frac{1}{D}\right) \sin\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &+ \frac{1}{P} \frac{D_3}{2 D_1} \frac{\sin\left(\frac{3}{P} - \frac{1}{D}\right) \cos\left(\frac{3}{P} - \frac{1}{D}\right)}{\sin\left(\frac{1}{P} - \frac{1}{D}\right) \cos\left(\frac{1}{P} - \frac{1}{D}\right)} \\ &\frac{\cos\left(\frac{3}{P} - \frac{1}{D}\right) \sin\left(\frac{3}{P} - \frac{1}{D}\right)}{\cos\left(\frac{1}{P} - \frac{1}{D}\right) \sin\left(\frac{1}{P} - \frac{1}{D}\right)} \end{aligned} \quad (19)$$

and similar expressions for the Σ^0 decays (replacing the factors $1=2$ for P in the expressions (18), (19)). At leading order,

$$A(\Lambda^0) = A(\Sigma^0) = \tan\left(\frac{1}{P} - \frac{1}{D}\right) \tan\left(\frac{1}{P} - \frac{1}{D}\right) ; \quad (20)$$

and

$$B(\Lambda^0) = B(\Sigma^0) = \cot\left(\frac{1}{P} - \frac{1}{D}\right) \tan\left(\frac{1}{P} - \frac{1}{D}\right) : \quad (21)$$

In the next section we will calculate the phase shifts $\frac{2I}{P}$, that are needed to estimate A and B .

III. LOW ENERGY INTERACTION

In order to describe the low energy interaction, a reliable way is to use effective Lagrangians, as it was done in a previous work [24]. A very important feature of this model is to allow the inclusion of spin 3/2 resonances in the intermediate states. In the low energy $\Lambda^0 P$ interactions, for example, the (1232) dominates almost completely the total cross section (when P is near the mass). Consequently, it is expected that in some reactions of Λ^0 scattering, the same behaviour will occur [24]. The lagrangians to be considered are

$$L = \frac{g}{2m} [\bar{\psi} \gamma_5 \psi] \phi \sim \quad (22)$$

$$L = g \bar{\psi} \gamma_5 \psi \left(Z + \frac{1}{2} \right) \sim g \phi \sim + H \psi :$$

(23)

$$L = \frac{0}{2} [\bar{\psi} \sim \psi] + \frac{0}{2} [\bar{\psi} (\frac{0}{4m}) i \sim \psi] : (\psi \sim \psi) : (24)$$

$$L = 0 \sim : (\psi \sim \psi) : \frac{0}{4m^2} (\psi \sim \psi) : (\psi \sim \psi) ; \quad (25)$$

where ψ , $\bar{\psi}$, \sim and \sim are the cascade, the resonance (1530), the pion and the rho mesons. Z is the ϕ -shell parameter [21], g and g' are the magnetic moments.

The lagrangians are almost the same as the N ones [21], because the N system is very similar to the Δ . N and Δ are particles with isospin 1/2, the only difference is that (1232) has isospin 3/2 and (1530), 1/2, so, a Δ matrix is included in (23).

The spin 3/2 propagator for a mass M particle, is then

$$G(p) = \frac{(p + M)}{p^2 - M^2} g \frac{1}{3} \left(\frac{p}{3M} + \frac{p}{3M} - \frac{2p \cdot p}{3M^2} \right) ; \quad (26)$$




FIG. 1: Diagrams to Interaction.

The contributing diagrams are shown in the Fig. 1 (we show only the direct diagrams, but in the calculations, the crossed diagrams are also included). The scattering matrix will have the general form

$$T^{ba} = \bar{u}(p) [A + \frac{(k + k^0)}{2} B] u(p) + [A + \frac{(k + k^0)}{2} A] i_{bac} \epsilon_{\mu\nu} g_{\mu\nu} ; \quad (27)$$

where k and k^0 are the initial and final momenta. Calculating the amplitudes from the diagrams, the contributions from Fig. 1 (a) (intermediate Δ) are

$$\begin{aligned} A^+ &= \frac{g^2}{m} \\ A &= 0 \\ B^+ &= g^2 \frac{1}{u - m^2} - \frac{1}{s - m^2} \\ B &= \frac{g^2}{2m} - g^2 \frac{1}{u - m^2} + \frac{1}{s - m^2} ; \quad (28) \end{aligned}$$

Fig 1 (d), the ρ exchange, gives

$$A^+ = B^+ = 0$$

$$\begin{aligned} A &= \frac{g^2}{m^2} \left(\frac{1}{1 - \frac{t=4m^2}{t-m^2}} \right) \\ B &= \frac{g^2}{m^2} \left(1 + \frac{1}{1 - \frac{t=4m^2}{t-m^2}} \right) ; \quad (29) \end{aligned}$$

The contribution from Fig. 1 (b), the interaction with the intermediate Δ , is

$$\begin{aligned} A^+ &= \frac{g^2}{3m} f \frac{r}{2} \hat{A} \\ &\quad \frac{m^2 + m}{m^2} (2m^2 - m^2 - m^2 + 2^2) \\ &\quad + \frac{4m}{m^2} [(m + m)Z + (2m + m)Z^2] k \cdot k^0 g \end{aligned} \quad (30)$$

$$A = \frac{g^2}{3m} f \frac{r}{2} \hat{A} + \frac{8m^2}{m^2} [(m + m)Z + (2m + m)Z^2] g \quad (31)$$

$$B^+ = \frac{g^2}{3m} f \frac{r}{2} \hat{B} - \frac{8m^2}{m^2} \frac{Z^2}{g} \quad (32)$$

$$\begin{aligned} B &= \frac{g^2}{3m} f \frac{r}{2} \hat{B} - \frac{4m}{m^2} [(2m^2 \\ &\quad + 2m^2 - m^2 - 2^2)Z + (2m^2 + 4m^2 - m^2)Z^2] \\ &\quad + \frac{(m + m)^2}{m^2} \frac{4m}{m^2} Z^2 k \cdot k^0 g ; \quad (33) \end{aligned}$$

with

$$\begin{aligned} \hat{A} &= \frac{(m + m)^2}{2m^2} [2m^3 - 2m^3 - 2m^2 m^2 \\ &\quad + 2m^2 m^2 + 2(2m^2 - m^2)] + \\ &\quad + \frac{3}{2} (m + m) t \\ \hat{B} &= \frac{1}{2m^2} [(m^2 - m^2)^2 - 2m^2 m^2 (m + m)^2 \\ &\quad + 6^2 m^2 (m + m)^2 - 2^2 (m + m)^2 + 4] + \frac{3}{2} t ; \end{aligned} \quad (34)$$

where t is the pion mass, and u and s are defined in the appendix A. One must remark that eq. (35) is different from the expression of [24], where there was a mistake. The correct expression is presented here.

In [24], [21] the ρ term (diagram 1.c) was simply considered as a parametrization

$$\begin{aligned} A &= a + bt \\ B &= 0 ; \quad (36) \end{aligned}$$

In fact, the ρ term represents the exchange of a scalar isoscalar system in the t -channel. This contribution is related to the scalar form factor of the baryon, and at large distances is dominated by triangle diagrams (Figure 2)

involving the exchange of 2 pions [27]. In the interaction, this contribution is associated with two triangle diagrams, with and intermediate states, as it was calculated in [26].




FIG. 2: The scalar form factor (grey blob) receives contributions from tree interactions (white blob) and triangle diagrams with spin 1=2 and 3=2 intermediate states.

The scalar form factor for a spin 1=2 baryon B is defined as $\langle B(p^0) | L_{sb} | B(p) \rangle = (t) u(p^0) u(p)$, where L_{sb} is the chiral symmetry breaking lagrangian.

The contribution of an intermediate particle of spin s and mass M to the scalar form factor is given by

$$\begin{aligned} \langle s(t; M) | u u \rangle &= \frac{g^2}{2m} T_a^y T_a \\ &= \frac{d^4 Q}{(2)^4} \frac{[u_s u]}{[(Q - q^2)^2 - M^2][(Q + q^2)^2 - M^2]} ; \end{aligned} \quad (37)$$

where Q is the average of the internal pion momenta, T is a vertex isospin matrix and

$$\begin{aligned} [u_{1=2} u] &= u \left((m + M) + \frac{(m - M)(m + M)^2}{s - M^2} \right) \\ &+ 1 + \frac{(m + M)^2}{s - M^2} \mathcal{Q} u ; \end{aligned}$$

$$\begin{aligned} [u_{3=2} u] &= u \left(\frac{1}{s - M^2} (m + M) (2 - t=2) \right. \\ &\quad \left. \frac{(2M + m)^4}{6M^2} + \frac{m^2 - M^2}{s - M^2} 1 - \frac{(m + M)}{6M^2} (m + M) (2M - m) + 2^2 \right. \\ &\quad \left. \frac{m(s - m^2)}{6M^2} + \frac{1}{s - M^2} (2 - t=2) + \frac{2m}{3} (m + M) \right. \\ &\quad \left. \frac{(m + M)^2}{3M} - \frac{4}{6M^2} + \frac{m^2 - M^2}{s - M^2} 1 \right. \\ &\quad \left. \frac{1}{6M^2} M^2 + 2mM - m^2 + 2^2 \frac{s - m^2}{6M^2} \right) \mathcal{Q} u ; \end{aligned} \quad (39)$$

Calculating (t) in the interaction, and using the loop integrals defined in appendix B, we obtain

$$\begin{aligned} 1=2(t) &= \frac{2}{(4)^2} \frac{g^2}{2m} (m + m) \frac{h}{cc}^{(000)} \\ &= \frac{m^2 - m^2}{2m} \frac{(000)}{sc} - \frac{m + m}{2m} \frac{(001)}{sc} ; \end{aligned} \quad (40)$$

$$3=2(t) = \frac{2}{(4)^2} \frac{g^2}{2m} \frac{1}{6m^2} (m + m)^2$$

$$\begin{aligned} & (2m - m) + 2^2 (m + m) + (2 - t=2) m \frac{(000)}{cc} \\ & 2^2 m \frac{(000)}{cc} + (m^2 - m^2) (m + m)^2 (2m - m) \\ & + 2^2 (m + m) (m^2 - m^2) \\ & + 6 (2 - t=2) m^2 (m + m)^4 (2m + m) \frac{(000)}{2m} \frac{sc}{sc} \\ & + (m + m)^2 (4m - m - m^2 - m^2) + 6m^2 (2 - t=2) \\ & 2^2 (m + m) (2m - m)^4 \frac{(001)}{2m} \frac{sc}{sc} : \end{aligned} \quad (41)$$

More details, as, for example regularization and the determination of $(t=0)$ can be found in [26].

The partial wave amplitudes are obtained summing the contributions from the diagrams of g. 1 and making a straightforward application of the expressions found in appendix A. This can be done, for example, in the center-of-mass frame, where \mathbf{p} is the momentum and $x = \cos \theta$, where θ is the scattering angle. One notes that the a_1 amplitudes are real, and, so, the corresponding S matrix is not unitary. To unitarize the amplitudes, we reinterpret them as elements of the K matrix [28], and then

$$a_1^U = \frac{a_1}{1 - i a_1} ; \quad (42)$$

where U means unitarized. Now the phase shifts are

$$\delta_1 = \tan^{-1} (f_1) ; \quad (43)$$

The parameters used are the same that were used in [24] and are $m = 1.318$ GeV, $m = 1.533$ GeV, $\mu = 1.25$, $\alpha = 0.349$, $g = 4$ and $\frac{2}{3}m^2 = 1 = (2f^2)$, with $f = 93$ MeV. The coupling constant g can be calculated comparing the resonant $\frac{1}{P}$ phase with the Breit-Wigner expression

$$\delta_1 = \tan^{-1} \frac{2 \frac{0}{2(m_r - p)} \frac{21}{3} \frac{7}{5}}{1} \quad (44)$$

where μ_0 is the center-of-mass momentum at the peak of the resonance. The obtained value is 4.54 GeV $^{\frac{1}{2}}$. The numerical results of the phase shifts at $\frac{1}{P} s = m$ are

$$\delta_P = 10.173^\circ \text{ and } \delta_D = 0.208^\circ \quad (45)$$

$$\delta_P = 0.106^\circ \text{ and } \delta_D = 0.078^\circ : \quad (46)$$

IV. SUMMARY AND CONCLUSIONS

We have calculated the strong P and D phase shifts for the decay at its mass including the contributions

from the diagrams of Fig. 1. The numerical values of the phases are shown in the expressions (45), (46). The respective values calculated in [18] are $\frac{1}{P} = 12.8^\circ$ and $\frac{1}{P} = 1.1^\circ$, that are greater in magnitude than the ones obtained in this work. One must remark that the calculations presented here have no heavy-baryon approximation and the π -exchange and the π -term are included, that are the sources of the differences. As we can see in eq. (45), (46), the D phases are much smaller than the P ones, as it is expected at low energies, and can even be neglected in a first approximation. We expect that the same pattern occurs for the weak phases.

In the other hyperon decays, the weak phases are of the order $\frac{1}{P} \sim 10^{-3}$ in the Weinberg-Higgs model and $\frac{1}{P} \sim 10^{-4}$ in the Kobayashi-Maskawa model [9]. In [18], using the KM model, the estimated value was $\frac{1}{P} \sim 10^{-3}$, which is significantly larger when compared with the other hyperons.

The asymmetry parameter A , eq. (20), in the decays $\Lambda \rightarrow p \pi^-$ depends on the factor $\tan(\frac{1}{P} - \frac{1}{D}) \sim -0.18$. In the $\Lambda \rightarrow p \pi^0$ decay, the factor is $\tan(\frac{1}{S} - \frac{1}{P}) \sim -0.12$, and using the results of [10], for the $\Sigma^- \rightarrow p \pi^-$ phases, in the $\Lambda \rightarrow p \pi^-$ decay, $\tan(\frac{1}{S} - \frac{1}{P}) \sim 0.05$. One notes that the greatest value happens in the $\Lambda \rightarrow p \pi^-$ decay (and the weak phases are also larger), so, we conclude that the A parameter must be the largest in this decay. On the other hand, considering the B parameter eq. (21), the $\Lambda \rightarrow p \pi^-$ decay shows a smaller term, from $\cot(\frac{1}{P} - \frac{1}{D})$, but the term that depends on the weak phases is larger. So, the B parameter is probably of the same order of the one that appears in the other hyperon decays. The B parameter seems to be the one where the CP violation would be most evident.

Acknowledgments

I wish to thank professor Y. Hamada and professor M. R. Robilotta, for many helpful discussions. This work was supported by FAPESP and CNPq.

APPENDIX A: BASIC FORMALISM

In this paper p and p^0 are the initial and final hyperon 4-momenta, k and k^0 are the initial and final pion 4-momenta, so the Mandelstam variables are

$$s = (p + k)^2 = (p^0 + k^0)^2 \quad (A1)$$

$$t = (p - p^0)^2 = (k - k^0)^2 \quad (A2)$$

$$u = (p^0 - k)^2 = (p - k^0)^2 \quad (A3)$$

With these variables, we can define

$$= \frac{s - u}{4m} \quad (A4)$$

$$0 = \frac{2 - t}{4m} \quad (A5)$$

$$r = \frac{m_r^2 - m^2 - k k^0}{2m} ; \quad (A6)$$

where m , m_r and k are, respectively, the hyperon mass, the resonance mass and the pion mass. The scattering amplitude for an isospin I state is

$$T_I = \bar{u}(p) [A_I + \frac{(k + k^0)}{2} B_I] u(p) ; \quad (A7)$$

where A_I and B_I are calculated using the Feynman diagrams. So the scattering matrix is

$$M_I^{ba} = \frac{T_I^{ba}}{8 \frac{1}{P} S} = f_I(\omega) + \sim \pi g(\omega) = f_I^I + \frac{(\sim k^0)(\sim k)}{k k^0} f_2^I ; \quad (A8)$$

with

$$f_1^I(\omega) = \frac{(E + m)}{8 \frac{1}{P} S} [A_I + (\frac{P}{S} - m) B_I] ; \quad (A9)$$

$$f_2^I(\omega) = \frac{(E - m)}{8 \frac{1}{P} S} [A_I + (\frac{P}{S} + m) B_I] ; \quad (A10)$$

where E is the hyperon energy, and

$$A^{\frac{1}{2}} = A^+ + 2A^- ; A^{\frac{3}{2}} = A^+ - A^- ; \quad (A11)$$

and similar expressions holds to B^I . The partial-wave decomposition is done with

$$a_l = \frac{1}{2} \int_{-1}^1 P_l(x) f_l(x) + P_{l+1}(x) f_{l+1}(x) dx ; \quad (A12)$$

In our calculation (tree level) a_l is real. With the unitarization, as explained in Section III, we obtain

$$a_l^U = \frac{1}{2i} [e^{2i\delta_l} - 1] = e^{i\delta_l} \sin(\delta_l) ! a_l ; \quad (A13)$$

APPENDIX B: LOOP INTEGRALS

The basic loop integrals needed in order to perform the calculations of Fig. 2 are

$$I_{cc} = \int \frac{d^4 Q}{(2\pi)^4} \frac{1}{[(Q - q)^2 - m^2][(Q + q)^2 - m^2]} ; \quad (B1)$$

$$I_{sc} = \int \frac{d^4 Q}{(2\pi)^4} \frac{1}{[(Q - q)^2 - m^2][(Q + q)^2 - m^2]} \frac{2m}{[s - M^2]} ; \quad (B2)$$

The integrals are dimensionless and have the following tensor structure

$$I_{cc} = \frac{i}{(4\pi)^2} \sum_{cc}^{(000)} ; \quad (B3)$$

$$I_{cc} = \frac{i}{(4)^2} \frac{q \cdot q}{2} \frac{(000)}{cc} + g \frac{(000)}{cc} ; \quad (B 4)$$

$$I_{sc} = \frac{i}{(4)^2} \frac{n}{sc} \frac{(000)}{sc}^O ; \quad (B 5)$$

$$I_{sc} = \frac{i}{(4)^2} \frac{P}{m} \frac{(001)}{sc} : \quad (B 6)$$

$$\frac{(00n)}{sc} = \left(\frac{2m}{\hbar} \right)^{n+1} \int_0^Z da \int_0^Z db \frac{(ab=2)^n}{D_{sc}} ; \quad (B 9)$$

with

Thus, the integrals that appear in the text are

$$\frac{(n00)}{cc} = \int_0^Z da (1=2-a)^n \ln \frac{D_{cc}}{2} ; \quad (B 7)$$

$$\frac{(000)}{cc} = \frac{1}{2} \int_0^Z da \frac{D_{cc}}{2} \ln \frac{D_{cc}}{2} ; \quad (B 8)$$

$$D_{cc} = a(1-a)q^2 + \frac{2}{m^2} ;$$

$$D_{sc} = a(1-a)(1-b)q^2 + \left[\frac{2}{m^2} ab \left(\frac{2}{m^2} + m^2 M^2 \right) + a^2 b^2 m^2 \right] ;$$

-
- [1] L. Wolfenstein, Phys. Lett. 13, 562 (1984).
[2] M. Kobayashi and T. Maskawa, Prog. Theor. Phys. 49, 652 (1973).
[3] F. Gilman and M. Wise, Phys. Lett. 93B, 129 (1980).
[4] S. Weinberg, Phys. Rev. Lett. 37, 657 (1976).
[5] J. H. Christenson, J. W. Cronin, V. L. Fitch and R. Turley, Phys. Rev. Lett. 13, 138 (1964).
[6] Y. Nir, hep-ph/0109090 (2001).
[7] S. Okubo, Phys. Rev. 109, 984 (1958).
[8] A. Pais, Phys. Rev. Lett. 3, 242 (1959).
[9] J. F. Donoghue, E. Golowich, W. A. Ponce and B. R. Holstein, Phys. Rev. D 21, 186 (1980); J. F. Donoghue, X. -G. He and S. Pakvasa, Phys. Rev. D 34, 833 (1986); X. -G. He, H. Steger and G. Valencia, Phys. Lett. B 272, 411 (1991); O. E. Overseth and S. Pakvasa, Phys. Rev. 184, 1163 (1969).
[10] A. N. Kamal, Phys. Rev. D 58, 077501 (1998).
[11] A. Datta, P. O'Donnell and S. Pakvasa, hep-ph/9806374 (1998).
[12] J. Tandean, A. W. Thomas and G. Valencia, Phys. Rev. D 64, 014005 (2001).
[13] U. G. Meißner and J. A. Oller, Phys. Rev. D 64, 014006 (2001).
[14] J. Tandean, and G. Valencia, Phys. Rev. D 67, 056001 (2003).
[15] X. G. He and G. Valencia, Phys. Rev. D 52, 5257 (1995).
[16] X. G. He, H. Murayama, S. Pakvasa and G. Valencia, Phys. Rev. D 61, 071701 (2000).
[17] K. B. Luk et al., hep-ex/0005004.
[18] J. Tandean, and G. Valencia, Phys. Lett. B 451, 382 (1999).
[19] R. Kosh and E. Pietarinen, Nucl. Phys. A 336, 331 (1980).
[20] T. Becher and H. Leutwyler, hep-ph/0103263 (2001).
[21] E. T. O'sypowski, Nucl. Phys. B 21, 615 (1970); M. G. Olsson and E. T. O'sypowski, Nucl. Phys. B 101, 136 (1975); H. T. Coelho, T. K. Das and M. R. Robilotta, Phys. Rev. C 28, 1812 (1983).
[22] B. Loiseau and S. Wysocki, Phys. Rev. C 63, 034003 (2001).
[23] N. Keiser, Phys. Rev. C 64, 045204 (2001).
[24] C. C. Barros and Y. Hamada, Phys. Rev. C 63, 065203 (2001).
[25] M. R. Robilotta, Phys. Rev. C 63, 044004 (2001).
[26] C. C. Barros and M. R. Robilotta, hep-ph/0209213, submitted to physical review C.
[27] T. Becher and H. Leutwyler, Eur. Phys. Journal C 9, 643 (1999); JHEP 106, 17 (2001).
[28] H. B. Tang and P. J. Ellis, Phys. Rev. C 56, 3363 (1997).
[29] Particle Data Group, C. Caso et al., Eur. Phys. J. C 3, 1 (1998).