

QCD sum rule analysis of the coupling constants g and g_1 C. Aydın and A. H. Yılmaz^y

Physics Department, Karadeniz Technical University, Trabzon, Turkey

(December 21, 2021)

Abstract

The coupling constants g and g_1 are calculated using QCD sum rules method by studying the three point and $!$ correlation functions. A comparison of the results with the values of the coupling constants that are deduced from the experimentally measured decay widths of $!$ and $!!$ decays is performed.

PACS numbers: 12.38.Lg; 13.40.Hq; 14.40.Aq

Typeset using REVTeX

 coskun@risc01.ktu.edu.tr
^yhakany@risc01.ktu.edu.tr

The method of QCD sum rules is one of the efficient tools for studying hadron physics. This method has been successfully applied to calculate many hadronic observables, such as decay constants and form factors [1,2]. On the other hand, radiative transitions between pseudoscalar (P) mesons have been an important area of study in low-energy hadron physics for more than three decades. These transitions have been analyzed within the frameworks of phenomenological quark models, potential models, bag models, and also by employing effective Lagrangian methods [4,5]. The radiative transitions $V \rightarrow P$ are characterized by the coupling constants g_V . Since low energy hadron physics is governed by nonperturbative QCD, it is very difficult to obtain the numerical values of these coupling constants from the first principles. For this reason, some specific nonperturbative methods have to be developed to be used as calculational tools. Among these methods QCD sum rules have proved to be very useful to extract the coupling constants. A recent review of QCD sum rules method is provided in [6] where more references can also be found.

In this work, we calculate the coupling constants g and g_1 associated with the radiative decays $\rho \rightarrow \pi \gamma$ and $\omega \rightarrow \pi \gamma$ by employing the traditional QCD sum rules method which provides a model independent way to calculate the coupling constants. The coupling constant g was previously calculated by T.M. Aliiev et al. [7] in the framework of light cone QCD sum rules. Our analysis, therefore, complements the results obtained in that paper.

In accordance with the general strategy of QCD sum rules method, we begin by considering the three point correlation function

$$\langle 0 | T \{ j^\mu(x) j^\nu(y) \} | 0 \rangle = \int d^4x d^4y e^{ip \cdot y} e^{-ip \cdot x} \langle 0 | T \{ j^\mu(x) j^\nu(y) \} | 0 \rangle \quad (1)$$

where the interpolating currents j^μ for vector meson and j^ν are $j^\mu = \frac{1}{\sqrt{2}} (\bar{u} \gamma^\mu u - \bar{d} \gamma^\mu d)$, $j^\nu = \frac{1}{\sqrt{2}} (\bar{u} \gamma^\nu u + \bar{d} \gamma^\nu d)$, respectively. We take ω mixing into account and use the interpolating current for ω meson as $j^\mu = \frac{1}{\sqrt{2}} (\bar{u} \gamma^\mu u + \bar{d} \gamma^\mu d) \cos \theta - (\bar{s} \gamma^\mu s) \sin \theta$ where θ is the mixing angle in the quark-flavour basis. The electromagnetic quark current is given as $j^\mu = e_u \bar{u} \gamma^\mu u + e_d \bar{d} \gamma^\mu d$, where e_u and e_d denote the quark charges.

The theoretical part of the sum rule for the coupling constant g_V is calculated by considering the perturbative contribution and the power corrections from operators of different dimensions to the three point correlation function. In the spirit of QCD sum rules techniques, we consider the three point correlation function in the Euclidian region defined by $p^2 = -Q^2 \leq -1 \text{ GeV}^2$, $p'^2 = -Q'^2 \leq -1 \text{ GeV}^2$. In this region, the perturbative contribution can be approximated by the lowest order quark loop diagram shown in Fig. 1. Moreover, we consider the power corrections from operators of different dimensions, proportional to terms $\langle \bar{q}q \rangle$, $\langle \bar{q} G q \rangle$ and $\langle \bar{q} q \bar{q} q \rangle$. Since the gluon condensate contribution proportional to $\langle G^2 \rangle$ is estimated to be negligible for light quark systems, it is not taken into account. We perform the calculations of the power corrections in the fixed point gauge [8]. Moreover, we work in the SU(2) flavour context with $m_u = m_d = m_q$ and we work in the limit $m_q = 0$. In this limit, the perturbative quark-loop diagram does not make any contribution, and only contributions result from the operators of dimensions $d=3$ and $d=5$ that are proportional to $\langle \bar{q}q \rangle$ and $\langle \bar{q} G q \rangle$, respectively. The relevant Feynman diagrams for the calculation of the power corrections are shown in Fig. 2 and Fig. 3.

We then calculate the three point correlation function $\Pi(p; p')$ using phenomenological considerations. This function satisfies a double dispersion relation. We choose the vector and pseudoscalar channels and by saturating this dispersion relation by the lowest lying meson states in these channels the physical part of the sum rule is obtained as

$$\Pi(p; p') = \frac{\langle 0 | j^\mu_V j^\nu_V | 0 \rangle \langle V(p) | j^\mu_V j^\nu_V | p' \rangle \langle j^\mu_V j^\nu_V \rangle}{(p^2 - m_V^2)(p'^2 - m_V^2)} + \dots \quad (2)$$

where the contributions from the higher states and the continuum are shown by dots. The overlap amplitudes for vector and pseudoscalar mesons are $\langle 0 | j^\mu_V j^\nu_V | 0 \rangle = \epsilon_V u_V^\mu u_V^\nu$ where u_V is the polarization vector of the vector meson $V = \rho, \omega$ and $\langle j^\mu_V j^\nu_V \rangle = \dots$. The matrix element of the electromagnetic current is given by

$$\langle V(p) | j^\mu_V j^\nu_V | p' \rangle = i \frac{e}{m_V} g_V K(q^2) \epsilon^{\mu\nu\alpha\beta} p_\alpha p'_\beta \quad (3)$$

where $q = p - p'$ and $K(q^2)$ is a form factor with $K(0) = 1$. This matrix element defines the coupling constant g_V through the effective Lagrangian

$$L = \frac{e}{m_V} g_V \quad \text{at } V \text{ at } A \quad (4)$$

describing the V -vertex [9].

After performing the double Borel transform with respect to the variables Q^2 and Q'^2 , we obtain the sum rule for the coupling constant g_V in the form

$$g_V = \frac{m_V}{\sqrt{V}} e_M^{\frac{m_V^2}{2}} e_M^{\frac{m^2}{2}} e_u \langle \bar{u}u \rangle - e_d \langle \bar{d}d \rangle \left(\frac{3}{2} + \frac{5}{16} m_0^2 \frac{1}{M^2} - \frac{3}{16} m_0^2 \frac{1}{M'^2} \right) \cos \quad (5)$$

where the relation $\langle \bar{q} G q \rangle = m_0^2 \langle \bar{q}q \rangle$ is used. In this expression the plus sign is for meson and the minus sign is for η' meson. In the numerical evaluation of the sum rule the values $m_0^2 = (0.8 \pm 0.02) \text{ GeV}^2$, $\langle \bar{u}u \rangle = \langle \bar{d}d \rangle = (-0.014 \pm 0.002) \text{ GeV}^3$ [6], and $m = 0.77 \text{ GeV}$, $m_\eta = 0.781 \text{ GeV}$, $m_{\eta'} = 0.547 \text{ GeV}$ are used [10]. The overlap amplitudes for vector meson states are calculated using the experimental leptonic decay widths of $V \rightarrow e^+ e^-$ decays [10] and the values $\Gamma = (0.17 \pm 0.03) \text{ GeV}^2$ and $\Gamma_{\eta'} = (0.15 \pm 0.02) \text{ GeV}^2$ are obtained. The overlap amplitude for meson state was determined earlier by QCD sum rules analysis as $\Gamma = (0.23 \pm 0.03) \text{ GeV}^2$ [11]. We use the value of the mixing angle as $\theta = 19^\circ \pm 2^\circ$ [11].

The dependence of the coupling constants g_V on the Borel parameters M^2 and M'^2 are analyzed by studying the independent variations of M^2 and M'^2 in the interval $0.6 \text{ GeV}^2 \leq M^2, M'^2 \leq 1.4 \text{ GeV}^2$ since these limits determine the allowed interval for the vector channel [12]. We show the variation of the coupling constant g and $g_{\eta'}$ as a function of the Borel parameters M^2 for different values of M'^2 in Fig. 4 and in Fig. 5, respectively. These figures indicate that the sum rule is quite stable with these reasonable variations of M^2 and M'^2 . We choose the middle value $M^2 = 1 \text{ GeV}^2$ for the Borel parameter in its interval of variation and obtain the coupling constants g_V as $g = 1.2 \pm 0.3$ and $g_{\eta'} = 0.4 \pm 0.06$ where the uncertainties result from the variations of M^2 and M'^2 and from the estimated values of the vacuum condensates.

If we use the effective Lagrangian given in Eq. 4, then the decay width for $V \rightarrow \eta'$ is

obtained as

$$\langle V \rightarrow \pi \pi \rangle = \frac{(m_V^2 - m_\pi^2)^3}{24 m_V^5} g_V^2 : \quad (6)$$

We then utilize the measured decay widths $\Gamma(V \rightarrow \pi \pi) = (57 \pm 10) \text{ keV}$ and $\Gamma(\rho \rightarrow \pi \pi) = (5.5 \pm 0.9) \text{ keV}$ [9] and obtain the coupling constants g_V as $g_V = 1.42 \pm 0.12$ and $g_\rho = 0.42 \pm 0.03$. Our results are, therefore, in good agreement with the coupling constants deduced from the experimental values of the respective decay widths. Moreover, our result for g_V is also consistent with the value $g_V = 1.42 \pm 0.2$ calculated by T.M. Aliev et al. [7] in the framework of light cone sum rules, thus our study employing traditional QCD sum rule method supplements the previous light cone QCD sum rules calculation.

ACKNOWLEDGMENT

We like to thank Profs. A. Gokalp and O. Yılmaz for suggesting this investigation to us and for helpful discussions during the course of our work.

REFERENCES

- [1] M .A .Shifm an, A .I.Vainstein, V .I.Zakharov, Nucl.Phys.B 147 (1979) 385 and 448.
- [2] L .J.Reinders, S.Yazaki and H .R .Rubinstein, Nucl.Phys.B 196 (1985) 125.
- [3] B .L .Io e, Nucl.Phys.B 188 (1981) 317; B 191 (1981) 591.
- [4] A .G okalp, O .Y ilmaz, Eur.Phys.J.C 24, (2002) 117
- [5] P .J.O 'Donnell, Rev.M od.Phys. 53 (1981) 673
- [6] P .Colangelo, A .Khodjam irian, in Boris Io e Festschrift, At the Frontier of Particle Physics / Handbook of the QCD , edited by M .Shifm an W orld Scienti c, Singapor.
- [7] [6] T M A liev, I K ank, A .O zpineci, hep-ph/0212187
- [8] A .V .Sm ilga, Sov.J.Nucl.Phys. 35 (1982) 271.
- [9] A .I.T itov, T .-S.H .Lee, H .Toki, O .Streltsova, Phys.Rev.C 60 (1999) 035205.
- [10] D .E.G room et al, Eur.Phys.J.C 15 (2000) 1;
- [11] S-L.Zhu, W -Y .P.Hwang, Z-S.Yang, Phys.Lett.B 420 (1998) 8.
- [12] V .L.E letsky, B .L .Io e, Ya.I.Kogan, Phys.Lett.B 122 (1983) 423.

FIGURES




FIG .1. Quark loop diagram for V vertex




FIG .2. Operators of dimension 3 corrections proportional to $\langle \bar{q}q \rangle$.




FIG .3. Operators of dimension 5 corrections proportional to $\langle \bar{q} G q \rangle$. The dot lines denote gluons.




FIG . 4. The coupling constant $g_{\rho\eta\gamma}$ as a function of the Borel parameter M^2 for different values of M'^2 .




FIG .5. The coupling constant $g_{\omega\eta\gamma}$ as a function of the Borel parameter M^2 for different values of M'^2 .