Non-renorm alization Theorem

Originating in a New Fixed Point of the Vector M an ifestation

Chihiro Sasaki^a

^aD epartm ent of P hysics, N agoya U niversity, N agoya, 464-8602, Japan.

We study the pion velocity at the critical temperature of chiral symmetry restoration in QCD. Starting from the premise that the bare elective eld theory is to be de ned from the underlying QCD, we incorporate the elects of Lorentz non-invariance into the bare theory by matching an elective eld theory to QCD at a suitable matching scale, and investigate how the Lorentz non-invariance existing in the bare theory in uences physical quantities. Using the hidden local symmetry model as the eld theory, where the chiral symmetry restoration is realized as the vector manifestation (VM), we not that the pion velocity at the critical temperature receives neither quantum nor (therm all) hadronic corrections at the critical temperature even when we start from the bare theory with Lorentz symmetry breaking. This is likely the manifestation of a new xed point in the Lorentz non-invariant form ulation of the VM.

1. Introduction

Chiral symmetry in QCD is expected to be restored under some extreme conditions such as large number of avor N_f and high temperature and/or density. In hadronic sector, the chiral symmetry restoration is described by various eld theories (EFTs) based on the chiral symmetry [1].

By using the hidden local symmetry (HLS) model [2] as an EFT and performing the W ilsonian matching which is one of them ethods that determine the bare theory from the underlying QCD [3], the vector manifestation (VM) in hot or dense matter was formulated in Refs. [4,5]. In the VM, the massless vector meson becomes the chiral partner of pion at the critical point [6] $^{#1}$. There, the intrinsic temperature or density dependences of the parameters of the HLS Lagrangian, which are obtained by integrating out the high energy modes (i.e., the quarks and gluons above the matching scale) in hot and/or dense matter, play the essential roles to realize the chiral symmetry restoration consistently.

In the analysis done in Ref. [7], it was shown that the e ect of Lorentz symmetry breaking to the bare parameters caused by the intrinsic temperature dependence through the W ilsonian matching are small [7,8]. Starting from the bare Lagrangian with Lorentz invariance, it was presented that the pion velocity approaches the speed of light at the critical temperature [7], although in low temperature region (T T_c) the pion velocity deviates from the speed of light due to hadronic corrections [8].

However there do exist the Lorentz non-invariant e ects in bare EFT anyway due to the intrinsic tem peraure and/or density e ects. Further the Lorentz non-invariance m ight be enhanced through the renorm alization group equations (RGEs), even if e ects of Lorentz sym m etry breaking at the bare level are sm all. Thus it is im portant to investigate how the Lorentz non-invariance at bare level in uences physical quantities.

In this paper, we pick up the pion velocity at the critical temperature and study the quantum and hadronic therm all ects based on the VM . The pion velocity is one of the important quantities since it controls the pion propagation in medium through the dispersion relation.

O urm ain result is that the pion velocity does not receive either quantum or hadronic corrections in the lim it T ! T_c :

$$v (T) = V_{\text{;bare}}(T);$$
 (1.1)

independently of the value of the bare pion velocity V_{jbare}. This non-renorm alization property on the pion velocity is protected by the VM. Equation (1.1) in plies that the Lorentz non-invariance at bare level is directly rejected on the physical pion velocity at the critical tem perature.

This paper is organized as follows: In section 2, we show the HLS Lagrangian with Lorentz noninvariance and the conditions satis ed at the critical tem perature for the bare parameters. In section 3, we show that the conditions for the bare parametes at the critical tem perature are satis ed in any energy scale and that this is protected as a xed point of the relevent RGEs. In section 4, we show the quantum and hadronic corrections to the pion velocity and derive our result (1.1). In section 5, we give a sum mary and discussions.

 $^{^{\}sharp \ 1}$ As studied in Ref. [12] in detail, the VM is de ned only as a lim it with bare parameters approaching the VM $\,$ xed point from the broken phase.

2. Hidden Local Sym m etry

Since Lorentz symmetry breaking e ects are included in the bare theory through the W ilsonian m atching, the HLS Lagrangian in hot and/or dense m atter is generically Lorentz non-invariant. Its explicit form was presented in Ref. [5]. In this section, we start from this Lagrangian with Lorentz non-invariance, and requiring that the axial-vector current correlator be equal to the vector current correlator at the critical point, we obtain the conditions for the bare param eters.

2.1. Lorentz N on-invariant H LS Lagrangian

In this subsection, we show the HLS Lagrangian at leading order including the e ects of Lorentz noninvariance.

The HLS model is based on the G_{global} H_{local} sym metry, where G = SU (N_f)_L SU (N_f)_R is the chiral sym metry and H = SU (N_f)_V is the HLS. The basic quantities are the HLS gauge boson V and two matrix valued variables $_{L}(x)$ and $_{R}(x)$ which transform as $_{L,R}(x)$! $_{L,R}^{0}(x) = h(x) _{L,R}(x)g_{L,R}^{y}$, where h(x) 2 H local and $g_{L,R}$ 2 [SU (N_f)_{L,R} lglobal. These variables are parameterized as ^{# 2}

$$L_{R}(x) = e^{i (x) = F'} e^{i (x) = F'};$$
 (2.1)

where = ${}^{a}T_{a}$ denotes the pseudoscalar N am bu-G oldstone bosons associated with the spontaneous sym m etry breaking of G_{global} chiral sym m etry, and = ${}^{a}T_{a}$ denotes the N am bu-G oldstone bosons associated with the spontaneous breaking of H_{local}. This is absorbed into the HLS gauge boson through the Higgs m echanism, and then the vector m eson acquires its m ass. F^t and F^t denote the tem poral components of the decay constant of and , respectively. The covariant derivative of L is given by

$$D_{L} = Q_{L} \quad \text{iV}_{L} + \text{i}_{L}L; \qquad (2.2)$$

and the covariant derivative of $_{\rm R}$ is obtained by the replacement of L with R in the above where V is the gauge eld of H $_{\rm local}$, and L and R are the external gauge elds introduced by gauging G $_{\rm global}$ symmetry. In terms of L and R , we denote the external axial-vector and vector elds as A = (R L)=2 and V = (R + L)=2.

In the HLS model it is possible to perform the derivative expansion system atically [10,11,12]. In the chiral perturbation theory (ChPT) with HLS, the vector m eson m ass is to be considered as sm all

compared with the chiral symmetry breaking scale

, by assigning O (p) to the HLS gauge coupling, g O (p) 10,11]. (For details of the ChPT with HLS, see Ref. [12].) The leading order Lagrangian with Lorentz non-invariance can be written as [5]

$$L = (F^{t})^{2}uu + F^{t}F^{s}(g uu)$$

$$tr[^{2}_{?} ^{2}_{?}]$$

$$+ (F^{t})^{2}uu + F^{t}F^{s}(g uu)$$

$$h i$$

$$tr^{*}_{k}^{k}$$

$$= \frac{1}{g_{L}^{2}}uug \frac{1}{2g_{T}^{2}}(g g 2uug)$$

$$tr V V ; (2.3)$$

where

$$^{*}_{?,k} = \frac{1}{2i} D_{R} ^{Y}_{R} D_{L} ^{Y}_{L}$$
 (2.4)

Here F^{s} denote the spatial pion decay constant and similarly F^{s} for the . The rest fram e of the medium is specified by u = (1;0) and V is the eld strength of V. g_{L} and g_{T} correspond in medium to the HLS gauge coupling g. The parametric and velocities are defined by [13]

$$V^{2} = F^{s} = F^{t}; \quad V^{2} = F^{s} = F^{t}:$$
 (2.5)

2.2. Conditions for the Bare Param eters

In this subsection follow ing R ef. [5] where the conditions for the current correlators with the bare param eters in dense m atter were presented, we show the Lorentz non-invariant version of the conditions satis ed at the critical tem perature for the bare param eters.

Concept of the matching in the W ilsonian sense is based on the following assumptions: The bare Lagrangian of the e ective eld theory (EFT) L bare is de ned at a suitable matching scale . Generating functional derived from L_{bare} leads to the same G reen's function as that derived from the generating functionalofQCD at . In otherwords, the bare param eters are obtained after integrating out the high energy modes, i.e., the quarks and gluons above . W hen we integrate out the high energy m odes in hot m atter, the bare param eters have a certain tem perature dependence, intrinsic tem perature dependence, converted from QCD to the EFT. The intrinsic tem perature dependence is nothing but the signature that hadrons have an internal structure constructed from quarks and gluons. In the following, we describe the chiral sym metry restoration based on the point of view that the bare HLS theory is de ned

^{# 2}The wave function renorm alization constant of the pion eld is given by the temporal component of the pion decay constant [9]. Thus we norm alize and by F^t and F^t respectively.

from the underlying QCD.W e note that the Lorentz non-invariance appears in the bare HLS theory as a result of including the intrinsic temperature dependence. Once the temperature dependence of the bare parameters is determ ined through the matching with QCD mentioned above, the parameters appearing in the hadronic corrections pick up the intrinsic them all ects through the RGEs.

The axial-vector and vector current correlators at bare level are constructed in terms of bare parameters and are divided into the longitudinal and transverse components:

$$G_{A;V} = P_{L} G_{A;V}^{L} + P_{T} G_{A;V}^{T}; \qquad (2.6)$$

where $P_{L,T}$ are the longitudinal and transverse projection operators, respectively. The axial-vector current correlator in the HLS around them atching scale

is well described by the following form s w ith the bare parameters [5,7]:

$$G_{A (H LS)}^{L}(p_{0};p) = \frac{p^{2}F_{,bare}^{t}F_{,bare}^{s}F_{,bare}^{s}}{[p_{0}^{2} V_{,bare}^{2}p^{2}]} 2p^{2}z_{2,bare}^{L};$$

$$G_{A (H LS)}^{T}(p_{0};p) = F_{,bare}^{t}F_{,bare}^{s}F_{,bare}^{s}$$

$$2 p_{0}^{2}z_{2,bare}^{L}p^{2}z_{2,bare}^{T}; (2.7)$$

where $z_{2,\text{pare}}^{L}$ and $z_{2,\text{pare}}^{T}$ are the coe cients of the higher order term s, and V _{jbare} is the bare pion velocity related to F^t_{jbare} and F^s_{jbare} as

$$V_{\text{;bare}}^{2} = \frac{F_{\text{;bare}}^{s}}{F_{\text{;bare}}^{t}} : \qquad (2.8)$$

Sim ilarly, two components of the vector current correlator have the following form s:

$$\begin{array}{l} G_{V \ (H \ LS)}^{\text{L}} \left(p_{0}; p \right) \\ = & \frac{p^{2} F_{;bare}^{\text{L}} F_{;bare}^{\text{s}} 1 \quad 2q_{;bare}^{2} z_{3;bare}^{\text{L}}}{h} \\ & 1 \\ \hline p_{0}^{2} \quad V_{;bare}^{2} p^{2} \quad M_{;bare}^{2} \\ & 2p_{2}^{\text{L}} z_{1;bare}^{\text{L}} + O \left(p^{4} \right); \\ G_{V \ (H \ LS)}^{\text{T}} \left(p_{0}; p \right) = & \frac{h}{p_{0}^{2}} \frac{F_{;bare}^{\text{T}} F_{;bare}^{\text{s}}}{V_{T;bare}^{2} p^{2} \quad M_{;bare}^{2}} \\ p_{0}^{2} \quad 1 \quad 2q_{;bare}^{2} z_{3;bare}^{\text{L}} \\ & V_{T;bare}^{2} p^{2} \quad 1 \quad 2q_{;bare}^{2} z_{3;bare}^{\text{T}} \\ & 2 \quad p_{0}^{2} z_{1;bare}^{\text{L}} \quad p^{2} z_{1;bare}^{\text{T}} + O \left(p^{4} \right); \end{array}$$

where $z_{1;2;\text{bare}}^{\text{L}}$ and $z_{1;2;\text{bare}}^{\text{T}}$ denote the coe cients of the higher order term s. In the above expressions, the bare vector m eson m ass in the rest fram e, M _{ibare}, is

$$M^{2}_{,bare} q^{2}_{,bare}F^{t}_{,bare}F^{s}_{,bare}$$
: (2.10)

W e de ne the bare param eters a_{bare}^{t} and a_{bare}^{s} as

$$a_{\text{bare}}^{t} = \frac{F_{;\text{bare}}^{t}}{F_{;\text{bare}}^{t}}; \quad a_{\text{bare}}^{s} = \frac{F_{;\text{bare}}^{s}}{F_{;\text{bare}}^{s}}; (2.11)$$

and the bare and transverse velocities as

$$V_{,bare}^{2} = \frac{F_{,bare}^{s}}{F_{,bare}^{t}}; V_{T,bare}^{2} = \frac{g_{L,bare}^{2}}{g_{T,bare}^{2}}:$$
 (2.12)

Now we consider the matching near the critical tem perature. At the chiral phase transition point, the axial-vector and vector current correlators must agree with each other: $G_{A\ (H\ LS)}^{\ L} = G_{V\ (H\ LS)}^{\ L}$ and $G_{A\ (H\ LS)}^{\ T} = G_{V\ (H\ LS)}^{\ T}$. These equalities are satis ed for any values of p_0 and p around the matching scale only if the following conditions are met:

This implies that at bare level the longitudinalm ode of the vector m eson becomes the real NG boson and couples to the vector current correlator, while the transverse m ode decouples.

3. Vector M an ifestation C ondition

In this section, we show that the conditions for the bare parameters for T ! T_c are satis ed in any energy scale and that this is protected by the xed point of the RGEs.

It was shown that the HLS gauge coupling g=0 is a xed point of the RGE forg at one-loop level [14,3]. The existence of the xed point g=0 is guaranteed by the gauge invariance. This is easily understood from the fact that the gauge eld is norm alized as V=g. In the present case without Lorentz symmetry, the gauge eld is norm alized by $g_{\rm L}$ as $V=g_{\rm L}$ and thus $g_{\rm L}=0$ becomes a xed point of the RGE for $g_{\rm L}$.

P rovided that $g_L = 0$ is a xed point, we can show that $a^t = a^s = 1$ is also a xed point of the coupled RGEs for a^t and a^s as follows: W e start from the bare theory de ned at a scale with $a_{bare}^t = a_{bare}^s = 1$ (and $g_L = 0$). The parameters a^t and a^s at () are calculated by integrating out the m odes in [;]. They are obtained from the two-point functions of A and V, denoted by ? and k. W e decompose these functions into

$$P_{2,k} = u u \frac{L}{2,k} + (g u u) \frac{s}{2,k} + P_{L} \frac{L}{2,k} + P_{T} \frac{T}{2,k};$$
 (3.1)

where u u , (g u u), P_L and P_T denote the tem poral, spatial, longitudinal and transverse projection operators, respectively. The parameters a^t



Figure 1. D lagram s for contributions to $_{?}$ at one-loop level.



Figure 2. D iagram s for contributions to $_{k}$ at one-loop level.

and a^s are defined by $a^t = \frac{t}{k} = \frac{t}{2}$; $a^s = \frac{s}{k} = \frac{s}{2}$ [8]. These expressions are further reduced to

where $\frac{t_{i}s}{2}$ (;) denotes the quantum correction obtained by integrating the modes out between [;]. We show the diagrams for contributions to $\frac{1}{2}$ and $\frac{1}{k}$ at one-loop level in Figs. 1 and 2.

The contributions (a) in Fig. 1 and (a) in Fig. 2 are proportional to $g_{L,bare}^2$. The contributions (c) in Fig.1 and (d) in Fig.2 are proportional to $(a_{bare}^t 1)$. Taking $g_{L,bare} = 0$ and $a_{bare}^t = a_{bare}^s = 1$, these contributions vanish. We note that (i.e., longitudinal vector m eson) is massless and the chiral partner of pion at the critical temperature. Then the contributions (b) and (c) in Fig. 2 have a symm etry factor 1=2 respectively and are obviously equal to the contribution (b) in Fig.1, i.e., $\binom{(b)}{2} = \binom{(b)+(c)}{k}$. Thus from Eq. (3.2), we obtain

$$a^{t}() = a^{t}_{bare} = 1;$$

 $a^{s}() = a^{s}_{bare} = 1:$ (3.3)

This implies that a^t and a^s are not renormalized at the scale (). Similarly, we include the corrections below the scale () in turn, and nd that a^t and a^s do not receive the quantum corrections. Eventually we conclude that $a^t = a^s = 1$ is a xed point of the RGEs for a^t and a^s .

From the above, we nd that $(g_L; a^t; a^s) = (0; 1; 1)$ is a xed point of the combined RGEs for $g_L; a^t$ and a^s. Thus the VM condition is given by

$$g_{L} ! 0;$$

 $a^{t} ! 1; a^{s} ! 1 \text{ for } T ! T_{c} : (3.4)$

The vector m eson m ass is never generated at the critical temperature since the quantum correction to M² is proportional to g_L^2 . Because of g_L ! 0, the transverse vector m eson at the critical point, in any energy scale, decouples from the vector current correlator. The VM condition for a^t and a^s leads to the equality between the and (i.e., longitudinal vector m eson) velocities:

$$V = V^{4} = F^{s}F^{t} = F^{s}F^{t^{2}}$$

= $a^{t} = a^{s^{T}}!^{T_{c}} 1:$ (3.5)

This is easily understood from a point of view of the VM since the longitudinal vectors eson become s the chiral partner of pion. We note that this condition V = V holds independently of the value of the bare pion velocity which is to be determ ined through the W ilsonian matching.

4. P ion Velocity at C ritical Tem perature

As we mentioned in Introduction, the intrinsic temperature dependence generates the e ect of Lorentz symmetry breaking at bare level. Then how does the Lorentz non-invariance at bare level in uence physical quantities? In order to make it clear, in this section we study the pion decay constants and the pion velocity near the critical temperature.

Following Ref. [7], we de ne the on-shell of the pion from the pole of the longitudinal component G_A^L of the axial-vector current correlator. This pole structure is expressed by tem poral and spatial com – ponents of the two-point function $_2$. The tem poral and spatial pion decay constants are expressed as follows [7,8]:

$$f^{t}(p;T)^{2} = \frac{t}{2} (V p;p;T);$$

$$f^{t}(p;T)f^{s}(p;T) = \frac{s}{2} (V p;p;T); \qquad (4.1)$$

where the on-shell condition p_0 ! V p was taken. W edivide the two-point function $_2$ into two parts, zero tem perature (vacuum) and non-zero tem perature parts, as $_2 = \frac{(vac)}{2} + \frac{1}{2}$. The quantum correction is included in the vacuum part $\frac{(vac)}{2}$, and the hadronic therm al correction is in $_2$. In the present perturbative analysis, we obtain the pion velocity as [8]

$$v^{2}(p;T) = \frac{f^{s}(p;T)}{f^{t}(p;T)}$$

= $V^{2} + e_{2}(V p;p)$

+
$$\frac{{}^{s}_{?} (V p; p; T) V^{2} {}^{t}_{?} (V p; p; T)}{{}^{F^{t}^{2}}};$$
 (4.2)

where $e_{?}$ (V p;p) denotes a possible nite renorm alization e ect. Note that the renorm alization condition on V is determined as $e_{?}$ (V p;p) $\mathbf{j}_{P=0} = 0$.

In the following, we study the quantum and hadronic corrections to the pion velocity for T ! T_c , on the assumption of the VM conditions (3.4). As we de ned above, the two-point function associated with the pion velocity v (p;T) is $_2$ (p₀;p;T). The diagram s contributing to $_2$ are shown in Fig. 1. As mentioned in section 3, diagram (a) is proportional to g_L and diagram (c) has the factor (a^t 1). Then these contributions vanish at the critical point. We consider the contribution from diagram (b) only.

First we evaluate the quantum correction to the vacuum part $2^{(vac)(b)}$. This is expressed as

$$\frac{(\text{vac})(b)}{2} \quad (p_0;p) = N_f \quad \frac{Z}{i(2)^n}$$

$$\frac{(k;p) \quad (k;p)}{[k_0^2 + V^2 k^2] M^2} \quad (k_0 \quad p_0)^2 + V^2 \tilde{j} \quad p_1^2]$$

$$(4.3)$$

where denotes the A vertex as

$$(k;p) = \frac{i^{p}}{2} \overline{a^{t}} g [u u + V^{2} (g u u)](2k p):$$
 (4.4)

W e note that the spatial component of this vertex ⁱ has an extra-factor V² as compared with the tem poralone. In the present analysis it is in portant to include the quadratic divergences to obtain the RGEs in the W ilsonian sense. In Refs. [15,3,12], the dim ensional regularization was adopted and the quadratic divergences were identied with the presence of poles of ultraviolet origin at n = 2 [16]. In this paper when we evaluate four dimensional integral, we rst integrate over k_0 from 1 to 1 . Then we carry out the integral over three-dim ensional m om entum κ with three-dim ensional cuto $_3$. In order to be consistent with ordinary regularization in four dimension [15,3,12], we use the following replacement associated with quadratic divergence:

$${}^{3} ! \frac{p}{2} \frac{1}{4} = \frac{1}{p} \frac{1}{2} ; \qquad (4.5)$$

$${}^{2} \frac{d^{n-1}k}{(2)^{n-1}} \frac{1}{k} ! \frac{2}{8 2} ;$$

$${}^{2} \frac{d^{n-1}k}{(2)^{n-1}} \frac{k^{i}k^{j}}{k^{3}} ! \frac{ij}{8 2} ; \qquad (4.6)$$

W hen we make these replacement, the present method of integral preserves chiral symmetry.

As shown in Appendix A, $\frac{(vac)^{t}}{2}$ and $\frac{(vac)^{s}}{2}$ are independent of the external momentum. A coord-ingly, the nite renum alization e ect e_{2} is also independent of the external momentum and then van-ishes:

$$e_{2}$$
 (V p;p) = 0: (4.7)

Thus in the following, we take the external momentum as zero. In that case, the tem poral and spatial components of $\binom{(vac)}{2}$ are expressed as $\binom{(vac)t}{2} = \binom{(vac)00}{2}$ and $\binom{(vac)s}{2} = \binom{(ij=3)}{2}$. Taking the VM limit (M ! 0 and V ! V), these components become

$$\begin{split} \lim_{V M} & \frac{(vac)00}{2} (p_0 = p = 0) \\ &= \frac{N_f}{4} \frac{Z}{(2 - 1)^n} \frac{dk_0 d^{n-1} k}{[k_0^2 + V^2 k^2]^2} \\ &= \frac{N_f}{4} \frac{Z}{(2 - 1)^{n-1}} \frac{d^{n-1} k}{V k} \frac{1}{V k} \\ &= \frac{N_f}{4} \frac{1}{V} \frac{2}{8 - 2}; \\ \lim_{V M} & \frac{(vac)ij}{2} (p_0 = p = 0) \\ &= \frac{N_f}{4} (V^2)^2 \frac{Z}{(2 - 1)^n} \frac{dk_0 d^{n-1} k}{(2 - 1)^n} \frac{4k^i k^j}{[k_0^2 + V^2 k^2]^2} \\ &= \frac{N_f}{4} V^4 \frac{Z}{(2 - 1)^{n-1}} \frac{d^{n-1} k}{(V k)^3} \frac{k^i k^j}{(V k)^3} \\ &= \frac{N_f}{4} V \frac{ij}{8 - 2}; \end{split}$$
(4.8)

Thus we obtain the tem poral and spatial parts as

$$\lim_{VM} \begin{array}{c} \lim_{2 \to 0} (vac)t (p_0 = p = 0) = \frac{N_f}{4} \frac{1}{V} \frac{1}{8} \frac{2}{2};$$

$$\lim_{VM} \begin{array}{c} (vac)s \\ 2 \end{array} (p_0 = p = 0) = \frac{N_f}{4} V \frac{2}{8^2}; \quad (4.9)$$

These quadratic divergences are renormalized by $(F^{t}_{;bare})^{2}$ and $F^{t}_{;bare}F^{s}_{;bare}$, respectively. Then RGEs for the parameters $(F^{t})^{2}$ and $F^{t}F^{s}$ are expressed as

$$\frac{d F^{t^{2}}}{d} = \frac{N_{f}}{(4)^{2}} \frac{1}{V}^{2}; \qquad (4.10)$$

$$\frac{d F^{t}F^{s}}{d} = \frac{N_{f}}{(4)^{2}} V^{2} : \qquad (4.11)$$

Both $(F^{t})^{2}$ and $F^{t}F^{s}$ scale following the quadratic running ². However, the coe cient of ² in the RGE for $(F^{t})^{2}$ is dimension that for $F^{t}F^{s}$.

W hen we use these RGEs, the scale dependence of the parametric pion velocity is

$$\frac{dV^2}{d} = \frac{d F^{t}F^{s} = (F^{t})^2}{d}$$

$$= \frac{1}{F^{t}} \frac{N_{f}}{(4)^{2}} V F^{t} F^{t} F^{s} \frac{1}{V} I^{2}$$

$$= 0: \qquad (4.12)$$

This in plies that the param etric pion velocity at the critical tem perature does not scale. In other words, the Lorentz non-invariance at bare level is not enhanced through the RGEs as long as we consider the pion velocity. As we noted below Eq. (4.4), the factor V² is in the spatial component of the vertex . If V were not equal to V, the coe cients of running in the right-hand-side of Eqs. (4.10) and (4.11) would change. However, since the VM conditions do guarantee V = V, the quadratic running caused from ² in (F^t)² and F^tF^s are exactly canceled in the second line of Eq. (4.12).

Next we study the hadronic therm alcorrection to the pion velocity at the critical tem perature. The tem poral and spatial parts of the hadronic therm al correction $_{?}$ contribute to the pion velocity, which have the same structure as those of the quantum correction $_{?}^{(vac)}$, except for a B ose E instein distribution function. Thus by the replacement of $^{2}=(4)^{2}$ with T²=12 in $_{?}^{(vac)t;s}$, hadronic corrections to the tem poral and spatial parts of $_{?}$ are obtained as follow s:

$$\lim_{V M} \quad \stackrel{t}{?} (p_0; p; T) = \quad \frac{N_f}{24} \frac{1}{V} T_c^2;$$

$$\lim_{V M} \quad \stackrel{s}{?} (p_0; p; T) = \quad \frac{N_f}{24} V T_c^2: \quad (4.13)$$

Substituting Eq. (4.13) into Eq. (4.2) with $e_2 = 0$ as shown in Eq. (4.7), we obtain the physical pion velocity in the VM as

$$v^{2} (p;T) {}^{T} ! {}^{T_{c}} V^{2} + \frac{ {}^{s} (V p;p;T_{c}) V^{2} ! (V p;p;T_{c}) }{ F^{t} {}^{2}} = V^{2} : (4.14)$$

Since the parametric pion velocity in the VM does not scale with energy [$\sec Eq.(4.12)$], V in the above expression is equivalent to the bare pion velocity:

 $v (p;T) = V_{bare}(T) \text{ for } T_c:$ (4.15)

This implies that the pion velocity in the limit T $\,!\,$ $T_{\rm c}$ receives neither hadronic nor quantum corrections due to the protection by the VM . This is our main result.

In order to clarify the reason of this nonrenorm alization property, let us recall the fact that only the diagram (b) in Fig.1 contributes to the pion velocity at the critical tem perature. A way from the 6

critical tem perature, the contribution of the mas-(i.e., the longitudinalm ode of massive vector sive meson) is suppressed owing to the Boltzmann factor exp[M = T], and then only the pion loop contributes to the pion velocity. Then there exists the $O(T^4)$ correction to the pion velocity [8]. Near the critical tem perature, on the other hand, becom es massless due to the VM since (i.e., the longitudinalvectorm eson) becom es the chiralpartner of the pion. Thus the absence of the hadronic corrections in the pion velocity at the critical tem perature is due to the exact cancellation between the contribution of pion and that of its chiral partner . Sim ilarly the quantum correction generated from the pion loop is exactly cancelled by that from the loop.

5. Sum m ary and D iscussions

In this paper, we started from the Lorentz non-invariant HLS Lagrangian at bare level and studied the pion velocity at the critical tem perature based on the VM . From the analysis of the quantum and hadronic therm al corrections to the pion velocity, we obtained the result that the pion velocity in the lim it T ! $T_{\rm c}$ is equal to the bare pion velocity. In other words, the pion velocity does not receive either quantum or hadronic corrections at the critical tem perature. This occurs due to the exact cancellation between the contribution of pion and that of the longitudinal vector m eson (i.e., the chiral partner of pion).

Now we consider the meaning of our result (4.15). Based on the point of view that the bare HLS theory is de ned from QCD, we presented the VM conditions realizing the chiral symmetry in QCD consistently, i.e., $(q_L; a^t; a^s)$! (0; 1; 1) for T ! T_c . This is the xed point of the RGEs for the parameters q_{I} ; a^{t} and a^{s} . As we showed in section 4, although both pion decay constants $(F^t)^2$ and F^tF^s scale following the quadratic running, $(F^{t})^{2}$ and $F^{t}F^{s}$ show a di erent running since the coe cient of 2 in Eq. (4.10) is di erent from that in Eq. (4.11). Nevertheless in the pion velocity at the critical tem perature, the quadratic running in (F^t)² is exactly cancelled by that in $F^{t}F^{s}$ [see second line of Eq. (4.12)]. There it was crucial for intricate cancellation of the quadratic running that the velocity of (i.e., longitudinal vector m eson) is equal to its chiral partner, ie., V ! V for T ! T_c. Note that this is not an extra condition but a consequence from the VM conditions for a^t and a^s; we started simply from the VM conditions alone and found that V does not receive quantum corrections at the restoration point. As we showed in Eq. (4.13), the hadronic correction to $(F^{t})^{2}$ is different from that to $F^{t}F^{s}$.

_+ _-

In the pion velocity, however, the hadronic correction from $(F^{t})^{2}$ is exactly cancelled by that from F^tF^s [see second line of Eq. (4.14)]. The VM conditions guarantee these exact cancellations of the quantum and hadronic corrections. This implies that $(g_L; a^t; a^s; V) = (0; 1; 1; any)$ form sa xed line for four RGEs of q_L ; a^t ; a^s and V . W hen one point on this xed line is selected through the matching procedure as done in Ref. [17], namely the value of V ;bare is xed, the present result in plies that the point does not move in a subspace of the parameters. This is likely the manifestation of a new xed point in the Lorentz non-invariant form ulation of the ${\tt VM}$. Approaching the restoration point of chiral sym metry, the physical pion velocity itself would ow into the xed point.

Several com m ents are in order:

We should distinguish the consequences within HLS/VM from those beyond HLS/VM .C learly the determ ination of the de nite value of the bare pion velocity is done outside HLS/VM . On the other hand, our main result (4.15) holds independently of the value of the bare pion velocity itself. Applying this result to the case where one starts from the bare HLS theory with Lorentz invariance, i.e., V $_{\rm ;bare}$ = 1, one nds that the pion velocity at T_c becomes the speed of light since v = V $_{\rm ;bare}$ = 1, as obtained in Ref. [7].

As a consequence of the relation (4.15), we can determ ine the tem poral and spatial pion decay constants at the critical tem perature when we take the bare pion velocity as nite. In the following, we study these decay constants and discuss their determ inations based on Eq. (4.15). Using Eq. (4.12), we solve the RGEs (4.10) and (4.11) and obtain a relation between two parametric pion decay constants as $F^{t}(0;T_{c})F^{s}(0;T_{c}) = V^{2}F^{t}(0;T)^{2}$. From this and (4.13), the tem poraland spatial pion decay constants with the quantum and hadronic corrections are obtained as

$$f^{t^{2}} = F^{t}(0;T_{c})^{2} \frac{N_{f}}{24} \frac{1}{V} T_{c}^{2};$$

$$f^{t}f^{s} = F^{t}(0;T_{c})F^{s}(0;T_{c}) \frac{N_{f}}{24}V T_{c}^{2}$$

$$= V^{2} f^{t^{2}}:$$
(5.1)

Since the order parameter $(f^t f^s)$ vanishes as expected at the critical temperature, we nd that $f^t f^s = V^2 f^{t^2} = 0$. Multiplying both side by $v^2 = V^2$, the above expression is reduced to

$$(5.2)^2 = V^4 (f^t)^2 = 0$$
:

Now, the spatial pion decay constant vanishes at the critical tem perature, $f^{\rm s}\left(T_{\rm c}\right)$ = 0 . In the case

of a vanishing pion velocity, f^t can be nite at the restoration point. On the other hand, when V is nite, Eq. (5.2) leads to $f^t(T_c) = 0$. Thus we not that both tem poral and spatial pion decay constants vanish simultaneously at the critical tem perature when the bare pion velocity is determined as nite.

In order to know the value of the (bare) pion velocity, we need to specify a method that determ ines the bare parameters of the e ective eld theory. As we stressed in subsection 22, the bare param eters of the HLS Lagrangian are determined by the underlying QCD. One possible way to determ ine them is the W ilsonian matching proposed in Ref. [3] which is done by m atching the axial-vector and vector current correlators derived from the HLS with those by the operator product expansion (OPE) in QCD at the matching scale . From the analysis perform ed on the basis of a W ilsonian m atching, the bare pion velocity at the critical tem perature is found to be nite, i.e., V ;bare 6 0 [17]. Thus, by combining Eq. (4.15) with estim ation of V ; bare, the value of the physicalpion velocity v (T) at the critical tem perature is obtained to be nite [17]. This is in contrast to the result obtaied from the chiral theory [18,19], where the relevent degree of freedom near T_c is only the pion. Their result is that the pion velocity becom es zero for T ! Tc. Therefore from the experim ental data, we may be able to distinguish which picture is correct, v 1 or v ! 0.

A cknow ledgm ent

The author would like to thank Professor M asayasu H arada, D octor Y oungm an K im and Professor M annque R ho form any useful discussions and com m ents.

Appendix

A. Two-point Function

In this appendix, we show that the temporal and spatial components of the two-point function (vac) are independent of the automalm on onter

 $\frac{(vac)}{2}$ are independent of the external m om entum in the VM limit.

From Eq. (4.3), $\frac{(vac)(b)}{2}$ is rew ritten as

$$\sum_{p}^{(\text{vac})} (p_{0}; p) = N_{f} \frac{a^{t}}{4} X X B^{(\text{vac})} (p_{0}; p; M; 0)$$

$$N_{f} \frac{a^{t}}{4} B^{(\text{vac})} (p_{0}; p; M; 0); \quad (A.1)$$

where we de ne

$$X = u u + V^{2} (g u u)$$
: (A.2)

$$+ \frac{2!^{p}}{2!^{p}} \frac{(!^{p} + p_{0})^{2}}{(!^{p} + p_{0})^{2}}$$

$$= \frac{d^{3}k}{(2)^{3}} \frac{(2k p ^{2}p^{2})^{2}}{p^{2}}$$

$$\left(\frac{1}{2!} \frac{1}{(! p_{0})^{2}} \frac{(!^{p})^{2}}{(!^{p} + p_{0})^{2}}\right)$$

$$+ \frac{1}{2!^{p}} \frac{1}{(!^{p} + p_{0})^{2}} \frac{1}{!^{2}}$$

$$+ \frac{p_{0}p k^{2} p}{p^{2}}$$

$$\left(\frac{1}{2!} \frac{2! p_{0}}{(! p_{0})^{2}} \frac{p_{0}}{(!^{p})^{2}}\right)$$

$$\frac{d^{3}k}{(2)^{3}} = \frac{1}{2!} \frac{(2! p)^{2}}{(! p)^{2} (!^{p})^{2}} + \frac{1}{2!^{p}} \frac{(2!^{p} + p_{0})^{2}}{(!^{p} + p_{0})^{2} !^{2}} + \frac{1}{2!^{p}} \frac{(2!^{p} + p_{0})^{2}}{(!^{p} + p_{0})^{2} !^{2}} + \frac{1}{2!} \frac{2! p}{(! p)^{2} (!^{p})^{2}} + \frac{1}{2!^{p}} \frac{2!^{p} + p_{0}}{(!^{p} + p_{0})^{2} !^{2}} ;$$

By using the expressions in Ref. [8], the componets $B^{(vac)t}$; $B^{(vac)s}$ and $B^{(vac)L}$ take the following form s:

B ^{(vac)t} (p₀;p;M ;0)

Ζ

$$\begin{split} B^{(vac)t}(p_{0};p;M;0) &= \overset{h}{B}^{(vac)t}(p_{0};p;M;0) \\ &+ (1 \quad V^{2})\frac{p^{2}}{p^{2}}B^{(vac)L}(p_{0};p;M;0); \\ B^{(vac)s}(p_{0};p;M;0) &= V^{4} \overset{h}{B}^{(vac)s}(p_{0};p;M;0) \\ &+ \frac{1 \quad V^{2}}{V^{2}}\frac{p_{0}^{2}}{p^{2}}B^{(vac)L}(p_{0};p;M;0) \overset{i}{:} (A.5) \end{split}$$

In term s of each components of B $^{\rm (vac)}$, the tem – poral and spatial parts of B are given by

$$!^{2} = V^{2}k^{2};$$

 $(!^{p})^{2} = V^{2}\tilde{K} p^{2} + M^{2}:$ (A.4)

with

$$B^{(vac)} (p_0; p; M; 0) = \begin{bmatrix} Z & \frac{dk_0}{i(2)} \\ \frac{d^3k}{i(2)} \end{bmatrix} \begin{bmatrix} \frac{d^3k}{i(2)^3} \\ \frac{(2k p)(2k p)}{k_0^2 !^2][(k_0 p_0)^2 (!^p)^2]} \end{bmatrix}; \quad (A.3)$$

d³k

=

In the above expression, we de ne the function B ^(vac) contributed to the diagram (b) in Fig. 1 as

$$\lim_{VM} B^{(vac)t}(p_{0};p;M;0) = \frac{d^{3}k}{(2)^{3}} \frac{1}{!}$$

$$\frac{I(k;p_{0};p) + V^{2}p}{[(! p_{0})^{2} (!^{p})^{2}][(! + p_{0})^{2} (!^{p})^{2}]};$$

$$\lim_{VM} B^{(vac)s}(p_{0};p;M;0) = \frac{Z}{\frac{d^{3}k}{(2)^{3}} \frac{1}{!} \frac{1}{V^{2}p^{2}}}$$

U sing Eqs. (A .7)-(A .9) with these functions, we obtain B^{~ (vac)t;s} as Z₂

I
$$(k; p_0; p)$$

= !² [4!² 4 (!^p)² 3p₀²] + p₀² [p₀² (!^p)²];
J $(k; p_0; p) = p_0 [3!^2 + p_0^2 (!^p)^2];$
K $(k; p_0; p) = !^2 + p_0^2 (!^p)^2$: (A 10)

where we de ne the functions as

$$\begin{split} \lim_{VM} B^{(vac)t}(p_{0};p;M;0) &= \frac{d^{3}k}{(2)^{3}} \frac{1}{!} \\ \frac{I(k;p_{0};p) + p}{[(! p_{0})^{2}} \frac{p}{(!^{p})^{2}} \frac{p}{[(! + p_{0})^{2}} \frac{k;p_{0};p}{(!^{p})^{2}}]; \\ \frac{I}{[(! p_{0})^{2}} \frac{p}{(!^{p})^{2}} \frac{p}{[(! + p_{0})^{2}} \frac{p}{(!^{p})^{2}}]; \\ \frac{I}{h} \\ \frac{1}{h} \\ \frac{1}{h} \\ \frac{1}{p^{2}} \frac{p}{k^{2}} \frac{p}{p} \frac{p}{k^{2}} \frac{p}{p} \frac{p}{k} \frac{p}{k;p_{0};p}; \\ + p^{2}_{0}p \frac{p}{k^{2}} \frac{p}{p} \frac{p}{k;p_{0};p}; \\ \frac{p}{k^{2}} \frac{p}{p} \frac{p}{k} \frac{p}{k;p_{0};p}; \\ \frac{p}{[(! p_{0})^{2}} \frac{p}{(!^{p})^{2}}][(! + p_{0})^{2}} \frac{p^{2}}{(!^{p})^{2}}]; \\ \frac{p}{[(! p_{0})^{2}} \frac{p}{(!^{p})^{2}} \frac{p}{[(! + p_{0})^{2}} \frac{p^{2}}{(!^{p})^{2}}]; \\ \frac{p}{[(! p_{0})^{2}} \frac{p}{(!^{p})^{2}}][(! + p_{0})^{2}} \frac{p^{2}}{(!^{p})^{2}}; \\ \end{array}$$

Now we take the VM lim it. Note that the VM condition for a^t and a^s im plies that \quad velocity becomes equal to the pion velocity, V ~!~ V for T !~ T_c as shown in Eq. (3.5). The components B (vac)t; B (vac)s and B $^{(vac)L}$ are calculated as follows: Z d^3k

$$+ \frac{1}{2!^{p}} \frac{2!^{p} + p_{0}}{(!^{p} + p_{0})^{2} !^{2}} ;$$

$$B^{(vac)L}(p_{0};p;M;0)$$

$$= \frac{2}{(2^{n})^{3}} \frac{p^{2}p}{p_{0}p^{2}} \frac{12}{p_{0}p^{2}} \frac{p}{p_{0}p^{2}} \frac{1}{(! p_{0})^{2} (!^{p})^{2}} + \frac{1}{2!^{p}} \frac{2!^{p} + p_{0}}{(!^{p} + p_{0})^{2} !^{2}} ; \quad (A.6)$$

$$\frac{1}{\begin{bmatrix} (! & p_0)^2 & (!^p)^2 \end{bmatrix} \begin{bmatrix} (! + p_0)^2 & (!^p)^2 \end{bmatrix}} \\ V^2 p & \mathcal{R} & p \end{bmatrix}^2 K (k; p_0; p) \\ + p_0^2 V^2 p & \mathcal{R} & p) J k; p_0; p \end{bmatrix} \overset{i}{:} (A.12)$$

The integrand of the functions $B^{(vac)t}$ and $B^{(vac)s}$ in the VM limit are same as those of $B^{(vac)t}$ and $B^{(vac)s}$ with V = 1 when we make the following replacement in $B^{(vac)t;s}$:

The functions B ^{(vac)t;s} with M ! 0 are independent of the external momentum p_0 and p [8] ^{# 3}. Thus we nd that the functions B ^{(vac)t} and B ^{(vac)s} in the VM lim it are obtained independently of the external momentum p_0 and p:

$$\lim_{V \to M} B^{(vac)t}(p_0;p;M;0) = \frac{1}{V} \frac{2}{8^2};$$

$$\lim_{V \to M} B^{(vac)s}(p_0;p;M;0) = V \frac{2}{8^2}; \quad (A.14)$$

REFERENCES

- T. Hatsuda and T. Kunihiro, Phys. Rept. 247, 221 (1994) [arX iv hep-ph/9401310];
 R. D. Pisarski, arX iv hep-ph/9503330;
 G.E.Brown and M.Rho, Phys. Rept. 269, 333 (1996) [arX iv hep-ph/9504250];
 G.E.Brown and M.Rho, Phys. Rept. 363, 85 (2002) [arX iv hep-ph/0103102];
 T.Hatsuda, H.Shiom i and H. Kuwabara, Prog. Theor. Phys. 95, 1009 (1996) [arX iv nucl-th/9603043];
 R. Rapp and J. Wambach, Adv. Nucl. Phys. 25, 1 (2000) [arX iv hep-ph/0003183.
- M.Bando, T.Kugo, S.Uehara, K.Yam awaki and T.Yanagida, Phys. Rev. Lett. 54, 1215 (1985); M.Bando, T.Kugo and K.Yam awaki, Phys.Rept.164, 217 (1988).
- M.Harada and K.Yam awaki, Phys. Rev. D 64, 014023 (2001) [arX iv hep-ph/0009163].
- M. Harada and C. Sasaki, Phys. Lett. B 537, 280 (2002) [arXiv:hep-ph/0109034].
- 5. M.Harada, Y.Kim and M.Rho, Phys.Rev.D 66,016003 (2002) [arXiv:hep-ph/0111120].
- M. Harada and K. Yam awaki, Phys. Rev. Lett. 86, 757 (2001) [arX iv hep-ph/0010207].

- 7. M. Harada, Y. Kim, M. Rho and C. Sasaki, Nucl. Phys. A 727, 437 (2003) [arXiv:hep-ph/0207012].
- 8. M. Harada and C. Sasaki, arX iv hep-ph/0304282; C. Sasaki, arX iv hep-ph/0304298.
- G. Meissner, J. A. Oller and A. Wirzba, Annals Phys. 297, 27 (2002) [arXiv:nucl-th/0109026].
- 10.H. Georgi, Phys. Rev. Lett. 63 (1989) 1917; Nucl. Phys. B 331, 311 (1990).
- 11.M. Tanabashi, Phys. Lett. B 316, 534 (1993) [arX iv hep-ph/9306237].
- 12.M. Harada and K. Yam awaki, Phys. Rept. 381, 1 (2003) [arX iv:hep-ph/0302103].
- R.D.Pisarskiand M.Tytgat, Phys.Rev.D 54, 2989 (1996) [arXiv:hep-ph/9604404].
- 14. M. Harada and K. Yam awaki, Phys.Lett. B 297, 151 (1992) [arXiv:hep-ph/9210208].
- 15. M. Harada and K. Yam awaki, Phys. Rev. Lett. 83, 3374 (1999) [arX iv hep-ph/9906445].
- 16.M. J. Veltman, Acta Phys. Polon. B 12, 437 (1981).
- 17.M. Harada, Y. Kin, M. Rho and C. Sasaki, Nucl. Phys. A. 730, 379 (2004) [arXiv:hep-ph/0308237].
- 18. D. T. Son and M. A. Stephanov, Phys. Rev. Lett. 88, 202302 (2002) [arX iv hep-ph/0111100].
- 19. D. T. Son and M. A. Stephanov, Phys. Rev. D 66, 076011 (2002) [arX iv hep-ph/0204226].

 $^{^{\}sharp 3}$ In Ref. [7] where V = 1 was taken, it was shown that the hadronic corrections $^{t;s}_{?}(p_0;p;T)$ at the VM lim it are independent of the external momentum p_0 and p. The structure of the integrand in the vacuum part is the same as that in the hadronic part except for the absence of the B ose-E instein distribution function. Thus the vacuum part is also independent of p_0 and p.