

Exclusive Double Diffractive Higgs Boson Production at LHC.

Petrov V. A. and Ryutin R. A.
Institute for High Energy Physics
142 281 Protvino, Russia

Abstract

Exclusive double diffractive (EDD) Higgs boson production is analyzed in the framework of the Regge-eikonal approach. Total and differential cross-sections for the process $p + p \rightarrow p + H + p$ are calculated. Experimental possibilities to find Higgs boson at LHC are discussed.

Keywords

Diffractive Higgs Boson Production { Pomeron { Regge-Eikonal model

1 Introduction

LHC collaborations aimed at working in low (TOTEM)- and high (CMS)- p_T regimes related to typical undulatory (diffractive) and corpuscular (point-like) behaviours of the corresponding cross-sections may offer a very exciting possibility to observe an interplay of both regimes [1]. In theory the "hard part" can be (hopefully) treated with perturbative methods whilst the "soft" one is definitely nonperturbative.

Below we give an example of such an interplay: exclusive Higgs boson production by diffractively scattered protons, i.e. the process $p + p \rightarrow p + H + p$, where $+$ means also a rapidity gap.

This process is related to the dominant amplitude of exclusive two-gluon production. Driving mechanism of the diffractive processes is the Pomeron. Data on the total cross-sections demands unambiguously for the Pomeron with larger-than-one intercept, therefore the need in "unitarisation".

As will be seen below the detection of the Higgs boson (in the bb mode) at LHC in the double diffractive regime looks fairly possible.

2 Calculations

In Figs. 1,2 we illustrate in detail the process $p + p \rightarrow p + H + p$. On-shell proton-gluon amplitudes in Fig. 2 are treated by the method developed in Ref. [2], which is based on the extension of Regge-eikonal approach, and successfully used for the description of the HERA data [3].

The amplitude of the process $p + p \rightarrow p + H + p$ consists of two parts (see Fig. 2). The "hard" part F is the usual gluon-gluon fusion process calculated in the Standard Model [4]. "Soft" amplitudes $T_{1,2}$ are obtained in the Regge-eikonal approach.

We use the following kinematics, which corresponds to the double Regge limit. It is convenient to use light-cone components $(+; -; ?)$. The components of momenta of the hadrons in Fig. 2 are

$$\begin{aligned}
 p_1 &= \left(\frac{\sqrt{s}}{2}; \frac{m^2}{2s}; 0 \right) \\
 p_2 &= \left(\frac{m^2}{2s}; \frac{\sqrt{s}}{2}; 0 \right) \\
 p_1^0 &= \left((1-x_1) \frac{\sqrt{s}}{2}; \frac{x_1^2 + m^2}{(1-x_1) 2s}; x_1 \right) \\
 p_2^0 &= \left(\frac{x_2^2 + m^2}{(1-x_2) 2s}; (1-x_2) \frac{\sqrt{s}}{2}; x_2 \right) \\
 q &= (q_+; q_-; q_\perp) \\
 q_1 &= q + p_1 \quad p_1^0 = q + p_1^0 \\
 q_2 &= q + p_2 \quad p_2^0 = q + p_2^0
 \end{aligned} \tag{1}$$

$x_{1,2}$ are fractions of protons momenta carried by gluons. For two-dimensional transverse

vectors we use boldface type. From the above notations we can obtain the relations:

$$t_{1,2} = \frac{s_{1,2}^2}{1 - s_{1,2}^2}, \quad \frac{s_{1,2}^2(1 + s_{1,2}^2) + s_{1,2}^2 m^2}{1 - s_{1,2}^2}, \quad s_{1,2}^2; \quad s_{1,2} \neq 0 \quad (2)$$

$$\begin{aligned} \cos \theta_0 &= \frac{1 - s_{1,2}^2}{j_{1,2} j_{2,2}} \\ M_H^2 &= s_{1,2} + t_1 + t_2 - 2 \sqrt{t_1 t_2} \cos \theta_0 \\ (p_1 + q)^2 &= m^2 + q^2 + \sqrt{2sq} = s_1 \\ (p_2 - q)^2 &= m^2 + q^2 - \sqrt{2sq} = s_2 : \end{aligned}$$

Physical region of diffractive events with two rapidity gaps is defined by the following kinematical cuts:

$$0.01 \text{ GeV}^2 \leq j_{1,2} \leq 1 \text{ GeV}^2; \quad (3)$$

$$m_{\text{in}} \leq \frac{M_H^2}{s_{\text{max}}} \leq m_{\text{ax}} = 0.1; \quad (4)$$

$$\begin{aligned} \sqrt{t_1} &= \sqrt{t_2} = \sqrt{t_1 + t_2} \\ &= s_{1,2} - M_H^2 = M_H^2 \end{aligned} \quad (5)$$

We can write the relations in terms of $y_{1,2}$ and y_H (hadron and Higgs boson rapidities correspondingly):

$$s_{1,2} = \frac{M_H^2}{s} e^{y_H}; \quad (6)$$

$$j_{1,2} = y = \ln \frac{\sqrt{s_{\text{max}}}}{M_H};$$

$$y_0 = 2.5 \text{ for } \sqrt{s} = 14 \text{ TeV}; M_H = 115 \text{ GeV};$$

$$j_{1,2} = \frac{1}{2} \ln \frac{(1 - s_{1,2}^2)s}{m^2 - t_{1,2}} \approx 9$$

The calculation of the amplitude is based on the nonfactorized scheme. The contribution of the diagram depicted in Fig. 2 is obtained by integrating over all internal loop momenta. It was shown in [5], that the leading contribution arises from the region of the integration, where momentum q is "Glauber-like", i.e. of the order $(k_+ m^2 = \sqrt{s}; k_- m^2 = \sqrt{s}; k_\perp)$, where k 's are of the order 1. The detailed consideration of the loop integral like

$$\int \frac{d^4 q}{(2\pi)^4} \frac{f(q; p_1; p_2; -1; -2)}{(q^2 + i0)(q_1^2 + i0)(q_2^2 + i0)} \quad (7)$$

shows that the main contribution comes from the poles

$$\begin{aligned} q_1^2 &= \sqrt{2s_{1,2}} q \quad q_1^2 = 0; \\ q_2^2 &= \sqrt{2s_{2,2}} q \quad q_2^2 = 0 : \end{aligned}$$

In this case

$$q = \frac{q_2^2}{2s}; \frac{q_1^2}{2s}; q; \quad (8)$$

where

$$q_1^2 = q^2 + t_1^2 + 2q t_1 \cos\left(\theta + \frac{0}{2}\right);$$

$$q_2^2 = q^2 + t_2^2 - 2q t_2 \cos\left(\theta - \frac{0}{2}\right)$$

In this region $q^2 = M_H^2$, and for the ggH -vertex F in the first order of the strong coupling we have the usual [4] expression

$$F = g \frac{2q_2 q_1}{M_H^2} F_{gg \rightarrow H}; \quad (9)$$

$$F_{gg \rightarrow H} = M_H^2 \frac{s}{2} \frac{G_F}{2} f(\theta); \quad (10)$$

$$f(\theta) = \frac{1}{8} \left(1 + \frac{1}{2} \left(1 - \frac{1}{4} L i_2 \left(\frac{2}{1 - \frac{1}{1-i0}} \right) + L i_2 \left(\frac{2}{1 + \frac{1}{1-i0}} \right) \right) \right);$$

where $s = M_H^2 = 4m_t^2$, G_F is the Fermi constant. NLO K -factor 1.5 for the $gg \rightarrow H$ process is included to the final answer.

Taking the general form for T -amplitudes that satisfy identities

$$q T^D = 0; q_1 T^D = 0; \quad (11)$$

and neglecting terms of the order $o(t_i)$, the following expression is found at $t_{ij} \ll 1 \text{ GeV}^2$:

$$T^D(p; q; q_1) = G \frac{p q_1 p q}{p q_1 p q} T_{gp \rightarrow gp}^D(s_i; t_i; q q_1) \quad (12)$$

$$G = g \frac{q_1 q}{q q_1};$$

$$p q_1 = p \frac{p q_1}{q q_1} q;$$

$$p q = p \frac{p q}{q q_1} q_1;$$

For $T_{gp \rightarrow gp}^D$ we use the Regge-eikonal approach [2, 6]. At small t_i it takes the form of the Born approximation, i.e. Regge factor:

$$T_{gp \rightarrow gp}^D(s_i; t_i; q q_1) = c_{gp} e^{-i \frac{s_i}{2} \frac{q q_1}{s_0} \frac{m^2}{m^2}} e^{b_0 t_i}; \quad (13)$$

$$b_0 = \frac{1}{4} \left(\frac{r_{pp}^2}{2} + r_{gp}^2 \right);$$

where $\alpha_p(0) = 1.203$, $\alpha_p(0) = 0.094 \text{ GeV}^{-2}$, $r_{pp}^2 = 2.477 \text{ GeV}^{-2}$ are found in [6]. Parameters $c_{gp} = 3.5$, $r_{gp}^2 = 2.54 \text{ GeV}^{-2}$ are defined from fitting the HERA data on elastic $J=$ production, which will be published elsewhere. The upper bound for the constant $c_{gp}^{\mu p} = 2.3(3.3)$ can be also estimated from the exclusive double diffractive di-jet production at Tevatron, if we take CDF cuts and the upper limit for the exclusive total di-jet cross-section [7]. The effective value $c_{gp} = 2.3$ corresponds to the case, when the Sudakov suppression factor is absorbed into the constant, and $c_{gp} = 3.3$ is obtained when taking into account this factor explicitly.

The full amplitude for Higgs boson production looks as follows:

$$T_{pp \rightarrow p H p} = \int \frac{d^4 q}{(2\pi)^4} \frac{8F(q_1; q_2) T^D(p_1; q; q_1) T^D(p_2; q; q)}{(q^2 + i0)(q_1^2 + i0)(q_2^2 + i0)} : \quad (14)$$

Factor 8 arises from the colour index contraction. Let $l^2 = q^2 = q_1^2 = q_2^2$, $y_H = \langle y_H \rangle = 0$ and contract all the tensor indices, then the integral (14) takes the form

$$T_{pp \rightarrow p H p} = c_{gp}^2 e^{b(t_1 + t_2)} \frac{s}{(2\pi)^2} \frac{M_H^2}{M_H^2} 8F_{gg \rightarrow H} I ; \quad (15)$$

$$b = \alpha_p(0) \ln \frac{s}{M_H^2} + b_0 ; \quad (16)$$

$$I = \int_0^{M_H^2} \frac{dl^2}{l^4} \frac{l^2}{s_0 m^2 + l^2} \quad ; \quad 1.88 \text{ GeV}^{-2} ; \quad (17)$$

where $M_H = 100 \text{ GeV}$, and $s_0 = m^2 = 1 \text{ GeV}^2$ is the scale parameter of the model that is used in the global fitting of the data on pp(pp) scattering for on-shell amplitudes [6]. It remains fixed in the present calculations.

If we take into account the emission of virtual "soft" gluons, while prohibiting the real ones, that could fill rapidity gaps, it results in the Sudakov-like suppression [8]:

$$F_s(l^2) = \exp \left[\frac{3}{2} \int_{l^2}^{M_H^2} \frac{dp_T^2}{p_T^2} s(p_T^2) \ln \frac{M_H^2}{4p_T^2} \right] ; \quad (18)$$

and to the new value of the integral (17):

$$I_s = \int_0^{M_H^2} \frac{dl^2}{l^4} F_s(l^2) \frac{l^2}{s_0 m^2 + l^2} \quad ; \quad 0.38 \text{ GeV}^{-2} ; M_H = 100 \text{ GeV} : \quad (19)$$

In this case the total cross-section becomes 24 times smaller, than without the factor F_s .

Unitarity corrections can be estimated from the elastic pp scattering by the method depicted in Fig.1, where

$$T_X = T_{pp \rightarrow p H p} ; \quad (20)$$

$$\begin{aligned}
V(s; q_T) &= 4s(2)^{2-2}(q_T) + 4s \int_0^Z d^2b e^{iq_T b} e^{i p p! p p} \int_0^1 \dots \\
T_X^{Unit:}(p_1; p_2; \dots) &= \frac{1}{16ss^0} \int_0^Z \frac{d^2 q_T}{(2)^2} \frac{d^2 q_T^0}{(2)^2} V(s; q_T) \dots \\
&\quad V(s; q_T^0); \\
1_T &= \dots \\
2_T &= \dots
\end{aligned}$$

and $p p! p p$ can be found in Ref. [6]. It reduces the integrated cross-section by the factor about 14.

3 Results and discussions

We have the following expression for the differential cross-section:

$$\begin{aligned}
\frac{d}{dt_1 dt_2 d_1 d_2} &= \frac{T_{pp! p p}^{Unit:}}{8s(2)^5} \dots \\
&= \dots + 2(t_1 + t_2) + (t_1 - t_2)^2 \dots
\end{aligned} \tag{21}$$

By partial integrating (21) we obtain the cross-sections $d = dt$ and $d = d$. The first result of our calculations is depicted in the Fig. 3. The antishrinkage of the diffraction peak with increasing Higgs boson mass is the direct consequence of the existence of the additional hard scale M_H , which makes the interaction radius smaller. The dependence is shown in Fig.4.

The second result is the total cross-section versus Higgs boson mass. We obtain the following results for the total cross-section of the EDD Higgs boson production:

c_{gp}	M_H (GeV)	$p+p! p+H+p$ (fb)			
		LHC		TeVatron	
		no Sud. suppr.	Sud. suppr.	no Sud. suppr.	Sud. suppr.
3.5	100 ! 500	110 ! 57	4.6 ! 0.14	12 ! 0.4	0.5 ! 0.001
2.3(3.3)	100 ! 500	20 ! 11	3.6 ! 0.11	2.2 ! 0.08	0.4 ! 0.0009

The above results for LHC energies are depicted in Figs.5-8. The form of the curve originates from the standard gluon-gluon-Higgs vertex and has a peak near $M_H \sim 2m_t$. For the case of Sudakov-like suppression the cross-section vanishes faster with M_H .

It is useful to compare our result with other studies. Results quite close to ours (with the normalization to the CDF data, $c_{gp} = 3.3$) were given in Ref. [8], where the value of the total cross-section is about 3 fb. In both cases the most important suppressing in the mass region $M_H > 100$ GeV is due to (perturbative) Sudakov factors, while the nonperturbative (absorptive) factors play relatively minor role.

Results of other authors were considered in details in [9]. The highest cross-section 2 pb for $M_H = 400$ GeV at LHC energies was obtained in Ref. [10]. They used a nonfactorized form of the amplitude and a "QCD inspired" model for $g p! g p$ amplitudes, taking into account the nonperturbative proton wave functions. Even if we multiply the result of

Ref. [10] by the suppressing factor, it will be larger than ours. This could serve as the indication of the role of nonperturbative effects. Our model is based on the Regge-Eikonal approach for the amplitudes, which is primarily nonperturbative, normalized to the data from HERA on $p \rightarrow J = p$ and improved by the CDF data on the exclusive di-jet production.

To estimate the signal to QCD background ratio for $b\bar{b}$ signal we use the standard expression for $gg \rightarrow b\bar{b}$ amplitude and assumptions [11]–[14]:

possibility to separate n gluon jets from b quark jets. If we cannot do it, it will increase the background by two orders of magnitude under the 50% efficiency.

suppression due to the absence of colour-octet b quark states

suppression of light fermion pair production, when $J_{tot} = 0$ (see also [15], [16])

cut $E_T > 50 \text{ GeV}$ ($\approx 6\theta$), since the cross-section of diffractive $b\bar{b}$ jet production strongly decreases with E_T .

The theoretical result of these numerical estimations is

$$\frac{\text{Signal}(pp \rightarrow pH p \rightarrow pbbp)}{\text{QCD background}} \approx 9.5 \cdot 10^4 \cdot \frac{M_H^3}{M^3} ; \quad (22)$$

where $\frac{M_H^3}{M^3} \approx 0.5 \cdot 10^3$ for Higgs boson masses $100 \leq M_H \leq 350 \text{ GeV}$. For $M_H \approx 115 \text{ GeV}$

$$\frac{\text{Signal}(pp \rightarrow pH p \rightarrow pbbp)}{\text{QCD background}} \approx 3 \cdot 10^8 \cdot \frac{\text{GeV}^3}{M^3} ; \quad (23)$$

where M is the mass resolution of the detector. Similar result was strictly obtained in [11], [12]. More exact estimations of the above ratio by fast Monte-Carlo simulations and the total efficiency of EDD Higgs boson production will be published in a forthcoming paper.

4 Conclusions

We see from the result that there is a real possibility to detect Higgs boson using the usual $b\bar{b}$ signal under the luminosity greater than 10^{32} in EDD events at LHC. Accuracy of the mass measurements could be improved by applying the missing mass method [17].

The low value of the exclusive Higgs boson production cross-section obtained in this paper is mainly due to the Sudakov suppression factor (18), the full validity of which is not obvious for us, because the confinement effects can strongly modify the "real gluon emission".

It is interesting that in spite of different models and quite different ways of account of absorptive effects in our paper and in Ref. [8], the final results (see Fig.8) appeared to be quite close.

Certainly, the cross-sections may be several times larger due to still not very well known non-perturbative factors.

It is possible to generalize our approach for the exclusive production of other particles like ϕ , ω , radion, K , K^* , gravitons, glueballs. In this case cross-sections can be larger than for EDD Higgs boson production, and some other important investigations like partial wave analysis and measurements of the diffractive pattern of the interaction could be done.

Acknowledgements

We are grateful to A. De Roeck, A. Prokudin, A. Rostovtsev, A. Sobol, S. Slabospitsky and participants of BLOIS2003 workshop and several CMS meetings for helpful discussions. This work is supported by the Russian Foundation for Basic Research, grant no. 02-02-16355

References

- [1] V. A. Petrov, Proc. of the 2nd Int. Symp. "LHC: Physics and Detectors". (Eds. A. N. Sissakian and Y. A. Kulchitsky), June 2000, Dubna. P. 223; V. A. Petrov, A. V. Prokudin, S. M. Troshin, N. E. Tyurin, J. Phys. G: Nucl. Part. Phys. 27 (2001) 2225.
- [2] V. A. Petrov, Proceedings of the VIIth Blois Workshop (Ed. M. Haguenaer et al., Editions Frontieres; Paris 1995).
- [3] V. A. Petrov and A. V. Prokudin, Phys. Atom. Nucl. 62 (1999) 1562.
- [4] A. Kniehl, Phys. Rept. 240 (1994) 211.
- [5] A. Berera, J. C. Collins, Nucl. Phys. B 474 (1996) 183.
- [6] V. A. Petrov, A. V. Prokudin, Eur. Phys. J. C 23 (2002) 135.
- [7] CDF Collaboration (K. Borras for the collaboration). FERMILAB-CONF-00-141-E, Jun 2000; K. Goulianos, talk given in the Xth Blois workshop, 2003, Helsinki, Finland; M. Gallinaro, hep-ph/0311192.
- [8] V. A. Khoze, A. D. Martin and M. G. Ryskin, Eur. Phys. J. C 14 (2000) 525; Eur. Phys. J. C 21 (2001) 99; V. A. Khoze, hep-ph/0105224;
- [9] V. A. Khoze, A. D. Martin, M. G. Ryskin, Eur. Phys. J. C 26 (2002) 229.
- [10] J.-R. Cudell and O. F. Hernandez, Nucl. Phys. B 471 (1996) 471.
- [11] V. A. Khoze, A. D. Martin, M. G. Ryskin, Eur. Phys. J. C 19 (2001) 477.

- [12] A . De Roeck, V . A . Khoze, A . D . Martin, R . Orava, M . G . Ryskin, Eur. Phys. J. C 25 (2002) 391.
- [13] V . A . Khoze, A . D . Martin, M . G . Ryskin, Eur. Phys. J. C 23 (2002) 311.
- [14] A . D . Martin, M . G . Ryskin and V . A . Khoze, Phys. Rev. D 56 (1997) 5867.
- [15] A . B ialas and V . Szerem eta, Phys. Lett. B 296 (1992) 191;
A . B ialas and R . Janik, Z. Phys. C 62 (1994) 487.
- [16] J. Pumplin, Phys. Rev. D 52 (1995) 1477.
- [17] M . G . Albrow and A . Rostovtsev, FERM ILAB-PUB-00-173. Aug. 2000.

Figure captions

Fig. 1: The full unitarization of the process $p + p \rightarrow p + X + p$.

Fig. 2: The process $p + p \rightarrow p + H + p$. Absorption in the initial and final pp-channels is not shown.

Fig. 3: t -distribution $d\sigma/dt = \sigma_{\text{tot}}$ of the process $p + p \rightarrow p + H + p$ for Higgs boson masses 100 and 500 GeV.

Fig. 4: η -distribution $d\sigma/d\eta = \sigma_{\text{tot}}$ of the process $p + p \rightarrow p + H + p$ for $M_H = 100$ GeV.

Fig. 5: The total cross-section (in fb) of the process $p + p \rightarrow p + H + p$ versus Higgs boson mass for $c_{\text{qp}} = 3.5$ without Sudakov-like suppression.

Fig. 6: The total cross-section (in fb) of the process $p + p \rightarrow p + H + p$ versus Higgs boson mass for $c_{\text{qp}} = 3.5$ with Sudakov-like suppression.

Fig. 7: The total cross-section (in fb) of the process $p + p \rightarrow p + H + p$ versus Higgs boson mass for $c_{\text{qp}} = 2.3$ without Sudakov-like suppression.

Fig. 8: The total cross-section (in fb) of the process $p + p \rightarrow p + H + p$ versus Higgs boson mass for $c_{\text{qp}} = 3.3$ with Sudakov-like suppression.




Figure 1:




Figure 2:




Figure 3:




Figure 4:




Figure 5:




Figure 6:




Figure 7:




Figure 8: