

CP asymmetry, branching ratios, and isospin breaking effects of $B \rightarrow K$ with the perturbative QCD approach

M. Matsumori and A. I. Sanda^y

Department of Physics, Graduate School of Science, Nagoya University.

Y.-Y. Keum^z

Institute of Physics, Academia Sinica, Taipei, Taiwan 115, Republic of China.

The main contribution to the radiative $B \rightarrow K$ mode is from penguin operators which are quantum corrections. Thus, this mode may be useful in the search for physics beyond the standard model. In this paper, we compute the branching ratio, direct CP asymmetry, and isospin breaking effects within the standard model in the framework of perturbative QCD, and discuss how new physics might show up in this decay.

PACS numbers: 13.20.He, 12.38.Bx, 13.40.Gp, 13.40.Hq

I. INTRODUCTION

The large CP violation in $B \rightarrow J/\psi K_s$ decay mode predicted by the standard model with Kobayashi-Masakawa (KM) scheme has been verified by B factories at KEK (High Energy Accelerator Research Organization) and Stanford Linear Accelerator Center (SLAC). The standard model predicts the CP asymmetries for $B \rightarrow J/\psi K_s$ and $B \rightarrow K_s$ to be equal to $\sin 2\beta$. However, recent experimental data from Belle showed that these asymmetries differ by nearly 2%; the averaged $\sin 2\beta$ from Belle and BaBar in $B \rightarrow J/\psi K_s$ system is $\sin 2\beta = 0.736 \pm 0.049$ [1] and in $B \rightarrow K_s$ decay mode is $\sin 2\beta = 0.06 \pm 0.33 \pm 0.09$ from Belle [2], and $\sin 2\beta = 0.50 \pm 0.25^{+0.07}_{-0.04}$ from BaBar [3]. Experimental error is still large, so the situation is inconclusive, but if this result continues to hold, it implies existence of new physics beyond the standard model.

In this paper, we want to concentrate on $B \rightarrow K$ decay mode. The decay mode has large branching ratio, so the experimental error on the CP asymmetry has been getting small and is now down to several percent.

CP Asymmetry is defined as

$$A_{CP} = \frac{\Gamma(B \rightarrow K^-) - \Gamma(B \rightarrow K^+)}{\Gamma(B \rightarrow K^-) + \Gamma(B \rightarrow K^+)}; \quad (1)$$

and in general, theoretical predictions of the CP asymmetries depend less on hadronic parameters than those of branching ratios as many uncertainties cancel in the ratio. So comparing predictions for CP asymmetries within the standard model with experimental data may be effective way to search for new physics. Many authors have pointed out for some time, that the CP asymmetry in this mode is very small. We can easily understand why the asymmetry is so small. In order to generate CP asymmetry, at least two amplitudes with nonvanishing relative weak and strong phases must interfere. This decay is mainly caused by an O_7 operator, and other contributions which interfere with this contribution are small and the Cabibbo-Kobayashi-Masakawa quark-mixing matrix (CKM) unitary triangle is crushed, making the CP asymmetry in this decay mode very small. However, if we were to look for new physics, we need to be able to give a quantitative estimate of the standard model contribution to the CP asymmetry. For this purpose, we include small contributions which interfere with O_7 , including also the long distance contributions, for example, $B \rightarrow K J/\psi K$.

Furthermore, the isospin breaking effect ϵ_{0+} is also very interesting because its size and sign are sensitive to the existence of physics beyond the standard model.

$$\begin{aligned} \epsilon_{0+} &= \frac{\Gamma(B^0 \rightarrow K^0) - \Gamma(B^+ \rightarrow K^+)}{\Gamma(B^0 \rightarrow K^0) + \Gamma(B^+ \rightarrow K^+)} \\ &= \frac{(\Gamma_{B^+ \rightarrow B^0})\Gamma(B^0 \rightarrow K^0) - \Gamma(B^+ \rightarrow K^+)}{(\Gamma_{B^+ \rightarrow B^0})\Gamma(B^0 \rightarrow K^0) + \Gamma(B^+ \rightarrow K^+)} \end{aligned} \quad (2)$$

In order to test the standard model, we need to know if the penguin contribution within the standard model can explain the experimental data. Experiments show $\epsilon_{0+} = +0.012 \pm 0.044 \pm 0.026$ in Belle [4] and

^Electronic address: mika@eken.phys.nagoya-u.ac.jp

^yElectronic address: sanda@eken.phys.nagoya-u.ac.jp

^zElectronic address: yykeum@phys.sinica.edu.tw

$\alpha_s = 0.050 \quad 0.045 \quad 0.028 \quad 0.024$ in BaBar [5]. More precise data will become available in the near future, so the theoretical prediction of its size and sign of this asymmetry should be pinned down.

In this paper, we calculate the branching ratio, direct CP asymmetry, and isospin breaking effects in $B \rightarrow K$ decay mode, based on the standard model. First, we briefly review the concept of the pQCD in Sect.II, and in Sect.III, we show the effective Hamiltonian which causes $B \rightarrow K$ decay. Then we present the factorization formulas for the $B \rightarrow K$ decay mode in Sect.IV, and in Sect.V, we mention about the long distance contributions. Next we will show the numerical results in Sect.VI, and Sect.VII is our conclusion. Finally in Appendix A, we present a brief review of pQCD.

II. OUTLINE OF PQCD

Theoretically, it is easy to analyze the inclusive B meson decay like $B \rightarrow X_s$ because we can estimate the decay width, for example, by inserting the complete set for all possible intermediate states. The experimental and theoretical branching ratio of $B \rightarrow X_s$ are

$$\text{Br}(B \rightarrow X_s)^{\text{exp}} = (3.52^{+0.30}_{-0.28}) \cdot 10^{-4}; \quad [6]$$

$$\text{Br}(B \rightarrow X_s)^{\text{th}} = (3.57 \pm 0.30) \cdot 10^{-4}; \quad [7]$$

and this good agreement strongly constraints new physics parameters. However, inclusive decays are experimentally difficult to analyze because all $B \rightarrow X_s$ candidates should be counted. If we can directly calculate the exclusive decay mode $B \rightarrow K$, we ought to obtain many interesting results to test the standard model or to search for new physics.

Perturbative QCD is one of the theoretical instrument for handling the exclusive decay modes. The concept of pQCD is the factorization between soft and hard dynamics. In order to physically understand the pQCD approach, we consider B^0 meson decays into K^0 meson and in the rest frame of the B^0 meson (Fig.1). The heavy b quark which has most of B^0 meson mass is nearly static in this frame and the other quark, which forms the B^0 meson together with the b quark, called the spectator quark, carries momentum of order $O(\Lambda) = O(M_B - m_b)$. This b quark decays into the light s quark and through the electromagnetic penguin operator and the decay products dash away back-to-back, with momentum of $O(M_B/2)$. (This process is depicted in Fig.1(a).) K^0 meson is composed of s quark and a spectator quark. In order for the fast moving s quark and slow moving spectator d quark to form a K^0 meson and nothing else, the spectator quark must be kicked by the gluon, so that the s and d quark have more or less parallel momenta in the direction of K^0 . (This process is depicted in Fig.1(b).) Since the invariant-mass square of this gluon is the order of $O(M_B)$, we can treat this decay process perturbatively.




FIG. 1: The left figure is no gluon exchange diagram. s and spectator d are not lines up to form an energetic K meson. In order to hadronize K meson, one gluon with large q^2 should be exchanged.




FIG. 2: Example of annihilation diagram which produces strong phase through the branch-cut.

There is also diagram shown in Fig.2. This can also be computed in the pQCD approach. The diagram can be cut along the dotted line indicating the presence of the physical intermediate state. This results in a strong interaction phase which can be computed. The direct CP asymmetry is caused by interfering some amplitudes which have relative weak and strong phase, and it can be written in the form proportioning to $\sin(\phi_1 - \phi_2) \sin(\phi_1 - \phi_2)$: in short, it depends on both weak and strong phases.

$$A(B \rightarrow f) = A_1 e^{i\phi_1} e^{i\phi_1} + A_2 e^{i\phi_2} e^{i\phi_2}$$

$$A(B \rightarrow f) = A_1 e^{i\phi_1} e^{i\phi_1} + A_2 e^{i\phi_2} e^{i\phi_2}$$

We can determine the strong phases by using the pQCD approach, then we can extract the information about the weak phases and examine the standard model. A more detailed review for the pQCD approach is in Appendix A.

III. KINEMATICS FOR $B \rightarrow K$ DECAY MODE

The effective Hamiltonian which induces flavor-changing $b \rightarrow s$ transition is given by [8]

$$H_{\text{eff}} = \frac{G_F}{2} \sum_{q=u,c} V_{qb} V_{qs}^* \left[C_1 (O_1^{(q)}) + C_2 (O_2^{(q)}) \right] + \sum_{i=3,8g} V_{tb} V_{ts}^* \left[C_i (O_i) + \text{h.c.} \right]; \quad (3)$$

$$\begin{aligned}
O_1^{(q)} &= (s_i q_j)_{V-A} (q_j b_i)_{V-A}; \\
O_2^{(q)} &= (s_i q_i)_{V-A} (q_j b_j)_{V-A}; \\
O_3 &= (s_i b_i)_{V-A} (q_j q_j)_{V-A}; \\
O_4 &= (s_i b_j)_{V-A} (q_j q_i)_{V-A}; \\
O_5 &= (s_i b_i)_{V-A} (q_j q_j)_{V+A}; \\
O_6 &= (s_i b_j)_{V-A} (q_j q_i)_{V+A}; \\
O_7 &= \frac{e}{4} s_i (m_s P_L + m_b P_R) b_i F^-; \\
O_{8g} &= \frac{g}{4} s_i (m_s P_L + m_b P_R) T_{ij}^a b_j G^a;
\end{aligned} \tag{4}$$

where $P_R^L = (1 \pm \gamma_5)/2$. We define the B meson and the K meson momenta P_1 and P_2 in the light-cone coordinates

$$p = (p^+, p^-, p_T) = \left(\frac{p^0 + p^3}{2}, \frac{p^0 - p^3}{2}, p^1, p^2 \right) \tag{5}$$

within the B meson rest frame as

$$P_1 = (P_1^+, P_1^-, P_{1T}) = \left(\frac{M_B}{2}, 1, 0_T \right); \tag{6}$$

$$P_2 = (P_2^+, P_2^-, P_{2T}) = \left(\frac{M_B}{2}, 0, 1, 0_T \right); \tag{7}$$

and photon and the K meson transverse polarization vectors as

$$\begin{aligned}
(\epsilon) &= (0; 0; \frac{1}{\sqrt{2}} (1; i)); \\
\epsilon_K(\epsilon) &= (0; 0; \frac{1}{\sqrt{2}} (1; i));
\end{aligned} \tag{8}$$

Throughout this paper, we keep only terms of order r_K in the computation of the numerator, where $r_K = M_K/M_B$.

The fractions of the momenta which have the spectator quarks in B and K meson are $x_1 = k_1^+ = P_1^+$ and $x_2 = k_2^+ = P_2^+$, so the momenta of these spectator quarks are expressed as follows,

$$k_1 = (k_1^+, k_1^-, k_{1T}) = \left(\frac{M_B}{2} x_1, 0; k_{1T} \right); \tag{9}$$

$$k_2 = (k_2^+, k_2^-, k_{2T}) = \left(0; \frac{M_B}{2} x_2; k_{2T} \right); \tag{10}$$

then the b and s quark momenta are $p_b = P_1 - k_1$ and $p_s = P_2 - k_2$, and we neglect the masses of the light quarks and identify the b quark mass with the B meson mass in calculations of the hard scattering amplitudes. The term proportional to $m_b = M_B - m_b$ is generated by higher order effects, so we included this effect in our error estimate.

From here, we extract the formulas for decay amplitudes caused by each operators,

$$\begin{aligned}
M &= \langle H | j H_{eff} | j \rangle i \\
&= \frac{G_F}{2} X_i V_{CKM} C_i(\gamma) \langle H | j O_i(\gamma) | j \rangle i
\end{aligned} \tag{11}$$

and they can be decomposed into scalar and pseudoscalar components as

$$M = (\epsilon_K)_K M^S + i (\epsilon_K)_K M^P; \tag{12}$$

IV. FORMULAS

In this section, we want to show the explicit formulas of the decay amplitudes caused by operators given in Sect.III.

A. O_7 contribution

If we define the common factor as

$$F^{(0)} = \frac{G_F}{2} e V_{cb} V_{cs} C_F M_B^5; \tag{13}$$

where C_F is color factor, and i as $V_{ib} V_{is} = V_{cb} V_{cs}$, the decay amplitude M_7 in Fig.3 can be expressed as follows.




FIG. 3: Feynman diagrams of electromagnetic penguin operator O_7 . A photon is emitted through the O_7 operator, and one hard gluon exchange is needed to form hadrons.




FIG. 4: Feynman diagrams of chromomagnetic penguin operator O_{sg} . A hard gluon is emitted through the O_{sg} operator and glued to the spectator quark line. Then photon is emitted by bremsstrahlung of external quark lines.

$$M_8^{S(d)} = F^{(0)}_{tQ_q} \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 \int_{b_2}^{b_2} db_2 \int_{B(x_1; b_1)}^{S_t(x_2)} S_t(x_2) \int_{s(t_8^d)}^{e^{[S_B(t_8^d) S_K(t_8^d)]}} C_8(t_8^d) H_8^{(d)}(A_8^{(2)}(b_1; E_8 b_1; E_8 b_2)) [6x_2 r_K^v(x_2) + (2 + x_2 - x_1) r_K^T(x_2)] \quad (22)$$

$$M_8^{P(d)} = F^{(0)}_{tQ_q} \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 \int_{b_2}^{b_2} db_2 \int_{B(x_1; b_1)}^{S_t(x_2)} S_t(x_2) \int_{s(t_8^d)}^{e^{[S_B(t_8^d) S_K(t_8^d)]}} C_8(t_8^d) H_8^{(d)}(A_8^{(2)}(b_1; E_8 b_1; E_8 b_2)) [(2 + x_2 - x_1) r_K^T(x_2) + 6x_2 r_K^a(x_2)] \quad t_8^d = \max(A_8^{(2)}(b_1; E_8 b_1; E_8 b_2)) \quad (23)$$

$$H_8^{(a)}(A_8 b_2; B_8 b_1; B_8 b_2) = K_0(A_8 b_2) (A_8 b_2) K_0(B_8 b_1) I_0(B_8 b_2) + (b_1 \otimes b_2) \quad (24)$$

$$H_8^{(b)}(A_8 b_1; C_8 b_1; C_8 b_2) = \frac{i}{2} K_0(A_8 b_1) (A_8 b_1) H_0^{(1)}(C_8 b_1) J_0(C_8 b_2) + (b_1 \otimes b_2) \quad (25)$$

$$H_8^{(c)}(A_8^{(2)}(b_1; D_8 b_1; D_8 b_2) = (A_8^{(2)}(b_1; D_8 b_1) K_0(D_8 b_1) I_0(D_8 b_2) + (b_1 \otimes b_2) + (A_8^{(2)}(b_1; D_8 b_1) H_0^{(1)}(D_8 b_1) I_0(D_8 b_2) + (b_1 \otimes b_2) \quad (26)$$

$$H_8^{(d)}(A_8^{(2)}(b_1; E_8 b_1; E_8 b_2) = (A_8^{(2)}(b_1; E_8 b_1) K_0(E_8 b_1) I_0(E_8 b_2) + (b_1 \otimes b_2) + (A_8^{(2)}(b_1; E_8 b_1) H_0^{(1)}(E_8 b_1) J_0(E_8 b_2) + (b_1 \otimes b_2) \quad (27)$$

$$A_8^2 = x_1 x_2 M_B^2; \quad B_8^2 = M_B^2 (1 + x_1); \quad C_8^2 = M_B^2 (1 - x_2); \quad A_8^{(2)} = (x_1 - x_2) M_B^2; \quad D_8^2 = x_1 M_B^2; \quad E_8^2 = x_2 M_B^2 \quad (28)$$

C. Loop contributions

1. Quark line photon emission

Next we want to mention about charm and up quark penguin contributions (Fig.5). The subtitle like "Quark line photon emission" means that a photon is emitted through the external quark lines. We define the c and u loop function in order that the $b \rightarrow sg$ vertex can be expressed as $(1 - \frac{5}{3}) I^a$. It has the gauge invariant form [13] and the explicit formula is as follows,

$$I^a = \frac{g T^a}{2} (k^2 g - k \cdot k) \int_0^1 dx x (1-x) \left[1 + \log \frac{m_i^2 - x(1-x)k^2}{t^2} \right] \\ = \frac{g T^a}{8} (k^2 g - k \cdot k) G(m_i^2; k^2; t) - \frac{2}{3}; \quad (29)$$




FIG. 5: Feynman diagrams of "Quark line photon emission". The charm or up loop go to gluon and attach to the spectator quark line. A photon is emitted through the external quark lines.

where k is the gluon momentum and m_i is the loop internal quark mass.

$$\begin{aligned}
 G(m_i^2; k^2; t) = & \left(k^2 \right) \frac{2}{3} \frac{h_5}{3} + \frac{4m_i^2}{k^2} \ln \frac{m_i^2}{t^2} + 1 + \frac{2m_i^2}{k^2} \left(1 - \frac{4m_i^2}{k^2} \right) \ln \frac{1 - 4m_i^2 = k^2}{1 - 4m_i^2 = k^2 + 1} \\
 & + \left(k^2 \right) (4m_i^2 - k^2) \frac{2}{3} \frac{h_5}{3} + \frac{4m_i^2}{k^2} \ln \frac{m_i^2}{t^2} - 2 + 1 + \frac{2m_i^2}{k^2} \left(\frac{4m_i^2}{k^2} - 1 \right) \arctan \frac{1}{\sqrt{4m_i^2 = k^2}} \\
 & + \left(k^2 - 4m_i^2 \right) \frac{2}{3} \frac{h_5}{3} + \frac{4m_i^2}{k^2} \ln \frac{m_i^2}{t^2} + 1 + \frac{2m_i^2}{k^2} \left(1 - \frac{4m_i^2}{k^2} \right) \ln \frac{1}{1 + \frac{1 - 4m_i^2 = k^2}{1 - 4m_i^2 = k^2}} + i
 \end{aligned} \quad (30)$$

The loop function G has the dependence of gluon momentum square of k^2 . But there is no singularity when we take the limit of $k \rightarrow 0$, so we can neglect k_T components of k^2 in the loop function G .

Then the "Quark line photon emission" contributions can be expressed as follows.

$$\begin{aligned}
 M_{1i}^{S(a)} &= M_{1i}^{P(a)} \\
 &= \frac{Q_b}{2} F^{(0)}_i \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 db_2 \int_{B_1}^{B_2} db_1 db_2 C_2(t_2^a) S_t(x_1) e^{[S_B(t_2^a) - S_K(t_2^a)]} H_2^{(a)}(A_2 b_2; B_2 b_1; B_2 b_2) \\
 &\quad G(m_i^2; A_2^2; t_2^a) \frac{2}{3} x_1 x_2 r_K \left(\frac{v}{K}(x_2) - \frac{a}{K}(x_2) \right) \quad (t_2^a = \max(A_2; B_2; 1=b_1; 1=b_2))
 \end{aligned} \quad (31)$$

$$\begin{aligned}
 M_{1i}^{S(b)} &= M_{1i}^{P(b)} \\
 &= \frac{Q_s}{2} F^{(0)}_i \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 db_2 \int_{B_1}^{B_2} db_1 db_2 C_2(t_2^b) S_t(x_2) e^{[S_B(t_2^b) - S_K(t_2^b)]} H_2^{(b)}(A_2 b_1; C_2 b_1; C_2 b_2) \\
 &\quad G(m_i^2; A_2^2; t_2^b) \frac{2}{3} x_2^2 r_K \left(\frac{v}{K}(x_2) + \frac{a}{K}(x_2) \right) + 3x_1 x_2 \frac{T}{K}(x_2) \quad t_2^b = \max(A_2; C_2; 1=b_1; 1=b_2)
 \end{aligned} \quad (32)$$

$$\begin{aligned}
 M_{1i}^{S(c)} &= M_{1i}^{P(c)} \\
 &= \frac{Q_q}{2} F^{(0)}_i \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 db_2 \int_{B_1}^{B_2} db_1 db_2 C_2(t_2^c) S_t(x_1) e^{[S_B(t_2^c) - S_K(t_2^c)]} H_2^{(c)}(A_2^0 b_2; D_2 b_1; D_2 b_2) \\
 &\quad G(m_i^2; A_2^0; t_2^c) \frac{2}{3} x_2 r_K \left(\frac{v}{K}(x_2) + \frac{a}{K}(x_2) \right) - x_1 \frac{T}{K}(x_2) \quad t_2^c = \max(A_2^0; D_2; 1=b_1; 1=b_2)
 \end{aligned} \quad (33)$$

$$\begin{aligned}
 M_{1i}^{S(d)} &= \frac{Q_q}{2} F^{(0)}_i \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 db_2 \int_{B_1}^{B_2} db_1 db_2 C_2(t_2^d) S_t(x_2) e^{[S_B(t_2^d) - S_K(t_2^d)]} H_2^{(d)}(A_2^0 b_1; E_2 b_1; E_2 b_2) \\
 &\quad G(m_i^2; A_2^0; t_2^d) \frac{2}{3} x_2 r_K [\beta(1+x_2) \frac{v}{K}(x_2) - (1-x_2) \frac{a}{K}(x_2)] + 3(x_2 - x_1) \frac{T}{K}(x_2)
 \end{aligned} \quad (34)$$




FIG. 6: The Feynman diagrams of "Loop line photon emission". The charm or up loop go to gluon and attach to the spectator quark line. A photon is emitted by bremsstrahlung of loop quark line.

$$M_{1i}^{P(d)} = \frac{Q_q}{2} F^{(0)}_{i0} \int_0^1 dx_1 dx_2 \int_0^1 db_1 db_2 db_3 \int_0^1 dt_2^d \int_0^1 dt_2^d S_t(x_2) e^{[S_B(t_2^d) - S_K(t_2^d)]} H_2^{(d)}(\vec{A}_2^{02}; \vec{b}_1; E_2 b_1; E_2 b_2) \quad (35)$$

$$G(m_i^2; A_2^{02}; t_2^d) = \frac{2i\hbar}{3} x_2 r_K \left[(1-x_2) \frac{v_K}{K}(x_2) + 3(1+x_2) \frac{a_K}{K}(x_2) \right] + 3(x_2-x_1) \frac{T_K}{K}(x_2)$$

$$t_2^d = \max(\vec{A}_2^{02}; E_2; 1=b_1; 1=b_2)$$

$$H_2^{(a)}(A_2 b_2; B_2 b_1; B_2 b_2) = H_8^{(a)}(A_8 b_2; B_8 b_1; B_8 b_2)$$

$$H_2^{(b)}(A_2 b_1; C_2 b_1; C_2 b_2) = H_8^{(b)}(A_8 b_1; C_8 b_1; C_8 b_2)$$

$$H_2^{(c)}(\vec{A}_2^{02}; D_2 b_1; D_2 b_2) = H_8^{(c)}(\vec{A}_8^{02}; D_8 b_1; D_8 b_2) \quad (36)$$

$$H_2^{(d)}(\vec{A}_2^{02}; E_2 b_1; E_2 b_2) = H_8^{(d)}(\vec{A}_8^{02}; E_8 b_1; E_8 b_2)$$

$$A_2^2 = x_1 x_2 M_B^2; \quad B_2^2 = (1+x_1) M_B^2; \quad C_2^2 = (1-x_2) M_B^2; \quad A_2^{02} = (x_1-x_2) M_B^2; \quad D_2^2 = x_1 M_B^2; \quad E_2^2 = x_2 M_B^2 \quad (37)$$

2. Loop line photon emission

Next we consider the "Loop line photon emission": a photon is emitted through the c or u quark loop line. We sum up Fig.6 (a) and 6 (b), the $b \rightarrow sg$ decay amplitude is expressed as

$$A(b \rightarrow sg) = (q)_a(k) s(p^0) I^a b(p); \quad (38)$$

where vertex function I is defined as follows [14, 15, 16],

$$I^a = F_1^a(k, q) (q-k) + q \frac{q \cdot k}{k^2} k - q \frac{q \cdot k}{L} L + F_2^a(k, q) k + k^2 \frac{q \cdot k}{L}; \quad (39)$$

$$F_1^a = \frac{i2}{3} \frac{p}{2} \overline{egT^a G_F} \int_0^1 dx \int_0^1 dy \frac{xy}{m_i^2 2xy(k-q) \cdot kx(1-x)}; \quad (40)$$

$$F_2^a = \frac{i2}{3} \frac{p}{2} \overline{egT^a G_F} \int_0^1 dx \int_0^1 dy \frac{x(1-x)}{m_i^2 2xy(k-q) \cdot kx(1-x)}; \quad (41)$$

where $L = (1-x)^2$, k is the gluon momentum and q is the photon one. Then the amplitudes can be expressed as follows:

$$M_{2i}^S = \frac{4}{3} F^{(0)}_{i0} \int_0^1 dx \int_0^1 dy \int_0^1 dx_1 dx_2 \int_0^1 db_1 db_2 \int_0^1 dt_2^d \int_0^1 dt_2^d S_t(x_2) e^{[S_B(t_2^d) - S_K(t_2^d)]} H_2(\vec{b}_1 A; \vec{b}_1 \vec{B}^2) \quad (42)$$

$$\frac{1}{xyx_2 M_B^2 m_i^2} xyx_2 (1-2x_2) r_K \frac{v_K}{K}(x_2) (1+2x_1) \frac{T_K}{K}(x_2) + r_K \frac{a_K}{K}(x_2) + x(1-x) x_2^2 r_K \left(\frac{v_K}{K}(x_2) + \frac{a_K}{K}(x_2) \right) + 3x_1 x_2 \frac{T_K}{K}(x_2)$$




FIG. 7: The energy scale of the loop contribution is $t = \max(A; \sqrt{B^2}; 1=b_1)$, where A^2 is gluon momentum, B^2 is the loop momentum, and b_1 is the transverse interval between the quark and antiquark pair in the B meson.

$$M_{2i}^P = \frac{4}{3} F^{(0)}_i \int_0^1 dx \int_0^1 dy \int_0^1 dx_1 dx_2 \int_{b_1}^{b_2} db_1 \int_{B^2}^{B^2} dA^2 C_2(t_2) s(t_2) e^{[S_B(t_2) - S_K(t_2)]} H_2(b_1 A; b_1 \sqrt{B^2})$$

$$\frac{1}{xyx_2 M_B^2 - m_i^2} xyx_2 r_K^v(x_2) (1 + 2x_1) r_K^T(x_2) + (1 - 2x_2) r_K^a(x_2)$$

$$+ x(1-x) [x_2^2 r_K^v(x_2) + r_K^a(x_2)] + 3x_1 x_2 r_K^T(x_2)$$

$$t_2 = \max(A; \sqrt{B^2}; 1=b_1) \quad (43)$$

$$H_2(b_1 A; b_1 \sqrt{B^2}) = K_0(b_1 A) - K_0(b_1 \sqrt{B^2}) \quad (B^2 > 0)$$

$$K_0(b_1 A) - \frac{i}{2} H_0(b_1 \sqrt{B^2}) \quad (B^2 < 0) \quad (44)$$

$$A^2 = x_1 x_2 M_B^2; \quad B^2 = x_1 x_2 M_B^2 - \frac{y}{1-x} x_2 M_B^2 + \frac{m_i^2}{x(1-x)} \quad (45)$$

In general, it is hard to estimate the u loop contributions accurately because of the nonperturbative hadronic uncertainties. For $k^2 < 1 \text{ GeV}$, nonperturbative correction to the u quark loop shown in Fig.7 is large and in fact uu pair may be better represented by resonances. On the other hand if k^2 is large, the perturbative computation is expected to be reliable.

In the pQCD approach, the factorization energy scale t is determined at each point of the integration, i.e. for each point $(x_1; x_2; b_1; b_2)$. Then these variables are integrated over the entire physical region. So for each point $(x_1; x_2; b_1; b_2)$, $s(t)$ can be determined. Thus we can observe the contribution to the amplitude as a function of $s(t)$. Figures 8 and 9 show the distribution of $s(t)$ for a diagram with the c quark loop, and u quark loop, respectively. We can see that the major part of the c loop contribution comes from a perturbative region, on the other hand u loop contribution includes also a nonperturbative region. Since M_{2u}^S and M_{2u}^P gets considerable contributions from the nonperturbative region, we introduce 100% theoretical error for these amplitudes.

D. Annihilation contributions

1. Tree annihilation

We now discuss the annihilation contributions caused by O_1 and O_2 operators. The diagrams are shown in Fig.10. The operators O_1, O_2 can be rewritten as

$$O_1 = (s_i u_j)_{V-A} (u_j b_i)_{V-A} = (s_i b_i)_{V-A} (u_j u_j)_{V-A}; \quad (46)$$

$$O_2 = (s_i u_i)_{V-A} (u_j b_j)_{V-A} = (s_i b_j)_{V-A} (u_j u_i)_{V-A}; \quad (47)$$

These annihilation contributions are tree processes: no hard gluons are needed because they are four Fermi interaction processes and do not include spectator quarks which should be line up to form hadrons. However, these contributions are small because it has $(V-A)(V-A)$ vertex: gets chiral suppression, and its CKM factor is $V_{ub}V_{us}$, $O(10^{-2})$ suppression compared to $V_{tb}V_{ts}$ and $V_{cb}V_{cs}$. Denoting $a_2(t) = C_2(t) + C_1(t) = 3$, the each decay amplitudes are as follows:

$$M_{2i}^{S(a)} = M_{2i}^{P(a)}$$

$$= F^{(0)}_u \frac{3}{4M_B^2} \frac{6Q_b f_K}{B} r_K \int_0^1 dx_1 \int_{b_1}^{b_2} db_1$$

$$a_2(t_a^a) S_t(x_1) e^{[S_B(t_a^a)]} H_2(b_1 A; b_1 \sqrt{B^2})$$

$$(t_a^a = \max(A; 1=b_1)) \quad (48)$$

$$M_{2i}^{S(b)} = F^{(0)}_u \frac{3}{4M_B^2} \frac{6Q_s f_B}{B} r_K \int_0^1 dx_2 \int_{b_2}^{b_2} db_2$$

$$a_2(t_a^b) S_t(x_2) e^{[S_K(t_a^b)]} \frac{i}{2} H_0^{(1)}(b_2 B_a)$$

$$h_i(2-x_2) r_K^v(x_2) + x_2 r_K^a(x_2) \quad (49)$$

$$M_{2i}^{P(b)} = F^{(0)}_u \frac{3}{4M_B^2} \frac{6Q_s f_B}{B} r_K \int_0^1 dx_2 \int_{b_2}^{b_2} db_2$$

$$a_2(t_a^b) S_t(x_2) e^{[S_K(t_a^b)]} \frac{i}{2} H_0^{(1)}(b_2 B_a)$$

$$[x_2 r_K^v(x_2) + (2-x_2) r_K^a(x_2)]$$

$$t_a^b = \max(B_a; 1=b_2) \quad (50)$$




FIG. 8: The horizontal line is $s(t)=$ and the vertical axis is the contribution from the each energy region in s to the total decay amplitude. The left figure is the real part and right one is imaginary part of the c loop contribution. These figures show that we can compute the c quark loop contribution perturbatively.




FIG. 9: The horizontal line is $s(t)=$ and the vertical axis is the distribution from the each energy region in s to the total decay amplitude. The left figure is the real part and right one is imaginary part of the u loop contribution. These figures show that nonperturbative contributions are important, so we cannot accurately compute the u quark loop contribution and have to take into account the hadronic uncertainty.




FIG. 10: Annihilation diagrams caused by O_1, O_2 operators. In these cases, no hard gluons are needed because they are four Fermi interactions and do not include spectator quarks which should be line up to form hadrons.

$$M_2^{S(d)} = F^{(0)}_u \frac{3}{4M_B^2} \frac{p_-}{6Q_u f_B} r_K \int_0^1 dx_2 \int_0^1 db_2 \int_0^1 dt_a^d S_t(x_2) e^{[S_K(t_a^d)]} i \frac{1}{2} H_0^{(1)}(b_2 D_a) [(1+x_2) \frac{v}{K}(x_2) (1-x_2) \frac{a}{K}(x_2)] \quad (52)$$

$$M_2^{P(d)} = F^{(0)}_u \frac{3}{4M_B^2} \frac{p_-}{6Q_u f_B} r_K \int_0^1 dx_2 \int_0^1 db_2 \int_0^1 dt_a^d S_t(x_2) e^{[S_K(t_a^d)]} i \frac{1}{2} H_0^{(1)}(b_2 D_a) [(1-x_2) \frac{v}{K}(x_2) (1+x_2) \frac{a}{K}(x_2)] \quad (53)$$

$$t_a^d = \max(D_a; 1-b_2)$$

$$A_a^2 = (1+x_1)M_B^2; \quad B_a^2 = (1-x_2)M_B^2; \quad C_a^2 = x_1 M_B^2; \quad D_a^2 = x_2 M_B^2 \quad (54)$$

2. QCD Penguin

Next we mention the QCD penguin annihilation caused by O_3, O_6 operators like in Fig.11. Here we define $a_4(t) = C_4(t) + C_3(t)=3$; $a_6(t) = C_6(t) + C_5(t)=3$. O_3, O_4 have the same expression of O_1, O_2 annihilation contributions. O_5, O_6 have a $(V-A)(V+A)$ vertex so they have chiral enhancement compared to $(V-A)(V-A)$ vertex, and its CKM factor are $V_{tb}V_{ts}$, then its' contributions are

$$M_2^{S(c)} = M_2^{P(c)} = F^{(0)}_u \frac{3}{4M_B^2} \frac{p_-}{6Q_u f_K} r_K \int_0^1 dx_1 \int_0^1 db_1 \int_0^1 dt_a^c S_t(x_1) e^{[S_B(t_a^c)]} \frac{1}{B}(x_1; b_1) K_0(b_1 C_a) \quad (51)$$

$$(t_a^c = \max(C_a; 1-b_1))$$




FIG. 11: Annihilation diagrams caused by $O_3 \sim O_6$ operators. QCD penguin annihilations include s in Wilson coefficient, so they are the same order of all contributions except for O_1, O_2 annihilations.

comparatively large and get main origins for isospin breaking effects.

$$\begin{aligned}
 M_4^{S(a)} &= M_4^{P(a)} \\
 &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_q f_K r_K dx_1 \int_0^1 b_1 db_1 \\
 &\quad a_4(t_a^a) S_t(x_1) e^{[S_B(t_a^a)]} \int_0^1 dx_2 \int_0^1 b_2 db_2 \\
 &\quad (t_a^a = \max(A_a; 1-b_1)) \quad (55)
 \end{aligned}$$

$$\begin{aligned}
 M_4^{S(b)} &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_s f_B r_K dx_2 \int_0^1 b_2 db_2 \\
 &\quad a_4(t_a^b) S_t(x_2) e^{[S_K(t_a^b)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^b = \max(B_a; 1-b_2)) \quad (56)
 \end{aligned}$$

$$\begin{aligned}
 M_4^{P(b)} &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_s f_B r_K dx_2 \int_0^1 b_2 db_2 \\
 &\quad a_4(t_a^b) S_t(x_2) e^{[S_K(t_a^b)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^b = \max(B_a; 1-b_2)) \quad (57)
 \end{aligned}$$

$$\begin{aligned}
 M_4^{S(c)} &= M_4^{P(c)} \\
 &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_q f_K r_K dx_1 \int_0^1 b_1 db_1 \\
 &\quad a_4(t_a^c) S_t(x_1) e^{[S_B(t_a^c)]} \int_0^1 dx_2 \int_0^1 b_2 db_2 \\
 &\quad (t_a^c = \max(C_a; 1-b_1)) \quad (58)
 \end{aligned}$$

$$\begin{aligned}
 M_4^{S(d)} &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_q f_B r_K dx_2 \int_0^1 b_2 db_2 \\
 &\quad a_4(t_a^d) S_t(x_2) e^{[S_K(t_a^d)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^d = \max(D_a; 1-b_2)) \quad (59)
 \end{aligned}$$

$$\begin{aligned}
 M_4^{P(d)} &= F^{(0)} \int_0^1 \frac{3}{4M_B^2} \bar{Q}_q f_B r_K dx_2 \int_0^1 b_2 db_2 \\
 &\quad a_4(t_a^d) S_t(x_2) e^{[S_K(t_a^d)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^d = \max(D_a; 1-b_2)) \quad (60)
 \end{aligned}$$

$$\begin{aligned}
 M_6^{S(b)} &= M_6^{P(b)} \\
 &= F^{(0)} \int_0^1 \frac{3}{2M_B^2} \bar{Q}_s f_B r_K dx_2 \int_0^1 b_2 db_2 a_6(t_a^b) \\
 &\quad S_t(x_2) e^{[S_K(t_a^b)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^b = \max(B_a; 1-b_2)) \quad (61)
 \end{aligned}$$

$$\begin{aligned}
 M_6^{S(d)} &= M_6^{P(d)} \\
 &= F^{(0)} \int_0^1 \frac{3}{2M_B^2} \bar{Q}_q f_B r_K dx_2 \int_0^1 b_2 db_2 a_6(t_a^d) \\
 &\quad S_t(x_2) e^{[S_K(t_a^d)]} \int_0^1 dx_1 \int_0^1 b_1 db_1 \\
 &\quad (t_a^d = \max(D_a; 1-b_2)) \quad (62)
 \end{aligned}$$

$$\begin{aligned}
 A_a^2 &= (1+x_1)M_B^2; & B_a^2 &= (1-x_2)M_B^2; \\
 C_a^2 &= x_1M_B^2; & D_a^2 &= x_2M_B^2 \quad (63)
 \end{aligned}$$

V. LONG DISTANCE CONTRIBUTIONS TO THE PHOTON QUARK COUPLING




FIG. 12: Vector-Meson-Dominance contributions mediated by ψ, ρ, ω .

Here we want to discuss the long distance contributions. In order to examine the standard model or search for new physics indirectly by comparing the experimental data with the values predicted within the standard model, we have to take into account these long distance effects: $B \rightarrow K$ ($J=0; 1; 2$) $\rightarrow K$ [17, 18] (Fig.12). It should be noted that $B \rightarrow D D K \rightarrow K$ is small compared to the $J=0$ intermediate state contribution.

These contributions are caused by O_1, O_2 operators, and the effective Hamiltonian describing these processes is

$$H_{\text{eff}} = \frac{G_F}{\sqrt{2}} \sum_{q=u;c} V_{qb} V_{qs}^* (C_1(t) O_1^{(q)}(t) + C_2(t) O_2^{(q)}(t)) + \text{h.c.} \quad (64)$$

If we use the vector-meson-dominance, the $B \rightarrow K$ decay amplitude can be expressed as inserting the complete set of




FIG. 13: (A), (B) are factorizable and (C), (D) are non-factorizable contributions to the hadronic matrix element for $\langle K | \mathcal{H}_{eff} | B \rangle$.

possible intermediate vector meson states like

$$\langle K | \mathcal{H}_{eff} | B \rangle = \sum_V \langle K | J_{em} | V \rangle \frac{i}{q_V^2 - m_V^2 - i\Gamma_V} \langle V | \mathcal{H}_{eff} | B \rangle; \quad (65)$$

where $V = \rho, \omega, \phi$. Now we concretely consider the $B \rightarrow K \gamma$ decay. Four diagrams contribute to the hadronic matrix element of $\langle K | \mathcal{H}_{eff} | B \rangle$ (see Fig.13), and first of all, we consider the leading contributions: the factorizable ones, Figs. 13(A) and 13(B).

A. Factorizable contribution

In this case, the $B \rightarrow K$ decay amplitude can be factorized as

$$\langle K | \mathcal{H}_{eff} | B \rangle = \frac{G_F}{2} V_{cb} V_{cs} a_1(t) \langle K | j_c | 0 \rangle \langle 0 | j_b | B \rangle; \quad (66)$$

and the definition of the decay constant is

$$\langle 0 | j_c | K \rangle = \frac{1}{\sqrt{2}} f_K q; \quad (67)$$

then the decay amplitude can be written as

$$M(B \rightarrow K \gamma(q)) = \frac{G_F}{2} V_{cb} V_{cs} a_1(t) \frac{1}{3} f_K q^2 \langle 0 | j_b | B \rangle; \quad (68)$$

where $a_1(t) = C_1(t) + C_2(t) = 3$. The conversion part of the meson into photon can be expressed as

$$\langle 0 | J_{em} | \gamma \rangle = \frac{2}{3} e m_V g_V(q^2); \quad (69)$$

then the total amplitude of $B \rightarrow K$ mediated by meson can be expressed as follows,

$$M(B \rightarrow K \gamma) = \frac{G_F}{2} V_{cb} V_{cs} a_1(t) \frac{2e g_V(0)^2}{3} \langle 0 | j_b | B \rangle \quad (70)$$

where the real photon momentum is $q^2 = 0$. In principle, we need to include the width of the vector meson in the propagator and write

$$\frac{i}{q^2 - m^2 - i\Gamma}; \quad (71)$$

but we have $\Gamma/m \approx 0.19$ and the effects of the width can be safely neglected.

The amplitudes for $B \rightarrow K \gamma \gamma$ can be computed in a similar manner. In this case, $\Gamma/m \approx 0.19$ and we can also neglect the width effect in the meson propagator. Differences with $B \rightarrow K \gamma$ are the value of decay constant $g_V(0)$ and the factor for the electromagnetic interaction.

$$M(B \rightarrow K \gamma \gamma) = \frac{G_F}{2} V_{ub} V_{us} a_1(t) \frac{e g_V(0)^2}{6} \langle 0 | j_b | B \rangle \quad (72)$$

However in the $B \rightarrow K \gamma$ case, the resonance peak is not so sharp, so the propagation of meson generates the strong phase: $\Gamma/m \approx 0.19$ and it introduces a strong phase.

$$M(B \rightarrow K \gamma) = \frac{G_F}{2} V_{ub} V_{us} a_1(t) \frac{e g_V(0)^2}{2(1 - i\Gamma/m)} \langle 0 | j_b | B \rangle \quad (73)$$

In order to estimate these long distance contributions, we have to know the decay constant g_V . The decay constants are experimentally determined by the $V \rightarrow e^+ e^-$ data [1]. The amplitude for $V \rightarrow e^+ e^-$ can be expressed as

$$M(V \rightarrow e^+ e^-) = Q e^2 m_V g_V(q^2) \quad (74)$$

where Q expresses the electric charge like that $Q = Q_c$ when $V = \rho$, $Q = (Q_u - Q_d)/\sqrt{2}$ in $V = \omega$ case, and in $V = \phi$ case, $Q = (Q_u + Q_d)/\sqrt{2}$. Then the decay width for $V \rightarrow e^+ e^-$ decay can be written like

$$\Gamma(V \rightarrow e^+ e^-) = \frac{4 Q^2 e^4 m_V^2 g_V^2(q^2)}{3 m_V}; \quad (75)$$

and the values of g_V are in Table I.

V	$\Gamma(V \rightarrow e^+ e^-)$ (GeV)	m_V (GeV)	g_V^2 (GeV ²)
J= (1S)	5.26 $\times 10^{-6}$	3.097	0.1642
(2S)	2.19 $\times 10^{-6}$	3.686	0.0814
(3770)	0.26 $\times 10^{-6}$	3.770	0.0099
(4040)	0.75 $\times 10^{-6}$	4.040	0.0306
(4160)	0.77 $\times 10^{-6}$	4.160	0.0323
(4415)	0.47 $\times 10^{-6}$	4.415	0.0209
	7.02 $\times 10^{-6}$	0.771	0.0485
!	0.60 $\times 10^{-6}$	0.783	0.0379

TABLE I: The coefficients g_V .

Branching Ratio

In order to see how much error is generated when we change some parameters in class(1), we introduce real parameter f_i 's as the fractional differences of the amplitudes from ones with a fixed hadronic parameter, where i and j express the flavor and electric charge. Note that the uncertainty in decay constants leads to an uncertainty in overall factor of the amplitude, i.e. they don't lead to an uncertainty in the phase of the amplitude. In the change wave function parameters on the other hand, the phase changes a little, but its effect is very small and we can introduce f_i 's as real parameters.

The decay widths of the B^- and B^0 meson decays can be expressed as

$$\Gamma(B^-) = |V_{tb}V_{ts}A_t^j(1 + f_t^j) + V_{cb}V_{cs}A_c^j(1 + f_c^j) + V_{ub}V_{us}A_u^j(1 + f_u^j)|^2 \quad (87)$$

$$\Gamma(B^0) = |V_{tb}V_{ts}A_t^j(1 + f_t^j) + V_{cb}V_{cs}A_c^j(1 + f_c^j) + V_{ub}V_{us}A_u^j(1 + f_u^j)|^2; \quad (88)$$

and we can see that the uncertainty to the branching ratio from input parameters comes from the error of the $O(1)$ amplitude, and it amounts to about $2f_t^j \sim 40\%$.

Direct CP Asymmetry

From eq.(87) and (88), the direct CP asymmetry can be expressed as follows,

$$A_{CP}^0 = \frac{2}{|V_{tb}V_{ts}f_t^j A_t^j(1 + f_t^j)|} \text{Im}(V_{tb}V_{ts}V_{cb}V_{cs}) \text{Im}(A_c^j A_c^j) (1 + f_c^j) + \text{Im}(V_{tb}V_{ts}V_{ub}V_{us}) \text{Im}(A_u^j A_u^j) (1 + f_u^j) \quad (89)$$

and the error for it is

$$\frac{\Delta A_{CP}}{A_{CP}} = \frac{\Delta A_{CP}}{A_{CP}} / \frac{(f_c^j/f_t^j) \text{Im}(A_c^j/A_t^j) + (f_u^j/f_t^j) \text{Im}(A_u^j/A_t^j)}{(1 + f_t^j) [\text{Im}(A_c^j/A_t^j) + \text{Im}(A_u^j/A_t^j)]} : \quad (90)$$

We can see that the uncertainties can cancel. We have checked that numerically the class(1) error for the CP asymmetry amounts to few percent and is small compared to other errors (see below).

Isospin Breaking

On the other hand, we want to show that the hadronic parameter uncertainties especially from f_B and c dependences of the isospin breaking effect can be large even though we take the ratio as the CP asymmetry. The decay width of the neutral and charged decay modes with the theoretical error can be written from eq.(87) and (88) as

$$\Gamma^0 = |V_{tb}V_{ts}A_t^0(1 + f_t^0) + V_{cb}V_{cs}A_c^0(1 + f_c^0) + V_{ub}V_{us}A_u^0(1 + f_u^0)|^2; \quad (91)$$

$$\Gamma^+ = |V_{tb}V_{ts}A_t^+(1 + f_t^+) + V_{cb}V_{cs}A_c^+(1 + f_c^+) + V_{ub}V_{us}A_u^+(1 + f_u^+)|^2; \quad (92)$$

and the isospin breaking effect is given by

$$\frac{\Gamma^0}{\Gamma^+} = \frac{A_t^0 f_t^0 (1 + f_t^0)^2}{A_t^+ f_t^+ (1 + f_t^+)^2} ; \quad (93)$$

where we neglected all terms except for those proportional to $A_t^j f_t^j$ because $A_c^j f_c^j = A_t^j f_t^j \sim O(10^{-4})$, and the CKM factor of the $A_u^j f_u^j$ is suppressed as $V_{ub}V_{us} = V_{tb}V_{ts} f_t^j \sim O(10^{-4})$. Then the error can be expressed as

$$\left(\frac{\Delta \Gamma^0}{\Gamma^0} \right) = \frac{\Delta \Gamma^0}{\Gamma^0} = \frac{4 A_t^0 f_t^0 A_t^+ f_t^+ (f_t^0/f_t^+)}{(A_t^0 f_t^0 - A_t^+ f_t^+) (A_t^0 f_t^0 + A_t^+ f_t^+)} : \quad (94)$$

We can easily imagine that the decay constant uncertainties are canceled as the direct CP asymmetry. However we observe that even though f_t^0/f_t^+ is small, there exist $A_t^0 f_t^0$ and $A_t^+ f_t^+$ in the denominator and it is also small, then the error enhancement can occur. Variation of f_B and c introduces $f_t^0/f_t^+ \sim 0.5\%$ while $(A_t^0 f_t^0 - A_t^+ f_t^+) = (A_t^0 f_t^0 + A_t^+ f_t^+) \sim 5\%$. This gives about 20% error for the isospin breaking. From the above argument, we can see that the error from f_B and c uncertainties remain somewhat large. Thus we estimate the class(1) error for the isospin breaking effect to be about 20%.

- Next we want to discuss the class(2) error. For class(2), we expect an error coming from the fact that we used the leading order term in $s(t)$. There are also errors coming from neglecting higher order decay amplitudes. But we have not checked the effect of class(2) errors as it requires actual computation of higher order amplitudes. We guess that the error is approximately 15% in the amplitude. Then the theoretical errors from class(2) are 30% in the branching ratio, about a few % in the direct CP asymmetry, and 20% in the isospin breaking effect.
- About the class(3) error, we change the α_s parameter in the range $\alpha_s = (1.2 \pm 0.2) = 0.20 \pm 0.09$ and $\alpha_s = (1.2 \pm 0.2) = 0.33 \pm 0.05$ [1], and numerically estimate how the physical quantities are affected by the changing of parameters. The major contributions to the branching ratio and isospin breaking effects come from the terms which are proportional to $V_{tb}V_{ts}$, so they are less sensitive to the error in α_s . On the other hand, direct CP asymmetry depends on $\text{Im}(V_{tb}V_{ts}V_{cb}V_{cs})$ and $\text{Im}(V_{tb}V_{ts}V_{ub}V_{us})$ as in eq.(89), thus the error from the α_s uncertainties amounts to about 15%.

4. The class(4) error comes from the u quark loop hadronic uncertainties. The terms which are proportional to $V_{ub}V_{us}$ are not very important to the computation of the branching ratio and isospin breaking effects, so for these quantities we can neglect the class(4) uncertainties.

However for CP asymmetry, c and u quark loops give comparable contributions as seen in eq.(89), thus the u quark loop contribution which is infected with nonperturbative correction, cannot be neglected. If we regard the u quark loop uncertainty as about 100% at the amplitude level for both real and imaginary part, the numerical error for the direct CP asymmetry amounts to about 75%.

In summary, we regard the error of the branching ratio, direct CP asymmetry, and isospin breaking effects as 50% (class(1); 40%, class(2); 30%), 75% (class(4); 75%), and 30% (class(1); 20%, class(2); 20%), respectively.




FIG. 15: CKM unitary triangle

B. Numerical Results

The numerical results for each decay amplitude M_i in the neutral decay (Table II) and charged decay (Table III) in unit of 10^{-6} GeV^{-2} are as follows.

The total decay amplitude can be expressed by using these components as

$$\begin{aligned} A(B \rightarrow K) &= M_t + M_c + M_u \\ &= (M_t^S + M_c^S + M_u^S) \\ &\quad + i(M_t^P + M_c^P + M_u^P); \end{aligned} \quad (95)$$

$$M_t = M_{7\gamma} + M_{8g} + M_{3\gamma};$$

$$M_c = M_{1c} + M_{2c} + M_{\gamma};$$

$$M_u = M_{1u+2u} + M_{2\gamma} + M_{\gamma};$$

where all components include CKM factors. If we express K and helicities as $1; 2$, the combinations which can contribute to the decay amplitude are $A_{1;2} = A_{+;+}; A_{-;-}$, if we take into account the fact that B meson is spinless and a real photon has helicities ± 1 . Then the total decay width of $B \rightarrow K$ is given by

$$\Gamma = \frac{1}{8 M_B} (M_t^S + M_c^S + M_u^S)^2 + (M_t^P + M_c^P + M_u^P)^2; \quad (96)$$

and the branching ratios for $B \rightarrow K$ become as follows:

$$\text{Br}(B^0 \rightarrow K^0) = (5.8 \pm 2.9) \times 10^{-5}; \quad (97)$$

$$\text{Br}(B^+ \rightarrow K^+) = (6.0 \pm 3.0) \times 10^{-5}; \quad (98)$$

Next we want to extract the direct CP asymmetry. We take into account up to $O(\alpha_s^4)$ about the CKM matrix components,

$$\begin{aligned} V_{CKM} &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \frac{\alpha_s}{4\pi} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^2}{(4\pi)^2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^3}{(4\pi)^3} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^4}{(4\pi)^4} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \\ &= \begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix} + \frac{\alpha_s}{4\pi} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^2}{(4\pi)^2} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^3}{(4\pi)^3} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} + \frac{\alpha_s^4}{(4\pi)^4} \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} \end{aligned} \quad (99)$$

and the unitary triangle related to this decay mode should be crushed (Fig. 15).

If we express each amplitudes M_i as $M_i = V_{ib}V_{is}^* V_{cb}V_{cs}^* e^{i\phi_i}$ where $\phi_i = \arg(V_{ib}V_{is}^* V_{cb}V_{cs}^*)$, in order to separate weak phase and strong phase ϕ_i , the decay amplitudes can be rewritten as

$$A(B \rightarrow K) = V_{cb}V_{cs}^* [V_{tb}V_{ts}^* e^{i\phi_t} + V_{cb}V_{cs}^* e^{i\phi_c} + V_{ub}V_{us}^* e^{i\phi_u}]; \quad (100)$$

$$A(B \rightarrow K) = V_{cb}V_{cs}^* [V_{tb}V_{ts}^* e^{i\phi_t} + V_{cb}V_{cs}^* e^{i\phi_c} + V_{ub}V_{us}^* e^{i\phi_u}]; \quad (101)$$

and the direct CP asymmetry can be expressed as

$$A_{CP} = \frac{(B \rightarrow K) - (B \rightarrow K)}{(B \rightarrow K) + (B \rightarrow K)} \frac{R_N}{R_D}; \quad (102)$$

$$\begin{aligned} R_N &= \frac{1}{2} [A_t A_c \sin(\phi_t - \phi_c) \text{Im}(V_{tb}V_{ts}^* V_{cb}V_{cs}^*) \\ &\quad + A_c A_u \sin(\phi_c - \phi_u) \text{Im}(V_{cb}V_{cs}^* V_{ub}V_{us}^*) \\ &\quad + A_u A_t \sin(\phi_u - \phi_t) \text{Im}(V_{ub}V_{us}^* V_{tb}V_{ts}^*)]; \end{aligned} \quad (103)$$

$$\begin{aligned} R_D &= A_t^2 |V_{tb}V_{ts}^*|^2 + A_c^2 |V_{cb}V_{cs}^*|^2 + A_u^2 |V_{ub}V_{us}^*|^2 \\ &\quad + A_t A_c \cos(\phi_t - \phi_c) \text{Re}(V_{tb}V_{ts}^* V_{cb}V_{cs}^*) \\ &\quad + A_c A_u \cos(\phi_c - \phi_u) \text{Re}(V_{cb}V_{cs}^* V_{ub}V_{us}^*) \\ &\quad + A_u A_t \cos(\phi_u - \phi_t) \text{Re}(V_{ub}V_{us}^* V_{tb}V_{ts}^*); \end{aligned} \quad (104)$$

then its' values are

$$A_{CP}(B^0 \rightarrow K^0) = (6.1 \pm 4.6) \times 10^{-3}; \quad (105)$$

$$A_{CP}(B^+ \rightarrow K^+) = (5.7 \pm 4.3) \times 10^{-3}; \quad (106)$$

Finally, we want to estimate the isospin breaking effects as eq.(2). This effect is caused by O_8 (Fig. 4), c and u loop contributions (Fig. 5), $O_1 - O_6$ annihilation (Figs. 10 and 11), and the long distance contributions mediated by photon in charged mode (Fig. 14). About the bremsstrahlung photon contributions emitted through quark lines, whether the spectator quark is u or d affects the strength and the sign for the coupling of photon and quark line, so they generate the isospin breaking effects. The most important contributions to the isospin breaking effects come from QCD penguin O_5, O_6 annihilation. These effects are additive to the dominant contribution O_7 in both neutral and charged decays (see Tables II and III). However its' size are different: the neutral mode's is larger than charged mode's. Then the sign of total isospin breaking effects becomes plus and its' value is as follows.

$$\delta_{0+} = (2.7 \pm 0.8) \times 10^{-2} \quad (107)$$

	$M_i^S = F^{(0)}$			$M_i^P = F^{(0)}$		
$V_{tb}V_{ts}$	$M_7^S = F^{(0)}$	$M_{8g}^S = F^{(0)}$	$M_{36}^S = F^{(0)}$	$M_7^P = F^{(0)}$	$M_{8g}^P = F^{(0)}$	$M_{36}^P = F^{(0)}$
	-218.67-3.86i	-2.19-0.55i	-11.56-5.70i	218.67+ 3.86i	2.27+ 0.59i	11.58+ 5.63i
$V_{cb}V_{cs}$	$M_{1c}^S = F^{(0)}$	$M_{2c}^S = F^{(0)}$	$M^S = F^{(0)}$	$M_{1c}^P = F^{(0)}$	$M_{2c}^P = F^{(0)}$	$M^P = F^{(0)}$
	-0.29-1.01i	6.42-12.63i	-13.29	-0.19+ 1.27i	-4.81+ 8.23i	15.09
$V_{ub}V_{us}$	$M_{1u+2u}^S = F^{(0)}$	$M_2^S = F^{(0)}$	$M_{+!}^S = F^{(0)}$	$M_{1u+2u}^P = F^{(0)}$	$M_2^P = F^{(0)}$	$M_{+!}^P = F^{(0)}$
	-0.63+ 0.22i	0	-0.03-0.06i	0.67-0.18i	0	0.03+ 0.07i

TABLE II: $B^0 \rightarrow K^0$ at $\alpha_s = 0.20$, $\alpha_s = 0.33$, $\mu_B = 0.40\text{ GeV}$.

	$M_i^S = F^{(0)}$			$M_i^P = F^{(0)}$		
$V_{tb}V_{ts}$	$M_7^S = F^{(0)}$	$M_{8g}^S = F^{(0)}$	$M_{36}^S = F^{(0)}$	$M_7^P = F^{(0)}$	$M_{8g}^P = F^{(0)}$	$M_{36}^P = F^{(0)}$
	-218.67-3.86i	-4.89-0.10i	-2.47+ 0.37i	218.67+ 3.86i	4.83-0.82i	2.86+ 0.14i
$V_{cb}V_{cs}$	$M_{1c}^S = F^{(0)}$	$M_{2c}^S = F^{(0)}$	$M^S = F^{(0)}$	$M_{1c}^P = F^{(0)}$	$M_{2c}^P = F^{(0)}$	$M^P = F^{(0)}$
	-0.66+ 2.15i	6.42-12.63i	-13.29	1.39-2.60i	-4.81+ 8.23i	15.09
$V_{ub}V_{us}$	$M_{1u+2u}^S = F^{(0)}$	$M_2^S = F^{(0)}$	$M_{+!}^S = F^{(0)}$	$M_{1u+2u}^P = F^{(0)}$	$M_2^P = F^{(0)}$	$M_{+!}^P = F^{(0)}$
	-0.75+ 0.51i	0.35+ 1.01i	-0.04+ 0.05i	0.79-0.18i	-0.75+ 1.16i	0.05-0.05i

TABLE III: $B^+ \rightarrow K^+$ at $\alpha_s = 0.20$, $\alpha_s = 0.33$, $\mu_B = 0.40\text{ GeV}$.

VII. CONCLUSION

In this paper, we calculated the branching ratio, direct CP asymmetry, and isospin breaking effect within the standard model using the pQCD approach. It is useful to compare our results with those existing in the literature. The decay amplitude can be obtained from the transition form factor

$$\langle K(P_2; K) | j_i q s | B(P_1) \rangle = i T_1^K(0) \epsilon_{K P q} \quad (108)$$

where $P = P_1 + P_2$, $q = P_1 - P_2$. Within the framework of pQCD, we obtain the value of the $B \rightarrow K$ transition form factor as $T_1^K(0) = 0.23 \pm 0.06$. The result can be compared with the ones extracted by another estimation. In the QCD Factorization, $T_1^K(0) = 0.27 \pm 0.04$ [25] and an updated phenomenological estimate of this quantity with the light-cone distribution amplitudes for the K meson is $T_1^K(0) = 0.27 \pm 0.02$ [26]. While the central value in the updated result is the same as before, the error is reduced by a factor of 2. In the light-cone QCD sum rule $T_1^K(0) = 0.38 \pm 0.06$ [27], the lattice QCD simulation $T_1^K(0) = 0.32^{+0.04}_{-0.02}$ [28] and $T_1^K(0) = 0.25^{+0.05}_{-0.02}$ [29], and the covariant light-front approach $T_1^K(0) = 0.24$ [30]. There are several estimates of the branching ratio by using the value of $T_1^K(0) = 0.38 \pm 0.06$ extracted from light-cone QCD sum rule. Comparing the results with experiments, this value of the form factor over estimates the branching ratios [31, 32, 33]. Also, it should be noted that $T_1^K(q^2)$ and other related form factors have been computed in the framework of pQCD [34]. They obtained the central value as $T_1^K(0) = 0.315$. The difference between our results and theirs is the K meson wave function. We take the new K wave function parameters computed in Ref.[12].

Note that we have also included the long distance contributions. If we neglect them, the branching ratios become $\text{Br}(B^0 \rightarrow K^0) = (5.2 \pm 2.6) \times 10^{-5}$ and

$\text{Br}(B^+ \rightarrow K^+) = (5.3 \pm 2.7) \times 10^{-5}$ to be pared with results shown in Eq.(97) and (98). The $B \rightarrow K \rightarrow K$ contribution to the total decay width amounts to about 12% and also it works additive to the branching ratios. We also emphasize that we can calculate the annihilation contributions with the pQCD approach, and these contribute to the total decay width which amount to about $2 \pm 10\%$.

This analysis predicts less than 1% direct CP asymmetry within the standard model. If we neglect the long distance contributions, the asymmetries become $A_{CP}(B^0 \rightarrow K^0) = (6.7 \pm 5.0) \times 10^{-3}$ and $A_{CP}(B^+ \rightarrow K^+) = (7.2 \pm 5.4) \times 10^{-3}$, and as to isospin breaking effect like $\alpha_{+} = + (2.6 \pm 0.3) \times 10^{-2}$. The long distance contributions do not seem to affect to these asymmetries.

The branching ratio of the neutral decay is similar to that of the charged decay, in spite of the difference of the lifetime between them. This effect is mainly caused by the 4-quark penguin operators O_5, O_6 . If we neglect these contributions, the isospin breaking is $\alpha_{+} = (1.2 \pm 0.4) \times 10^{-2}$, so we can see that they generate about 4% isospin breaking effect. This result is similar to the conclusion of Ref.[35].

$B \rightarrow K$ decay, as we have first mentioned, is an attractive decay mode to test the standard model and search for new physics. In order to look for the new physics, we have to reduce the experimental errors. The error to the direct CP asymmetry must get smaller than 1%. That is to say, we need at least 20 times more data. This is not possible without the super B factory.

Acknowledgments

We acknowledge useful discussion with the pQCD group members, especially Hsiang-nan Li. AIS acknowledges support from the Japan Society for the Promotion of Science, Japan-US collaboration program, and a grant from Ministry




FIG. 16: $O(\alpha_s)$ corrections to the hard scattering H .

of Education, Culture, Sports, Science and Technology of Japan.

APPENDIX A: BRIEF REVIEW OF PQCD

1. Divergences in perturbative diagrams

Here we want to review the k_T factorization [36]. At a higher order, in addition to many gluon exchanges must be considered. In order to understand the factorization procedure, we refer to the diagrams of Fig. 16.

They describe the $O(\alpha_s)$ radiative corrections to the hard scattering process H . In general, individual higher order diagrams have two types of infrared divergences: soft and collinear. Soft divergence comes from the region of a loop momentum where all its momentum components in the light-cone coordinate vanish:

$$l = (l^+, l_\perp, l_-) = (0; 0; 0): \quad (A1)$$

Collinear divergence originates from the gluon momentum region which is parallel to the massless quark momentum,

$$l = (l^+, l_\perp, l_-) = (M_B; 0; 0): \quad (A2)$$

In both cases, the loop integration corresponds to $\int d^4l = \int d^2l_\perp \int dl^+ \int dl_-$, so logarithmic divergences are generated. It has been shown order by order in perturbation theory that these divergences can be separated from the hard kernel and absorbed into meson wave functions using eikonal approximation [37].

Furthermore, there are also double logarithm divergences in Fig. 16(a) and 16(b) when soft and collinear momentum overlap. These large double logarithms can be summed by using renormalization group equation. This factor is called the Sudakov factor and also factorized into the definition of meson wave function [38, 39, 40]. The explicit expression for Sudakov factor is given by [39] (see Appendix B).

There are also ultraviolet divergences, and also another type of double logarithm which comes from the loop correction for the weak decay vertex correction. These double logarithms can also be factored out from the hard part and grouped into the quark jet function. These double logarithms also




FIG. 17: An electron which is scattered by the electromagnetic interaction (a) is observed with many soft photons. Similarly, a quark which is scattered by the strong interaction (b) is not observed as a single gluon: accompanied by many soft gluons, and they form hadron jets.

should be resummed as the threshold factor [41, 42]. This factor decreases faster than any other power of x as $x \rightarrow 0$, so it removes the endpoint singularity. Thus we can factor out the Sudakov factor, the threshold factor, and the ultraviolet divergences from the hard part and group them into the meson wave function (Appendix B). Then the redefinition of wave functions including these loop corrections get factorization energy scale dependence t .

Thus the amplitude can be factorized into a perturbative part including a hard gluon exchange, and a nonperturbative part characterized by the meson distribution amplitudes. Then the total decay amplitude can be expressed as the convolution:

$$\int_0^1 dx_1 dx_2 \int_0^1 d^2b_1 d^2b_2 C(t) \int_K (x_2; b_2; t) H(x_1; x_2; b_1; b_2; t) \int_B (x_1; b_1; t); \quad (A3)$$

here $\int_K (x_2; b_2; t)$, $\int_B (x_1; b_1; t)$ are meson distribution amplitudes that contain the soft divergences which come from quantum correction and $H(x_1; x_2; b_1; b_2; t)$ is the hard kernel including a finite piece of quantum correction, where b_1, b_2 are the conjugate variables to transverse momentum, and x_1, x_2 are the momentum fractions of spectator quarks.

2. Physical interpretation of Sudakov factor

In order to understand the Sudakov factor physically, first we consider QED. When a charged particle is accelerated, infinitely many photons must be emitted by the bremsstrahlung (Fig. 17(a)). A similar phenomenon occurs when a quark is accelerated: infinitely many gluons must be emitted. According to the feature of strong interaction, gluons cannot exist freely, so hadronic jets are produced. Then we observe many hadrons in the end if gluonic bremsstrahlung occurs. Thus the amplitude for an exclusive decay $B \rightarrow K$ is proportional to the probability that no bremsstrahlung gluon is emitted. This is the Sudakov factor and it is depicted in Fig. 19. As seen in Fig. 19, the Sudakov factor is large for small b and Q . Large b implies that the quark and antiquark pair is separated, which

-
- [1] S. Eidelman et al. [Particle Data Group], Phys. Lett. B 592 (2004) 1.
- [2] K. Abe et al. [BELLE Collaboration], arXiv:hep-ex/0409049.
- [3] B. Aubert et al. [BABAR Collaboration], arXiv:hep-ex/0408072.
- [4] M. Nakao et al. [BELLE Collaboration], Phys. Rev. D 69, 112001 (2004) [arXiv:hep-ex/0402042].
- [5] B. Aubert et al. [BABAR Collaboration], Phys. Rev. D 70, 112006 (2004) [arXiv:hep-ex/0407003].
- [6] J. Alexander et al. [Heavy Flavor Averaging Group (HFAG) Collaboration], arXiv:hep-ex/0412073.
- [7] A. J. Buras, A. Czamecki, M. Misiak and J. Urban, Nucl. Phys. B 631, 219 (2002) [arXiv:hep-ph/0203135].
- [8] G. Buchalla, A. J. Buras and M. E. Lautenbacher, Rev. Mod. Phys. 68, 1125 (1996) [arXiv:hep-ph/9512380].
- [9] M. Bauer and M. Wirbel, Z. Phys. C 42, 671 (1989).
- [10] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Lett. B 504, 6 (2001) [arXiv:hep-ph/0004004].
- [11] P. Ball, V. M. Braun, Y. Koike and K. Tanaka, Nucl. Phys. B 529, 323 (1998) [arXiv:hep-ph/9802299].
- [12] P. Ball and M. Boglione, Phys. Rev. D 68, 094006 (2003) [arXiv:hep-ph/0307337].
- [13] M. Bander, D. Silverman and A. Soni, Phys. Rev. Lett. 43, 242 (1979).
- [14] J. Liu and Y. P. Yao, Phys. Rev. D 42, 1485 (1990).
- [15] H. Simma and D. Wyler, Nucl. Phys. B 344, 283 (1990).
- [16] H. N. Li and G. L. Lin, Phys. Rev. D 60, 054001 (1999) [arXiv:hep-ph/9812508].
- [17] E. Golwisch and S. Pakvasa, Phys. Rev. D 51, 1215 (1995) [arXiv:hep-ph/9408370].
- [18] N. G. Deshpande, X. G. He and J. Trampetic, Phys. Lett. B 367, 362 (1996).
- [19] R. L. Anderson et al., Phys. Rev. Lett. 38, 263 (1977).
- [20] E. Paul, BONN-HE-81-26 Invited talk given at 10th Int. Symp. on Lepton and Photon Interactions at High Energy, Bonn, West Germany, Aug 24-29, 1981
- [21] K. Abe et al. [Belle Collaboration], Phys. Lett. B 538, 11 (2002) [arXiv:hep-ex/0205021].
- [22] B. Aubert et al. [BABAR Collaboration], Phys. Rev. Lett. 87, 241801 (2001) [arXiv:hep-ex/0107049].
- [23] M. Verderi, Workshop on the Unitarity Triangle in 2005, WG5 Session2, San Diego, 2005 (to be published)
- [24] A. L. Kagan, Phys. Lett. B 601, 151 (2004) [arXiv:hep-ph/0405134].
- [25] M. Beneke, T. Feldmann and D. Seidel, Nucl. Phys. B 612, 25 (2001) [arXiv:hep-ph/0106067].
- [26] A. Ali, E. Lunghi and A. Y. Parkhomenko, Phys. Lett. B 595, 323 (2004) [arXiv:hep-ph/0405075].
- [27] P. Ball and V. M. Braun, Phys. Rev. D 58, 094016 (1998) [arXiv:hep-ph/9805422].
- [28] L. Del Debbio, J. M. Flynn, L. Lellouch and J. Nieves [UKQCD Collaboration], Phys. Lett. B 416, 392 (1998) [arXiv:hep-lat/9708008].
- [29] D. Becirevic, talk presented at the Flavor Physics & CP Violation conference, Ecole Polytechnique, Paris France, June 2003
- [30] H. Y. Cheng and C. K. Chua, Phys. Rev. D 69, 094007 (2004) [arXiv:hep-ph/0401141].
- [31] A. Ali and A. Y. Parkhomenko, Eur. Phys. J. C 23, 89 (2002) [arXiv:hep-ph/0105302].
- [32] S. W. Bosch, arXiv:hep-ph/0208203.
- [33] S. W. Bosch and G. Buchalla, Nucl. Phys. B 621, 459 (2002) [arXiv:hep-ph/0106081].
- [34] C. H. Chen and C. Q. Geng, Nucl. Phys. B 636, 338 (2002) [arXiv:hep-ph/0203003].
- [35] A. L. Kagan and M. Neubert, Phys. Lett. B 539, 227 (2002) [arXiv:hep-ph/0110078].
- [36] M. Nagashima and H. N. Li, Phys. Rev. D 67, 034001 (2003) [arXiv:hep-ph/0210173].
- [37] H. N. Li and H. L. Yu, Phys. Rev. D 53, 2480 (1996) [arXiv:hep-ph/9411308].
- [38] J. C. Collins and D. E. Soper, Nucl. Phys. B 193, 381 (1981).
- [39] J. Botts and G. Sterman, Nucl. Phys. B 325, 62 (1989).
- [40] H. N. Li and G. Sterman, Nucl. Phys. B 381, 129 (1992).
- [41] H. N. Li, Phys. Rev. D 66, 094010 (2002) [arXiv:hep-ph/0102013].
- [42] T. Kurimoto, H. N. Li and A. I. Sanda, Phys. Rev. D 65, 014007 (2002) [arXiv:hep-ph/0105003].
- [43] J. C. Collins and G. Sterman, Nucl. Phys. B 185 (1981) 172.
- [44] Y. Y. Keum, H. N. Li and A. I. Sanda, Phys. Rev. D 63 (2001) 054008 [arXiv:hep-ph/0004173].