N eutrinos as a D iagnostic of H igh Energy A strophysical P rocesses

Luis A. Anchordoqui,¹ Haim Goldberg,¹ Francis Halzen,² and Thomas J. Weiler³

¹Department of Physics, Northeastern University, Boston, MA 02115

²Department of Physics, University of W isconsin, M adison W I 53706

 ^3D epartm ent of P hysics and A stronom y, V anderbilt U niversity, N ashville TN 37235

A leading candidate for the extragalactic source of high energy cosm ic rays is the Ferm i engine m echanism, in which protons con ned by m agnetic elds are accelerated to very high energy through repeated scattering by plasm a shock fronts. In the process of acceleration, collisions of trapped protons with the am bient plasm a produce pions which decay to electrom agnetic energy and neutrinos. For optically thin sources, a strong connection between the em erging cosm ic rays and secondary neutrinos can be established. In this context, we show the feasibility of using the G lashow resonance as a discriminator between the pp and p interactions in Ferm i engines as sources of neutrinos. In particular, we demonstrate how three years of observation at the km³ LeeC ube facility can serve as a liter for the dom inance of the pp interaction at the source.

Neutrinos can serve as unique astronom ical messengers. Except for oscillations induced by transit in a vacuum higgs eld, neutrinos propagate without interactions between source and Earth, providing powerful probes of high energy astrophysics [1]. The deployment of the km³ IceC ube facility at the South Pole [2] will greatly increase the statistics required for the realization of such a program.

The avor composition of neutrinos originating at astrophysical sources can also serve as a probe of new physics in the electroweak sector [3]. Speci cally, Ice-Cube will have the capability to clearly identify neutrino species [4], and consequently will be able to measure deviations of avor composition from standard expectations. In addition, in resonant scattering $_{e}e$! W ! anything [5], the detector can simultaneously discrim inate between $_{e}$ and $_{e}$: In this Letter, we show that the signal for $_{e}$ at the G lashow resonance, when norm alized to the total + — ux, can be used to di erentiate between the two primary candidates (p and pp collisions) for neutrino-producing interactions in optically thin sources of cosm ic rays.

It is helpful to envision the cosm ic ray engines as machines where protons are accelerated and (possibly) perm anently con ned by the magnetic elds of the acceleration region. The production of neutrons and charged pions and subsequent decay produces both neutrinos and $\cos m$ ic rays: the form er via $^+$! e^+ $_e$ — (and the conjugate process { m ore on this below), the latter via neutron di usion from the region of the con ned protons. If the neutrino-em itting source also produces high energy cosm ic rays, then pion production must be the principal agent for the high energy cuto on the proton spectrum [6, 7]. Conversely, since the protons must undergo su cient acceleration, inelastic pion production needs to be sm all below the cuto energy; consequently, the plasm a must be optically thin. Since the interaction time for protons is greatly increased over that of neutrons because of magnetic con nem ent, the neutrons escape before interacting, and on decay give rise to the observed cosm ic ray ux. A desirable property of this low -dam ping scenario is that a single source will produce

cosm ic rays with a sm ooth spectrum across a wide range of energy.

For optically thin sources, the neutrino power density scales linearly with the cosm ic ray power density $_{-e_R}$ [8]. The actual value of the neutrino ux depends on what fraction of the proton energy is converted to pions (which then decay to neutrinos and photons). To quantify this, here we de ne as the ratio of pion energy to the em erging nucleon energy at the source. Since there is about one cosm ic ray (neutron) produced per proton collision, a significant consequence of the neutron leakage m odel is that

is expected to be approximately equal to the fraction of collision energy carried o by pions in a single collision. Following W axm an and Bahcall [8], the expected total (all species) neutrino ux is

$$E^{2}$$
 total 2 10⁸ , GeV cm 2 s 1 sr 1 ; (1)

where for no source evolution $_z$ 0:6; whereas for an evolution / $(1 + z)^3$ as seen in the star-form ation rate, $_z$ 3 \mathfrak{P}].

Depending on the relative ambient gas and photon densities, pion production proceeds either through inelastic pp scattering [10], or photopion production predom inantly through the resonant process $p ! ! ! ! n + orp^{0}$ [8]. For the rst of these, pion production is well-characterized by a central rapidity (Feynman) plateau, yielding 0.6 [1]. For resonant photoproduction, the inelasticity is kinem atically determ ined by requiring equal boosts for the decay products of the ⁺ [12], giving 0.25 [13].

The rst production mechanism (described above) involves the dominance of inelastic pp collisions in generating charged pions. The nearly isotopically neutral mix of pions will create on decay a neutrino population in the ratio N = N- = 2N = 2N-: In propagation to Earth, a distance longer than all oscillation lengths, avor-changing amplitudes are replaced by probabilities [14], resulting in equal uxes for all six neutrino species, so that N $\frac{E}{e}$ arth = $\frac{1}{6}$ N $\frac{total}{+}$: Here, we have incorporated maximal \$ mixing and the known smallness of $j_h e j_3 i j_r^2$, where $_3$ ' (+)= $\frac{1}{2}$ is the third neutrino eigenstate.

The second mechanism proceeds via photoproduction of pions by trapped protons on the thermal photon background, leaving the isotopically asymmetric process p ! + ! + n as the dominant source of neutrinos. At production, $N = N - = N_e = N_e$. After oscillation, the e ux at Earth is

$$N \frac{E_{e}^{\text{Earth}}}{=} = N - P (-! - e)$$
$$= \frac{1}{3} \sin^{2} \cos^{2} N \frac{\text{total}}{+} : (2)$$

U sing the most recent SNO value of the solar mixing angle ' 32:5 [15], we obtain $N \frac{E}{e}^{arth} = \frac{1}{15} N \frac{total}{+}$: Because of averaging over various therm all environments, this relation holds across the neutrino spectrum [8].

The preceding discussion indicates the feasibility of di erentiating between the two processes as follows: (i) obtain a norm alization for the total ux from data outside the G lashow resonance region; (ii) m easure the event rate in the resonance region, and com pare with the ratios given above. In what follows, we will assess the potential of LoeC ube to obtain a statistically signi cant signal-tobackground ratio.

The cross section for G lashow resonant scattering reads

$$= \frac{1}{2m_{e}E} M f (2m_{e}E m_{W}^{2}); \qquad (3)$$

where $M = g^2 m_W^2 = 2$; m_e and m_W are the electron and W m asses, and $g^2 = 4$ (m_W) = $\sin^2 w$ (m_W) ' 0:43: Now, for an incoming ux of antineutrinos \mathcal{F}_e (E) the event rate (assuming 100% detection e ciency) reads

$$\frac{dN}{dt} = N_{e} \qquad dE \quad J_{-e} (E) (E)$$

$$= N_{e} \qquad \frac{g^{2}}{4m_{e}} J_{-e} \quad \frac{m_{W}^{2}}{2m_{e}} ; \qquad (4)$$

where N_e ' 6 10^{38} is the number of target electrons in the e ective volume V 2 km³ of the IœC ube experiment, and ' 2 is the solid angle aperture [16]. If neutrinos are produced via pp inelastic collisions, according to the previous discussion (with = 0:6 and $_z = 3$), the antineutrino ux on Earth is $E^2 J_{-e} (E) =$ 6 $10^9 \text{ GeV cm}^2 \text{ s}^1 \text{ sr}^1$, so that the event rate is dN =dt = 4:6 yr¹: For antineutrinos produced in photopion interactions (= 0:25), the expected rate is a factor of 6 sm aller, dN =dt = 0:8 yr¹:

C an the expected resonant signal be di erentiated from the continuum background? In order to estimate the background, we note that all but the muon decays of the W devolve into an electrom agnetic shower in the detector. For the hadronic decays (which constitute 70% of the total), the interaction mean free path of charged pions in ice is orders of magnitude smaller than the pion decay length at TeV energies, allowing a progressive channeling of nearly all the energy into electrom agnetic modes through 0 decay. For the e $-_{e}$ and - decay modes,

there will be a single electrom agnetic burst triggered by the electron interaction or by the decay, each how ever m an ifesting only half the energy. As a conservative procedure, we will use in our signal estim ate only the hadronic events with full energy in the resonant band, giving (in the case of pp interactions) a signal dN = dt = 32 yr¹: The background will then constitute the non-resonant scattering of e and e within the resonant band. This is calculated as an integral of the ux times the charged current cross section [17] over the resonant acceptance bin $(10^{6:7} \text{ GeV}; 10^{6:9} \text{ GeV})$ of the AMANDA detector [18]. The process being o -resonance, the Earth is transparent and so we take = 4 . The num ber of target nucleons is twice the number of electrons. This procedure yields a background of 0.6 events yr 1 . Thus, within 3 years ofdata accumulation (1/5 of the total lifetim e of the experim ent), we expect about 2 background events, with a signal of 9.6 events, well above the 99% CL for discovery (corresponding to 6.69 events [19]).

E lectron antineutrinos can also be produced through neutron -decay. This contribution turns out to be negligible. To obtain an estim ate, we sum over the neutron-em itting sources out to the edge of the universe at a distance $1=H_0$ [20]

$$-_{e} = \frac{m_{n}}{8_{0}H_{0}} \frac{Z_{E_{n}^{max}}}{\frac{m_{n}E_{-}}{2_{0}}} \frac{dE_{n}}{E_{n}} F_{n} (E_{n}) ; \qquad (5)$$

where F_n (E_n) is the neutron volum e em issivity and m_n the neutron m ass. Here, we have assumed that the neutrino is produced monoenergetically in the neutron rest frame, i.e., $_0 m_N (1 m_e^2 = {}^2m_N) = 2 0.55 \text{ MeV}$, where m_N ' 1:30 MeV is the neutron proton m ass difference. An upper limit can be placed on F_n by assum - ing an extrem e case in which all the cosm ic rays escaping the source are neutrons, i.e., $_{e_R} = dE_n E_n F_n (E_n)$: W ith the production rate of ultrahigh energy protons $_{e_R}^{(10^{10};10^{12}]}$ 5 10⁴ erg M pc ³ yr ¹ [21], and an assumed injection spectrum F_n / E_n^2 ; we obtain

$$E^{2} - 3 10^{11} \text{ GeV cm}^{2} \text{ s}^{1} \text{ sr}^{1}$$
; (6)

about 3 orders of m agnitude sm aller than the charge pion contribution with = 0.6: (N ote that oscillations will reduce the $-_{e}$ ux on E arth by 40% from this value.)

Finally, we note that in the case that pions are produced via p ! * ! N , the event rate for the G lashow resonance cannot be separated from back-ground. Thus, the presence or absence of a signal at the G lashow resonance can be used as a liter for pp-dom inance at the source.

In summary, we have shown how several years of data collection at IceC ube can potentially isolate the dominance of inelastic pp interactions as neutrino progenitors in cosm ic ray sources. The analysis depends on the sources being optically thin, which can be ascertained if the observed di use neutrino ux is in agreement with Eq. (1). Then a signal at the G lashow resonance will provide a strong m arker for pp-dom inance.

<u>Note added</u>: The previous version of this paper contained a brief discussion of how the ux of neutrinos from optically thin sources can be used as a marker of the cosmic ray Galactic/extra-galactic transition. This topic has received detailed discussion in a separate paper [22].

- [1] T.K.Gaisser, F.Halzen and T.Stanev, Phys.Rept.258, 173 (1995) [Erratum -ibid. 271, 355 (1996)]; F.Halzen and D.Hooper, Rept. Prog. Phys. 65, 1025 (2002)
- [2] F.Halzen, astro-ph/0311004.
- [3] See e.g., J.F. Beacom, N.F. Bell, D.Hooper, S.Pakvasa and T.J.W eiler, Phys. Rev. Lett. 90, 181301 (2003); Phys.Rev.D 69,017303 (2004); J.F.Beacom, N.F.Bell, D.Hooper, J.G. Learned, S.Pakvasa and T.J.W eiler, Phys.Rev.Lett.92,011101 (2004); H.Athar, M.Jezabek and O.Yasuda, Phys.Rev.D 62, 103007 (2000).
- [4] J. F. Beacom, N. F. Bell, D. Hooper, S. Pakvasa and T. J.W eiler, Phys. Rev. D 68, 093005 (2003).
- [5] S.L.G lashow, Phys. Rev. 118, 316 (1960).
- [6] P. L. Bierm ann and P. A. Strittm atter, Astrophys. J. 322, 643 (1987).
- [7] For proton energies $< 10^{11}~{\rm G\,eV}$, the H illas criterion is easily satis ed.
- [8] E.W axm an and J.N.Bahcall, Phys. Rev.D 59, 023002 (1999).
- [9] The assumption of optically thin sources also guarantees that the muon decays well before it interacts. Note that the condition on neutron leakage requires that the neutron interaction length is larger than the accelerating region. Since the muon interaction length (mostly from pair production and brem sstrahlung) is in turn larger than that of the neutron, the muon will decay without losing energy, either inside or outside the source, giving rise to secondary neutrinos, each with 1/4 of the pion energy.
- [10] See e.g., L.A. Anchordoqui, H.G oldberg, F.Halzen and T.J.W eiler, Phys. Lett. B 600, 202 (2004).
- [11] J. A lvarez-M uniz, R. Engel, T. K. Gaisser, J. A. Ortiz

This work has been partially supported by US NSF (Grant Nos. OPP-0236449, PHY-0140407, PHY-0244507), US DOE (Grant Nos. DE-FG 02-95ER 40896, DE-FG 05-85ER 40226), NASA-ATP 02-000-0151, and the W isconsin A lum ni Research Foundation.

- and T.Stanev, Phys.Rev.D 66,033011 (2002).
- [12] F.W. Stecker, Phys. Rev. Lett. 21, 1016 (1968).
- [13] We note that for the case of pion production in inelastic pp collisions, the average fraction of the total pion energy carried by charged pions is 2/3, com pared to 1/2 in the photopion case discussed in [8]. For the pp case, this will enhance the neutrino ux by 33%; however, for the sake of sim plicity we will om it this consideration.
- [14] S. Eidelm an et al. Particle Data G roup Collaboration], Phys.Lett.B 592,1 (2004).
- [15] S.N.Ahm ed et al, Phys.Rev.Lett. 92, 181301 (2004).
- [16] Note that, o -resonance, the IceC ube detector will have nearly full solid angle acceptance at PeV energies.
- [17] R.Gandhi, C.Quigg, M.H.Reno and I.Sarœvic, Phys. Rev. D 58, 093009 (1998). The neutral current interaction cross section is a factor of 2.5 sm aller than the charged current process. Additionally, since 80% of the energy is carried o by the neutrino, the background event rate at the G lashow resonance would require prim ary neutrino energies at about 10^{7:5} GeV, where the ux is 25 tim es sm aller.
- [18] M. Ackermann et al., [AMANDA Collaboration], astro-ph/0405218.
- [19] G.J.Feldm an and R.D.Cousins, Phys. Rev. D 57, 3873 (1998).
- [20] L. A. Anchordoqui, H. Goldberg, F. Halzen and T.J.W eiler, Phys.Lett.B 593, 42 (2004).
- [21] E.W axm an, A strophys. J. 452, L1 (1995).
- [22] M. Ahlers, L. A. Anchordoqui, H. Goldberg, F. Halzen, A. Ringwald and T. J. W eiler, arX is astro-ph/0503229.