

Statistical mechanics of confined quantum particles

Vishnu M .Bannur and K .M .Udayanandan
Department of Physics,
University of Calicut, Kerala-673 635, India.

October 27, 2021

Abstract

We develop statistical mechanics and thermodynamics of Bose and Fermi systems in relativistic harmonic oscillator (RHO) confining potential, which may be applicable in quark gluon plasma (QGP), astrophysics, Bose-Einstein condensation (BEC) etc. Detailed study of QGP system is carried out and compared with lattice results. Further, as an application, our equation of state (EoS) of QGP is used to study compact stars like quark star.

PACS Nos : 05.30.-d, 12.38.Mh, 12.38.Gc, 26.60.+c, 05.70.Ce, 12.39.Ki

Keywords : Relativistic harmonic oscillator, Equation of state, quark gluon plasma, quark star

1 Introduction :

There are many systems like QGP, quark stars, BEC etc. where particles are confined by harmonic oscillator type potentials and are at thermodynamic equilibrium. So we need to study statistical mechanics of such a system. Taking QGP, a new form of matter made of quarks and gluons, as an example we discuss this problem. It is generally accepted that hadrons are confined state of quarks and gluons and at high temperature and density, we get what is called QGP [1]. In relativistic heavy ion collisions (RHICs), quark stars etc., QGP consists of quarks and gluons confined in a volume large compared to the size of hadrons. In the case of early universe, just before hadronization, again we have QGP. In all above cases, EoS of QGP is needed to explain their evolution and study various properties. Since quantum chromodynamics (QCD) is nonlinear, it is difficult to solve analytically at low energy and only at high energy one can do perturbative calculations. That is why one goes to numerical simulations of QCD on lattice and EoS of QGP without and with quark flavors has been derived [2]. Many phenomenological models like hadron thermal loop (HTL) [3], quasi-particle QGP (qQGP) [4, 5], strongly interacting QGP (sQGP) [6], strongly coupled QGP (SCQGP) [7] and various confinement models [8, 5, 9] were proposed to explain lattice results and there is no satisfactory model. Models which fit well with lattice results, generally, involve large number of parameters [4, 5] and models with detailed theory like HTL [3] don't fit lattice results well, so on. However, recently we found that SCQGP model [7] as well as a new qQGP model [10] fit surprisingly well with lattice results with minimum number of parameters compared to other models. Both models are purely phenomenological and based on the collective properties of plasma.

Here we approach the problem from the known fact that RHQ model of hadrons quite successfully explains many properties of hadrons [11]. QGP formed in RHICs and compact stars may be viewed as a large hadron with large number of quarks and gluons, confined in a RHQ confining potential. Hence, a study of statistical mechanics and thermodynamics of such a system will give EoS of QGP. It is just a generalization of widely studied Bag Model of QGP. Instead of sharp bag wall, here we have a smooth confining potential. As a first step,

here we neglect effects like Color-Coulomb interactions, hadron formations etc. and see how the EoS of QGP differs from ideal EoS. Earlier, similar calculations has been done [9], but without proper counting of degeneracy. They have taken only accidental degeneracy where as here we have taken full degeneracy arising from coalesce of continuum levels to oscillator levels as a result of RHO interaction, just like in Landau's theory of diamagnetism [12].

2 EoS of QGP in RHO Model :

From the earlier study of hadrons in RHO model, we have the single particle energy levels [11],

$$\epsilon_n^2 = C_g^2 (2n + 1); \quad (1)$$

for gluons and

$$\epsilon_n^2 = m_q^2 + C_q^2 \frac{1}{n + m_q} (2n + 1); \quad (2)$$

for quarks. C_g and C_q are spring constants of gluon and quark RHO confining potentials respectively. $n = 1; 2; \dots$ is the oscillator quantum numbers and m_q is the mass of quark. Let us first consider gluon plasma without quarks which is a Bose system. Following the standard procedure of statistical mechanics, pressure or logarithm of partition function is given by,

$$\frac{P V}{T} = \sum_n g_n g_f \ln(1 - e^{-\epsilon_n}); \quad (3)$$

where g_n is the full degeneracy arising from the "coalescing together" of an almost continuous set of the zero-potential levels and is given by,

$$g_n = \frac{V C_g^3}{6^2} ((2n + 3)^{3=2} - (2n + 1)^{3=2}); \quad (4)$$

V is the volume and β is the inverse temperature. $g_f = 16$ is the internal degrees of freedom of gluons. Since gluons are bosons, the fugacity $z = 1$. For fermions like quarks $z \neq 1$ and pressure is given by,

$$\frac{P V}{T} = \sum_n g_n g_f \ln(1 + z e^{-\epsilon_n}); \quad (5)$$

where g_f is the internal degrees of quarks. Above expressions for pressure are exact and one need to do infinite sum which may be done numerically. However, for high temperature, we may use Euler-Maclaurin series to get an approximate expression as, after some algebra,

$$\begin{aligned} \frac{P}{T} &= \frac{g_I}{2} \sum_{l=1}^{\infty} \frac{1}{l^{5/2}} \left[K_{3/2} \left(\frac{C_g l}{T} \right) + \frac{3}{4} \frac{C_g l}{T} K_{\frac{1}{2}} \left(\frac{C_g l}{T} \right) \right] \\ &+ \frac{C_g^3 g_I}{12} \left(\frac{3}{3} - 1 \right) \ln \left(1 - e^{-\frac{C_g}{T}} \right) \end{aligned} \quad (6)$$

for gluon plasma, in terms of modified Bessel functions $K_n(x)$. But we know that at extremely high temperature, confinement effects may be negligible and we expect to get EoS of relativistic free gas which we indeed get as shown below. As $T \rightarrow \infty$,

$$\begin{aligned} P &\rightarrow \frac{g_I}{90} T^4 \left(1 + \frac{45}{4} \frac{C_g^2}{T^2} + \dots \right) \\ &\rightarrow \frac{g_I}{90} T^4 = P_s \end{aligned} \quad (7)$$

Note that in our quantum calculation with proper choice of g_n , Eq. (4), as expected, we get T^4 , rather than T^7 as in [9]. Further, we also know that as $T \rightarrow \infty$, we approach classical limit and from our earlier classical calculations of this problem, we indeed got T^4 law [13]. Next let us compare our EoS with the results of lattice simulation of QGP or lattice gauge theory (LGT).

3 Comparison with LGT :

We can see, from Eq. (7), that our EoS of QGP is temperature dependent modification of normal T^4 law. Similar modification is seen in LGT calculations and only at $T \rightarrow \infty$, one expects T^4 law, which is termed as non-ideal effect. As an example, let us consider pure gauge QGP or gluon plasma and the pressure is, Eq. (3),

$$\frac{P}{T} = \frac{C_g^3 g_I}{6} \sum_n^{\infty} \frac{1}{n^{5/2}} \left[(2n+3)^{3/2} - (2n+1)^{3/2} \right] \ln \left(1 - e^{-\frac{C_g}{T} n} \right); \quad (8)$$

which may written as

$$p = \frac{P}{P_s} = \frac{a^3 g_I}{6} \frac{1}{t^{5/2}} \sum_n^{\infty} \frac{1}{n^{5/2}} \left[(2n+3)^{3/2} - (2n+1)^{3/2} \right] \ln \left(1 - e^{-\frac{a}{t} n} \right); \quad (9)$$

where P_s is the Stefan-Boltzmann limit pressure and $t = T = T_c$. T_c is the critical temperature of the transition of hadrons to QGP and the parameter a is defined as, $a = C_g = T_c$. Above equation may be compared with LGT results by varying the parameter a as discussed in section V. Knowing P or partition function, all other thermodynamic quantities like energy density, entropy, sound speed etc. may be derived. Further extension of our calculations for QGP with finite number of quark flavors, without or with finite chemical potential, is straight forward [14].

4 Quark star:

As another example, let us consider what are called quark stars [15]. We know that, at the last stages of a star, it undergoes various changes like white-dwarf star, neutron star and then, probably, quark star. Sometimes more complex compact star with different layers of matter with quark matter inside and then mixture of quarks and hadrons, neutrons, so on. Here we take a very simple compact star, just cold quark star, where the degenerate pressure of quarks in RHOModel balances the gravitational collapse. We estimate the limiting mass for stability and mass-radius relation, using Newtonian description. A detailed study using Tolman-Oppenheimer-Volkoff (TOV) equation with complex compact star in our model is a separate topic. With these simplifications, we get the degenerate pressure from the energy of the system ,

$$E_0 = \frac{C_q^4 V g_f}{6^2} \sum_{n=1}^{N_F} ((2n+3)^{3=2} - (2n+1)^{3=2}) \frac{m_q^2}{m_q^2 + 2n+1}; \quad (10)$$

where we have approximated the single particle energy level of quarks, Eq. (2), for simplicity, as [9]

$$\epsilon_n^2 = m_q^2 + C_q^2 (2n+1); \quad (11)$$

Here g_f is the internal degrees of quarks and $m_q = \frac{m_q}{C_q}$. N_F is the quantum number corresponding to Fermi level, given by the relation,

$$N = \frac{C_q^3 V g_f}{6^2} \sum_{n=1}^{N_F} ((2n+3)^{3=2} - (2n+1)^{3=2}); \quad (12)$$

where N is the total number of quarks and in the case of cold star approximation, quarks fill the single particle RHO levels, according to Pauli's exclusion principle, upto the level n_F . Following the same procedure as that of white-dwarf star calculations we may relate the n_F to the radius, R , of the star as

$$R = 4 \frac{9 M}{2 m_q g_f C_q^3 ((2n_F + 3)^{3=2} \frac{3^{3=2}}{3} \frac{3^{3=2}}{3})^5} ; \quad (13)$$

where M is the mass of star. Degenerate pressure is obtained from E_0 ,

$$P = \frac{\partial E_0}{\partial V} = \frac{C_q^4 g_f}{6^2} \frac{1}{3} \frac{(2n_F + 3)^{3=2} \frac{3^{3=2}}{3} \frac{3^{3=2}}{3}}{(2n_F + 3)^{1=2}} \frac{\partial}{\partial n_F} \sum_{n=1}^{n_F} \frac{1}{(2n+3)^{3=2} (2n+1)^{3=2}} \frac{1}{m_q^2 + 2n + 1} ; \quad (14)$$

To proceed further, we approximate the summation over n by integration using Euler-Maclaurin series. The error introduced by this approximation is small, which may be checked numerically by evaluating summation and Euler-Maclaurin formula. This degenerate pressure balances the gravitational pressure inward, which we take, the Newtonian formula,

$$\frac{M^2}{4 R^4} ; \quad (15)$$

where γ is the gravitational constant and β is a geometrical factor. After some lengthy calculations for large n_F , we get

$$(\tilde{R}^2 + 1) (2 - \tilde{R}^2)^2 = \tilde{M}^{4=3} ; \quad (16)$$

where \tilde{R} and \tilde{M} are related to radius, R , and mass, M , of the star respectively as,

$$\tilde{R} = R C_q \left(\frac{2 g_f m_q}{9 M} \right)^{1=3} ; \quad (17)$$

and

$$\tilde{M} = \frac{M}{m_q} \left(\frac{m_q}{m_q} \right)^{3=2} \frac{g_f}{3} \frac{32}{9} ; \quad (18)$$

It is interesting to compare above mass-radius relation with familiar white-dwarf star mass-radius relation, proceeding with similar normalization of mass and radius,

$$(\tilde{R}^2 + 1) (2=3 - \tilde{R}^2)^2 = \tilde{M}^{4=3} ; \quad (19)$$

We see that the only change in our case is that the factor $2=3$ is replaced by 2 in mass-radius relation and C_q plays the role of m_e , the electron mass. From the Eq. (16), we can estimate the limiting mass, M_0^{QS} , similar to the Chandrasekhar's limit (M_0^W) in the case of white-dwarf star, above which the star will collapse and we get,

$$M_0^{QS} = \left(\frac{m_p}{m_q}\right)^2 \frac{32}{g_f} M_0^W ; \quad (20)$$

where m_p is the mass of proton. It is generally assumed [15] that quark star contains up, down and strange-antistrange quarks and hence $g_f = (2 \times 3 \times 2 + 1 \times 2 \times 3 \times 2) = 24$. The quark mass m_q in RHO is a constituent mass which ranges from 300 MeV to 600 MeV for up quark to strange quark and on an average we may approximate $m_q = m_p=2$. Therefore, as a rough estimate, from Eq. (20), limiting mass of quark star may be of the order of 4 times the solar mass, where as the M_0^W is of the order of solar mass. Further numerical calculation of Eq. (16), without the approximation of $n_F \ll 1$ which is used to get the Eq. (16), we obtained a plot of mass-radius relation as shown Fig. 2.

5 Results :

In Fig. 1, we plotted our results for pressure in terms of $p = P_s$ as function of $t = T/T_c$ along with LGT results. By varying the parameter a , so as to get a similar behaviour as LGT results as function of t , we get $a = 2:3$. However, the curve lies slightly above the lattice data. We may correct our pressure by subtracting a constant such that both the curve matches at large t values and we get good fit except near to $T = T_c$. The value of β here is 0.1. It is not unreasonable to subtract by β because we have neglected the mutual interactions between particles, mainly, attractive Color-Coulombic interactions. We know that attractive mutual interactions among particles reduces the pressure. From the expression for a , we may estimate $T_c = C_g/a = 200 \text{ MeV}$ for $C_g = 2.35 \text{ fm}^{-1}$ [9]. Once pressure or partition function is obtained all other thermodynamic functions like μ , c_s^2 etc. may be easily evaluated. Similar comparison of our results and LGT results may be carried for other systems with flavored QGP [14].

In Fig. 2, we plotted mass-radius relations of quark star. Mass is expressed in terms of solar mass and radius in kilometers (km) by solving, numerically, Eq. (14) after equating to the gravitational pressure. We have taken $m_q = 2.5$ which is equivalent to taking $C_q = 1 \text{ fm}^{-1}$ [9].

6 Conclusions :

We developed statistical mechanics and thermodynamics of a system of quantum particles in RHO confining potential and applied to QGP. We compared the results with LGT and conclude that the overall QCD confining effect, modelled by RHO, explains the LGT data with small correction to take into account of the attractive mutual interactions among particles. We applied our model to study quark star and obtained the limiting mass and mass-radius relation and found to agree of the order of magnitude. To get an accurate result one need to study complex compact stars using TOV equations. However, Boson or Fermion system in RHO confining potential is a new general problem which may be applicable to many fields like BEC, RHICs, astrophysics and so on.

References

- [1] Proceedings of 5th International Conference on Physics and Astrophysics of Quark Gluon Plasma, Kolkata, India (2005), www.veccalernet.in/icpaqgp.
- [2] G. Boyd, J. Engels, F. Karsch, E. Laermann, C. Legeland, M. Lutgen eier and B. Petersson, Phys. Rev. Lett. 75, 4169 (1995); Nucl. Phys. B 469, 419 (1996); F. Karsch, Nucl. Phys. A, 698, 199 (2002); E. Laermann and O. Philipsen, Ann. Rev. Nucl. Part. Sci. 53, 163 (2003).
- [3] A. Rebhan, hep-ph/0010252, 2000.
- [4] A. Peshier, B. Kampfer, O. P. Pavlenko and G. So, Phys. Lett. B 337, 235 (1994); Phys. Rev. D 54, 2399 (1996); P. Levai and U. Heinz, Phys. Rev. C 57, 1879 (1998); R. A. Schneider and W. Weise, Phys. Rev. C 64, 055201 (2001).

- [5] D . H . R ischke, nucl-th/0305030, 2003.
- [6] E . Shuryak, hep-ph/0405066, (2004).
- [7] V . M . Bannur, Eur. Phys. J. C 11, 169 (1999); hep-ph/0504072.
- [8] V . M . Bannur, Phys. Lett. B 362, 7 (1995).
- [9] S . B . K hadkikar, J . C . Parikh and P . C . V inodkum ar, Mod. Phys. Lett. A 8, 749 (1993).
- [10] V . M . Bannur, hep-ph/0508069.
- [11] S . B . K hadkikar and S . K . G upta, Phys. Lett. B 124, 523 (1983); Phys. Rev. D 36, 307 (1987); P . C . V inodkum ar, J . N . Pandya, V . M . Bannur and S . B . K hadkikar, Eur. Phys. J. A 4, 83 (1999).
- [12] R . K . P atria, Statistical M echanics, Butterworth H einem ann, O xford (1997).
- [13] K . M . U dayanandan, V . M . Bannur and P . C . V inodkum ar, submitted to J. Phys. G .
- [14] K . M . U dayanandan and V . M . Bannur, in preparation.
- [15] N . K . G lendening, Compact stars, Springer, Berlin (2000); R . X . X u, astro-ph/0211348; S . Banerjee, S . K . G hosh and S . Raha, J. Phys. G : Nucl. Part. Phys., 26, L1 (2000).

Figure 1: Plots of $P = P_s$ as a function of $T = T_c$ from our model and lattice results (symbols) for pure gauge.

Figure 2: Mass-radius plot of quark star, mass in terms of solar mass and radius in kilometers.




Fig. 1




Fig. 2