

The crystallography of strange quark matter

K. Rajagopal and R. Sharma

Center for Theoretical Physics, Massachusetts Institute of Technology, Cambridge, MA 02139
 Nuclear Science Division, MS 70R 319, Lawrence Berkeley National Laboratory, Berkeley, CA 94720

Abstract. Cold three-flavor quark matter at large (but not asymptotically large) densities may exist as a crystalline color superconductor. We explore this possibility by calculating the gap parameter and free energy for possible crystal structures within a Ginzburg-Landau approximation, evaluating to order μ^{-6} . We develop a qualitative understanding of what makes a crystal structure stable, and find two structures with particularly large values of μ and the condensation energy, within a factor of two of those for the CFL phase known to characterize QCD at asymptotically large densities. The robustness of these phases results in their being favored over wide ranges of density and though it also implies that the Ginzburg-Landau approximation is not quantitatively reliable, previous work suggests that it can be trusted for qualitative comparisons between crystal structures. We close with a look ahead at the calculations that remain to be done in order to make contact with observed pulsar glitches and neutron star cooling.

We explore properties of quark matter at zero temperature and at densities that may occur within neutron star cores. Absent interactions, each flavor of quark will fill momentum eigenstates up to a Fermi surface. However, this configuration is unstable to the formation of Cooper pairs in the presence of any attractive interaction between quarks. In QCD, the interaction between pairs of quarks that are antisymmetric in color is attractive, and hence the ground state features a diquark condensate that is dominantly antisymmetric in color. This is color superconductivity. We consider condensates antisymmetric in Dirac indices and, consequently, in flavor also, implying that quarks in a pair have different flavors.

At asymptotic densities ($\mu_s \rightarrow \infty$), where the three light quarks can be treated as massless, quark matter exists in the CFL phase [1] in which quarks of all three colors and all three flavors form Cooper pairs with zero total momentum, yielding chiral, chiral and chiral condensates, and in which all fermionic excitations are gapped, with a gap parameter $\mu_0 \approx 100 \text{ MeV}$. However, at densities relevant for neutron star phenomenology, meaning quark chemical potentials at most 500 MeV , the strange quark mass M_s cannot be neglected. In neutral unpaired quark matter in weak equilibrium, M_s introduces splitting between the Fermi surfaces for quarks of different flavor which can be taken into account to lowest order in $M_s^2 = \mu^2$ by treating the quarks as if they were massless but with chemical potential splittings $\mu_2 = (\mu_u - \mu_s) = 2$ and $\mu_3 = (\mu_d - \mu_u) = 2$ given by $\mu_2 = \mu_3 = M_s^2 = (8)$. Note that the splitting between unpaired Fermi surfaces increases with decreasing density. In the CFL phase, the Fermi momenta are not given by these optimal values for unpaired quark matter; instead, the system pays a free energy price μ^2 to equalize all Fermi momenta and gains a pairing energy benefit μ_0^2 . As a function of decreasing density, there comes a point (at which $\mu_0 = 4$ [2]) when the system can lower its energy by breaking pairs. Below this density the CFL phase can certainly not be the ground state of matter. Within a spatially homogeneous pairing ansatz, the phase that results when CFL Cooper pairs start to break is the gapless CFL (gCFL) phase of Ref. [2]. However, this phase turns out to be unstable to the formation of counter-propagating currents, presumably leading to inhomogeneity, and therefore cannot be the ground state of matter at any density [3]. Within the range of densities in which the gCFL phase has lower free energy than the CFL and unpaired phases, the true ground state of dense matter must have lower free energy still. We have recently proposed a candidate phase (actually, two candidate phases) for the ground state of matter over much of this

z Speaker.

density regime and at still lower densities [4]. In these proceedings we describe this proposal. For further references, in particular to other suggested resolutions of the gCFL instability, see Ref. [4].

Crystalline color superconductivity [5, 6] naturally permits pairing between quarks living at split Fermi surfaces by allowing Cooper pairs with nonzero net momentum. In three-flavor quark matter, this allows pairing to occur even with the Fermi surfaces split in the free-energetically optimal way as in the absence of pairing [7, 9, 4]. This is the origin of the advantage that crystalline color superconducting phases have over the CFL and gCFL phases at large values of the splitting. For example, by allowing u quarks with momentum $p + q_3$ to pair with d quarks with momentum $p + q_3$, for any p , we can pair u and d quarks along rings on their respective Fermi surfaces [5, 6]. In coordinate space, this corresponds to a condensate of the form $\text{hudi} \exp 2iq_3 \cdot r$. The net free energy gained due to pairing is then a balance between increasing μ_3 yielding pairing on larger rings while exacting a greater kinetic energy cost. The optimal choice turns out to be $\mu_3 = \mu_1$ with $\mu_1 = 1.1997$, corresponding to pairing rings on the Fermi surfaces with opening angle 67.1° [5]. It is possible to cover larger areas of the Fermi surfaces by allowing Cooper pairs with the same μ_3 but various q_3 , yielding $\text{hudi} \exp 2iq_3^a \cdot r$ with the q_3^a chosen from some specified set fq_3g . This is a condensate modulated in position space in some crystalline pattern, with the crystal structure defined by fq_3g . In this two-flavor context, a Ginzburg-Landau analysis reveals that the best fq_3g contains eight vectors pointing at the corners of a cube, say in the $(1; 1; 1)$ directions in momentum space, yielding a face-centered cubic structure in position space [6].

We use the following ansatz for the three-flavor crystalline color superconducting condensate [4]:

$$\text{h}_{ij} = C \prod_{I=1}^3 \exp(iq_I^a \cdot r) \quad (1)$$

This is antisymmetric in color $(i; j)$, spin, and flavor $(i; j)$ indices and is thus a generalization of the CFL condensate to crystalline color superconductivity. We set $\mu_1 = 0$, neglecting hdsi pairing because the d and s Fermi surfaces are twice as far apart from each other as each is from the intervening u Fermi surface. Hence, I can be taken to run over 2 and 3 only. fq_2g and fq_3g define the crystal structures of the hdsi and hudi condensates respectively. We only consider crystal structures in which all the vectors in fq_2g are equivalent to each other, and same for fq_3g , as this justifies our simplifying assumption that the hdsi and hudi condensates are each specified by a single gap parameter (μ_2 and μ_3 respectively), avoiding having to introduce one gap parameter per q . We furthermore only consider crystal structures which are exchange symmetric, meaning that fq_2g and fq_3g can be exchanged by some combination of rigid rotations and reflections applied simultaneously to all the vectors in both sets. This simplification, together with $\mu_2 = \mu_3$ (an approximation corrected only at order $M_s^4 = \mu_s^3$), guarantees that we find solutions with $\mu_2 = \mu_3$.

We analyze and compare candidate crystal structures by evaluating the free energy $\mathcal{F}(\mu_2; \mu_3)$ for each crystal structure in a Ginzburg-Landau expansion in powers of the μ 's. This approximation is controlled if $\mu_2; \mu_3 \ll \mu_0$, with μ_0 the gap parameter in the CFL phase at $M_s^2 = \mu_s = 0$. The terms in the Ginzburg-Landau expansion must respect the global $U(1)$ symmetry for each flavor, meaning that each μ_I can only appear in the combination μ_I^2 . (The $U(1)$ symmetries are spontaneously broken by the condensate, but not explicitly broken.) Therefore, $\mathcal{F}(\mu_2; \mu_3)$ is given to sextic order by

$$\mathcal{F}(\mu_2; \mu_3) = \frac{\mu_2^2}{2} P_2 \mu_2^2 + P_3 \mu_3^2 + \frac{1}{2} \mu_2^4 + \mu_3^4 + \mu_2^2 \mu_3^2 + \frac{1}{3} \mu_2^6 + \mu_3^6 + \mu_2^4 \mu_3^2 + \mu_2^2 \mu_3^4 \quad (2)$$

where we have chosen notation consistent with that used in the two-flavor study of Ref. [6], which arises as a special case of (2) if we take μ_2 or μ_3 to be zero. P_I is the number of vectors in the set fq_Ig . The form of the Ginzburg-Landau expansion (2) is model-independent, whereas the expressions for the coefficients $P_I, \mu_I, \mu_{IJ}, \mu_{IJK}$ for a specific crystal structure are model-dependent. For exchange symmetric crystal structures, $\mu_2 = \mu_3, \mu_2^2 = \mu_3^2, \mu_2^4 = \mu_3^4$ and $\mu_2^2 \mu_3^2 = \mu_2^4 \mu_3^2$.

Because setting one of the q_i to zero reduces the problem to one with two-flavor pairing only, we can obtain \mathcal{F} , and \mathcal{F}' via applying the two-flavor analysis described in Ref. [6] to either f_{q_2g} or f_{q_3g} separately. Using \mathcal{F} as an example, we learn that

$$\mathcal{F} = \mathcal{F}(q_i; \mu_i) = \mu_i + \frac{\mu_i}{2q_i} \log \frac{q_i + \mu_i}{q_i - \mu_i} - \frac{1}{2} \log \frac{\Delta_{2SC}^2}{4(q_i^2 - \mu_i^2)}; \quad (3)$$

Here, $q_i = \sum_j v_j$ and Δ_{2SC} is the gap parameter for the 2SC (2-flavor, 2-color) BCS pairing obtained with $\mu_i = 0$ and μ_j nonzero with the other two gap parameters set to zero. Assuming that $\mu_i = 0$, it is given by $\Delta_{2SC} = 2^{1/3} \mu_i$ [10]. In the Ginzburg-Landau approximation, in which the μ_i are assumed small, we must first minimize the quadratic contribution to the free energy, and only then investigate the quartic and sextic contributions. Minimizing \mathcal{F} fixes the length of all the vectors in the set $f_{q_i g}$, yielding $q_i = \mu_i$ with $\mu_i = 1:1997$ the solution to $\frac{1}{2} \log [(q_i + \mu_i)/(q_i - \mu_i)] = 1$ [5]. Upon setting $q_i = \mu_i$, (3) becomes

$$\mathcal{F}(\mu_i) = \frac{1}{2} \log \frac{\Delta_{2SC}^2}{4(\mu_i^2 - 1)}; \quad (4)$$

Furthermore, the only dimensionful quantities on which the quartic and sextic coefficients can depend are then the μ_i [6, 4], meaning that for exchange symmetric crystal structures and with $\mu_2 = \mu_3 = \mu$ we have $\mathcal{F} = \mathcal{F}(\mu) = \mathcal{F}(\mu)$, $\mathcal{F}'_{32} = \mathcal{F}'_{32} = \mathcal{F}'(\mu)$, $\mathcal{F}'_{322} = \mathcal{F}'_{322} = \mathcal{F}'(\mu)$ and $\mathcal{F}'_{322} = \mathcal{F}'_{322} = \mathcal{F}'(\mu)$ where the barred quantities are dimensionless numbers which depend only on f_{q_2g} and f_{q_3g} that must be evaluated for each crystal structure. Doing so requires evaluating one loop Feynman diagrams with 4 or 6 insertions of μ_i 's. Each insertion of μ_i (μ_i) adds (subtracts) momentum $2q_i^a$ for some a , meaning that the calculation consists of a bookkeeping task (determining which combinations of 4 or 6 q_i^a 's are allowed) that grows rapidly in complexity with the complexity of the crystal structure and a loop integration that is nontrivial because the momentum in the propagator changes after each insertion. See Ref. [4], where this calculation is carried out explicitly for 11 crystal structures in a mean-field NJL model upon making the weak coupling (μ_0 and μ both much less than μ_0) approximation. Note that in this approximation neither the NJL cutoff nor the NJL coupling constant appear in any quartic or higher Ginzburg-Landau coefficient, and they appear in \mathcal{F} only within μ_0 . Hence, the details of the model do not matter as long as one thinks of μ_0 as a parameter, kept fixed.

It is easy to show that for exchange symmetric crystal structures any extremum of $\mathcal{F}(\mu_2; \mu_3)$ in $(\mu_2; \mu_3)$ -space must either have $\mu_2 = \mu_3 = \mu$, or have one of μ_2 and μ_3 vanishing [4]. It is also possible to show that the three-flavor crystalline phases with $\mu_2 = \mu_3 = \mu$ are electrically neutral whereas two-flavor solutions in which only one of the μ_i 's is nonzero are not [4]. We therefore analyze only solutions with $\mu_2 = \mu_3 = \mu$. In marked contrast to the two-flavor results of Ref. [6] which (ignoring the requirement of neutrality) show that many two-flavor crystal structures have negative \mathcal{F} and hence sextic order free energies that are unbounded from below, we find that $\mathcal{F}(\mu; \mu)$ is positive for large μ for all the crystal structures that we investigate [4]. This allows us to minimize $\mathcal{F}(\mu; \mu)$ with respect to μ , thus evaluating \mathcal{F} and \mathcal{F}' .

We begin with the simplest three-flavor "crystal" structure in which f_{q_2g} and f_{q_3g} each contain only a single vector, making the u and d condensates each a single plane wave. This simple condensate yields a qualitative lesson which proves helpful in winnowing the space of multiple plane wave crystal structures [4]. For this simple "crystal" structure, all the coefficients in the Ginzburg-Landau free energy can be evaluated analytically [4]. The terms that occur in the three-flavor case but not in the two-flavor case, namely \mathcal{F}'_{32} and \mathcal{F}'_{322} , describe the interaction between the two condensates, and depend on the angle θ between q_2 and q_3 . For any angle θ , both \mathcal{F}'_{32} and \mathcal{F}'_{322} are positive. And, both increase monotonically with θ and diverge as $\theta \rightarrow \pi$ [4]. This divergence tells us that choosing q_2 and q_3 precisely antiparallel exacts an infinite free energy price in the combined Ginzburg-Landau and weak-coupling limit in which $\mu_0 \rightarrow 0$; $\mu \rightarrow 0$, meaning that in this limit if we chose $\mu = \mu$ we find $\mathcal{F} = 0$. Away from the Ginzburg-Landau limit, when the pairing rings on the Fermi surfaces widen into bands, choosing $\mu = \mu$ exacts a finite price meaning that \mathcal{F} is nonzero but smaller than that for any other choice of μ . The high cost of choosing q_2 and q_3 precisely antiparallel can be understood qualitatively as arising from the fact that in this case the ring of states on the u -quark Fermi surface that "want to" pair with d -quarks coincides precisely with the ring

that wants to pair with s-quarks [9]. This simple two plane wave ansatz has been analyzed upon making the weak-coupling approximation but without making the Ginzburg-Landau approximation [9]. All the qualitative lessons learned from the Ginzburg-Landau approximation remain valid and we learn further that the Ginzburg-Landau approximation always underestimates [9].

The analysis of the simple two plane wave "crystal" structure, together with the observation that in more complicated crystal structures with more than one vector in f_{q_2g} and f_{q_3g} the Ginzburg-Landau coefficient $\chi_2(\chi_{322})$ is given in whole (in part) by a sum of many two plane wave contributions, yields one of two rules for constructing favorable crystal structures for three-avor crystalline color superconductivity [4]: f_{q_2g} and f_{q_3g} should be rotated with respect to each other in a way that best keeps vectors in one set away from the antipodes of vectors in the other set. The second rule is that the sets f_{q_2g} and f_{q_3g} should each be chosen to yield crystal structures which, seen as separate two-avor crystalline phases, are as favorable as possible. The 11 crystal structures analyzed in Ref. [4] allow one to make several pairwise comparisons that test these two rules. There are instances of two structures which differ only in the relative orientation of f_{q_2g} and f_{q_3g} and in these cases the structure in which vectors in f_{q_2g} get closer to the antipodes of vectors in f_{q_3g} are disfavored. And, there are instances where the smallest angle between a vector in f_{q_2g} and the antipodes of a vector in f_{q_3g} are the same for two different crystal structures, and in these cases the one with the more favorable two-avor structure is more favorable. These considerations, together with explicit calculations, indicate that two structures, which we denote "2Cube45z" and "CubeX", are particularly favorable.

In the 2Cube45z crystal, f_{q_2g} and f_{q_3g} each contain eight vectors pointing at the corners of a cube. If we orient f_{q_2g} so that its vectors point in the $(1; 1; 1)$ directions in momentum space, then f_{q_3g} is rotated relative to f_{q_2g} by 45° about the z-axis. In this crystal structure, the h_{ud} and h_{us} condensates are each given by the most favored two-avor crystal structure [6]. The relative rotation maximizes the separation between any vector in f_{q_2g} and the nearest antipodes of a vector in f_{q_3g} .


We arrive at the CubeX structure by reducing the number of vectors in f_{q_2g} and f_{q_3g} . This worsens the two-avor free energy of each condensate separately, but allows vectors in f_{q_2g} to be kept farther away from the antipodes of vectors in f_{q_3g} . We have not analyzed all structures obtainable in this way, but we have found one and only one which has a condensation energy comparable to that of the 2Cube45z structure. In the CubeX structure, f_{q_2g} and f_{q_3g} each contain four vectors forming a rectangle. The eight vectors together point toward the corners of a cube. The 2 rectangles intersect to look like an "X" if viewed end-on. The color, flavor and position space dependence of the CubeX condensate is given by

$$\chi_2 = \frac{1}{2} \sum_{ij} \cos \frac{2}{a} (x + y + z) + \frac{1}{2} \sum_{ij} \cos \frac{2}{a} (x - y + z) + \frac{1}{2} \sum_{ij} \cos \frac{2}{a} (x + y - z) + \frac{1}{2} \sum_{ij} \cos \frac{2}{a} (x - y - z); \quad (5)$$

where $a = \frac{p}{3} = q = 4.536 \text{ fm} = (1.764 M_s^2)$ is the lattice spacing. For example, with $M_s^2 = 100; 150; 200 \text{ MeV}$ the lattice spacing is $a = 72; 48; 36 \text{ fm}$. We depict this condensate in Fig. 1.

In Figs. 2 and 3, we plot χ_2 and χ_3 versus M_s^2 for the most favorable crystal structures that we have found, namely the CubeX and 2Cube45z structures described above. We have taken the CFL gap parameter $\mu_0 = 25 \text{ MeV}$ in these figures, but they can easily be rescaled to any value of μ_0 [4]. Fig. 2 shows that the gap parameters are large enough that the Ginzburg-Landau approximation is at the edge of its domain of reliability. However, results obtained for simpler crystal structures suggest that the Ginzburg-Landau calculation underestimates χ_2 and the condensation energy and that, even when it breaks down, it is a good qualitative guide to the favorable structure [9]. We therefore trust the result, evident in Fig. 3, that these crystalline phases are both impressively robust, with one or other of them favored over a wide swath of M_s^2 and hence density. We do not trust the Ginzburg-Landau calculation to discriminate between these two structures, particularly given that although we have a qualitative understanding of why these two are favorable we have no qualitative argument for why one should be favored over the other.

Figure 1. The CubeX crystal structure of Eq. (5). The figure extends from 0 to $a=2$ in the x , y and z directions. Both $\rho_2(x)$ and $\rho_3(x)$ vanish at the horizontal plane. $\rho_2(x)$ vanishes on the darker vertical planes, and $\rho_3(x)$ vanishes on the lighter vertical planes. On the upper (lower) dark cylinders and the lower (upper) two small corners of dark cylinders, $\rho_2(x) = +3/3$ ($\rho_2(x) = -3/3$). On the upper (lower) lighter cylinders and the lower (upper) two small corners of lighter cylinders, $\rho_3(x) = -3/3$ ($\rho_3(x) = +3/3$). The largest value of $j_1(x)$ is 4, occurring along lines at the centers of the cylinders. The lattice spacing is a when one takes into account the signs of the condensates; if one looks only at $j_1(x)$, the lattice spacing is $a=2$.



We are confident that 2Cube45z is the most favorable structure obtained by rotating one cube relative to another. We are not as confident that CubeX is the best possible structure with fewer than $8+8$ vectors. Regardless, the 2Cube45z and CubeX crystalline phases together make the case that three-avor crystalline color superconducting phases are the ground state of cold quark matter over a wide range of densities. If even better crystal structures can be found, this will only further strengthen this case.

Fig. 3 shows that over most of the range of $M_s^2 = \mu$ where it was once considered a possibility, the gCFL phase can be replaced by a much more favorable three-avor crystalline color superconducting phase. It also shows that it is hard to find a crystalline phase with lower free energy than the gCFL phase at the lowest values of $M_s^2 = \mu$ (highest densities) in the "gCFL window". This narrow window where the gCFL curve remains the lowest in Fig. 3 is thus the most likely place in the QCD phase diagram to find the gCFL phase augmented by current-carrying meson condensates described in Refs. [11]. Except within this window, crystalline color superconducting phases with the CubeX or the 2Cube45z crystal structures provide an attractive resolution to the instability of the gCFL phase.

The three-avor crystalline color superconducting phases with CubeX and 2Cube45z crystal structures are the lowest free energy phases that we know of, and hence candidates for the ground state of QCD, over a wide range of densities. One or other is favored over the CFL, gCFL and unpaired phases for $2.9 \mu_0 < M_s^2 = \mu < 10.4 \mu_0$, as shown in Fig. 3. For $\mu_0 = 25 \text{ MeV}$ and $M_s = 250 \text{ MeV}$, this translates to $240 \text{ MeV} < \mu < 847 \text{ MeV}$. With these choices of parameters, then, the lower part of this range of μ (higher part of the range of $M_s^2 = \mu$) is certainly superseded by nuclear matter. And, the high end of this range extends far beyond the $\sim 500 \text{ MeV}$ characteristic of the quark matter at the densities expected at the center of compact stars. If compact stars have quark matter cores, then, it is reasonable to include the possibility that the entire quark matter core could be in a crystalline color superconducting phase on the menu of options that must ultimately be winnowed by confrontation with astrophysical observations. If μ_0 is larger, say $\sim 100 \text{ MeV}$, the entire quark matter core could be in the CFL phase.

Now that we have two candidates for the crystal structure of the three-avor crystalline color superconducting phase of cold quark matter, favorable over a very wide range of intermediate densities, and a qualitative guide to the scale of μ and μ_0 , we can look ahead toward the calculation of astrophysically relevant observables. The heat capacity of the CubeX and 2Cube45z phases should be only quantitatively suppressed relative to that of unpaired quark matter, but their neutrino emissivity may be more significantly suppressed [4]. Both can now be calculated, yielding an estimate of the effects of a crystalline quark matter core on the rate at which a neutron star cools by neutrino emission. A crystalline color superconducting core, being both crystalline and superfluid, could be a region within which rotational vortices are pinned and hence (some) pulsar glitches originate [5]. Or, the presence of crystalline quark matter within neutron stars




Figure 2. Gap parameter versus $M_s^2 = \mu$ for the CFL gap parameter, the three gap parameters $\mu_1 < \mu_2 < \mu_3$ describing uds , hus and hud pairing in the gCFL phase, and the gap parameter in the crystalline color superconducting phases with CubeX and 2Cube45z crystal structures.




Figure 3. Free energy relative to that for neutral unpaired quark matter versus $M_s^2 = \mu$, for the same phases as in Fig. 2. The gCFL phase free energy is a dashed line as a reminder that this phase is unstable: the free energy of the ground state of matter must lie below this dashed line.

could be ruled out if it predicts glitch phenomenology in qualitative disagreement with that observed. The two microphysical properties of crystalline quark matter that must be estimated before glitch phenomenology can be addressed are the vortex pinning force and the shear modulus. Glitches occur if rotational vortices are pinned and immobile while the spinning pulsar's angular velocity slows over years, with the glitch triggered by the catastrophic unpinning of long-immobile vortices. Immobilizing vortices requires sufficient pinning force and shear modulus. Estimating the pinning force will require analyzing how the CubeX and 2Cube45z phases respond when rotated. The shear modulus is related to the coefficients in the effective theory that describes the phonon modes of the crystal [12]. Analyzing the phonons in the three-flavor crystalline color superconducting phases with the 2Cube45z and CubeX crystal structures is thus a priority. These are the prerequisites to determining whether observations of pulsar glitches can be used to rule out (or in) the presence of quark matter in a crystalline color superconducting phase within neutron stars.

We thank the LBNL Nuclear Theory Group for hospitality. Research supported in part by the U.S. DOE under contract # DE-AC02-05CH11231 and cooperative research agreement # DF-FC02-94ER40818.

- [1] M. G. Alford, K. Rajagopal and F. Wilczek, Nucl. Phys. B 537, 443 (1999).
- [2] M. Alford, C. Kouvaris and K. Rajagopal, Phys. Rev. Lett. 92, 222001 (2004); Phys. Rev. D 71, 054009 (2005).
- [3] R. Casalbuoni, R. Gatto, M. M. Annarelli, G. Nardulli and M. Ruggieri, Phys. Lett. B 605, 362 (2005) [Erratum -ibid. B 615, 297 (2005)]; K. Fukushima, Phys. Rev. D 72, 074002 (2005); this instability was analyzed previously in a gapless paired phase of two-flavor quark matter in M. Huang and I. A. Shovkovy, Phys. Rev. D 70, 051501 (2004); Phys. Rev. D 70, 094030 (2004); I. Giannakis and H. C. Ren, Phys. Lett. B 611, 137 (2005).
- [4] K. Rajagopal and R. Shama, arXiv:hep-ph/0605316.
- [5] M. G. Alford, J. A. Bowers and K. Rajagopal, Phys. Rev. D 63, 074016 (2001); R. Casalbuoni, R. Gatto, M. M. Annarelli and G. Nardulli, Phys. Lett. B 511, 218 (2001); R. Casalbuoni and G. Nardulli, Rev. Mod. Phys. 263, 320 (2004); this is the QCD analogue of a phenomenon first described in A. I. Larkin and Yu. N. Ovchinnikov, Zh. Eksp. Teor. Fiz. 47, 1136 (1964) [Sov. Phys. JETP 20, 762 (1965)]; P. Fulde and R. A. Ferrell, Phys. Rev. 135, A550 (1964).
- [6] J. A. Bowers and K. Rajagopal, Phys. Rev. D 66, 065002 (2002).
- [7] R. Casalbuoni, R. Gatto, N. Ippolito, G. Nardulli and M. Ruggieri, Phys. Lett. B 627, 89 (2005).
- [8] M. Cimiale, G. Nardulli, M. Ruggieri and R. Gatto, arXiv:hep-ph/0602180.
- [9] M. M. Annarelli, K. Rajagopal and R. Shama, arXiv:hep-ph/0603076.
- [10] T. Schafer, Nucl. Phys. B 575, 269 (2000).
- [11] A. K. Ryjevski, hep-ph/0508180; T. Schafer, Phys. Rev. Lett. 96, 012305 (2006); A. Geethold and T. Schafer, hep-ph/0603257.
- [12] R. Casalbuoni, E. Fabiano, R. Gatto, M. M. Annarelli and G. Nardulli, Phys. Rev. D 66, 094006 (2002).