

Axino dark matter from Q-balls in Affleck-Dine baryogenesis and the $\Omega_b = \Omega_{DM}$ coincidence problem

Leszek Roszkowski¹ and Osamu Seto²

(1) Department of Physics and Astronomy, University of Sheffield, Sheffield, UK,

(2) Department of Physics and Astronomy, University of Sussex, Brighton BN1 9QJ, UK

(Dated: December 2, 2021)

We show that the $\Omega_b = \Omega_{DM}$ coincidence can naturally be explained in a framework where axino is cold dark matter which is predominantly produced in nonthermal processes involving decays of Q-balls formed in Affleck-Dine baryogenesis. In this approach, the similarity of Ω_b and Ω_{DM} is a direct consequence of the (sub-)GeV scale of the mass of the axino, while the reheating temperature T_R must be low, some 10^2 GeV, or less.

1. Introduction.

The origin of nonbaryonic cold dark matter (DM) and of baryon asymmetry in the Universe are among the longest lasting puzzles in cosmology as well as in particle physics today. In particular, the question of why the observed values of baryon density Ω_b and of dark matter Ω_{DM} are so close to each other, $\Omega_{DM} = \Omega_b = 5.65 \pm 0.58$ [1], remains a mystery.

A standard paradigm is that the nonbaryonic cold dark matter is made up of some weakly interacting massive particle (WIMP) which freezes out of thermal equilibrium in the early Universe. Perhaps the most popular WIMP candidate is the lightest neutralino of the Minimal Supersymmetric Standard Model (MSSM) as the lightest supersymmetric particle (LSP). It remains stable due to the conservation of R-parity. This economical scenario does not, however, explain the proximity of Ω_b and Ω_{DM} .

The same is generally true for conventional mechanisms of baryogenesis or leptogenesis. This may indicate that the observed baryon-to-DM density ratio is just a pure accident, or else a result of some underlying, and as yet unknown, more fundamental theory. An alternative approach is to try to identify a physical mechanism which would simultaneously produce both baryon asymmetry and DM in the proportions consistent with observations. It is clear that this basically necessitates abandoning standard paradigms for producing both types of species in the Universe. This may be one important lesson to learn from these considerations.

A number of attempts at explaining baryon-to-DM ratio have been suggested in the literature. For instance, recently a right handed sneutrino [2] and a sneutrino condensate as an Affleck eld [3] have been proposed.

A few years ago, Enqvist and McDonald (EMD) proposed [4, 5] an attractive solution based on a variant of Affleck-Dine (AD) baryogenesis [6]. In that scenario, an AD condensate forms during inflation and develops a large vacuum expectation value (VEV) along a D - at direction in the MSSM. D - at directions are configurations

of scalar elds for which the D -part of the potential vanishes. Prevalent in theories with many scalar elds like the MSSM, they are of much interest to cosmology [7].

In the standard AD scenario, after the end of inflation, the scalar eld condensate slowly rolls towards the origin and, after a few dozens of coherent oscillations, produces a nonzero baryon number in presence of baryon number-violating couplings of the elds making up the flat direction. Originally, Kusenko and Shaposhnikov argued that the AD condensate can instead fragment into non-topological solitons called Q-balls [8]. If their baryonic charge is large enough, as in models with gauge mediated SUSY breaking, Q-balls remain effectively stable until today, and contribute to the DM density, despite severe astrophysical constraints [9]. On the other hand, EMD demonstrated that, under nontrivial but natural conditions (that we summarize below), in a large class of supergravity (SUGRA) models with gravity mediated SUSY breaking (GRMSB) Q-balls subsequently decay into baryonic matter and neutralino WIMPs assumed to be the LSP [4].

In the EMD scenario, the baryon-to-DM ratio can easily be estimated to be in the right ballpark, as we shall see below. This otherwise attractive framework suffers, however, from a serious problem: neutralino production in Q-ball decays is in fact too efficient, and density can only agree with observations for low neutralino mass $m_{\tilde{\chi}^0_1} \sim 1$ GeV, well below LEP limits [10]. Moreover, this puts into a potential jeopardy the AD mechanism in a large class of GRMSB supergravity models.

In this Letter, we suggest a way out from the above problems of the EMD scenario which at the same time preserves its successful features, in particular, an explanation of the $\Omega_b = \Omega_{DM}$ ratio. We propose that the DM is not made up of the neutralino but instead of an axino, a superpartner of the axion. The axino is a neutral Majorana, chiral fermion. It arises in SUSY models incorporating a Peccei-Quinn solution to the strong CP problem in QCD. Unlike for the neutralino or gravitino, its mass is strongly model dependent and can be much smaller than the (gravity mediated) SUSY breaking scale [11, 12, 13]. Similarly to the axion, its interactions are suppressed by the PQ scale $f_a \sim 10^{11}$ GeV, well below the sensitivity of LEP. The axino has a number of properties which make

Present address: Instituto de Física Teórica, Universidad Autónoma de Madrid, Cantoblanco, Madrid 28049, Spain.

it a promising candidate for cold dark matter [14, 15]. Earlier papers considered warm axino relics [13, 16]. As we will show below, axinos are naturally produced at low temperatures of a few GeV, consistent with the Q-ball scenario of EMD but still before the period of Big Bang Nucleosynthesis (BBN).

2. The Enqvist-Mcdonald scenario.

We now briefly present the main features of the EMD variant of AD baryogenesis. It is assumed that the AD eld is a D - at direction in the M SSM. Its potential is, in general, lifted by soft supersymmetric (SUSY) breaking terms and nonrenormalizable terms [17, 18].

The potential of the AD eld, including in aton-induced terms, reads

$$V(\phi) = (m^2 - qH^2) \phi + K \ln \frac{j \bar{j}}{2} + j \bar{j} + c_2 H + A m_{3=2} \frac{n}{n M_P^{n-3}} + H \phi + \frac{2 j \bar{j}^{n-2}}{M^{2n-6}}; \quad (1)$$

where m is the soft SUSY breaking mass for the AD eld and a radiative correction is given by $K \ln j \bar{j}$. A at direction dependent constant, K , takes values from 0.01 to 0.1 [19, 20]. ϕ denotes a renormalization scale and qH^2 , with $c_1 = 1$, is the negative mass-squared term induced by the energy density of the in aton [18]. Terms proportional to A and c_2 are the trilinear terms from low energy SUSY breaking and those induced by the in aton, respectively, while $m_{3=2}$ denotes the gravitino mass. The nonrenormalizable terms in Eq. (1) come from the superpotential $W = \phi^n / (n M_P^{n-3})$, where ϕ is the Yukawa coupling and M is some large scale acting as a cut-off. In SUGRA, it is natural to assume $M = M_P = 2.4 \times 10^{18}$ GeV which is the reduced Planck mass.

Since during in ation the Hubble parameter $H = m_{3=2}$, the AD eld settles down at the minimum of the potential (1) which is given by

$$j \bar{j} = \frac{c_1}{n-1} \frac{H M_P^{n-3}}{1} = \frac{H M_P^{n-3}}{n-1} = \frac{H M_P^{n-3}}{n-1} = \frac{H M_P^{n-3}}{n-1}; \quad (2)$$

It is clear that the AD eld can naturally develop a very large VEV, which is possible in nonminimal Kahler potentials [18], or if large enough trilinear term A is induced by the in aton [21].

We have neglected in Eq. (1) thermal mass terms $h^2 T^2 j \bar{j}$, where h denotes couplings of the AD eld to other particles [22]. They would play a role if the AD eld VEV were relatively small. We have also neglected two loop thermal effects due to the running of gauge coupling which generate a term $T^4 \ln j \bar{j} = T^2$, where $j \bar{j} = O(10^2)$ [23]. They will not be important below.

As H decreases, the AD eld traces the instantaneous minimum after in ation, begins to oscillate when $H_{osc}^2 \sim m^2$ and, after a few dozen turns, produces a nonzero baryon number and then fragments into Q-balls.

The baryon number density for the AD eld is given by $n_b = i q (-\dot{\phi})$ where q is the baryonic charge for the AD eld. By using the equation of motion of the AD eld, the charge density can be rewritten as

$$n_b(t) = \frac{1}{a(t)^3} \int_{t_{osc}}^t dt^0 a(t^0)^3 \frac{2 q m_{3=2}}{M_P^{n-3}} \text{Im}(A^n); \quad (3)$$

with $a(t)$ being the scale factor. When the AD eld starts to oscillate around the origin, the baryon number density is induced by the relative phase between A and c_2 . With the entropy density after reheating $s = 4 \pi^2 g T^3 = 90$, we can express the baryon asymmetry as

$$\frac{n_b}{s} = \frac{T_R n_b}{4 M_P^2 H^2 t_{osc}} = \frac{q A \text{Im}_{3=2} T_R j_{osc}}{2 H_{osc}^3 M_P^{n-1}} \sin \phi; \quad (4)$$

Here, t_{osc} denotes the time of the start of the oscillation, and $\sin \phi$ is the effective CP phase.

From now on, we consider the case of $n = 6$ because a promising AD eld for our scenario, a odd direction belongs to this class. Let us first evaluate n_b/s . The baryon asymmetry, Eq. (4), for the relevant case is estimated as

$$\frac{n_b}{s} = \frac{q A \text{Im}_{3=2} T_R}{2 M_P^{3=2} m_{3=2} M_P^{1=2}} = 10^{10} \frac{q A \text{Im}_{3=2}}{3=2} \frac{m_{3=2}}{100 \text{ GeV}} \frac{10^3 \text{ GeV}^{3=2}}{m} \frac{T_R}{100 \text{ GeV}} \quad (5)$$

which is of the right order. (The previously made assumption $M = M_P$ is crucial, for otherwise Q-balls would decay too early or would evaporate.) The low reheating temperature T_R after in ation is required to explain the appropriate baryon asymmetry. In this case, the AD condensate fragments into Q-balls [19, 20].

The growth of perturbations of the AD eld and its subsequent fragmentation into Q-balls crucially depends on the logarithmic correction to the ϕ^2 mass term in $V(\phi)$, Eq. (1). An essential requirement is that $V(\phi)$ is flatter than quadratic, or that $K < 0$ [20]. This can be achieved in SUGRA models with a nonminimal Kahler potential [19].

In order to discuss the evolution of Q-balls, first we briefly summarize their relevant properties in GRM SB models. The radius of a Q-ball, R , is estimated as $R^2 = 2/(K \text{Im}^2)$ [19]. The charge is roughly given by $Q = \frac{4}{3} R^3 n_b(t_i) = \frac{4}{3} R^3 (H_i = H_{osc})^2 n_b t_{osc}$; where the suffix i represents the time when the spatial inhomogeneity becomes nonlinear, which can be evaluated as [24]

$$Q = 6 \times 10^3 \frac{2 q A \text{Im}_{3=2} T_R}{3=2} \frac{m_{3=2} M_P^{3=2}}{m^{5=2}} = 10^{20} \frac{q A \text{Im}_{3=2}}{3=2} \frac{m_{3=2}}{100 \text{ GeV}} \frac{1 \text{ TeV}^{5=2}}{m} \quad (6)$$

Unless $Q > 0$ (10^{18}), Q -balls will evaporate before decaying [25]. For Q as in Eq. (6), Q -ball decay temperature is $T_d \sim 1 \text{ GeV}$ to 1 MeV [4, 26]. For example [26],

$$T_d < 2 \text{ GeV} \quad \frac{0.03}{k_B} \quad \frac{m}{1 \text{ TeV}} \quad \frac{10^{20}}{Q} \quad \frac{1}{Q}$$

which is lower than the typical freeze out temperature of WIMPs, $T_f \sim m = 24$. Thus, the LSPs generated in Q -ball decays do not subsequently thermalize. Nor will the baryon asymmetry be washed out by sphaleron effects since $T_d < T_{ew}$ [4]. Note that Q -balls decay prior to BBN and thus do not spoil its successful predictions.

In the EMD scheme, T_R also must be rather low. (This justifies neglecting thermal effects in Eq. (1).) In fact, unless $T_R < 10^{3.5} \text{ GeV}$, Q -balls could thermalize [4]. In order to preserve the $b \{ DM \}$ relation, one needs to suppress the neutralino population from freeze-out. For this to happen it would be sufficient to assume $T_R < T_f$.

It is easy to see why in the EMD scenario, the ratio $b = \frac{n_b}{n_{DM}}$ should be less than 1. The Q -ball is basically a huge "bag" of squarks. It decays predominantly via $q \rightarrow q + \tilde{\chi}$. Thus, for one unit of a baryon number, at least $N \sim 3$ units of nonbaryonic number density are created. (This number can be larger than 3 if one takes into account additional decays of squarks into heavier charginos and neutralinos which then cascade decay into the lightest neutralino, which are model dependent.) In other words, the LSP number density n after Q -ball decay is given by $n = N f_B n_b$, where f_B is the fraction of baryon asymmetry carried by the AD field that is transferred into Q -balls. From lattice calculations, $f_B \sim 1$ [27]. Assuming that the LSPs subsequently do not undergo any significant self-annihilation, and since in general $h^2 = m Y = m n/s$, this can be recast into

$$\frac{b}{m} = \frac{m_n Y_b}{m Y} = \frac{m_n}{m} \frac{1}{f_B N}; \quad (7)$$

where m_n denotes the mass of a nucleon and m the mass of the neutralino. It is clear that Eq. (7) implies $b = 1$ to be less than one but not > 1 . In the EMD scenario, not only both types of matter are simultaneously produced but also a right ratio of their abundances is predicted.

Unfortunately, this attractive picture runs into a serious problem of over-producing neutralinos, as noticed already by EMD themselves [5, 28]. Since $Y_m \sim 3.9 \times 10^{-10}$, $h^2 = 0.11 \text{ GeV}$, one can rewrite Eq. (7) as

$$m \sim 1.5 \text{ GeV} \quad \frac{3}{N} \quad \frac{1}{f_B} \quad \frac{0.86}{n_b/s} \quad \frac{10^{10}}{h^2} \quad \frac{1}{0.11}; \quad (8)$$

In order to remain consistent with the values of n_b and h^2 derived from observations, the neutralino mass has to be $O(1 \text{ GeV})$ which, in the MSSM, is excluded by LEP [10]. Here, we have neglected a possible contribution to the LSP density from freeze-out. If it were significant, the problem would become only worse. Moreover, the condition (8) puts into question an attractive AD mechanism in a large class of SUGRA models.

To circumvent these problems, one has to review assumptions in the above discussions, namely:

- (i) LSPs produced in Q -ball decay do not annihilate;
- (ii) The LSP is the lightest neutralino of the MSSM.

If we relax assumption (i), the neutralino LSP with the mass of $O(10^2 \text{ GeV})$, consistent with LEP, becomes acceptable. Indeed, allowing for subsequent LSP self-annihilation, the LSP density will be reduced by $h v_i T_d$ [29]. If the cross section for the LSP (self-)annihilation is large enough, e.g., when the LSP is Higgsino or Wino-like [29], the relation between b and DM is lost. One interesting exception is when the energy density of universe is dominated by Q -ball itself [30].

Alternatively, if we lift assumption (ii), the $b \{ DM \}$ relation may be preserved. One way is to consider, e.g., models with the Higgs sector supplemented by a singlet. If its fermionic partner, the singlino, is the LSP then, for some specific choices of parameters [31], it could be possible to circumvent the LEP bound and perhaps also to suppress the LSP abundance from freeze-out. In the rest of the paper, we will investigate axino LSP as DM.

3. Axino dark matter from Q -balls.

In general, axinos, like gravitinos, can be produced in both thermal processes (TP) and in nonthermal processes (NTP), e.g., in late decays. TP consists of the scatterings and the decays of particles in the thermal bath. NTP is given by the decay of the Next-to-LSP (NLSP) relic (which, for simplicity, we assume to be the neutralino) from freeze-out or from the decay of Q -balls in our scenario.

The relevant Boltzmann equations can be written as

$$\dot{n} + 3Hn = h v_i (n^2 - n_{eq}^2) + Q n; \quad (9)$$

$$\dot{n}_a + 3Hn_a = h v_{ij} n_i n_j + h v_{i1} n_i + n; \quad (10)$$

where $h v_i$ is the usual neutralino freeze-out term, Q denotes the contribution to $\dot{n}$ from Q -balls decay, i is the decay rate of the neutralino, $h v_{(i+j) \rightarrow a} \equiv i_{ij}$ and $h v_{(i) \rightarrow a} \equiv i_i$ are the scattering cross section and the decay rate for the thermal production of axinos.

The total NLSP abundance is given by

$$Y = N f_B \frac{n_b}{s} + Y^{TP} \quad (11)$$

where $Y^{TP} \sim H = s_{\tilde{\chi} = m} = T_f = h v_{ann}$, as usual. Since $n_a = n$, owing to R-parity conservation, the resulting number density of axino is given by

$$Y_a = Y_a^{NTP} + Y_a^{TP}; \quad (12)$$

with

$$Y_a^{NTP} = 1 - 10^{10} N \quad \frac{f_B}{1} \quad \frac{n_b/s}{1 - 10^{10}} + Y^{TP}; \quad (13)$$

where Y_a^{TP} denotes the axinos produced by thermal processes. Since typically $Y^{TP} \sim 10^{11}$, one can see that nonthermal production of axinos due to the thermal relic

NLSPs decay can easily be negligible compared to that from Q -ball production, and its contribution to $a h^2$ is further suppressed by the ratio $m_a = m$. The thermally produced axino $\tilde{\chi}_a^{\text{TP}}$ also can be subdominant, say $\tilde{\chi}_a^{\text{TP}} < 10^{11}$, for $T_R < 100 \text{ GeV}$ [15, 32]. Hence, the axino dark matter density is estimated as

$$\frac{a h^2}{0.11}, \quad \frac{m_a}{1.5 \text{ GeV}}, \quad \frac{N f_B}{3}, \quad \frac{b h^2}{0.02} : \quad (14)$$

One can see that the baryon asymmetry and the dark matter abundance are readily linked.

As mentioned above, axino mass is strongly model-dependent; in particular, it critically depends on how the visible and hidden sectors are coupled [11, 12, 13]. At tree level, either $m_a = 0$ ($m_{3=2}$) or $O(m_{3=2}^2 = f_a) = O(\text{keV})$. However, in the latter case, trilinear terms can generate a substantial 1-loop correction of order $f_0^2 = 8^{-2} A$, where f_0 is the Yukawa coupling of the heavy quark to a singlet field containing the axion, which gives m_a in the range of a few tens of GeV or less [12, 13].

The final check point is the compatibility with successful predictions of BBN. However, this is not really a problem for the axino LSP because its interactions

are less suppressed than those of the gravitino, roughly by $(M_P = f_a)^2$ and, so long as the NLSP is heavier than about 150 GeV, axinos are produced before the time of BBN [15]. In contrast, the gravitino LSP would be produced in late NLSP neutralino decays, which faces strong constraints from BBN [33, 34].

4. Conclusions.

We have shown that the framework with cold dark matter axino LSP produced in Q -ball decays can explain the abundance of dark matter and the baryon asymmetry simultaneously and may be an answer to the $b - D_M$ coincidence. In this approach, the similarity between b and D_M is explained by basically only the axino mass of order of (sub-)GeV. The essential property of Q -ball decays is that one can predict the number of SU(5) particles per one baryonic charge from Q -ball N. A characteristic feature is low reheat temperature T_R of 10^2 GeV .

Acknowledgments. The work of O.S. is supported by PPARC. We thank the European Network of Theoretical Astroparticle Physics (ENTAP), part of ILIAS, under contract number RII3-CT-2004-506222 and the EC 6th Framework Programme MRTN-CT-2004-503369 for partial financial support.

-
- [1] D. N. Spergel et al, astro-ph/0603449.
 - [2] D. Hooper, J. March-Russell and S. M. West, Phys. Lett. B 605, 228 (2005).
 - [3] J. McDonald, JCAP 0701, 001 (2007).
 - [4] K. Enqvist and J. McDonald, Nucl. Phys. B 538, 321 (1999).
 - [5] K. Enqvist and J. McDonald, Phys. Lett. B 440, 59 (1998).
 - [6] I. A. Heck and M. Dine, Phys. Lett. B 145, 181 (1984).
 - [7] K. Enqvist and A. Mazumdar, Phys. Rept. 380, 99 (2003).
 - [8] A. Kusenko and M. E. Shaposhnikov, Phys. Lett. B 418, 46 (1998).
 - [9] A. Kusenko, L. C. Loveridge and M. E. Shaposhnikov, JCAP 0508, 011 (2005).
 - [10] W. M. Yao, et al. [Particle Data Group], J. Phys. G 33 (2006) 1.
 - [11] E. J. Chun, J. E. Kim and H. P. Nilles, Phys. Lett. B 287, 123 (1992); J. Chun and A. Lukas, Phys. Lett. B 357, 43 (1995).
 - [12] P. Moxhay and K. Yamamoto, Phys. Lett. B 151, 363 (1985).
 - [13] T. Goto and M. Yamaguchi, Phys. Lett. B 276, 103 (1992).
 - [14] L. Covi, J. E. Kim and L. Roszkowski, Phys. Rev. Lett. 82, 4180 (1999).
 - [15] L. Covi, H. B. Kim, J. E. Kim and L. Roszkowski, JHEP 0105, 033 (2001).
 - [16] J. E. Kim, A. Masiero and D. V. Nanopoulos, Phys. Lett. B 139, 346 (1984); S. A. Bionetto, F. Gabbiani and A. Masiero, Phys. Lett. B 222, 433 (1989) and Phys. Rev. D 49, 3918 (1994); K. Rajagopal, M. S. Turner and F. Wilczek, Nucl. Phys. B 358, 447 (1991).
 - [17] K. W. Ng, Nucl. Phys. B 321, 528 (1989).
 - [18] M. Dine, L. Randall and S. Thomas, Nucl. Phys. B 458, 291 (1996).
 - [19] K. Enqvist and J. McDonald, Phys. Lett. B 425, 309 (1998).
 - [20] K. Enqvist, A. Jokinen and J. McDonald, Phys. Lett. B 483, 191 (2000).
 - [21] S. Kasuya and M. Kawasaki, Phys. Rev. D 74, 063507 (2006).
 - [22] R. Allahverdi, B. A. Campbell and J. R. Ellis, Nucl. Phys. B 579, 355 (2000).
 - [23] A. Anisimov and M. Dine, Nucl. Phys. B 619, 729 (2001).
 - [24] S. Kasuya and M. Kawasaki, Phys. Rev. D 62, 023512 (2000).
 - [25] R. Banerjee and K. Jedamzik, Phys. Lett. B 484, 278 (2000).
 - [26] A. G. Cohen, S. R. Coleman, H. Georgi and A. Manohar, Nucl. Phys. B 272, 301 (1986).
 - [27] S. Kasuya and M. Kawasaki, Phys. Rev. D 61, 041301 (2000).
 - [28] K. Enqvist and J. McDonald, Nucl. Phys. B 570, 407 (2000).
 - [29] M. Fujii and K. Hamaguchi, Phys. Lett. B 525, 143 (2002); Phys. Rev. D 66, 083501 (2002).
 - [30] M. Fujii, T. Yanagida, Phys. Lett. B 542, 80 (2002).
 - [31] R. Flores, K. A. Olive and D. Thomas, Phys. Lett. B 245, 509 (1990).
 - [32] L. Covi, L. Roszkowski, R. Ruiz de Austri and M. Smal, JHEP 0406, 003 (2004).
 - [33] L. Roszkowski, R. Ruiz de Austri and K.-Y. Choi, JHEP 0508, 080 (2005).
 - [34] O. Seto, Phys. Rev. D 73, 043509 (2006).