

Heavy meson semileptonic decays in two dimensions in the large N_c

Jorge M ondejar^a

^aDept. d'Estructura i Constituents de la Matèria, U.B. Barcelona,
Diagonal, 647, E-08028 Barcelona, Spain

We study QCD in $1+1$ dimensions in the large N_c limit using light-front Hamiltonian perturbation theory in the $1/N_c$ expansion. We use this formalism to exactly compute hadronic transition matrix elements for arbitrary currents at leading order in $1/N_c$, which we use to write the semileptonic differential decay rate of a heavy meson and its moments. We then compare with the results obtained using an effective field theory approach based on perturbative factorization, with the intention of better understanding quark-hadron duality. A very good numerical agreement is obtained between the exact result and the result using effective theories.

1. Introduction

The results presented here are part of a work done in collaboration with Antonio Pineda and Joan Rósp [1].

Asymptotic freedom can be seen as the first example of factorization between high and low energies, since it dictates that Green functions at high Euclidean energies (Q^2) can be described by perturbation theory up to corrections suppressed by powers of $1/N_c$ over Q . Therefore, the use of the operator product expansion (OPE) in processes where the relevant momentum scale is large and Euclidean is safe. This is quite restrictive, since, in most of the cases, it can only be tested with experiment through dispersion relations, which involve measurements up to arbitrarily high energies. What one usually does is to try to directly apply the same perturbative factorization techniques to observables living in the Minkowski regime. In practice this means to perform the analytic continuation of approximate perturbative results obtained in the Euclidean region to the Minkowski region, but such calculations do not come from first principles. This problem affects the OPE and effective field theories that are built using perturbative factorization techniques aiming to factorize high from low energies, and it is usually stated as duality violations. We will follow here the definition of [2] for duality violations.

One can quantify the discrepancy between the

exact result and the one using perturbative factorization in the large N_c limit of QCD [3]. In this case one finds a clear discrepancy between both results in the physical cut of the Green functions, where one has infinitely narrow resonances on the one hand and an smooth function on the other. This can be further quantified in the 't Hooft model [4], which we will consider in what follows.

The specific observable we use to illustrate this discussion is the differential semileptonic inclusive decay of a heavy meson: $H_Q \rightarrow X l$. The duality violations in this case are maximal, but if we consider the Mellin moments of the differential decay rate we find that there is a very good agreement between the exact and the perturbative result.

In sec. 2 we analyze QCD₁₊₁ in the light front. In sec. 3 we compute the hadronic differential decay rate and its moments. In sec. 4 we construct an effective theory to compute these quantities at one loop, and compare the two results. Finally, in section 5 we present our conclusions.

2. QCD₁₊₁ in the light front

The QCD Lagrangian is given by

$$L = \frac{1}{4} G^a_{ij} G^{aj} + \sum_i \bar{\psi}_i (i \not{D} - m_i) \psi_i; \quad (1)$$

where $D = \partial + igA$ and i labels the flavor.

in the past, we will follow here the work of Bigi et al. [5]. The authors considered the flavour changing weak interaction

$$L_{\text{weak}}^V = \frac{G}{2} c_Q \bar{l}_a \gamma_\mu l_b : \quad (11)$$

The total decay width can be written as

$$\Gamma_{H_Q} = \frac{G^2}{M_{H_Q}^2} \int \frac{d^2q}{(2\pi)^2} \text{Im} \langle q | \text{Im} T | q \rangle ; \quad (12)$$

where $\langle q |$ and $T | q \rangle$ are defined as

$$\langle q | = i h_0 \int \bar{l}_a(x) \bar{l}_b(x) \bar{l}_b(0) \bar{l}_a(0) \delta^2(x) ; \quad (13)$$

$$T | q \rangle = i h_{H_Q} \int \bar{Q}(x) c(x) c(0) Q(0) \delta^2(x) ;$$

and their Fourier transform as

$$\langle q | = \int d^2x e^{iqx} \langle q | (x) ; \quad (14)$$

$$T | q \rangle = \int d^2x e^{iqx} T | q \rangle (x) ;$$

The leptonic tensor can be easily calculated; to calculate the hadronic tensor we just have to insert a complete set of intermediate states: in the large N_c limit this set will consist exclusively of mesons, so we can use the transition matrix elements we found previously. In the end, the differential decay rate reads (the details can be found in [1])

$$\frac{d\Gamma_{\text{hadr}}}{dx} = \sum_{M_n} \frac{G^2 M_{H_Q}^2 M_n^2}{4 M_{H_Q}^2} \quad (15)$$

$$\int_0^1 dz \langle n | T_{H_Q}(z) | n \rangle^2 x \left(1 + \frac{M_n^2}{M_{H_Q}^2} \right) ;$$

where

$$x = \frac{q^+}{P_{H_Q}^+} ; \quad (16)$$

The moments are defined as

$$M_N = \int_0^1 dx x^N \langle n | T_{H_Q}(z) | n \rangle ; \quad (17)$$

We can insert (15) in this expression right away, but since our interest here is to compare the result

with that of an effective theory, we will give an alternative expression for the moments which is valid up to $N = 2$ (again, the details of the calculation, and the approximations made, can be found in [1]),

$$M_N^{\text{hadr}} = \frac{G^2 M_{H_Q}^2 m_{Q;R}^2}{4 M_{H_Q}^2} \int_0^1 \frac{dx}{x^2} x^N \langle n | T_{H_Q}(x) | n \rangle ; \quad (18)$$

where $m_{x;R}^2 = m_x^2 - x^2$. This formula has a precision of $O(x^2 = m_Q^2)$ when $N = 1$ which reduces to $O(x = m_Q)$ when $N = 2$.

4. Effective theory

To construct the effective theory we switch to a partonic picture, in which the decay of the initial meson becomes just the decay of the heavy quark into a light quark. The decay is given by the imaginary part of the diagrams shown in Figure 1.




Figure 1. Diagrams involved in the construction of the effective theory at one loop. The dotted lines represent the outgoing leptons.

We work in the kinematics in which the momentum carried away by the leptons is

$$q^+ = x p_Q^+ ; \quad q^- = 0 ; \quad (19)$$

This means that the light outgoing quark resulting from the decay has the following momentum :

$$p_c^+ = p_Q^+ (1 - x) ; \quad p_c^- = p_Q^- ; \quad (20)$$

The important point is that $p_c = p_Q = m_Q$, which in our light front quantization frame means that the propagating light quark in the diagram has a very large "energy", and that the process happens at very short "times". So, we can safely integrate out the light quark from our lagrangian. The resulting lagrangian will be essentially that of HQET, plus a local vertex in x^+ , the imaginary part of which will represent the decay process. The construction of the effective theory amounts then to computing in full QCD the diagrams in Figure 1, and then matching to an effective vertex in which the intermediate light quark will have disappeared. The effective vertex is

$$L_I = \frac{G^2}{2} (\bar{\psi}^+) \frac{m_{Q;R}}{i\partial^+} Q_+ \frac{1}{i\partial^+} \frac{1}{\frac{m_{C;R}^2}{m_{Q;R}^2} i\partial^+} \frac{m_{Q;R}}{i\partial^+} Q_+ (\psi^+)^y; \quad (21)$$

where ψ is a pseudoscalar massless field that represents the (massless) outgoing leptons. The differential decay rate is then given by

$$\frac{d^{\text{pert}}}{dx} = \frac{1}{M_{H_Q}} \frac{1}{2(2-x)^2} \frac{G^2}{2} (M_{H_Q} x)^2 2 \text{Im} T_{\text{eff}} \quad (22)$$

$$= \frac{G^2 M_{H_Q} m_{Q;R}^2 m_{C;R}^2}{4 m_{Q;R}^2} \frac{m_{Q;R}^2}{M_{H_Q}^2} \frac{1}{x} \frac{2}{M_{H_Q}} \frac{B}{\partial^+} \frac{x}{1 - \frac{m_{C;R}^2}{m_{Q;R}^2}} \quad (23)$$

Comparing with (15) we see the maximal duality violation: one result is a sum of deltas whereas the other is a smooth function. However, if we compute the moments, we find

$$M_N^{\text{pert}} = \frac{G^2 M_{H_Q} m_{Q;R}^2}{4 M_{H_Q}^2} \frac{1}{\int_0^1 dx x^{N-2} \frac{2}{M_{H_Q}} (x); \quad (23)$$

the same expression as (18), with the same precision. Thus, up to the precision we are working with, there are no duality violations for the moments. Actually, a numerical analysis (see [1]) shows that the numerical agreement is very good up to high moments. In [5] it is shown for the inclusive decay width that differences appear at

$O(1/m_Q^9)$ (they are strongly suppressed, but they are there). In [1] it is also shown that, although very good results are found for the moments, things don't look so well if we average over smaller ranges. Namely, for

$$\int_{x_n}^{x_n+x} \frac{d}{dx} dx; \quad (24)$$

where x_n is any x which satisfies the delta in (15), we find that, first, we have to re-tune the value of x for the hadronic and perturbative results to agree at leading order, but even then, they differ at $O(1/m_Q^2)$.

5. Conclusions

To sum up, we have computed the differential semileptonic inclusive decay of a heavy meson and its moments, through both a hadronic calculation and a perturbative calculation (using an effective theory), and found that, whereas a comparison between the two results is impossible for the differential decay rate, the moments show no duality violations with the precision we have worked with; however, other observables, like the one defined in (24) don't work so well. So, in conclusion, effective theories can only be a good approximation for inclusive observables on which one averages over a large fraction of the final bound states. They are not suited for point-to-point comparisons, or comparisons between arbitrarily smeared functions.

REFERENCES

1. J. M ondejar, A. Pineda and J. Rojo, J. High Energy Phys. 09, 060 (2006) [arXiv:hep-ph/0605248].
2. M. A. Shifman, arXiv:hep-ph/0009131.
3. G. 't Hooft, Nucl. Phys. B 72, 461 (1974).
4. G. 't Hooft, Nucl. Phys. B 75, 461 (1974).
5. I. I. Y. Bigi, M. A. Shifman, N. Uraltsev and A. I. Vainshtein, Phys. Rev. D 59, 054011 (1999) [arXiv:hep-ph/9805241].