Resolving the octant 23 degeneracy by neutrino oscillation experiments HiroshiNunokawa^a

^a Departam ento de F sica, Pontif cia Universidade Catolica do Rio de Janeiro, C.P. 38071, 22452-970, Rio de Janeiro, Brazil

W e discuss how and to what extent the degeneracy associated with $_{23}$, if it's not maximal, can be resolved by future oscillation experiments which utilize conventional neutrino beam from accelerator and/or reactor neutrinos.

1. Introduction

Physics of neutrinos will soon be entering into the era of precision physics. Proposed projects such as T2K [3] and NO A [4] experiment will be able to measure the mixing parameters responsible for atm ospheric neutrino oscillation, $\sin^2 2_{23}$ and m $_{23}^2$ with a few percent uncertainty. Moreover, these experiments can probe $\sin^2 2_{13}$ down to 10⁻² or smaller. If non-zero $_{13}$ will be established, the future phases of these experiments will be aimed to determ ine the CP violating phase as well as the mass hierarchy [3,4] (see also [5]).

It has been known that in order to perform precise determ inations of neutrino mixing param – eters including the CP phase and mass hierarchy, one must confront with the problem of so called parameter degeneracy. There are 3 types of such degeneracy, octant [6], intrinsic [7] and sign m² [8] degeneracy. Here we focus only on the octant degeneracy, which is decoupled from the other ones with a good approximation in our experimental set up. We consider only experiments based on the conventional neutrino beam from accelerator and reactor neutrinos.

Suppose that $_{23}$ is di erent from =4. D isappearance ! experiment can determ ine quite precisely the value of $\sin^2 2_{23}$ with 1% uncertainty [3] but this does not allow us to know in which octant $_{23}$ lives, and leads, at rst approximation, ignoring $_{13}$, to the following 2 degenerate solutions, $\sin^2 _{23} = 1$ 1 $\sin^2 2_{23} = 2$. For example, $\sin^2_{23} = 0.4 \text{ or } 0.6 (0.45 \text{ or } 0.55)$ if $\sin^2 2_{23} = 0.96 (0.99)$. How and to what extent this can be resolved is the topics of this talk.

2. C om bining A ccelerator and R eactor

Let us rst discuss the possibility to resolve this degeneracy by combining accelerator and reactor neutrinos [1], based on the suggestion in [10]. As a concrete example, we consider the second phase of the T2K experiment [3] with upgraded beam powerof4MW and Hyper-Kamiokande (HK) detector, and high statistics second generation reactor experiment, e.g. Angra project [9]. We x the mixing parameters relevant for solar neutrinos as $m_{21}^2 = 8.0$ 10 ⁵ eV² and $sin^2_{12} =$ 0.31, and the atmospheric m $^2_{23}$ = 2.5 10 3 eV^2 . The exposures for accelerator are assumed to be 2 (6) years of neutrino (anti-neutrino) running with HK whose ducial volume is 0.54 Mt, whereas for the reactor we assume an exposure of 10 GW kt yr, which is de ned as a product of reactor power, detector volum e and running time.

In Fig. 1 we show the process of how the octant degeneracy can be resolved by combining the results from accelerator and reactor. Figs. 1(a) and (b) show that by combining the ! and

! $_{\rm e}$ modes from accelerator, we can get 2 separated allowed regions where one of them corresponds to the fake solution. Then if we add the result of the measurement of $_{13}$ form reactor experiment, which is shown in Fig. 1(c), we can eliminate the fake solution and end up with the true one as shown in Fig. 1(d).

Talk presented at NOW 2006, Otrant, Lecce, Italy, September 9-16, 2006, based on [1,2]





Figure 2. Region of parameters where ₂₃ degeneracy can be resolved. Upper (lower) panel corresponds to conservative (optim istic) choice of the system atic uncertainties. Taken from [1].

The idea is to detect the \solar" term de ned as $\cos^2_{23} \sin^2 2_{12}$ (m $_{21}^2 L=4E$)², which is possible if the detector is placed far enough. Since this term is proportional to \cos^2_{23} , in principle, one can tell in which octant $_{23}$ lies. In Fig. 3, we show the expected number of m uon and electron events as a function of reconstructed neutrino energy at Kam ioka and Korea for $\sin^2_{23} = 0.4$ and 0.6. We observe that at Korea for the electron events, appreciable di erence exist between the data corresponding to these 2 degenerate solutions, which can tell us the correct octant.

In Fig. 4 we show the region of parameters where the octant degeneracy can be resolved by the 2 detector method. Let us note that resolving power (sensitivity) does not depend much on the value of $_{13}$ which is very dimensity from the



Figure 1. (a) Region allowed by the disappearance (2 approximately horizontal bands) and by the appearance (the other curved band) accelerator measurement. (b) The regions that remain allowed when results shown in (a) are combined. (c) The regions allowed by reactor measurement. (d) The regions allowed after combining all the results from accelerator and the reactor experiments. Taken from [1].

We show in Fig. 2 the parameter region of $\sin^2 2_{13}$ and \sin^2_{23} where the octant degeneracy can be resolved. The degeneracy can be resolved for $\sin^2 2_{23} < 0.96$ at 2 CL for $\sin^2 2_{13} > 0.05$ (see upper panel). We con rm ed the strong dependence of sensitivity on $_{13}$ expected in [10].

3. A ccelerator based 2 detector m ethod

Next we discuss another possibility based on [11]. In the second phase of T 2K experiment (HK with 4 MW beam), in order to improve the sensitivity to them asshierarchy determ ination, it was suggested to place the second identical detector at K orea with baseline of 1050 km, in addition to the one at K am ioka. Here we assume 2 detectors are not only identical but also receive the neutrino beam with the same energy spectrum by choosing the same o axis angle (2.5 degree in this case). It was shown that this experimental



Figure 3. Expected number of electron (upper panels) and muon (lower panels) events as a function of the reconstructed neutrino energy for $\sin^2_{23} = 0.40$ (open circles) and 0.60 (lled circles). Taken from [2].



Figure 4. Region of parameters where 23 degeneracy can be resolved. Light gray (dark gray) area corresponds to 2 (3) signi cance. 0.27 M ton detectors both in K am ioka and K orea and 4 years running with neutrino beam and another 4 years with anti-neutrino beam are assumed. In left (right) 2 panels, the sensitivity is de ned so that the experiment can resolve the octant degeneracy for any (half) values of the CP phase . Taken from [2].

previous case shown in Fig. 2 where the sensitiv-

3

ity is signi cantly worse (better) when $_{13}$ is small (large). We conclude that in this method, the octant degeneracy can be resolved for $\sin^2 2_{23} < 0.97$ at 2 CL even for very small values of $_{13}$.

4. Sum m ary

The expected sensitivities by the 2 m ethods we discussed show very di erent dependence on $_{13}$. If $_{13}$ is larger, close to the present lim it, the m ethod of com bining accelerator and reactor would give better sensitivity whereas for sm aller $_{13}$, the accelerator based 2 detector m ethods would be better, and therefore, these 2 m ethods are com plem entary to each other. See R efs. [1,2] for details.

A cknow ledgm ent. The author would like to thank K. Hiraide, T. Kajita, H. Minakata, T. Nakaya, S. Nakayam a, H. Sugiyam a, W. J. C. Teves and R. Zuaknovich Funchal for collaborations. The author acknow ledges also CNPq and FAPERJ for nancial support.

REFERENCES

- K. Himaide et al., Phys. Rev. D 73 (2006) 093008 [arX iv hep-ph/0601258].
- T.Kajita et al., arX iv hep-ph/0609286, to be published in Phys.Rev.D.
- Y. Itow et al., arX iv hep-ex/0106019.
 For an updated version, see: http://neutrino. kek.jp/hfnu/bi/biv2.030528.pdf
- D. S. Ayres et al. NOvA Collaboration], arX iv:hep-ex/0503053.
- 5. 0.M ena et al, arX iv hep-ph/0609011.
- G.Fogliand E.Lisi, Phys. Rev. D 54 (1996) 3667 [arX iv hep-ph/9604415].
- J.Burguet-Castellet al, Nucl. Phys. B 608 (2001) 301 [arX iv hep-ph/0103258].
- H.M inakata and H.Nunokawa, JHEP 0110 (2001) 001 [arX iv hep-ph/0108085].
- J.C.Anjos et al, Nucl. Phys. Proc. Suppl. 155 (2006) 231 [arX iv hep-ex/0511059].
- 10. H. M inakata et al, Phys. Rev. D 68 (2003) 033017 [Erratum - ibid. D 70 (2004) 059901] [arX iv hep-ph/0211111].
- 11. M. Ishitsuka et al, Phys. Rev. D 72 (2005) 033003 [arX iv hep-ph/0504026].