

DEBYE SCREENING IN THE HOT NON-ABELIAN PLASMA

N.O.AGASIAN and YU.A.SIMONOV

Institute of Theoretical and Experimental Physics, Moscow, 117218, Russia
E-mail: agasian@itep.ru

The Debye mass m_D is computed nonperturbatively in the deconfined phase of QCD, where chromomagnetic confinement is known to be present. The latter defines m_D to be $m_D = c_D \frac{P_s}{s}$, where $c_D = 2.06$ and $s = s(T)$ is the spatial string tension. The resulting magnitude of $m_D(T)$ and temperature dependence are in good agreement with lattice calculations.

Keywords: quark-gluon plasma; background perturbation theory.

1. Introduction

The screening of electric fields in QCD was originally considered in analogy to QED plasma, where the Debye screening mass was well understood¹, and the perturbative leading order (LO) result for QCD was obtained long ago², $m_D^{(LO)} = \frac{N_c}{3} + \frac{N_f}{6} g^2 T$. Another difficulty of the perturbative approach is that the gauge-invariant definition of the one-gluon Debye mass is not available. Recently the calculations of $m_D(T)$ was computed on the lattice for $N_f = 0; 2; 3; 4; 5$ using the free-energy asymptotics.

The purpose of our talk is to provide a gauge-invariant and a nonperturbative method, which allows to obtain Debye masses in a rather simple analytic calculational scheme. In what follows we use the basically nonperturbative approach of Field Correlator Method (FCM)^{6;7} and Background Perturbation Theory (BPT) for nonzero T ^{8;9;10} to calculate $m_D(T)$ in a series, where the first and dominant term is purely nonperturbative¹¹

$$m_D(T) = M_0(T) + \text{BPT series} \quad (1)$$

Here $M_0(T)$ is the gluon lump mass due to chromomagnetic confinement in 3d, which is computed¹¹ to be $M_0(T) = c_D \frac{P_s(T)}{s(T)}$, with $s(T)$ being the spatial string tension and $c_D = 2$ for $N_c = 3$. The latter is simply expressed in FCM through chromomagnetic

correlator^{12;13}, and can be found from the lattice data^{14;15}. Therefore $M_0(T)$ is predicted for all T and can be compared with lattice data.

2. Background Perturbation Theory

Theory for the thermal Wilson loop

One starts with the correlator of Polyakov loops $\langle L(0)L^\dagger(r) \rangle$

$$\langle L(0)L^\dagger(r) \rangle = \frac{1}{N_c^2} \exp \left(-\frac{F_1(r;T)}{T} \right) + \frac{N_c^2}{N_c^2} \exp \left(-\frac{F_8(r;T)}{T} \right) :$$

The first term contributing to the free energy F_1 of the static $Q\bar{Q}$ -pair in the singlet color state, the second to the octet free energy F_8 .

A convenient way to define Debye mass was suggested in^{3;4}

$$F_1(r;T) = F_1(1;T) - \frac{4}{3} \frac{s(T)}{r} e^{-m_D(T)r}$$

Accordingly one ends for F_1 with the thermal Wilson loop of time extension $\tau = 1/T$ and space extension r ,

$$\exp(-F_1(r;T)) = \langle W(r;T) \rangle = \frac{1}{N_c} \int \mathcal{D}A \exp(i \int_c A dz) \quad (2)$$

arXiv:hep-ph/0612233v2 9 Jul 2007

Eq.(2) is the basis of our approach. In what follows we shall calculate however not F_1 , which contains all tower of excited states over the ground state of heavy quarks QQ , but rather the static potential $V_1(r;T)$, corresponding to this ground state. As a result the Debye mass m_D is calculated gauge-invariantly in terms of the thermal Wilson loop.

Separating, as in BPT, the field A into NP background B and valence gluon field a

$$A = B + a \quad (3)$$

one can assign gauge transformations as follows

$$B \rightarrow U^+ (B + \frac{i}{g} \partial) U; \quad a \rightarrow U^+ a U \quad (4)$$

As a next step one inserts (3) into (2) and expands in powers of ga , which gives an expansion for the Wilson loop

$$\begin{aligned} \langle W(r; \tau) \rangle &= \langle W^{(0)}(r; \tau) \rangle_{i_B} + \\ &+ \langle W^{(2)}(r; \tau) \rangle_{i_B, a} + \dots; \end{aligned} \quad (5)$$

and $\langle W^{(2)} \rangle$ can be written as

$$\begin{aligned} \langle W^{(2)} \rangle_{i_B, a} &= \frac{(ig)^2}{N_c} \int dx dy \\ \text{trP} \left(\int_{xy}^Y \right) & \text{ha}(x) a(y) i_a \left(\int_{xy}^{\tilde{a}^Y} \right) i_B : \end{aligned} \quad (6)$$

Here $\left(\int_{xy}^Q \right)$ and $\left(\int_{xy}^{\tilde{a}^Y} \right)$ are parallel transporters along the pieces of the original Wilson loop $W(r; \tau)$, which result from the dissection of the Wilson loop at points x and y . Thus the Wilson loop $W^{(2)}(r; \tau)$ is the standard loop $W^{(0)}(r; \tau)$ augmented by the adjoint line connecting points x and y . It is easy to see using (4), that this construction is gauge invariant.

For OGE propagator one can write the path integral Fock-Feynman-Schwinger (FFS) representation for nonzero T

$$G(x; y) = \text{ha}(x) a(y) i_a$$

$$\begin{aligned} &= \int_0^Z ds \int_0^Z (D^4 z)_{xy}^w \exp(-K) \text{adj}(C_{xy}) \\ &P_F \exp(2ig \int_0^Z F(z) dz) ; \quad (7) \\ &\text{adj}(C_{xy}) = P \exp(ig \int_{C_{xy}} B dz) \end{aligned}$$

where the open contour C_{xy} runs along the integration path in (7) from the point x to the point y and $K = \frac{1}{4} \int_0^R (z)^2 dz$. The $(D^4 z)_{xy}^w$ is a path integration measure with boundary conditions $z(\tau=0) = x$ and $z(\tau=s) = y$ and with all possible windings in the Euclidean temporal direction (this is marked by the superscript w).

We must now average over B the geometrical construction obtained by inserting (7) into (6), i.e.

$$\langle \left(\int_{xy}^Y \right) \text{adj}(C_{xy}) \left(\int_{xy}^{\tilde{a}^Y} \right) i_B \rangle = \langle W_{xy}(r; \tau) \rangle_{i_B} \quad (8)$$

For $T > T_c$ one can apply the nonabelian Stokes theorem, which yields the area law⁶ for distances $r \gg r_g$, r_g gluon correlation length, $r_g \approx 0.2 \text{ fm}$

$$\langle W_{xy}(r; \tau) \rangle_{i_B} = \exp(-S_{adj}^{H_{g1}}); \quad (9)$$

where S_{g1}^H is the area of the space-like projection of gluon-deformed piece of surface S_{g1} , and S_{adj}^H is the adjoint spatial string tension, $S_{adj}^H = (9-4) s$ for $SU(3)$. s is the fundamental spatial string tension.

As the result one obtains exactly the form containing the gluon loop Green's function

$$\begin{aligned} \langle W^{(2)} \rangle_{i_B, a} &= (ig)^2 C_2(f) \\ &\int_0^Z dx_4 \int_0^Z dy_4 G_{44}(r; t_4); \end{aligned} \quad (10)$$

where $t_4 = x_4 - y_4$, $C_2(f) = (N_c^2 - 1)/2N_c$ and $G_{44}(r; t_4)$ is

$$G_{44}(r; t_4) = \int_0^Z ds \int_0^Z (D^4 z)_{xy}^w$$

$$\exp(-K) \exp\left(\frac{H}{\text{adj}} S_{gl}^H\right); \quad (11)$$

Thus the gluon Green's function in the confined phase becomes a gluelump Green's function, where the adjoint source trajectory is the projection of the gluon trajectory on the Wilson loop plane. Gluelump is the system when a gluon is connected by the string to the Wilson loop plane. String is made of chromomagnetic field only. Thus the problem reduces to the calculation of the gluelump Green's function

$$G(x; y) = \langle \text{Tr} \int_{\mathcal{X}} \exp(-H_n(r)) \rangle$$

The lowest eigenvalue of M_0 Hamiltonian yields the static QQ potential

$$V_1^{(1)}(r; T) = \frac{N_c^2}{2N_c} \frac{1}{r} e^{M_0 r}$$

As a result one finds $m_D = M_0$

$$m_D = (2.82 - 5.08 \frac{e}{s}) \frac{p}{s} - 2.06 \frac{p}{s}$$

This is the central result of our talk. Note, that no free parameters are used, $\frac{e}{s}$ is taken from glueball spectrum and BFKL.

3. Numerical results

One can now compare our prediction for $m_D(T) = c_D \frac{p}{s(T)}$ with the latest lattice data⁵. At temperatures $T > T_{dr}$ (1.2 - 1.5) T_c , (where T_{dr} is the temperature of dimensionally reduction and physical justification for resorting to dimensionally reduced regime at $T' = \frac{p}{s(T)}$ was given in^{13,16}) the spatial string tension is chosen in the form^{14,15}

$$\frac{p}{s(T)} = c g^2(T) T; \quad (12)$$

with the two-loop expression for $g^2(T)$

$$g^2(t) = 2b_0 \ln \frac{t}{L} + \frac{b_1}{b_0} \ln^2 \frac{t}{L}; \quad (13)$$

where $t = T = T_c$.

The measured¹⁵ spatial string tension in pure glue QCD corresponds to the values of $c = 0.566 - 0.013$ and $L = T_c =$




Fig. 1. Chromoelectric Debye mass $m_D = T$ for 2-flavor QCD (upper lines) and quenched QCD ($N_f = 0$) (lower lines) versus $T = T_c$. Solid lines are calculated using $m_D(T) = 2.09 \sqrt{s(T)}$, where $\sqrt{s(T)}$ corresponds to Eq.(12) with $N_f = 2$ for upper solid line and $N_f = 0$ for lower solid line. Dashed lines are calculated using Eq.(14), $N_f = 2$ (upper line, $N_f = 0$ (lower line). The lattice data are from Okazaki and F. Zantow, Phys. Rev. D 71 114510 (2005).

0.104 - 0.009: On the Fig. 1 are shown lattice data³ and theoretical curves for the Debye mass in quenched ($N_f = 0$) and 2-flavor QCD. Solid lines correspond to our theoretical prediction, $m_D(T) = c_D \frac{p}{s(T)}$, with $c_D = 2.09$ and for $\frac{p}{s(T)}$ we exploit the parameters $c = 0.564; L = 0.104$. The upper solid line is for the Debye mass in 2-flavor QCD, and the lower { for quenched QCD. We note that in computing $m_D(T)$ using (12), (13) all dependence on N_f enters only through the Gell-Mann coefficients b_0 and b_1 . For comparison we display in the Fig. 1 dashed lines for $m_D(T) = T$, calculated with a perturbative-like ansatz⁵.

$$m_D^{Latt}(T) = A_{N_f} \left(1 + \frac{N_f}{6} g(T) T \right); \quad (14)$$

Comparison of our theoretical prediction with the perturbative-like ansatz shows that both agree reasonably with lattice data in the temperature interval $T_c < T < 5T_c$; the agreement is slightly better for our results. At the same time, in (14) a fitting constant is used $A_{N_f} = 1.5$, which is necessary even at $T = T_c$. We expect the accuracy of the first approximation of our approach is around 10%, taking also into account the bias in the

definition of σ_s for the gluon loop. The temperature region near T_c needs additional care because i) the behaviour of $\sigma_s(T)$ deviates from the lattice data and ii) contribution of chromoelectric fields above T_c which was neglected above.

4. Conclusions

We have studied Debye screening in the hot nonabelian theory. For that purpose the gauge-invariant definition of the free energy of the static $Q\bar{Q}$ -pair in the singlet color state was given in terms of the thermal Wilson loop. Due to the chromomagnetic confinement persisting at all temperatures T , the hot QCD is essentially nonperturbative. To account for this fact in a gauge-invariant way the BPTH was developed for the thermal Wilson loop using path-integral FFS formalism. As a result one obtains from the thermal Wilson loop the screened Coulomb potential with the screening mass corresponding to the lowest gluon loop mass. Applying the Hamiltonian formalism to the BPTH Green's functions with the einbein technique the gluon loop mass spectrum was obtained. As a result, we have derived the leading term of the BPTH for the Debye mass which is the purely nonperturbative,

$$m_D(T) = \sigma_s(T); \quad \sigma_s(T) = 2:$$

Acknowledgments

The authors are grateful to F. Karsch for supplying us with numerical data and we thank B.L. Ioé, A.B. Kaidalov, D.Kharzeev and V.A. Novikov for useful discussions. Our gratitude is also to M.A. Tusov for help and useful advices.

The work is supported by the Federal Program of the Russian Ministry of Industry, Science, and Technology No.40.052.1.1.1112, and by the grant of RFBR No. 06-02-17012, and by the grant for scientific schools NS-843.2006.2.

References

1. I. A. Khiezer and S. Peletninski, *ZhETF* 38, 1829 (1960); J.-P. Blaizot, E. Iancu and R. Parwani, *Phys. Rev. D* 52, 2543 (1995).
2. E. Shuryak, *ZhETF* 74, 408 (1978); J. Kapusta, *Nucl. Phys. B* 148, 461 (1979); D. Gross, R. Pisarski and L. Yaffe, *Rev. Mod. Phys.* 53, 43 (1981).
3. O. Kaczmarek and F. Zantow, *Phys. Rev. D* 71, 114510 (2005).
4. M. Doring, S. Ejiri, O. Kaczmarek, F. Karsch and E. Laermann, *hep-lat/0509150*.
5. O. Kaczmarek and F. Zantow, *hep-lat/0512031*.
6. H.G. Dosch, *Phys. Lett. B* 190, 177 (1987); H.G. Dosch and Yu.A. Simonov, *Phys. Lett. B* 205, 339 (1988); Yu.A. Simonov, *Nucl. Phys. B* 307, 512 (1988); A. Di Giacomo, H.G. Dosch, V.I. Shevchenko and Yu.A. Simonov, *Phys. Rept.* 372, 319 (2002).
7. Yu.A. Simonov, *JETP Lett.* 54, 249 (1991); *JETP Lett.* 55, 605 (1992).
8. Yu.A. Simonov, *Phys. At. Nucl.* 58, 309 (1995); N.O. Agasian, *JETP Lett.* 57, 208 (1993); *JETP Lett.* 71, 43 (2000).
9. Yu.A. Simonov and J.A. Tjon, *Ann. Phys.* 300, 54 (2002).
10. E.L. Gubankova and Yu.A. Simonov, *Phys. Lett. B* 360, 93 (1995).
11. N.O. Agasian and Yu.A. Simonov, *Phys. Lett. B* 639, 82 (2006).
12. Yu.A. Simonov, *JETP Lett.* 54, 249 (1991).
13. N.O. Agasian, *Phys. Lett. B* 562, 257 (2003).
14. F. Karsch, E. Laermann and M. Lutgen eier, *Phys. Lett. B* 346, 94 (1995).
15. G. Boyd et al., *Nucl. Phys. B* 469, 419 (1996).
16. N.O. Agasian and K. Zarembo, *Phys. Rev. D* 57, 2475 (1998).