

Recent Models of Neutrino Masses and Mixing

Ambar Ghosal

Saha Institute of Nuclear Physics,
1/AF Bidhannagar, Kolkata 700 064, India

April 16, 2024

1 Introduction

Neutrino physics is now playing a major role to probe physics beyond standard model. Confirmation of tiny but non-zero neutrino mass through neutrino oscillation experiments have thrown light on the structure of leptonic sector. On neutrino mass, there are some direct experimental upper bounds on the individual neutrino masses from the decay of other particles, nuclei and also on the sum of the masses of neutrino from cosmological observations whereas neutrino oscillation experiments put bounds on neutrino mass-squared differences on neutrinos which oscillate into each other. Furthermore, there is an additional bound on the effective mass of the Majorana type neutrino mass coming from the $\beta\beta$ decay. This bound essentially restricts 11 element of the neutrino mass matrix.

2 Experimental Bounds

2.1 Direct bounds

A direct bound on m_e is obtained from the hadronic decay of π^+ as $m_e < 18 \text{ MeV}$ (95% C.L.) and from the μ^+ decay a bound on m_e is coming as $m_e < 170 \text{ KeV}$ (95% C.L.), and from the tritium beta decay a bound on m_e is coming as $m_e < 2.2 \text{ eV}$ (95% C.L.). Future KATRIN experiment [1] will bring down this limit to $m_e < 0.2 \text{ eV}$.

2.2 Indirect Bounds

Bounds on solar neutrino mass splitting if ν_e is oscillated to ν_μ , combining SNO + KamLAND + SAGE + GALLEX [3, 4] neutrino experimental results give $m_{21}^2 = (7.3 - 8.5) \times 10^{-5} \text{ eV}^2$ at 2 σ range with the best fit value $m_{21}^2 = 7.9 \times 10^{-5} \text{ eV}^2$ [2, 5] and the bound on solar neutrino mixing angle θ_{12} is $\sin^2 \theta_{12} = 0.3$ with the 2 σ range is $0.26 - 0.36$ [2, 5, 6]. The global analysis of Atmospheric + K2K [7, 8] experimental results give the best-fit

MINOS data. The best fit value of $\sin^2 \theta_{23} = 0.5$ with the 2σ range $0.38 - 0.64$. The CHOOZ [9] experimental upper bound on θ_{13} is given by $\theta_{13} < 12^\circ$ (at 3σ) with best fit value $\sin^2 \theta_{13} = 0.0$. An additional bound on the sum of the neutrino masses is coming from the observation of WMAP [10] experimental result on cosmic microwave background anisotropies. Combining analysis of 2dF Galaxy Redshift survey, CBI and ACBAR [11], WMAP has determined the amount of critical density contributed due to relativistic neutrinos which in turn gives an upper bound on the total neutrino mass as $m_{\Sigma} = 0.7$ eV at 95% C.L. Another bound is due to β decay which gives the bound on effective Majorana neutrino mass as $\langle m_{ee} \rangle = (0.05 - 0.84)$ eV at 95% C.L. [14] (relaxing the uncertainty of the nuclear matrix elements up to 50% and the contribution to this process due to particles other than Majorana neutrino is negligible).

3 Models with $\theta_{23} = 45^\circ$ and $\theta_{13} = 0^\circ$

At the lowest order, models with maximal atmospheric mixing angle, ($\theta_{23} = 45^\circ$) and vanishing value of CHOOZ mixing angle ($\theta_{13} = 0$) driven by some symmetry in addition with the standard model obtained through the choice of following neutrino mass matrix assuming the charged lepton mass matrix diagonal. Consider the following neutrino mass matrix

$$m_\nu = \begin{pmatrix} 0 & a & b \\ b & c & d \\ b & d & c \end{pmatrix} \quad (1)$$

Upon diagonalization, the above mass matrix gives three non-degenerate eigenvalues and two out of the three mixing angles are independent of model parameters and they are ($\theta_{23} = 45^\circ$), ($\theta_{13} = 0$), however, the solar mixing angle depends upon the model parameters.

The above mass matrix can easily be obtained invoking $\mu - \tau$ reflection symmetry [15, 21] and regarding that symmetry a point should be noted. If we invoke $\nu_{\mu L} \leftrightarrow \nu_{\tau L}$ and $\nu_{\mu R} \leftrightarrow \nu_{\tau R}$ then a diagonal charged lepton mass matrix will lead to unacceptable relation $m_\mu = m_\tau$. Thus it is necessary to build up a concrete model keeping track of the contribution from the charged lepton sector.

There is another approach to generate the above mixing angles through non-diagonal charged lepton mass matrix and keeping neutrino mass matrix diagonal. Examples of such types are $S_3 \supset S_3$ [16] permutation symmetry which gives rise to 'democratic type' charged lepton mixing matrix,lopsided structure of charged lepton mass matrix using GUT model [18].

Apart from the discrete symmetries few continuous U(1) symmetries are important in the context of neutrino mass and mixing pattern. Among them U(1) _{$L_e - L_\mu - L_\tau$} [19] is the most popular one. The neutrino mass matrix in the basis of diagonal charged lepton mass matrix this symmetry gives

$$m_\nu = \begin{pmatrix} 0 & 0 & p \\ 0 & p & q \\ p & q & 0 \end{pmatrix} \quad (2)$$

consequence of those flavor symmetries, however, $\theta_{12} = 45^\circ$ goes beyond the 5° value of the present experimental solar neutrino data. There is another $U(1)$ symmetry which can generate $\theta_{13} = 0$, $\theta_{23} = 45^\circ$, namely, $U(1)_{L-L}$ symmetry [20], which generates quasi-degenerate mass spectrum of neutrinos. In addition it generates $\theta_{12} = 0$ which can be cured under soft breaking of the symmetry.

4 Models with $\theta_{23} \neq 45^\circ$ and $\theta_{13} \neq 0^\circ$

In this section, we discuss a specific model which results the above mentioned mixing angles, however, still arbitrary solar mixing angle. The model is based on $U(1)_{L-L}$ symmetry as follows:

$$l_{2L} \sim l_{3L} \quad \text{and} \quad \nu_R \sim \nu_R \quad (3)$$

Neutrino masses are generated through higher dimensional operators, such as $\bar{l}_L / M$ where Φ is the usual Higgs doublet. The resulting charged lepton and neutrino mass matrices are [21]

$$m_E = \begin{pmatrix} 0 & a & b & b \\ \bar{a} & c & d & e \\ c & e & d & 0 \end{pmatrix}; m_\nu = \begin{pmatrix} 0 & p & q & q \\ \bar{p} & q & r & r \\ q & r & 0 & r \end{pmatrix} \quad (4)$$

Considering both the mass matrices are complex upon diagonalization the resulting mixing angles are coming out as

$$\sin^2 \theta_{atm} \approx 0.99; \sin^2 2\theta_{13} = 0 (10^{-5}); \sin^2 \theta_{solar} \approx 0.85 \quad (5)$$

for a specific choice of model parameters with acceptable charged lepton and neutrino mass values. There are several other types of models exist in the literature [22] which results to the above value of mixing angles. For all those types of models, solar mixing angles depends on the choice of model parameters. Thus, our search for a more predictive model driven by some symmetry is still in progress.

5 Tri-bimaximal Mixing

Experimental values of the three neutrino mixing angles are found to be remarkably close to the conjectured tri-bimaximal mixing ansatz proposed by Harrison, Perkins and Scott (HPS) [23] is given by

$$U_{tb} = \begin{pmatrix} 0 & q\frac{1}{2} & q\frac{1}{3} & 1 \\ \bar{q}\frac{1}{2} & q\frac{3}{2} & q\frac{1}{3} & 0 \\ \bar{q}\frac{1}{6} & q\frac{1}{3} & q\frac{1}{2} & \bar{q}\frac{1}{2} \\ q\frac{1}{6} & q\frac{1}{3} & q\frac{1}{2} & \bar{q}\frac{1}{2} \end{pmatrix} : \quad (6)$$

which gives $\theta_{12} = 35.2^\circ$, $\theta_{23} = 45^\circ$, $\theta_{13} = 0^\circ$ with a suitable normalization

with the standard model the HPS ansatz is realized. In the next section we will present a brief introduction to A_4 symmetry. To understand the very different mass pattern of charged lepton and neutrino A_4 symmetry has been proposed by E Ma and others [24]. However, to realize the HPS mixing within the framework of an $SU(2)_L \times U(1)_Y \times A_4$ model has been done by Altarelli and Feruglio [25].

6 A_4 Symmetry

A_4 is a non-abelian discrete symmetry group generated due to even permutation of four objects. It can also be viewed as an equivalent representation of the symmetry of a Tetrahedron. Number of elements of A_4 is 12 and those elements can be divided in four distinct classes as follows :

Identity transformation : E (1)

Four three fold axis of rotation : C_3 (4)

Four three fold axis of rotation : C_3^2 (4)

Three two fold axis of rotation : C_2 (3)

where $C_n = e^{2\pi i/n}$

Obviously, A_4 has four irreducible representation with dimension $\sum_{i=1}^4 n_i^2 = 12$ and the only solution exists as

$$n_1 = n_2 = n_3 = 1; n_4 = 3 \quad (7)$$

Those irreducible representations are labeled as $1, 1^0, 1^{\bar{0}}$ and 3 . Multiplication between different representations are given by

$$\begin{aligned} 3 \otimes 3 &= 1 + 1^0 + 1^{\bar{0}} + 3 + 3 \\ 1 \otimes 1 &= 1, \quad 1 \otimes 1^0 = 1^0, \quad 1 \otimes 1^{\bar{0}} = 1^{\bar{0}}, \quad 1^0 \otimes 1^0 = 1^{\bar{0}}, \quad 1^0 \otimes 1^{\bar{0}} = 1, \quad 1^{\bar{0}} \otimes 1^{\bar{0}} = 1^0 \\ 3 \otimes 1 &= 3, \quad 3 \otimes 1^0 = 3, \quad 3 \otimes 1^{\bar{0}} = 3. \end{aligned}$$

A_4 can also be generated by the two elements S and T satisfying the relation $S^2 = (ST)^3 = T^3 = 1$ where $S = (4321)$ and $T = (2314)$. Under such representation the above mentioned decomposed states can be viewed as

$$\begin{aligned} 1 \quad S &= 1 \quad T = 1 \\ 1' \quad S &= 1 \quad T = e^{i4\pi/3} = \omega^2 \\ 1'' \quad S &= 1 \quad T = e^{i2\pi/3} = \omega \end{aligned}$$

A_4 has two subgroups G_S : reflection subgroup generated by S isomorphic to Z_2 , G_T : rotational subgroup generated by T and isomorphic to Z_3 . If the flavor symmetry associated to A_4 is broken by the VEV of a triplet $\Phi = (\phi_1; \phi_2; \phi_3)$ there are two interesting breaking patterns generated. We will see in the next section G_S and G_T play a very crucial role to generate appropriate neutrino and charged lepton mass matrices which after diagonalization lead to HPS mixing matrix exactly.

7 Altarelli-Feruglio Model

Lepton	SU (2) _L	A ₄	
($\bar{l}_i; l_i$)	2	3	
$\bar{l}_i^c$	1	1	
Scalar			VEV
h_u	2	1	$\langle h_u^0 \rangle = v_u$
h_d	2	1	$\langle h_d^0 \rangle = v_d$
	1	1	$\langle \phi^0 \rangle = u$
s	1	3	$\langle \phi_s^0 \rangle = (v_s; v_s; v_s)$
t	1	3	$\langle \phi_t \rangle = (v_t; 0; 0)$

Table 1: List of fermion and scalar fields used in AF model.

interaction is given by [25]

$$L_1 = y_e e^c (\bar{t} l) h_d + y^c (\bar{t} l) \phi h_d + y^c (\bar{t} l) \phi h_d + x_a (\bar{h}_u h_u) =^2 + x_b (\bar{s} h_u h_u) =^2 + h.c \quad (8)$$

where x_a, x_b, y_e, y^c, y are Yukawa couplings and Λ is the scale of new physics. After spontaneous breaking of A₄ due to $\langle \phi_s \rangle \neq 0$ and $\langle \phi_t \rangle \neq 0$ and SU (2)_L → U (1)_Y due to $\langle h_u \rangle \neq 0$ and $\langle h_d \rangle \neq 0$, we obtain the charged lepton and neutrino mass matrices as

$$m_1 = \begin{pmatrix} m_e & 0 & 0 \\ 0 & m & 0 \\ 0 & 0 & m \end{pmatrix}; m = \begin{pmatrix} a+2d & d & d \\ d & 2d & a \\ d & a & 2d \end{pmatrix} \quad (9)$$

where $m_e = y_e v_d \frac{v_t}{\Lambda}$, $m = y v_d \frac{v_t}{\Lambda}$, $m_e = y v_d \frac{v_t}{\Lambda}$, $a = 2x_a v_u^2 \frac{u}{\Lambda^2}$ and $d = 2x_b v_u^2 \frac{v_s}{\Lambda^2}$. Upon diagonalization of m , the three non degenerate eigenvalues obtained as

$$m_1 = a + d; \quad m_2 = a; \quad m_3 = d - a \quad (10)$$

and the mixing matrix comes out same as the tribimaximal pattern. It is to be noted that choice of VEV is not very natural in the present model that causes vacuum alignment problem, however, that has been overcome through either by supersymmetrization [25, 26] of the model or through the introduction of another extra space dimension [27].

8 Beyond AF Model

Although AF model successfully generates tribimaximal mixing, however, due to the following reasons it is our endeavor to go beyond AF model:

HPS mixing gives $\theta_{12} = 35.26^\circ$ which is at the 1 σ range of the solar angle whereas the best fit value is 33.9° ,

θ_{13} may be non-zero according to CHOOZ experimental bound,

Motivated with the above, AF model has been modified through the inclusion of a $SU(2)_L$ singlet charged scalar $\phi = (\phi_1; \phi_2; \phi_3)$ and due to the soft breaking of A_4 symmetry in the scalar sector. A trilinear A_4 symmetry breaking term is incorporated and due to radiative Zee mechanism the following mass matrix of the neutrino is obtained [28]

$$m = \begin{pmatrix} 0 & a + \frac{2d}{3} & \frac{d}{3} & \frac{d}{3} \\ \frac{d}{3} & \frac{2d}{3} & a - \frac{d}{3} + \frac{2d}{3} \\ \frac{d}{3} & a - \frac{d}{3} + \frac{2d}{3} & \frac{2d}{3} \\ \frac{d}{3} & \frac{2d}{3} & a - \frac{d}{3} + \frac{2d}{3} \end{pmatrix} \quad (11)$$

where $\phi = f m^2 \frac{G_{12} v_u}{v_d} F(m^2; m_{hd}^2)$ with

$$F(M_1^2; M_2^2) = \frac{1}{16 M_1^2 (M_1^2 - M_2^2)} \ln \frac{M_1^2}{M_2^2} : \quad (12)$$

Here we consider two cases: i) Here we assume a, d, ϕ are all real and $j_1 < j_2 < j_3$, we obtain three eigenvalues as

$$m_1 = a + d + \phi; m_2 = a; m_3 = d - a \quad (13)$$

and the three mixing angles are

$$\sin \theta_{12} = \frac{1}{\sqrt{3}} + \frac{\phi}{3}; \sin \theta_{23} = \frac{1}{\sqrt{2}} - \frac{\phi}{2}; \sin \theta_{13} = \frac{\phi}{3} \quad (14)$$

where

$$\phi_1 = \frac{\phi}{d\sqrt{3}}; \phi_2 = \frac{1}{3} - \frac{\phi}{4a} - \frac{\phi}{2(2a-d)}; \phi_3 = \frac{1}{3} - \frac{\phi}{a} + \frac{\phi}{2(2a-d)} :$$

Assuming a constraint relation $d = 2a$, from Eq. (12) we obtain,

$$\begin{aligned} m_1 &= \frac{1}{2} \sqrt{\frac{2}{3}} m_{atm}^2 + \frac{\phi}{2} \sqrt{\frac{2}{3}} m_{atm}^2 R \\ m_2 &= \frac{1}{2} \sqrt{\frac{2}{3}} m_{atm}^2 \\ m_3 &= \frac{3}{2} \sqrt{\frac{2}{3}} m_{atm}^2 - \frac{\phi}{2} \sqrt{\frac{2}{3}} m_{atm}^2 R \end{aligned}$$

and

$$\begin{aligned} \sin \theta_{13} &= \frac{5}{3} \sqrt{\frac{2}{3}} R \\ \sin \theta_{12} &= \frac{1}{\sqrt{3}} - \frac{2}{5} \sqrt{\frac{2}{3}} \sin \theta_{13} \\ \tan^2 \theta_{23} &= 1 + \frac{4}{5} \sqrt{\frac{2}{3}} \sin \theta_{13} \end{aligned}$$

$$\theta_{12} = 31.13^\circ - 33.5^\circ; \quad \theta_{12} = 43.5^\circ - 46^\circ; \quad \theta_{12} = 3.5^\circ - 1.5^\circ$$

Thus the mass eigenvalues are normally ordered. While there is hierarchy between $m_{1,2}$ and m_3 , the masses m_1 and m_2 are nearly quasi-degenerate. The above mass values also give

$$|m_{ee} - j'| \approx 0.01 \text{ eV}$$

which is far beyond the detection limits of planned ν_μ experiments in near future. Furthermore, the smaller value of θ_{13} is correlated with the larger value of θ_{12} and θ_{23} . For example, with $d = 2.25a$, we obtain $\theta_{13} = 6^\circ$, but $\theta_{12} = 29.2^\circ$ which is below the range allowed at 99.9% C.L. and hence ruled out.

ii) One Complex and two real parameters:

In this section we consider the above model with one complex and two real parameters which in addition to the three mixing angles and three mass eigenvalues give rise to CP violation in the leptonic sector which has been parametrized through the J_{CP} parameter. We express mixing angles θ_{13} and θ_{23} in terms of a single model parameter and constrained the allowed parameter space. Utilizing all those constraints, we explore the extent of J_{CP} . We also describe the mass pattern obtained in this case [29].

We consider the parameter δ' is complex and the neutrino mass matrix comes out as

$$m = \begin{pmatrix} a + \frac{2de^i}{3} & \frac{de^i}{3} & \frac{de^i}{3} \\ \frac{de^i}{3} & \frac{2de^i}{3} & a - \frac{de^i}{3} \\ \frac{de^i}{3} & a - \frac{de^i}{3} + \frac{2de^i}{3} \end{pmatrix} \quad (15)$$

Diagonalizing the neutrino mass matrix through

$$U^Y m U = \text{diag}(de^i + a; a; de^i - a) \quad (16)$$

up to the first order of δ' and the matrix U becomes

$$U = U_{tb} + O(\delta') \quad (17)$$

Explicitly we get,

$$\begin{aligned} \sin \theta_{12} = |J_{12}| &= \frac{1}{3} + \frac{1}{3} \frac{(2 + \cos^2 \theta)}{\cos^2 (\theta + 2)} \\ \sin \theta_{13} = |J_{13}| &= \frac{1}{3} \frac{1 + \cos^2 (\theta^2 + 8)}{2(2)} \cos^2 \theta \quad (18) \\ \tan^2 \theta_{23} &= 1 + \frac{2}{3(2)} \end{aligned}$$

where $\theta = \delta'$; $d = a \cos \theta$. It is possible to express the parameter δ' as

From the above equation we get $m_{32}^2 > 0$ since $a^2 > 0$. Also if we express $\cos^2 \theta_{12}$ in the similar way, we get for the best fit value of observables $\theta_{12} < 2^\circ$ for $\cos^2 \theta_{12} < 1$. The model gives for $\theta_{12} = 34.0^\circ$, $\theta_{23} > 45.8^\circ$ and $\theta_{13} = 3.0^\circ$. If we allow 1 deviation of θ_{23} ($\theta_{23} = 48^\circ$) the maximum possible allowed value obtained from the experiment (CHOOZ : $0^\circ < \theta_{13} < 12^\circ$ at 3 σ).

Keeping all those constraints in view we explore the parameter space of CP violation parameter J_{CP} . The parameter J_{CP} is defined as

$$J_{CP} = \frac{1}{8} \sin 2\theta_{12} \sin 2\theta_{23} \sin 2\theta_{13} \cos \theta_{13} \sin \delta = \frac{\text{Im}[h_{12}h_{23}h_{31}]}{m_{21}^2 m_{31}^2 m_{32}^2} \quad (20)$$

where $h = m m^{-1/2}$. From the analytical expression of J_{CP} it has been found that 1 $^\circ$ deviation of θ_{23} (46° at $\delta = 2.03$) from its best fit value predicts $J_{CP} = 7.1 \times 10^{-5}$. If we allow 1 deviation of θ_{23} ($\theta_{23} = 48^\circ$ at $\delta = 2.41$) the value of $J_{CP} = 2.6 \times 10^{-3}$ is larger and which can be probed through the upcoming base line experiments.

It is also possible to constrain δ and for the best-fit values of θ_{12} , m_{21}^2 , m_{32}^2 we obtain $\delta = 3.6^\circ$ for $\theta_{23} = 48^\circ$ at $\delta = 2.41$. Moreover, the sum of the three neutrino masses vary as $0.07 < m_1 + m_2 + m_3 < 0.076$ due to the variation of parameter δ , $2.41 < \delta < 2$ ($48^\circ > \theta_{23} > 45.8^\circ$) which is also consistent with the cosmological bounds on the sum of the neutrino mass.

9 Open Problems

There are still few open problems which needs further attention

1. Embedding within a GUT model, so that the whole scenario of quarks and leptons could be explained. One of the major ongoing problem is to achieve quark-lepton complementarity relation which gives $\theta_c + \theta_{12} = 45.1^\circ \pm 2.4^\circ$ [1-30]. An attempt has been made within the context of supersymmetric $SU(4)_C \times SU(2)_L \times SU(2)_R$ model [31], however, a successful scenario with $SU(2)_L \times U(1)_Y \times A_4$ model is still yet unanswered.

2. See-saw realization of the AF model has been done [25, 32], however, connection between CP violating phases with leptogenesis has not yet been studied.

10 Acknowledgement

Author thanks the organizers of the XV IIAE-BRNS High Energy Physics Symposium held at Indian Institute of Technology, Kharagpur, India during December 2006, for the invitation to give this talk. Author also acknowledges Biswajit Adhikary for many helpful discussions.

References

- [3] B. Ahamin et al. [SNO Collaboration], Phys. Rev. C 72, 055502 (2005) [arXiv:nuclex/0502021]. S. Fukuda et al. [Super-Kamiokande Collaboration], Phys. Lett. B 539, 179 (2002); Q. R. Ahmad et al. [SNO Collaboration], Phys. Rev. Lett. 89, 011301 (2002); Q. R. Ahmad et al. [SNO Collaboration], Phys. Rev. Lett. 89, 011302 (2002); B. T. Cleveland et al., Astrophys. J. 496, 505 (1998); J. N. Abdurashitov et al. [SAGE Collaboration], J. Exp. Theor. Phys. 95, 181 (2002) [Zh. Eksp. Teor. Fiz. 122, 211 (2002)]; W. Hampel et al. [GALLIE Collaboration], Phys. Lett. B 447, 127 (1999); C. Cattadori, Talk at Neutrino 2004, Paris, France, June 14-19, 2004.
- [4] T. Araki et al. [KamLAND Collaboration], Phys. Rev. Lett. 94, 081801 (2005).
- [5] S. Goswami, A. Bandyopadhyay and S. Choubey, Nucl. Phys. Proc. Suppl. 143, 121 (2005); A. Bandyopadhyay et al., Phys. Lett. B 608, 115 (2005); G. L. Fogli et al., arXiv:hep-ph/0506083; M. Maltoni et al., New J. Phys. 6, 122 (2004).
- [6] A. Strumia and F. Vissani, Nucl. Phys. B 726, 294 (2005) [arXiv:hep-ph/0503246].
- [7] Y. Ashie et al. [Super-Kamiokande Collaboration], Phys. Rev. D 71, 112005 (2005).
- [8] E. Aliu et al. [K2K Collaboration], Phys. Rev. Lett. 94, 081802 (2005).
- [9] M. Apollonio et al., Eur. Phys. J. C 27, 331 (2003) [arXiv:hep-ex/0301017].
- [10] D. N. Spergel et al. [WMAP Collaboration], Astrophys. J. Suppl. 148, 175 (2003) [arXiv:astro-ph/0302209].
- [11] O. Elgaroy et al., Phys. Rev. Lett. 89, 061301 (2002) [arXiv:astro-ph/0204152], C. L. Kuo et al. [ACBAR collaboration], Astrophys. J. 600, 32 (2004) [arXiv:astro-ph/0212289].
- [12] S. R. Elliott and J. Engel, J. Phys. G 30, R183 (2004) [arXiv:hep-ph/0405078].
- [13] G. L. Fogli, E. Lisi, A. Marrone, A. Melchiorri, A. Palazzo, P. Serra and J. Silk, Phys. Rev. D 70, 113003 (2004) [arXiv:hep-ph/0408045].
- [14] H. V. Klapdor-Kleingrothaus, A. Dietz, H. L. Hamey and I. V. Krivosheina, Mod. Phys. Lett. A 16, 2409 (2001) [arXiv:hep-ph/0201231].
- [15] W. Grimus and L. Lavoura, JHEP 0107, 045 (2001) [arXiv:hep-ph/0105212]. W. Grimus and L. Lavoura, Acta Phys. Polon. B 32, 3719 (2001) [arXiv:hep-ph/0110041]. W. Grimus, S. Kaneko,

- [arXiv:hep-ph/0209294]. I. Aizawa, M. Ishiguro, T. Kitabayashi and M. Yasue, Phys. Rev. D 70, 015011 (2004) [arXiv:hep-ph/0405201]. A. Ghosal and D. Majumdar, Mod. Phys. Lett. A 21, 1067 (2006) [arXiv:hep-ph/0505173]. A. Ghosal and D. Majumdar, Phys. Rev. D 66, 053004 (2002) [arXiv:hep-ph/0209280].
- [16] H. Fritzsch and Z. Z. Xing, Phys. Lett. B 372, 265 (1996) [arXiv:hep-ph/9509389], *ibid* B 598, 237 (2004).
- [17] M. Fukugita, M. Tanimoto and T. Yanagida, Phys. Rev. D 57, 4429 (1998) [arXiv:hep-ph/9709388].
- [18] K. S. Babu and S. M. Barr, Phys. Lett. B 381, 202 (1996) [arXiv:hep-ph/9511446], *ibid* B 525, 289 (2002) J. Sato and T. Yanagida, Phys. Lett. B 430, 127 (1998) [arXiv:hep-ph/9710516]. G. Altarelli and F. Feruglio, Phys. Lett. B 451, 388 (1999) [arXiv:hep-ph/9812475].
- [19] S. T. Petcov, Phys. Lett. B 110, 245 (1982); for more recent studies see, e.g., R. Barbieri et al, JHEP 9812, 017 (1998); A. S. Joshipura and S. D. Rindani, Eur. Phys. J. C 14, 85 (2000); R. N. Mohapatra, A. Perez-Lorenzana and C. A. de Sousa Pires, Phys. Lett. B 474, 355 (2000); Q. Sha and Z. Tavartkiladze, Phys. Lett. B 482, 145 (2000). L. Lavoura, Phys. Rev. D 62, 093011 (2000); W. Grimus and L. Lavoura, Phys. Rev. D 62, 093012 (2000); T. Kitabayashi and M. Yasue, Phys. Rev. D 63, 095002 (2001); A. Aranda, C. D. Carone and P. Meade, Phys. Rev. D 65, 013011 (2002); K. S. Babu and R. N. Mohapatra, Phys. Lett. B 532, 77 (2002); H. J. He, D. A. Dicus and J. N. Ng, Phys. Lett. B 536, 83 (2002) H. S. Goh, R. N. Mohapatra and S. P. Ng, Phys. Lett. B 542, 116 (2002); G. K. Leontaris, J. Rizos and A. Psallidas, Phys. Lett. B 597, 182 (2004). JHEP 0009, 007 (2000); hep-ph/0410279. S. T. Petcov and W. Rodejohann, Phys. Rev. D 71, 073002 (2005) [arXiv:hep-ph/0409135].
- [20] S. Choubey and W. Rodejohann, Eur. Phys. J. C 40, 259 (2005) [arXiv:hep-ph/0411190]. B. Adhikary, Phys. Rev. D 74, 033002 (2006) [arXiv:hep-ph/0604009]. T. Ota and W. Rodejohann, Phys. Lett. B 639, 322 (2006) [arXiv:hep-ph/0605231]. W. Rodejohann and M. A. Schmidt, Phys. Atom. Nucl. 69, 1833 (2006) [arXiv:hep-ph/0507300].
- [21] A. Ghosal, Mod. Phys. Lett. A 19, 2579 (2004).
- [22] A. S. Joshipura, arXiv:hep-ph/0512252. R. N. Mohapatra and W. Rodejohann, Phys. Rev. D 72, 053001 (2005) [arXiv:hep-ph/0507312].
- [23] P. F. Harrison, D. H. Perkins and W. G. Scott, Phys. Lett. B 530, 167 (2002) [arXiv:hep-ph/0202074], *ibid* 530, 167 (2002); P. F. Harrison and

- Valle, Phys. Lett. B 552, 207 (2003). E. M a, Phys. Rev. D 70, 031901 (2004); E. M a, Phys. Rev. D 72, 037301 (2005).
- [25] G. Altarelli and F. Feruglio, Nucl. Phys. B 741, 215 (2006) [arXiv:hep-ph/0512103].
- [26] G. Altarelli, arXiv:hep-ph/0610164.
- [27] G. Altarelli, F. Feruglio and Y. Lin, arXiv:hep-ph/0610165. G. Altarelli and F. Feruglio, Nucl. Phys. B 720, 64 (2005) [arXiv:hep-ph/0504165].
- [28] B. Adhikary, B. Brahmachari, A. Ghosal, E. M a and M. K. Parida, Phys. Lett. B 638, 345 (2006) [arXiv:hep-ph/0603059].
- [29] B. Adhikary and A. Ghosal, arXiv:hep-ph/0609193.
- [30] J. Ferrandis and S. Pakvasa, Phys. Rev. D 71, 033004 (2005) [arXiv:hep-ph/0412038]. S. K. Kang, C. S. Kim and J. Lee, Phys. Lett. B 619, 129 (2005) [arXiv:hep-ph/0501029].
- [31] S. F. King and M. Malinsky, arXiv:hep-ph/0610250.
- [32] X. G. He and A. Zee, arXiv:hep-ph/0607163.