

Precision bounds on m_H and m_t

F. del Aguila

Departamento de Física Teórica y del Cosmos
 Universidad de Granada, E-18071 Granada, Spain

M. Martinez

PPE Division, CERN
 CH-1211 Geneva 23, Switzerland
 and

M. Quiros^y

Theory Division, CERN
 CH-1211 Geneva 23, Switzerland

Abstract

We perform a fit to precise electroweak data to determine the Higgs and top masses. Penalty functions taking into account their production limits are included. We find $m_H = 65^{+245}_4$ GeV and $m_t = 122^{+25}_{-20}$ GeV. However whereas the top 2 distribution behaves properly near them in minimum, the Higgs 2 distribution does not, indicating a statistical fluctuation or new physics. In fact no significant bound on the Higgs mass can be given at present. However, if the LEP accuracy is improved and the top is discovered in the preferred range of top masses, a meaningful bound on the Higgs mass could be obtained within the standard model framework.

CERN-TH.6389/92

IEM-TH-52/92

February 1992

Work partially supported by CICYT under contracts AEN 90-0139, AEN 90-0683.

^yPermanent address: Instituto de Estructura de la Materia, CSIC, Serrano 123, E-28006 Madrid, Spain.

Introduction

The standard model of electroweak interactions [1] describes impressively well all experimental data [2]. This description depends on two unknown parameters, the top mass, m_t , and the Higgs mass, m_H . The present experimental accuracy of the electroweak measurements justifies to look for indirect bounds on these masses. In this paper we present the χ^2 distributions resulting from the most precise and recent electroweak data as functions of m_t and m_H . Similar analyses can be found in the literature [3, 4, 5]. We pay particular attention to the Higgs penalty function describing the Higgs production limit. It plays an important role due to the low sensitivity of the data on m_H and to the fact that the minimum of the χ^2 , if only indirect data are used, is (at present) well below the direct production limit. Actually, the present sensitivity of the data on m_H will prove to be too low to obtain stringent, statistically significant, limits. However the sensitivity could be greatly improved in the near future, provided that LEP errors get reduced and the top is discovered.

Data

The experimental values used in the fits are gathered in Table 1. The detailed way LEP excludes Higgs production can be found in Ref. [6]. For our discussion it is enough to realize that the final result of the search is that no event with a given signature is found, whereas some events should be present if the Higgs boson would have been produced. Of course, the number of expected events decreases with increasing m_H . Therefore, assuming a Poisson distribution for the number of expected events, values below m_H can be excluded (for instance) at the 95% c.l. if the number of expected events for m_H is larger than or equal to 3. The number of expected events at LEP, N , as a function of m_H is shown in Fig.1a (this plot is obtained from the compilation in Ref. [7]). The confidence level, c , corresponding to seeing no events when N events are expected is equal to $1 - e^{-N}$; and this confidence level is transformed into the corresponding number of standard deviations, x , for a single sided $N(0;1)$ distribution using the relation

$$\frac{c + 1}{2} = \frac{1}{\sqrt{2\pi}} \int_1^x e^{-t^2/2} dt;$$

In this way x can be written as a function of m_H . This function is plotted in Fig.1b, and the best linear fit,

$$x = 15.81 - 0.24 m_H;$$

is superimposed on it. Thus, the corresponding contribution to χ^2 is given by the penalty function

$$\chi_H^2 = \begin{cases} \frac{65.0 - m_H (\text{GeV})}{4.1} & \text{if } m_H \leq 65 \text{ GeV;} \\ 0 & \text{if } m_H > 65 \text{ GeV;} \end{cases}$$

In the case of the top quark the detailed way TEVATRON excludes top production can be found in Ref. [8]. In this case, however, the minimum of the χ^2 , is well above the direct production limit, and then the corresponding penalty function plays no important

role in the fit. The strategy for the top search is also different. What is studied in the top case is the presence of an excess of events in some distributions (basically in the transverse mass distribution) due to top production. These distributions are fitted with a linear superposition of the standard model predictions with no top quark plus the top production. The result of this fit for each top mass is a probability density distribution $L(\tau)$ for the variable τ defined as the ratio of the number of fitted to the number of expected ($N_t(m_t)$) top events for such a mass. No top events corresponds to $\tau = 0$. To exclude a top mass (for instance) with a 95% c.l. we associate to each mass m_t the value of $\tau_0(m_t)$ which defines for the probability density corresponding to this mass an area, $\int_0^{\tau_0(m_t)} L(\tau) d\tau$, equal to the 95% of the total area or probability, $\int_0^{\tau_1} L(\tau) d\tau$. Then $\tau_0(m_t) = 1$ gives the m_t limit below which the top mass is excluded. To make this computation for different confidence levels we have assumed that $\tau_0(m_t) = \tau_1(m_t)$ is independent of m_t in the region of interest. (Note that this is equivalent to assume that the shape of the top distribution is independent of the top mass in this region.) To translate the c.l. to which a given top mass is excluded into a number of standard deviations we proceed as in the case of the Higgs mass. The corresponding fit gives the approximate penalty function

$$\chi_t^2 = \begin{cases} \frac{108 - m_t(\text{GeV})}{9} & \text{if } m_t \leq 108 \text{ GeV;} \\ 0 & \text{if } m_t > 108 \text{ GeV;} \end{cases}$$

The χ^2 also includes neutral current data: neutrino-quark data, q [9], the latest neutrino-electron results, e [10], and the parity violation in atoms data, eH [11, 12]; the W mass limits: M_W [13] and $\frac{M_W}{M_Z}$ [14]; the LEP results presented at the Joint International Lepton-Photon Symposium and Europhysics Conference on High-Energy Physics in August 1991 [15] (with the correlation matrix given in Ref. [16]); and the strong coupling constant, α_s [17]. We take for α_s the latest (most precise) ALEPH measurement using hadronic Z decays. The definite α_s value is important to fix the χ^2 value at the minimum but modifies little the relative χ^2 distribution and does not affect the conclusions.

Theory

We use the on-shell scheme of Ref. [18] with the electromagnetic coupling constant, α , and the Fermi constant, G_F , [9] as input parameters. For the calculation of the observables in the Z physics sector two independent electroweak libraries have been used: the one from G. Burgers and W. Hollik [19] and the one from the Dubna-Zeuthen group [20]. In spite of the fact that they use a completely different calculational scheme, their results are in very good agreement [21]. We have upgraded the first one to incorporate:

The missing relevant pieces of two-loop corrections [22];

The dominant QCD corrections to m_t dependent terms [23, 24];

The updated calculation of the QCD corrections to the decay $Z \rightarrow b\bar{b}$ [24]; and,

$O(\alpha_s^3)$ corrections to the decay widths [25].

Similar upgrades have been implemented in the Dubna-Zeuthen library by their authors. A comparison of the present versions of both programs shows an even better agreement than the one previously quoted. While the numbers and plots shown in this study have been obtained with the first library, we have explicitly checked that the changes are negligible if the Dubna-Zeuthen library is used instead. In our fits we use the explicit expressions for all the observables, in particular for the neutral current ones [26]. We have checked that reducing the full experimental information for the latter to the equivalent electroweak mixing angle value, our fits and conclusions change little.

Interpretation

Our results are summarized in Figs 2-8. Fig 2a shows the contribution to the χ^2 distribution of neutral current and pp collider data; Fig 3a that of LEP data and α_s ; Fig 4a their sum; and Fig 5a that of all the data including the penalty functions. Thus Figs 2a-4a give partial contributions to Fig 5a. At each point in the $m_t - m_H$ plane we minimize with respect to M_Z and α_s (G_F and $\sin^2 \theta_W$ being fixed). Thus, the total number of degrees of freedom is $20 - 2 = 18$. (The two degrees of freedom corresponding to the penalty functions are taken care by the two parameters, m_t and m_H , with respect to which we do not minimize.) Figs 2b-5b show the same fits but replacing the central experimental values of the different observables by their predicted (standard model) values near the minimum of the χ^2 distribution. In particular we have taken $m_H = 70$ GeV, $m_t = 130$ GeV. The corresponding standard model predictions for the observables used in the fits are given in Table 2. Comparing Figs 2a-4a and Fig 5a we see that the Higgs penalty function moves the minimum from $m_H = 10$ to 70 GeV. However, the main observation to be elaborated below when comparing Fig 3a and Fig 3b is the apparent change in the LEP distribution, and then in the global one. This tells us that some experimental central values manifest a statistical fluctuation or indicate new physics. This comparison also proves the actual lack of sensitivity of the data on m_H . To understand this change it is illuminating to rewrite (in obvious notation)

$$\begin{aligned} \chi^2(m_H; m_t; \dots) &= \sum_i \frac{(\chi_i^{\text{exp}} - \chi_i^{\text{th}}(m_H; m_t; \dots))^2}{\sigma_i^2} = \sum_i \frac{(\chi_i^{\text{exp}} - \chi_i^0 + \delta_i)^2}{\sigma_i^2} \\ &= \sum_i \frac{(\chi_i^{\text{exp}} - \chi_i^0)^2}{\sigma_i^2} - 2 \sum_i \frac{(\chi_i^{\text{exp}} - \chi_i^0) \delta_i}{\sigma_i^2} + \sum_i \frac{\delta_i^2}{\sigma_i^2}; \end{aligned}$$

where χ_i^0 are the standard model predictions at the χ^2 minimum and δ_i are the differences between the predicted values at a given point and the values at the minimum. The last expression shows that a statistical fluctuation or a relatively large change in the central value of any observable contributes to the χ^2 with a relatively large constant term $\frac{(\chi_i^{\text{exp}} - \chi_i^0)^2}{\sigma_i^2}$ and with a linear term of pronounced slope $-\frac{2(\chi_i^{\text{exp}} - \chi_i^0)}{\sigma_i^2}$. However, what gives the sensitivity of the observable is $\frac{\delta_i}{\sigma_i^2}$. Comparing Figs 2a-5a and Figs 2b-5b

we see that indirect LEP data are very insensitive to m_H but some observables have a large discrepancy between their measured central values and the predicted ones. The global fit to all data (Fig. 5a) gives

$$m_H = 65^{+245}_4 \text{ GeV};$$

$$m_t = 122^{+25}_{20} \text{ GeV};$$

Although the fits in Figs. 2a-5a give the actual experimental information, those in Figs. 2b-5b give the actual sensitivity of the data on m_H and m_t . Hence present indirect electroweak data are not sensitive to m_H . Actually, the present LEP discrepancy can be traced back to the value of the ratio R_b and the bottom forward/backward asymmetry A_{FB}^{bb} . This is to say that the data show a higher sensitivity than expected on m_H due to the pronounced slope terms (introduced above) resulting from R_b and A_{FB}^{bb} . If this discrepancy is a statistical fluctuation and it disappears when more data are available, the subsequent fit will approach Figs. 2b-5b. However, it can also happen that the central values of other observables fluctuate, in which case our discussion would apply to them. At any rate this will be always the case (if new physics is discarded) when the preferred m_H value from indirect LEP data lies below the production limit. In Figs. 6a,b we show the χ^2 distribution as a function of m_H (minimization with respect to m_t is understood) for the global fits in Figs. 5a,b. Plotting the same distribution for the different observables it can be explicitly proven that only R_b and A_{FB}^{bb} show the pronounced slope behaviour for the real data, as can be seen in Fig. 6a, whereas the other observables show a small variation, which we do not plot. Fig. 6b confirms that the sensitivity of R_b and A_{FB}^{bb} on m_H is very low, as it is the sensitivity of the other observables and of the full set. In fact we plot χ^2 in Fig. 6a, subtracting from each χ^2 contribution its value at the global minimum: 10.79, 0.88, 2.73 for the global, R_b , A_{FB}^{bb} contributions, respectively. Using higher values for R_b would arrange the R_b deviation from the standard model prediction. This would translate into a modification of the m_t value at the minimum in order to conserve the agreement between the measured and the predicted Z values. However A_{FB}^{bb} would still constrain the fit and keep unchanged the value of m_H at the minimum and the conclusions. Lower values of R_b make worse the disagreement between the measured and the predicted R_b values. It is worth to note that although the top mass limit is significative, it is strongly correlated to the Higgs mass as can be seen in Fig. 5. For instance if the Higgs mass were known to be $300 \pm 30 \text{ GeV}$, the present data would imply $m_t = 139^{+21}_{25} \text{ GeV}$ (see Fig. 5a).

Conclusions

As stressed above, today's data show a cautious m_H sensitivity due to a statistical fluctuation or to new physics manifested by the experimental values of R_b and A_{FB}^{bb} (Figs. 2a-6a). As a matter of fact, the actual sensitivity of the considered observables on m_H is very low (Figs. 2b-6b). The Higgs penalty function plays an important role in the fits, pushing up the Higgs mass value at the minimum. What is more interesting is the expected improvement of the m_H sensitivity in the near future [27]. Figs. 2-5 show the correlation between m_H and m_t which should translate, once the top quark is discovered and its mass

determined, into a definite bound on m_H . In fact, if present LEP accuracy is improved as assumed in Table 2, and the top mass is in the preferred range given above, a meaningful bound on the Higgs mass (within the standard model framework) could be obtained. This fact is illustrated in Figs. 7 and 8. In Fig. 7 we plot the χ^2 -distribution using the standard model predictions as central values and the set of LEP improved errors quoted in Table 2. We minimize at each point with respect to M_Z and α_s . Fig. 7 shows that fixing m_t a relatively strong upper bound on m_H can be deduced. This is explicitly shown in Fig. 8, where we fix the top mass, $m_t = 130 \pm 1$ GeV. In this case $m_H < 315$ GeV at 95% c.l.

Acknowledgements

One of us (M.M.) would like to thank J. Steinberger for triggering the interest for this study as well as for many suggestions and A. Blondel and G. Rolandi for discussions. We have benefited from discussions with J. Ellis, with W. Hollik about the upgrade of his libraries and also with D. Schaile concerning the actual interpretation of the χ^2 distribution. We acknowledge W. Hollik and the Dubna-Zeuthen group for making us available their computer programs.

References

- [1] S.L. Glashow, Nucl. Phys. 22 (1961) 579; A. Salam, in Elementary particle theory: relativistic groups and analyticity (Nobel Symposium No.8), ed. N. Svartholm (Almqvist and Wiksell, Stockholm, 1968); S. Weinberg, Phys. Rev. Lett. 19 (1967) 1264; S.L. Glashow, J. Iliopoulos and L. Maiani, Phys. Rev. D 2 (1970) 1285; G. 't Hooft, Nucl. Phys. B 33 (1971) 173; B 35 (1971) 167.
- [2] See for instance: H. Burhardt and J. Steinberger, Annu. Rev. Nucl. Part. Sci. 41 (1991) 55.
- [3] J. Ellis and G.L. Fogli, Phys. Lett. B 249 (1990) 543; J. Ellis, G.L. Fogli and E. Lisi, CERN preprint, CERN/TH.6273/91 (Sep.1991).
- [4] G. Passarino, Phys. Lett. B 255 (1991) 127.
- [5] D. Schaile, CERN preprint, CERN/PPE/91/187 (Nov.1991).
- [6] D. Decamp et al., ALEPH Collaboration, Phys. Lett. B 246 (1990) 306; P. Abreu et al., DELPHI Collaboration, Z. Phys. C 51 (1991) 25; B. Adeva et al., L3 Collaboration, Phys. Lett. 257B (1991) 450; M. Z. Akray et al., OPAL Collaboration, Phys. Lett. 253B (1991) 511.
- [7] M. Davier, Proceedings of the International Lepton{Photon Symposium and Europhysics Conference on High Energy Physics (Eds. S. Hegerty, K. Potter and E. Quercigh), Geneva 1991 (to appear).
- [8] F. Abe et al., CDF Collaboration, Phys. Rev. Lett. 64 (1990) 142; Phys. Rev. D 43 (1991) 664; Phys. Rev. Lett. 68 (1992) 447.
- [9] Particle Data Group, Phys. Lett. B 239 (1990) 1.
- [10] R. Beyer, CHARM II Collaboration, Proceedings of the International Lepton{Photon Symposium and Europhysics Conference on High Energy Physics (Eds. S. Hegerty, K. Potter and E. Quercigh), Geneva 1991 (to appear).
- [11] P. Langacker, Phys. Lett. B 256 (1991) 277.
- [12] M.C. Noecker, B.P. Masterson and C.E. Wieman, Phys. Rev. Lett. 61 (1988) 310; S.A. Blundell, W.R. Johnson and J. Sapirstein, Phys. Rev. Lett. 65 (1990) 1411; V. Dzubaj, V. Flambaum and O. Sushkov, Phys. Lett. A 141 (1989) 147.
- [13] F. Abe et al., CDF Collaboration, Phys. Rev. Lett. 65 (1990) 2243.
- [14] J. Alitti et al., UA2 Collaboration, CERN/PPE/91-163 (Sep.1991).
- [15] J. Carter, Proceedings of the International Lepton{Photon Symposium and Europhysics Conference on High Energy Physics (Eds. S. Hegerty, K. Potter and E. Quercigh), Geneva 1991 (to appear).

- [16] The LEP Collaborations: ALEPH, DELPHI, L3 and OPAL, CERN preprint, CERN-PPE/91-232, to appear in Phys. Lett. B .
- [17] For some recent compilations see for instance: S. Bethke, CERN preprint, CERN/PPE/91-36 (1991); T. Hebbeker, Proceedings of the International Lepton{Photon Symposium and Europhysics Conference on High Energy Physics (Eds. S. Hegerty, K. Potter and E. Quercigh), Geneva 1991 (to appear). The actual measurement used in this study comes from the ALEPH Collaboration (in preparation).
- [18] D.A. Ross and J.C. Taylor, Nucl. Phys. B 51 (1973) 25; A. Sirlin, Phys. Rev. D 22 (1980) 971; M. Consoli and W. Hollik, in Z Physics at LEP 1, CERN 89{08 (Ed. G. Altarelli, R. Kleiss and C. Verzegnassi), Vol.1 (1989) 7; W. Hollik, Fortschr. Phys. 38 (1990) 3.
- [19] Program GAMMAZ, G. Burgers and W. Hollik; G. Burgers, F. Jegerlehner, B. Kniehl and J. Kuhn, Proceedings of the Workshop of Z physics at LEP 1, CERN Report 89-08 Vol.1 (1989) 55; M. Consoli, W. Hollik and F. Jegerlehner, ibid. Vol.1 (1989) 7.
- [20] Program DIZET, Dubna-Zeuthen radiative correction group, D. Bardin et al., Z. Phys. C 44 (1989) 493; Nucl. Phys. B 351 (1991) 1; Phys. Lett. B 255 (1991) 290.
- [21] D. Bardin, W. Hollik and T. Riemann, Z. Phys. C 49 (1991) 485.
- [22] S. Fanchiotti and A. Sirlin, NYU preprint, in memoriam M.A.B. Beg, (Feb.1990); G. Degrossi, S. Fanchiotti and A. Sirlin, Nucl. Phys. B 351 (1991) 49; G. Degrossi and A. Sirlin, Nucl. Phys. B 352 (1991) 342; F. Halzen and B. Kniehl, Nucl. Phys. B 353 (1991) 567; F. Halzen, B. Kniehl and M.L. Stong, MAD/PH/658 (Sep.1991) .
- [23] A. Djouadi and C. Verzegnassi, Phys. Lett. B 195 (1987) 265; A. Djouadi, Nuovo Cim. 100A (1988) 357.
- [24] B.A. Kniehl and J.H. Kuhn, Phys. Lett. B 224 (1989) 229; Nucl. Phys. B 329 (1990) 547; K.G. Chetyrkin and J.H. Kuhn, Phys. Lett. B 248 (1990) 359.
- [25] L.R. Surgulazde and M.A. Samuel, Phys. Rev. Lett. 66 (1991) 560.
- [26] W.J. Marciano and A. Sirlin, Phys. Rev. D 22 (1980) 2695; W.J. Marciano, S. Sarantakos and A. Sirlin, Nucl. Phys. B 217 (1983) 84; W.J. Marciano and A. Sirlin, Phys. Rev. D 27 (1983) 552; ibid. D 29 (1984) 75.
- [27] A. Blondel, F.M. Renard and C. Verzegnassi, Phys. Lett. B 269 (1991) 419.

Table captions

Table 1 Experimental values of the observables used in the fits.

Table 2 Standard model predictions for the observables used in the fits and for $m_H = 70 \text{ GeV}$ and $m_t = 130 \text{ GeV}$. The improved error column gives a guess of future LEP errors.

Figure captions

Fig.1a [b] Number of expected events [standard deviations] at LEP as a function of m_H .

Fig.2a [b] Level contours for the contribution to the χ^2 -distribution of neutral current and pp collider data in Table 1 [standard model predictions in Table 2 with the present experimental errors in Table 1]. Minimization with respect to M_Z and α_s is understood.

Fig.3a [b] The same as in Fig.2a [b] but for LEP observables including α_s .

Fig.4a [b] The same as in Fig.2a [b] but for all observables in Table 1.

Fig.5a [b] The same as in Fig.2a [b] but for all observables in Table 1 plus the top and Higgs penalty functions given in the text.

Fig.6a [b] χ^2 -distributions as functions of m_H for the global fit in Fig.5a [b] (solid curve). The R_V (dashed curve) and the A_{FB}^{bb} (dotted curve) contributions are also shown. Minimization with respect to M_Z , α_s and m_t is understood.

Fig.7 Level contours for the total χ^2 -distribution assuming the standard model predictions as central values and the set of improved errors given in Table 2. Minimization with respect to M_Z and α_s is understood.

Fig.8 χ^2 -distribution as a function of m_H for the global fit in Fig.7 and $m_t = 130 \pm 1 \text{ GeV}$. Minimization with respect to M_Z , α_s and m_t is understood.

Q u a n t i t y		Ex p e r i m e n t a l V a l u e	C o r r e l a t i o n M a t r i x			
q	g_L^2	0.2977 0.0042	1.			
	g_R^2	0.0317 0.0034		1.		
	χ_L^2	2.50 0.03			1.	
	χ_R^2	$4.59^{+0.44}_{-0.27}$				1.
e	g_A^e	0.503 0.018	1.	0.05		
	g_V^e	0.025 0.020		1.		
eH	$C_{1u} + C_{1d}$	0.144 0.007	1.			
	$C_{1u} - C_{1d}$	0.60 0.09		1.		
	$C_{2u} - C_{2d}$	0.05 0.11			1.	
pp	M_W	79.91 0.39 GeV				
	$M_W = M_Z$	0.8813 0.0041				
LEP	M_Z	91.175 0.021 GeV	1.	0.09	0.01	0.00
	χ_Z^2	2487 10 MeV		1.	{0.25	{0.07
	χ_h^2	41.36 0.23 nb			1.	0.18
	R_γ	20.92 0.11				1.
	From $A_{FB}^{l_2}$: $\frac{g_V}{g_A}(M_Z)$	0.0048 0.0012	1.			
	From $A^{pol}(\gamma)$: $\frac{g_V}{g_A}(M_Z)$	0.072 0.017		1.		
	From bb-asymmetry: $A_{FB}^{bb}(M_Z)$	0.132 0.022			1.	
	From qq-asymmetry: $\sin^2 \theta_w(M_Z)$	0.2303 0.0035				1.
	s	0.125 0.005				

Table 1

Q uantity		Standard m odel prediction	Im proved errors
q	g_L^2	0.3012	0.0042
	g_R^2	0.0302	0.0034
	ϵ_L	2.46	0.03
	ϵ_R	5.18	0.44
e	g_A^e	0.505	0.018
	g_V^e	0.048	0.020
eH	$C_{1u} + C_{1d}$	0.146	0.007
	$C_{1u} - C_{1d}$	0.54	0.09
	$C_{2u} - C_{2d}$	0.09	0.11
pp	M_W	80.14 G eV	0.06 G eV
	$M_W = M_Z$	0.8790	0.0041
LEP	M_Z	91.175 G eV	0.005 G eV
	σ_{had}^0	2489 M eV	4 M eV
	R_γ	41.41 nb	0.08 nb
	R_γ	20.81	0.02
	From $A_{FB}^{l,2}$: $\frac{g_V}{g_A}(M_Z)$	0.0048	0.0006
	From $A^{\text{pol}}()$: $\frac{g_V}{g_A}(M_Z)$	0.069	0.009
	From bb-asymmetry: $A_{FB}^{bb}(M_Z)$	0.097	0.010
	From qq-asymmetry: $\sin^2 \theta_W(M_Z)$	0.2320	0.0035
	s	0.125	0.005

Table 2