

Electroweak Symmetry Breaking via a Technicolor Nambu–Jona-Lasinio Model

Markus A. Luty

Theoretical Physics Group
Lawrence Berkeley Laboratory
1 Cyclotron Road
Berkeley, California 94720

Abstract

We consider a theory of gauge fields and fermions which we argue gives rise to dynamics similar to that of the Nambu–Jona-Lasinio (NJL) model when a gauge coupling constant is appropriately re-tuned. We discuss the application of this model to dynamical electroweak symmetry breaking by a top-quark condensate. In this model, custodial symmetry is violated solely by perturbatively weak interactions, and the top–bottom mass splitting is due to the enhanced sensitivity to custodial symmetry violation near the critical point. We also consider models in which electroweak symmetry is broken by new strongly-interacting fermions with NJL-like dynamics. We argue that these models require additional re-tuning in order to keep corrections to the electroweak ρ -parameter acceptably small.

1. Introduction

In the last few years, there has been a revived interest in the Nambu{Jona-Lasinio (NJL) model [1] as an effective description of electroweak symmetry breaking via a top-quark condensate [2][3]. Numerous variations on this basic theme have also been explored [4]. More recently, several groups have constructed renormalizable models which are supposed to give rise to top-quark condensates and NJL-like dynamics. [5]. In this paper, we will consider a model of asymptotically free gauge fields and fermions which we argue has dynamics similar to that of the NJL model. In this model, all symmetry breaking is driven by strongly-coupled $SU(N)$ gauge groups with fermions in the fundamental representation. We can therefore use our understanding of chiral symmetry breaking in QCD-like theories to analyze the model. The only new dynamical assumption is that the chiral symmetry breaking phase transition is second order. We will consider models in which the electroweak symmetry is broken by top-quark condensates, as well as models in which electroweak symmetry is broken by condensates of new fermions.

The plan of this paper is as follows: In section 2, we will review some basic results concerning the NJL model, and discuss the sense in which the NJL model can be viewed as a low-energy effective lagrangian. In section 3, we will present a model of asymptotically-free gauge fields and fermions and give qualitative arguments that this model gives rise to dynamics similar to the NJL model. In section 4, we flesh out the dynamical picture of the previous section by analyzing the model using the Schwinger{Dyson equations in ladder approximation. In section 5, we apply the model to electroweak symmetry breaking, and section 6 contains our conclusions.

2. The NJL model

We begin by treating the NJL model as a toy model with no reference to electroweak physics. The NJL model we will consider has F "flavors" and N "technicolors." The lagrangian is

$$\mathcal{L}_{\text{NJL}} = \bar{\psi} i \not{\partial} \psi + \frac{G}{N} (\bar{\psi}_{Lja} \psi_{Rka}) (\bar{\psi}_{Rkb} \psi_{Ljb}); \quad (1)$$

where $j,k = 1, \dots, F$ are flavor indices and $a,b = 1, \dots, N$ are technicolor indices. This theory has a global chiral symmetry

$$SU(N)_{TC} \times U(F)_L \times U(F)_R; \quad (2)$$

in an obvious notation. The theory is nonrenormalizable, and must therefore be interpreted as an effective theory equipped with an energy cutoff. In order to describe physics at

scales above Λ , we must construct a more fundamental theory which reduces to the NJL model at low energies.

In the limit $N \rightarrow \infty$ (with F held finite), this model can be solved exactly. The qualitative features are well known: if G is less than a critical value G_{crit} , then the chiral symmetry of eq. (2) is unbroken, and the effective theory below the scale Λ simply describes massless fermions interacting through the contact interaction of the form eq. (1). However, if $G > G_{\text{crit}}$, then the chiral symmetry is partially broken: the condensate

$$\langle \bar{\psi}_{Lj} \psi_{Rk} \rangle \neq 0 \quad (3)$$

results in the symmetry-breaking pattern

$$U(F)_L \times U(F)_R \rightarrow U(F)_{L+R}; \quad (4)$$

where $U(F)_{L+R}$ is the vector-like subgroup of $U(F)_L \times U(F)_R$.

If G is tuned to be very close to G_{crit} ,

$$G = G_{\text{crit}} (1 + O(m^2)) \quad \text{with} \quad m^2 \propto G^{-1}; \quad (5)$$

then the effective theory near the scale Λ is precisely a $U(F)_L \times U(F)_R$ linear sigma model coupled to the fermions. The effective lagrangian at the scale Λ is

$$\begin{aligned} \mathcal{L}_e = & \bar{\psi} \not{\partial} \psi + y \bar{\psi}_{Lj} \psi_{Rk} + \text{h.c.} \\ & + \text{tr} \left[\frac{1}{2} (\partial_\mu \phi)^2 + \frac{1}{2} \text{tr} \left(\frac{1}{2} \partial_\mu \vec{\phi}^2 \right) + \frac{1}{2} \text{tr} \left(\frac{1}{2} \partial_\mu \vec{\phi}^2 \right) + \dots \right]; \end{aligned} \quad (6)$$

where the ellipses indicate terms containing higher-dimension operators. If $G < G_{\text{crit}}$, then $m^2 > 0$ and the theory is in the unbroken phase. The fermions are then massless, and there are $2F^2$ physical scalars with equal masses of order m . If $G > G_{\text{crit}}$, then $m^2 < 0$ and the theory is in the broken phase. The fermions then have masses of order m , and there are F^2 physical scalars with masses of order m and F^2 massless Nambu-Goldstone bosons. Because the underlying theory eq. (1) has only one coupling constant G , there will be non-trivial relations among the parameters of the effective low-energy lagrangian eq. (6).

An important feature of the NJL model in the large- N limit is that the low-energy dynamics is continuous as a function of G near the critical point. An analogous assumption in the context of our renormalizable model will play an important role in what follows.

Before presenting the renormalizable model, we discuss the sense in which the NJL model can arise as an effective theory from a more fundamental theory. When we write

an effective field theory which matches onto a more fundamental theory at a scale Λ , we must include all possible interaction terms among the low-energy fields consistent with the symmetries of the underlying theory. The coefficients of these operators are determined by matching the predictions of the low-energy theory with that of the underlying theory. There will be an infinite number of operators in the low-energy effective lagrangian, but only a few of these, the so-called "relevant" operators, will be important for describing the physics at energies $E \ll \Lambda$. The effects of the remaining "irrelevant" operators at low energies will be suppressed by powers of E/Λ . This is the reason for the power of the effective-lagrangian approach.

For weakly-coupled theories, dimensional analysis is sufficient to classify operators as relevant or irrelevant: operators with engineering dimension 4 or less are relevant, and all others are irrelevant. As the NJL model illustrates, the classification becomes non-trivial when couplings become strong. We therefore ask if there are any other operators which are relevant if they are added to the NJL lagrangian eq. (1) when G is near its critical value.

A clue to the answer to this question can be obtained from the fact that the low-energy limit of the fine-tuned NJL model is a linear sigma model. This equivalence can be made explicit by rewriting the NJL interaction in terms of an auxiliary scalar field σ :

$$L_{\text{NJL}}^0 = \bar{\psi} i \not{\partial} \psi + \frac{G}{4} (\bar{\psi} \psi)^2 + \frac{1}{2} (\partial_\mu \sigma)^2 - L_{\text{ja}} - \bar{\psi} \psi \sigma + \text{h.c.} : \quad (7)$$

This is the effective lagrangian at the scale Λ , in which σ has no kinetic or self-interaction terms. σ can be explicitly integrated out to recover the NJL lagrangian eq. (1). At scales below Λ , σ acquires both kinetic and self-interaction terms as in eq. (6), and can serve as an interpolating field for the physical scalar states.

In this formulation of the NJL model, it is clear that all of the the dimension-four operators appearing in eq. (6) are relevant. If these are added to the effective lagrangian at the scale Λ and the field σ is integrated out, the result will be a non-local effective action containing only the fermion fields. If we expand this action in powers of derivatives of fermion fields, it seems clear that there should be a small number of higher-dimension operators that become relevant in the fine-tuned limit, and whose effects at low energies precisely reproduce the effects of the operators of eq. (6).

One such operator was considered in ref. [6]. A more complete analysis was performed in ref. [7], which explicitly identified operators that, when added to the effective lagrangian at the scale Λ , reproduced the entire parameter space of the linear sigma model. It may therefore seem that the predictions of refs. [2][3][4] based on the NJL model are vacuous, but this is not so. The reason is that as long as Λ is many orders of magnitude above

the weak scale, and the linear sigma model has a large Yukawa coupling at scales just below Λ , the logarithmic evolution of the parameters of the effective lagrangian drives the parameters of the theory to an approximate infrared fixed point at the weak scale [8]. (This is true for all models which reduce to the standard model with one or two Higgs doublets in the low-energy limit.) The role of the NJL-like dynamics in these models is therefore to explain why the top-quark coupling is singled out to be strong enough to lie in the basin of attraction of the approximate infrared fixed point.

The situation is clearly quite different if the compositeness scale is taken to be near the weak scale. In this case, the couplings do not run over a sufficient range to be affected by the approximate infrared fixed point, and all the relevant operators are a priori equally important. The construction of explicit renormalizable models is clearly especially interesting in this case.

If we consider some fundamental theory which gives rise to the NJL interaction eq. (1), it is clear that all of the relevant operators discussed above will be generated. Therefore, in attempting to construct a renormalizable model which captures the physics of the NJL model, strictly speaking we can only expect to find a model with a fermion bilinear order parameter whose low-energy limit is a linear sigma model. Nonetheless, in the model we construct, we will find that the dynamics is very similar to the NJL model eq. (1).

3. A Renormalizable Model

The NJL model described in the previous section bears a superficial resemblance to a model in which the four-fermion coupling in eq. (1) is replaced by $SU(N)_{TC}$ gauge interactions. In the $SU(N)_{TC}$ gauge model, the gauge coupling becomes strong at some scale Λ_{TC} , and a condensate of the form eq. (3) forms, giving rise to $F^2 - 1$ NGB's. (Recall that the NJL model of eq. NJL has $F^2 - 1$ NGB's.) We can try to press the analogy further by using Fierz identities to rewrite the NJL interaction as

$$(\bar{\psi}_{Lja} \psi_{Rka})(\bar{\psi}_{Rkb} \psi_{Ljb}) = (\bar{\psi}_{Lj} T_A \psi_{Lj})(\bar{\psi}_{Rk} T_A \psi_{Rk}) + O(1/N); \quad (8)$$

where the T_A are $SU(N)_{TC}$ generators normalized so that $\text{tr} T_A T_B = \delta_{AB}/2$. Eq. (8) is the operator corresponding to the most attractive channel for one gauge boson exchange in the $SU(N)_{TC}$ theory. However, the two models are clearly qualitatively different: The NJL model contains propagating fermions at low energies, while the fermions are confined in the $SU(N)_{TC}$ gauge model. Also, the NJL model can be re-tuned to make the scalar and fermion masses small compared to the cutoff scale Λ , while the $SU(N)_{TC}$ gauge model has no adjustable parameter; all dimensionful quantities in the $SU(N)_{TC}$ model are of order Λ_{TC} raised to the appropriate power.

We can, however, construct a class of models whose low-energy dynamics interpolates continuously between that of the NJL model and the $SU(N)_{TC}$ gauge model. The models we will construct consist entirely of gauge fields coupled to fermions, so the theories are well behaved at high energies. Chiral symmetry breaking in these models is triggered by asymptotically-free gauge couplings, and the fine-tuning in the NJL limit can be understood in terms of the fine-tuning of gauge couplings (defined at a suitable subtraction point) in the underlying gauge theory.

The basic idea underlying the model is very simple. We assume that there is another "interloping" sector which breaks the $SU(N)_{TC}$ group completely, so that all the $SU(N)_{TC}$ gauge bosons are massive. If this symmetry breaking occurs at a scale where the $SU(N)_{TC}$ coupling is weak, the effective theory below this scale will consist of massless fermions interacting through contact interactions such as those of eq. (8), and $h^{-1} = 0$. On the other hand, if the $SU(N)_{TC}$ coupling gets strong at a scale higher than the scale at which the interloping sector becomes strong, then the $SU(N)_{TC}$ interactions will break the chiral symmetry and $h^{-1} = \frac{3}{TC}$. In this case, the low-energy effective theory contains only the Nambu-Goldstone bosons resulting from the chiral symmetry breaking. If we assume that the transition between the two limits just described is smooth, then we can obtain a model in which $h^{-1} = \frac{3}{TC}$ by tuning the model between the two limits described above. We expect the dynamics of such a model to be similar to the fine-tuned NJL model discussed in the previous section.

Actually, since we are ultimately interested in obtaining a condensate for quarks carrying only color indices, we will consider models where the technicolor group is broken in the pattern

$$SU(N)_{TC} \rightarrow SU(N)_C \times SU(N)_0; \quad (9)$$

where $SU(N)_C$ is weakly coupled at the symmetry-breaking scale. In this case, the surviving $SU(N)_0$ group is weakly coupled, and we will later take $N = 3$ and identify it with QCD.

To make these ideas specific, consider a theory with a gauge group

$$SU(N)_{TC} \times SU(K) \times SU(K)_T \times SU(N)_C; \quad (10)$$

and fermion content

$$\begin{aligned} L_j; R_j & (N; 1; 1; 1); & j = 1; \dots; F; \\ L & (N; K; 1; 1); \\ R & (1; K; 1; N); \\ R & (N; 1; K; 1); \\ L & (1; 1; K; N); \end{aligned} \quad (11)$$

For simplicity, we will impose a discrete symmetry to set the coupling constants of the two $SU(K)$ groups equal. The first three group factors in eq. (10) are "strong" gauge groups which will form fermion condensates, while $SU(N)_C$ is weakly coupled in the energy regime in which we are interested. Note that the fermion content of each gauge group consists of vector-like fermions in the fundamental representation. Thus, all gauge anomalies are cancelled in this theory. More importantly, this means that as long as only one of the gauge couplings is strong, we can analyze the symmetry breaking pattern by appealing to our understanding of QCD-like theories.

The coupling strength of the gauge groups $SU(N)_{TC}$ and $[SU(K) \times SU(K)]_I$ can be characterized by the values of the gauge coupling constants g_{TC} and g_I evaluated at some fixed scale μ_0 . Alternatively, we can characterize the gauge couplings by mass scales Λ_{TC} and Λ_I at which the respective gauge couplings become strong enough to trigger chiral symmetry breaking. There are several possibilities for the low-energy dynamics of this model, depending on the relative magnitudes of these parameters.

Suppose first that $g = g_{TC}$ at the scale μ_0 , so that $\Lambda_I \gg \Lambda_{TC}$. Then $SU(N)_{TC}$ is weakly coupled at the scale Λ_I , and the $SU(K)_{I^0}$ interactions give rise to the condensates

$$\langle \bar{h}_{L,R} i; \bar{h}_{L,R} i \rangle \sim \Lambda_I^3: \quad (12)$$

This condensate results in the chiral symmetry breaking pattern

$$\begin{aligned} SU(N)_L \times SU(N)_R &\rightarrow SU(N)_{L+R}; \\ SU(N)_L \times SU(N)_R &\rightarrow SU(N)_{L+R}; \end{aligned} \quad (13)$$

giving rise to $2(N^2 - 1)$ potential NGB's. (Each of the $SU(K)$ factors acts like a copy of QCD.) The condensate eq. (12) results in the gauge symmetry breaking pattern

$$SU(N)_{TC} \times SU(N)_I \rightarrow SU(N)_{I^0}: \quad (14)$$

$N^2 - 1$ potential NGB's are therefore eaten by the broken gauge bosons, which acquire masses of order $g_{TC} \Lambda_I$. The remaining $N^2 - 1$ potential NGB's transform in the adjoint representation of the unbroken $SU(N)_{C^0}$ gauge group, and therefore acquire masses of order $g_C \Lambda_I$.

Through all of this, the fermions remain massless, and $\langle \bar{h} i \rangle = 0$. The effective theory at scales $\mu \gg g_C \Lambda_I$ therefore consists of the massless fermions subject to various contact interactions suppressed by inverse powers of Λ_I . This is qualitatively similar to the NJL model far into the unbroken phase.

Now suppose that $g_C = g_I$ at the scale μ_0 , so that $T_C = I$. In this case, the $SU(N)_{TC}$ gauge coupling becomes strong at a scale where the $[SU(K) \times SU(K)]_I$ gauge coupling is weak. This results in the formation of the condensate

$$h \bar{\psi}_{L,R} + \bar{\psi}_{L,R} i \gamma_5 : \quad (15)$$

(There are other possibilities for the alignment of the condensate, but a calculation along the lines of ref. [9] which treats the $SU(K)$ and electroweak gauge groups perturbatively shows that this vacuum alignment is chosen for sufficiently weak $SU(K)$ coupling. This is consistent with the lore that condensates will align so as to preserve the maximal gauge symmetry.) This condensate results in the chiral symmetry-breaking pattern

$$SU(F+K)_L \times SU(F+K)_R \rightarrow SU(F+K)_{L+R}; \quad (16)$$

giving rise to $(F+K)^2 - 1$ potential NGB's. This gives rise to the gauge symmetry breaking pattern

$$[SU(K) \times SU(K)]_I \rightarrow SU(K)_{I^0}; \quad (17)$$

$K^2 - 1$ NGB's resulting from the condensate eq. (15) are therefore eaten by the broken gauge bosons, which acquire masses $\sim g_{TC}$.

In a basis where the first F entries correspond to ψ_L and the last K entries correspond to ψ_R , we can classify the potential NGB's associated with the broken axial generators as follows:

$$\begin{pmatrix} X & 0 \\ 0 & 0 \end{pmatrix} \quad F^2 - 1 \text{ physical NGB's}, \quad (18)$$

$$\begin{pmatrix} 0 & 0 \\ 0 & Y \end{pmatrix} \quad K^2 - 1 \text{ eaten NGB's}, \quad (19)$$

$$\begin{pmatrix} 0 & E \\ E^Y & 0 \end{pmatrix} \quad 2FK \text{ massive pseudo-NGB's}, \quad (20)$$

$$\begin{pmatrix} K & 1 & 0 \\ 0 & F & 1 \end{pmatrix} \quad 1 \text{ physical "axion"}. \quad (21)$$

The fate of the NGB's of eqs. (18) and (19) should be obvious. The NGB's of eq. (20) can be viewed as $2F$ NGB's transforming in the fundamental representation of the unbroken $SU(K)_{I^0}$ gauge group. These NGB's therefore acquire masses of order g_{TC} . The "axion" of eq. (21) can be associated with the charge

$$\begin{aligned} L &: +K; & R &: -K; \\ L &: -F; & R &: +F; \\ L &: +F; & R &: -F; \end{aligned} \quad (22)$$

The current associated with this charge has no $SU(N)_{TC} [SU(K) \times SU(K)]$ anomalies, but it does have a $SU(N)_C$ anomaly. Therefore, the axion will acquire a mass of order $\frac{2}{C} = \frac{TC}{C}$.

The R and L fermions remain massless at this stage, and transform in the fundamental representation of the unbroken $SU(K)_{I^0}$ gauge group. At the scale Λ_{I^0} , the $SU(K)_{I^0}$ gauge group becomes strong, resulting in the formation of the condensate

$$h_{LR}^{-1} \sim \Lambda_{I^0}^3 \quad (23)$$

This results in the symmetry-breaking pattern

$$SU(N)_L \times SU(N)_R \rightarrow SU(N)_{L+R}; \quad (24)$$

giving rise to $N^2 - 1$ potential NGB's. These potential NGB's transform in the adjoint representation of the unbroken $SU(N)_C$ gauge group, and therefore acquire masses of order $g_C \Lambda_{I^0}$. Thus, $h_{LR}^{-1} \sim \Lambda_{TC}^3$ and the low-energy theory consists entirely of the F^2 NGB's resulting from the symmetry breaking at the scale Λ_{TC} . This is qualitatively similar to the NJL model far into the broken phase.

We now consider the nature of the transition between the two limits just described. If we assume that the order parameter h_{LR}^{-1} is continuous across the transition (i.e. that the transition is second order), then we can tune the $[SU(K) \times SU(K)]$ coupling so that $h_{LR}^{-1} \sim \Lambda_{TC}^3$, where $\Lambda_{TC} \sim \Lambda_{I^0}$. In this case, both $SU(N)_{TC}$ and $[SU(K) \times SU(K)]$ are becoming strongly coupled near the scale Λ , but we can analyze this model using continuity arguments.

In the netuned limit, it seems plausible that the fermions are propagating degrees of freedom with mass m . The reason is that in the unbroken phase, there are poles at zero momentum transfer in Green's functions coming from an intermediate state consisting of a single fermion. If the transition is second order, we expect the position of these poles to be continuous across the transition, and the fermions to be massive propagating states.

Also, in the netuned limit, we expect the decay constant f of the NGB's to be finite. To see this, define

$$J_{AB}(q) = \int d^4x e^{iq \cdot x} \langle 0 | T J_A(x) J_B(0) | 0 \rangle \quad (25)$$

where

$$J_A = \bar{\psi} \gamma_5 T_A \psi \quad (26)$$

are the spontaneously broken axial currents. From Goldstone's theorem, we know that $J_{AB}(q)$ has a pole at $q^2 = 0$:

$$J_{AB}(q) \sim (q \cdot q - q^2) \frac{f_{AB}^2}{q^2} \quad \text{as } q^2 \rightarrow 0; \quad (27)$$

where f_{AB}^2 is the matrix of NGB decay constants. From the Jackiw-Johnson sum rule (eq. (54) below), we know that the NGB decay constants vanish if the fermion self-energy vanishes. Therefore, if we assume that the transition is second order, then we expect the decay constants will be of order m . (This is made explicit in the approximations to the Jackiw-Johnson sum rule which we will consider below.)

We now turn to the low-energy effective lagrangian for this model. At sufficiently low energies $\ll f$, the effective lagrangian describes the interactions among the NGB's (and possibly the fermions if their masses are smaller than f). In this effective lagrangian, the $SU(N)_F$ symmetry is realized nonlinearly on the NGB fields. As a result, this effective lagrangian breaks down for processes involving momentum transfers $p^2 \sim f^2$. One signal for this fact is that an infinite number of operators are important for such processes. We therefore conclude that the theory must contain new particles and interactions at scales $\lesssim f_{TC}$. The simplest possibility is that the symmetry is realized linearly in the effective lagrangian at scales $\ll f$, implying that the theory contains light scalars with the quantum numbers of the field in eq. (6). (It is difficult to imagine a reasonable alternative if one accepts the assumption that the condensate can be re-tuned to be small.)

In fact, we may conjecture that the effective lagrangian at scales below f_{TC} is in the same universality class as the NJL model. To make this statement precise, we write the effective lagrangian as

$$L_e = L_{NJL} + L \quad (28)$$

where L_{NJL} is the NJL lagrangian of eq. (1) and L contains all remaining terms. We conjecture that the four-fermion coupling in L_{NJL} is close to its critical value, and that the coefficients of the operators in L are sufficiently small that the theory is in the same universality class as the NJL model. Similar assumptions have been discussed elsewhere for the case of top-condensate models [12] and "strong extended technicolor" models [13].

4. Schwinger-Dyson Analysis

We can flesh out our picture of the NJL limit by considering the Schwinger-Dyson equations for the theory we have described. The full Schwinger-Dyson equations are clearly intractable, and in order to make progress we must truncate these equations and

hope that what remains captures the essential physics we wish to discuss. We do not pretend that this analysis justifies the dynamical assumptions made in the last section, but we will see that these assumptions are incorporated in a simple and natural way in our analysis. Also, we can use the truncated Schwinger-Dyson equations to make crude quantitative estimates for various quantities of interest. This section can be skimmed by a reader interested mainly in the application of the model discussed above to electroweak symmetry breaking.

4.1. Fermion Self-Energy

$$\text{If we define} \quad \begin{aligned} &0 \quad 1 \\ &\text{@} \quad A ; \end{aligned} \quad (29)$$

the full Schwinger-Dyson equation for the propagator is

$$iS^{-1}(p) = \sum_{r;r^0}^X \int \frac{d^4k}{(2\pi)^4} g_r \text{Tr}_A S(k) \text{Tr}_{r^0B}(k;p) \text{Tr}_{r^0B} G^{AB}(k-p); \quad (30)$$

where $r;r^0$ run over the fermion representations of the gauge group, Tr_A is the gauge boson vertex function, and G^{AB} is the gauge boson propagator. We restrict the fermion propagator to have the block-diagonal form

$$S = \begin{pmatrix} S & \\ & S \end{pmatrix} : \quad (31)$$

This is consistent with the full Schwinger-Dyson equations, and is also the correct form in both of the limits considered in the previous section. The Schwinger-Dyson equation for S then involves only the exchange of $SU(N)_{TC}$ gauge bosons. We will assume in what follows that the $[SU(K) \times SU(K)]_{\frac{1}{2}}$ coupling can be chosen so that the mass M of the $SU(N)_{TC}$ gauge bosons has any desired value.

We will follow a venerable tradition and approximate the full gauge boson vertex function by its tree-level value (with a running coupling evaluated at an appropriate momentum scale), approximate the gauge boson propagators by their asymptotic forms, and neglect fermion wave-function renormalization. (See [10] for more details.) The resulting integral equation for the fermion self-energy is then

$$(p^2) = \int \frac{dk^2}{4\pi^2} G(\max\{k^2; p^2; M^2\}) \frac{k^2 - (k^2)}{k^2 + (k^2)}; \quad (32)$$

where M is the mass of the $SU(N)_{TC}$ gauge bosons,

$$G(p^2) = \frac{3g^2(p^2)C_2}{p^2} \quad (33)$$

can be viewed as the strength of the one-gauge-boson-exchange interaction, and C_2 is the Casimir of the fermion representation.

For $p^2 \ll M^2$, the right-hand side of eq. (32) is independent of p^2 , and we have

$$(p^2) = m \quad \text{for } p^2 \ll M^2: \quad (34)$$

This situation is rather reminiscent of the NJL model. In fact, we can write

$$\begin{aligned} m &= m \frac{G(M^2)}{4} \int_0^{M^2} dk^2 \frac{k^2}{k^2 + m^2} + m \\ &= m \frac{G(M^2)}{4} M^2 - m^2 \ln \frac{M^2}{m^2} + m; \end{aligned} \quad (35)$$

where

$$m \int_{M^2}^{\infty} \frac{dk^2}{4} G(k^2) \frac{k^2}{k^2 + m^2} \quad (36)$$

depends only on the behavior of (p^2) with $p^2 > M^2$. If we naively identify $G(M^2)$ with the NJL coupling G and identify M with the NJL cutoff, then eq. (35) is similar to the NJL gap equation with the addition a "counterterm" $-m$. To understand the relation between eq. (35) and the NJL gap equation, we note that the NJL coupling G includes the effects of integrating out all field models with $p^2 > \Lambda^2$, while $G(M^2)$ has no such interpretation. The role of the term $-m$ is to include the effects of the high-energy gauge bosons.

For $p^2 > M^2$, eq. (32) is the same as the gap equation for massless gauge bosons, which has been studied by many authors. The integral equation can be converted to a differential equation:

$$p^2 (p^2) - \frac{G^0(p^2)}{G^0(p^2)} p^2 (p^2) - \frac{G^0(p^2)}{4} (p^2) = 0; \quad (37)$$

where the prime denotes differentiation with respect to p^2 . This equation has been linearized in (p^2) , which is completely justified in the present case, since we are interested in the regime where $p^2 \gg M^2 \gg m^2(p^2)$. Using the known asymptotically-free behavior of $G(p^2)$ for large p^2 , we can write the ultraviolet boundary condition

$$(p^2) \sim \frac{3}{p^2} G(p^2) \quad \text{as } p^2 \rightarrow \infty; \quad (38)$$

for some constant C_2 . Eq. (37) has an infrared boundary condition

$$\begin{aligned} G^0(M^2) &= m \int_0^{M^2} \frac{dk^2}{4k^2} \frac{k^2}{k^2 + m^2} \\ &= m \frac{G^0(M^2)}{4} - M^2 - m^2 \ln \frac{M^2}{m^2} : \end{aligned} \quad (39)$$

It is easily verified by direct substitution that eq. (37) with these boundary conditions is equivalent to eq. (32).

In this formalism, we can explicitly see the fine-tuning necessary to obtain $m = M$. Because eq. (37) is linear in $F(p^2)$, once we impose the ultraviolet boundary condition eq. (38), $F(p^2)$ is completely determined up to an overall constant. We can therefore write

$$F(p^2) = m \frac{F(p^2)}{F(M^2)}; \quad (40)$$

where $F(p^2)$ is the solution to eq. (37) satisfying the boundary condition

$$F(p^2) \rightarrow G(p^2) \text{ as } p^2 \rightarrow 1 : \quad (41)$$

The infrared boundary condition can then be written

$$\frac{M^2 R(M^2)}{4} - \frac{M^2 G^0(M^2) F(M^2)}{4 F^0(M^2)} = 1 + O(m^2/M^2); \quad (42)$$

The left-hand side depends only on M and is of order 1 for $M \gg M_{TC}$. We see that M must be equal to a critical value M_{crit} to an accuracy $O(m^2/M^2)$ if we want $m = M$. This in turn means that the $[SU(K) \times SU(K)]$ gauge coupling (evaluated at the scale μ_0) must be fine-tuned to an accuracy $O(m^2/M^2)$.

To show that eq. (42) has a solution for some M^2 , we note that the quantity $R(p^2)$ defined above satisfies the differential equation

$$R^0(p^2) = G^0(p^2) - R^2(p^2); \quad (43)$$

with ultraviolet boundary condition

$$R(p^2) \rightarrow G(p^2) \text{ as } p^2 \rightarrow 1 : \quad (44)$$

Thus, the left-hand side of eq. (42) is small for $M \gg M_{TC}$. Because $R^0(p^2) < G^0(p^2)$, we have $R(p^2) < G(p^2)$, and thus

$$M^2 R(M^2) > 3g^2(M^2)C_2 : \quad (45)$$

Therefore, as long as $g^2(M^2)$ is sufficiently large, the left-hand side of eq. (42) will be greater than unity and the differential equation has a solution. This is just the condition that the gauge coupling becomes sufficiently strong to trigger chiral symmetry breaking.

Note also that $\chi(0)$ is continuous as a function of M in this formalism.

4.2. Light Scalars

We now address the question of the existence of light scalars in this model. Our strategy is to compute $\pi\pi \rightarrow \pi\pi$ scattering in the color-singlet scalar and pseudo-scalar channels. The light scalars will manifest themselves as poles in the s channel. We will compute the scattering amplitude in the ladder approximation shown in fig. 1.

We will make several further simplifications. Since the phase transition is second order in this formalism, it is sufficient to establish the existence of the scalars in the unbroken phase where $\chi = 0$. Also, we will compute the amplitude in the kinematic regime shown in fig. 1. The Mandelstam variables are then $s = 0, t = u = p^2$. A light scalar with mass $m_s \ll M$ in the s channel will result in the behavior

$$T(p^2) \sim \frac{1}{m_s^2} + \dots \quad (46)$$

In this kinematic regime, only a single form-factor contributes to the amplitude, and we can write the following simple integral equation for the scattering amplitude $T(p^2)$:

$$T(p^2) = G(\max(p^2; M^2)g) + \int_0^Z \frac{dk^2}{4} G(\max(k^2; p^2; M^2)g) T(k^2) \quad (47)$$

Since we are in the unbroken phase, this equation holds for all flavor channels and for both the scalar and pseudoscalar channels.

For $p^2 \ll M^2$, the right-hand side of eq. (47) is independent of p^2 , and we have

$$T(p^2) = T \quad \text{for } p^2 \ll M^2 \quad (48)$$

Again, this situation is very similar to the NJL model, where the scattering amplitude considered here is independent of t and u .

For $p^2 > M^2$, the amplitude satisfies the differential equation

$$T'(p^2) - \frac{G'(p^2)}{G(p^2)} T(p^2) - \frac{G'(p^2)}{4} T(p^2) = 0; \quad (49)$$

where the prime again denoted differentiation with respect to p^2 . This is the same differential equation satisfied by χ above, but with different boundary conditions and with the

crucial difference that it holds for arbitrarily large T . Eq. (49) has the ultraviolet boundary condition

$$T(p^2) \rightarrow cG(p^2) \text{ as } p^2 \rightarrow 1; \quad (50)$$

where c is a dimensionless constant, and an infrared boundary condition

$$T^0(M^2) = G^0(M^2) \left(1 + \frac{T(M^2)M^2}{4} \right); \quad (51)$$

Once we impose the ultraviolet boundary condition, we can write the solution as

$$T(p^2) = T \frac{F(p^2)}{F(M^2)}; \quad (52)$$

where $F(p^2)$ is the function introduced in eq. (40) above. The infrared boundary condition can then be written

$$\begin{aligned} \frac{1}{T} &= \frac{F^0(M^2)}{G^0(M^2)F(M^2)} - \frac{M^2}{4} \\ &= \frac{1}{R(M^2)} - \frac{M^2}{4} \end{aligned} \quad (53)$$

where $R(p^2)$ is the function defined in eq. (42). Eq. (53) shows explicitly that a solution exists. Of more interest is the fact that eq. (53) shows that as we approach the re-tuning condition eq. (42), T becomes singular.^y This shows that there are light scalars in the theory near the critical point in the ladder approximation.

4.3. Decay Constants

We now discuss the decay constants which result from the fermion condensation. They are given by the Jackiw-Johnson sum rule [11]

$$f_{AB}^2 = \frac{Z}{(2)^4} \text{tr} \int \frac{d^4k}{(2)^4} \text{tr} \left[\gamma_5 (Z_J T_A) S(k) e_B(k; k) S(k) \right] + f^2; \quad (54)$$

where e_A is the current vertex function with the NGB pole removed, $S(k)$ is the propagator of the fermions in the current, Z_J is the renormalization constant of the current operator, and f^2 is a counterterm.

^y In an exact treatment, we certainly expect the same critical value of M for the chiral symmetry-breaking transition and the appearance of light scalars, but a priori we have no right to expect the critical values of M to coincide exactly in our approximation, since it is not clear that the gap equation eq. (32) and the partial summation we have performed to compute T are part of a single consistent approximation.

If we make the same approximations made in deriving eq. (32) we obtain $f_{AB}^2 = f_{AB}^2$, with

$$f^2 = \frac{N}{16\pi^2} \int_0^{\Lambda^2} dk^2 \frac{k^2 (k^2 - 2k^2 + k^2)}{[k^2 + k^2 (k^2)]^2} \quad (55)$$

$$= \frac{N}{16\pi^2} m^2 \ln \frac{M^2}{m^2} + \int_{M^2}^{\Lambda^2} dk^2 \frac{k^2 (k^2)}{k^2} : \quad (56)$$

The integral in eq. (56) is of order m^2 , so the first term dominates for $m \ll M$. If we approximate (k^2) for $k^2 > M^2$ by its asymptotic form eq. (38), we obtain

$$f^2 = \frac{N m^2}{16\pi^2} \ln \frac{M^2}{m^2} + \frac{1}{2} : \quad (57)$$

To summarize, we see that the gap-equation analysis incorporates many of the features we expected on the basis of qualitative arguments of section 3: the transition is second order, the fine-tuning is manifest, the decay constants are small compared to the compositeness scale, and the light scalars are present.

5. Application to Electroweak Symmetry Breaking

5.1. "Top-Mode" Model

In order to apply the model described above to electroweak symmetry breaking, we take $N = 3$ and identify $SU(3)_C$ with ordinary color. We also take $F = 2$ and identify

$$= \begin{pmatrix} t \\ b \end{pmatrix} : \quad (58)$$

The low energy theory contains two Higgs doublets; in the notation of eq. (6), we have

$$= \begin{pmatrix} H_t^0 & H_b^+ \\ H_t^- & H_b^0 \end{pmatrix} ; \quad (59)$$

where H_t and H_b are Higgs doublet fields.

This theory may seem to be a phenomenological disaster, since it appears that custodial symmetry of the technicolor interactions implies $m_t = m_b$, and the theory contains an axion (see eq. (22)) with a decay constant near the weak scale. Such axions are ruled out by a combination of laboratory experiments and astrophysical considerations [16]. However, the model actually does not suffer from these problems, as we will describe below.

In the standard model, we know that the custodial $SU(2)$ symmetry is broken by the weak hypercharge. In fact, the hypercharge assignments are

$$Y(\ell_L) = \frac{1}{3}; \quad Y(t_R) = \frac{4}{3}; \quad Y(b_R) = -\frac{2}{3}; \quad (60)$$

Thus, $U(1)_Y$ gauge boson exchange will mediate an attractive force between ℓ_L and \bar{t}_R and a repulsive force between ℓ_L and \bar{b}_R . Although this force is weak, when the theory is tuned near the critical point, the contribution from $U(1)_Y$ gauge boson exchange can result in $\tilde{m}_b = 0$ while $\tilde{m}_t \neq 0$. The phenomenon of the amplification of small perturbations near a critical point has been discussed in the context of technicolor theories [14] and in top-condensate models [15], where it has been dubbed "critical instability."

To see how this phenomenon emerges from the gap equation, write

$$\begin{aligned} Y &= Y_L P_L + Y_R P_R \\ &= \frac{1}{3} 1_2 P_L + \frac{1}{3} 1_2 + \frac{2}{3} P_R; \end{aligned} \quad (61)$$

where $P_{L,R}$ are the left- and right-handed helicity projection operators. The self-energy is a diagonal 2×2 matrix

$$= \begin{pmatrix} \tilde{m}_t & 0 \\ 0 & \tilde{m}_b \end{pmatrix}; \quad (62)$$

The gap equation has the same form as eq. (32) with the replacement

$$G(p^2) = \frac{3}{4} \frac{g_{TC}^2(p^2) C_2 + g_Y^2(p^2) Y_L Y_R}{p^2}; \quad (63)$$

In the approximations we are making, the Schwinger-Dyson equations for the fermion masses split into separate equations for the top- and bottom-quark self-energies. Suppose now that the theory is tuned so that $\tilde{m}_t = M$. Then

$$\begin{aligned} G_t(M^2) &= G_{\text{crit}} + O(\tilde{m}_t^2 - M^2); \\ G_b(M^2) &= G_{\text{crit}} - \frac{2g_Y^2(M^2)}{M^2} + O(\tilde{m}_t^2 - M^2); \end{aligned} \quad (64)$$

Thus, $\tilde{m}_b = 0$ as long as

$$g_Y > \frac{m_t}{M}; \quad (65)$$

(Note that we can consider the possibility that the scale M is chosen so that the hierarchy $\tilde{m}_t = \tilde{m}_b$ emerges entirely as a result of $U(1)_Y$ custodial symmetry breaking. However, this requires $M \sim 1 \text{ TeV}$, and results in a top mass $m_t \sim 500 \text{ GeV}$, as we will see below.)

We now consider the low-energy theory for the case where $\mu_b = 0$. Taking into account the breaking of $SU(2)_R$ due to hypercharge, the symmetry breaking pattern can be written

$$SU(2)_W \times U(1)_L \times U(1)_{BR} \rightarrow U(1)_{EM} \times U(1)_{BR}; \quad (66)$$

resulting in 3 potential NGB's which are eaten by the massive electroweak gauge bosons. The effective theory below the scale Λ_{TC} contains two Higgs doublets with

$$\langle H_t \rangle = v; \quad \langle H_b \rangle = 0; \quad (67)$$

H_b is an unbroken doublet with mass $\sim M$, which can be far above the electroweak scale. In this model as discussed so far, all fermions other than the top quark are exactly massless.

The axion of eq. (21) is not present, but this is only because there is an unbroken axial $U(1)$ symmetry in the low-energy theory as a result of the massless b quark. Clearly, this symmetry, as well as the flavor symmetries of the remaining quarks and leptons, must be broken somehow if the theory is to account for the observed fermion masses and mixings. This is exactly the situation in technicolor theories, where the technifermion condensate does not give mass to ordinary quarks and leptons in the absence of additional "extended technicolor" interactions which connect the technifermions with the quarks and leptons.

Thus, as in technicolor, we assume that the effective theory at the scale Λ_{TC} contains four-fermion "extended technicolor" (ETC) interactions of the form

$$\frac{1}{M_E^2} (\bar{\psi} \psi) (\bar{f} f) \quad (68)$$

where the ψ 's represent third-generation quark fields and the f 's represent quark or lepton fields which are $SU(N)_{TC}$ singlets. We will not be concerned here with the dynamics which gives rise to these operators, or with exhibiting a completely realistic theory. Our aim is simply to show that there is in principle no obstacle to making the theory realistic.

Four-fermion interactions of the form of eq. (68) can give rise to masses for all the quarks and leptons, but they necessarily preserve a $U(1)$ symmetry which counts the number of quarks from the first two generations.^y If this symmetry were exact, elements of the Cabibbo-Kobayashi-Maskawa matrix involving the third generation would vanish identically, and so this $U(1)$ symmetry must be broken somehow. This $U(1)$ symmetry can be broken by dimension-7 interactions such as

$$\frac{1}{M_E^3} (\bar{Q}_L \psi_L) (\bar{\psi}_L \psi_R); \quad (69)$$

^y I am indebted to R. Sundrum for pointing this out to me.

where Q_L is a left-handed quark field from the first two generations. Alternatively, we can imagine that all quarks carry $SU(3)_{TC}$ rather than $SU(3)_C$, and the fact that the top quark is the only fermion which condenses is explained by critical instability, due for example to additional higher-dimension operators. In this case, all masses and mixings can be obtained from operators of the form of eq. (68). Thus, there seems to be no obstacle in principle to incorporating light fermions into the theory.

The scale $M_E > M_{TC}$ is associated with new interactions, such as massive gauge boson exchange. It may seem that much of the motivation for writing a renormalizable model is lost once we introduce these non-renormalizable terms. However, unlike the four-fermion coupling which drives the symmetry breaking in the NJL model, these four-fermion interactions are perturbatively weak, and no exotic dynamics is required to generate them. The present model can be viewed as a valid effective theory for scales up to M_E , which can be well above the compositeness scale, while the NJL model cannot be extended above scales $\sim \frac{1}{\sqrt{4\pi G}}$.

Through the standard ETC mechanism, the four-fermion interactions of eq. (68) will give rise to fermion masses of order

$$m_f \sim \frac{1}{M_E^2} \int^M \frac{d^4 k}{(2\pi)^4} \text{tr} \frac{1}{\not{k}} m_t \frac{1}{\not{k}} \sim \frac{M^2 m_t}{4\pi^2 M_E^2} : \quad (70)$$

This can be viewed as due to the presence of a condensate

$$\langle \bar{\psi} \psi \rangle \sim \frac{M^2 m_t}{4\pi^2} : \quad (71)$$

Alternatively, using the NJL model as a guide, we expect that the ETC interactions eq. (68) will give rise to Yukawa couplings between the ordinary fermions and the composite Higgs field of order

$$Y \sim \frac{M^2}{4\pi^2 M_E^2} \frac{m_t}{v} \quad (72)$$

This gives rise to the fermion masses eq. (70) when the Higgs field acquires a vacuum expectation value.

We now consider the case where M is far above the weak scale. In this case, the effective lagrangian at low scales is indistinguishable from the standard model, and we expect the renormalization group analysis of ref. [3] to be valid. This scenario necessitates a severe re-tuning of the order $m_t^2 = \frac{2}{3} M_{TC}^2$, and results in a top-quark mass of 230 GeV for $M_{TC} \sim 10^{15}$ GeV (200 GeV for $M_{TC} \sim 10^{19}$ GeV). Such values for the top quark mass are disfavored by a global analysis of the radiative effects in the standard model, which gives $m_t < 201$ GeV at the 95% confidence level [17].

Note that if M is far above the weak scale, eq. (70) shows that realistic fermion masses can easily be generated with M_E sufficiently large to suppress flavor-changing neutral currents which can arise from four-fermion operators of the form

$$\frac{1}{M_E^2} (\bar{f}f)(\bar{f}f): \quad (73)$$

Of course, this feature is obtained only at the expense of the fine-tuning of the $SU(N)_{TC}$ breaking scale.

If M is close to the weak scale, the leading-log effects which form the basis of the predictions of ref. [3] are not expected to be important. Using eq. (57) and noting that $v = f = \frac{p}{\sqrt{2}} = 246 \text{ GeV}$, we find e.g.

$$m_t \simeq 440 \text{ GeV} \quad \text{for } M \simeq 10 \text{ TeV}: \quad (74)$$

While the approximations used to obtain this result are rather crude, it is clear that this is well outside the limits on the top mass coming from α_s -parameter constraints. It is of course possible that other sources of custodial symmetry breaking (for example from higher-dimensional operators from the ETC sector) can cancel the effects of the top quark on the α_s parameter, although there is no good reason to expect this to happen.

Since the "top-mode" model described above seems to have phenomenological troubles for all reasonable choices of parameters, we now turn to models in which other fermions are responsible for the electroweak symmetry breaking.

5.2. Technicolor-like Models

In this subsection, we consider models in which the fermions are new strongly-interacting fermions. In this case, the direct connection between the top-quark mass and the electroweak scale is lost, but raising the compositeness scale may make these theories more phenomenologically attractive than technicolor theories (at the expense of fine-tuning). We will argue that because of the critical instability mechanism discussed above, we cannot have the electroweak symmetry broken by new fermions without additional fine-tuning.

There are several strong constraints on the new fermions, which we will generically denote by T . We assume that the new fermions occur in weak doublets so that the condensate of these fermions is an $I_3 = \frac{1}{2}$ order parameter. The masses of the T fermions are constrained by demanding that they give rise to the correct value of the electroweak

scale v . Using the approximation eq. (57) to the Jackiw-Johnson sum rule, we obtain

$$\sum_j m_j^2 = \frac{8}{3} \frac{v^2}{M^2} \ln \frac{M^2}{\bar{m}^2} + \frac{1}{2} \quad (75)$$

$$(400 \text{ GeV})^2 \quad \text{for } M = 10 \text{ TeV}; \quad (76)$$

where the sum runs over all T fermions and \bar{m} is their average mass. (This formula will have large corrections if $\ln(M^2/\bar{m}^2) \ll 1$.) This shows that in any such model, we expect new heavy weak-doublet fermions which will be accessible in future experiments.

In order that they do not contribute significantly to the β -parameter, T fermions in the same weak doublet must be nearly degenerate in mass. In particular, we must avoid large mass splittings induced by $U(1)_Y$ couplings through critical instability, as discussed above. Of course, we can always add additional interactions to cancel this effect, but the strength of these interactions must be re-tuned, and we want to see whether re-tuning can be avoided.

To avoid the critical instability mechanism described in the last section, we must take the weak doublet T fermions to carry zero $U(1)_Y$ charge. The $SU(2)_W \times U(1)_X$ singlet T fermions must carry $U(1)_Y$ charge so that the condensate $\langle \bar{T}_L T_R \rangle$ preserves $U(1)_{EM}$. Thus, the only possibility is to take the extra fermions to transform under $SU(3)_{TC} \times SU(2)_W \times U(1)_X$ as

$$\begin{aligned} T_L &= (3; 2; 0); \\ U_R &= (3; 1; 1); \\ D_R &= (3; 1; -1); \end{aligned} \quad (77)$$

When $SU(3)_{TC}$ becomes strong, they give rise to a condensate

$$\langle \bar{U}_L U_R \rangle; \langle \bar{D}_L D_R \rangle \neq 0; \quad (78)$$

where we have defined

$$T_{L,R} = \begin{pmatrix} U \\ D \end{pmatrix}_{L,R} : \quad (79)$$

The theory has an approximate discrete custodial symmetry

$$\begin{aligned} T_{L,R} &\xrightarrow{T} \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix} T_{L,R} \quad \times T_{L,R}; \\ B &\xrightarrow{T} B; \\ W &\xrightarrow{T} X W X; \end{aligned} \quad (80)$$

where B and W are the $U(1)_Y$ and $SU(2)_W$ gauge fields, respectively. If the T fields were the only fermions in the theory, this custodial symmetry would be exact, and we would have $m_U = m_D$. In the full theory, this symmetry is violated by the $U(1)_Y$ gauge couplings to the quarks and leptons and by ETC operators. In particular, there must be strong ETC operators to give rise to the large top mass, so that there will in general be custodial-symmetry violating four-fermion couplings among the T fermions with strength $1/M_E^2 \sim 1/M^2$, which will give rise to a large splitting $m_U - m_D$. (Note that we cannot explain the large top-quark mass by taking the third-generation quarks to transform under $SU(N)_{TC}$, since the effect of $U(1)_Y$ would give $m_t = m_T$.)

We have argued that additional fine-tuning is required in "non-minimal" models incorporating NJL-like dynamics to break electroweak symmetry in order to avoid an unacceptably large value for the β -parameter. If we accept this additional fine-tuning, however, there seems to be no obstacle to making such theories fully realistic.

6. Conclusions

We have considered a renormalizable model which we argued can break the electroweak symmetry via NJL-like dynamics. The crucial dynamical assumption is that the chiral symmetry-breaking transition in the model is second order. The fine-tuning needed to obtain a large hierarchy between the weak scale and the compositeness scale is manifested in the fine-tuning of a gauge coupling constant. We have seen that the minimal "top-mode" standard model can be obtained from such a model, as well as models in which new fermions are responsible for electroweak symmetry breaking. In the latter case, we have argued that additional fine-tuning is required to avoid unacceptably large corrections to the electroweak β -parameter.

7. Acknowledgements

I would like to thank R. Rattazzi and especially R. Sundrum for discussions on the topic of this paper. This work was supported by the Director, Office of Energy Research, Office of High Energy and Nuclear Physics, Division of High Energy Physics of the U.S. Department of Energy under Contract DE-AC03-76SF00098.

8. References

- [1] G. Jona-Lasinio and Y. Nambu, Phys. Rev. 122, 345 (1961).
- [2] Y. Nambu, EFI-88-39 (1988), EFI-88-62 (1988), and EFI-89-08 (1989) (unpublished); V. A. Miransky, M. Tanabashi, and K. Yamawaki, Mod. Phys. Lett. A 4, 1043 (1989); Phys. Lett. 221B, 177 (1989); W. Marciano, Phys. Rev. Lett. 62, 2793 (1989).
- [3] W. A. Bardeen, C. T. Hill, and M. Lindner, Phys. Rev. D 41, 1647 (1990).
- [4] For example, see M. A. Luty, Phys. Rev. D 41, 2893 (1990); M. Suzuki, Phys. Rev. D 41, 3457 (1990); C. T. Hill, M. A. Luty, and E. A. Paschos, Phys. Rev. D 43, 3011 (1991); R. N. Mohapatra and K. S. Babu, Phys. Rev. Lett. 66, 556 (1991); P. H. Frampton, O. Yasuda, Phys. Rev. D 44, 3709 (1991); M. Suzuki, Phys. Rev. D 44, 3628 (1991); W. A. Bardeen, M. Carena, T. E. Clark, K. Sasaki, and C. Wagner, Nucl. Phys. B 369, 33 (1992).
- [5] C. T. Hill, Phys. Lett. 266B, 419 (1991); D. E. Clague and G. G. Ross, Nucl. Phys. B 364, 43 (1991); M. Lindner and D. Ross, Nucl. Phys. B 370, 30 (1992); S. P. Martin, Florida Univ. preprint UFIT-HEP-91-24; R. Bonish, Phys. Lett. 268B, 394 (1991).
- [6] M. Suzuki, Mod. Phys. Lett. A 5, 1205 (1990)
- [7] A. Hasenfratz, P. Hasenfratz, K. Jansen, J. Kuti, and Y. Shen, Nucl. Phys. B 365, 79 (1991).
- [8] C. T. Hill, Phys. Rev. D 24, 691 (1981); C. T. Hill, C. Leung, and S. Rao, Nucl. Phys. B 262, 517 (1985).
- [9] S. Chadha and M. E. Peskin, Nucl. Phys. B 185, 61 (1981), and references therein.
- [10] For reviews with various points of view, see T. Appelquist, Yale preprint YCTP-P 23-91, to be published in the proceedings of the Fourth Mexican Summer School of Particles and Fields; M. Peskin in Recent Advances in Field Theory and Statistical Mechanics, edited by R. Stora and J. B. Zuber (Elsevier, Amsterdam, 1982); A. G. Cohen and H. Georgi, Nucl. Phys. B 314, 7 (1989).
- [11] R. Jackiw and K. Johnson, Phys. Rev. D 8, 2386 (1973); For a recent discussion, see T. Appelquist, M. B. Einhorn, T. Takeuchi, and L. C. R. Wijewardhana, Phys. Rev. D 41, 3192 (1990).

[12] E . A . Paschos and V . I. Zakharov, Phys. Lett. 272B , 105 (1991).

[13] R . S. Chivukula, A . G . Cohen, and K . Lane, Nucl. Phys. B 343, 554 (1990).

[14] T . Appelquist and O . Shapira, Phys. Lett. 249B , 83 (1990); T . Appelquist, J. Terning, and L . C . R . Wijewardhana, Phys. Rev. D 44, (1991).

[15] M . Bando, T . Kugo, and K . Suehiro, Prog. Theor. Phys. 85, 1299 (1991).

[16] For reviews, see R . Peccei in CP Violation, edited by C . Jarlskog (World Scientific, Singapore, 1989); J. Kim, Phys. Rep. 150, 1 (1987).

[17] Particle Data Group, private communication.

Figure Captions

Fig. 1: Ladder approximation to the $\pi\pi$ scattering amplitude.