

University of Wisconsin – Madison

MAD/PH/693

FERMILAB-PUB-92/61-T

February 1992

Test of the Diophantine-Hall-Raby Ansatz for Fermion Mass Matrices

V. Barger, M. S. Berger, T. Han and M. Zralek^y

Physics Department, University of Wisconsin, Madison, WI 53706, USA

Fermi National Accelerator Laboratory, P.O. Box 500, Batavia, IL 60510, USA

^yPhysics Department, University of Silesia, Katowice, Poland

ABSTRACT

By evolution of fermion mass matrices of the Fritzsch and the Georgi-Jarlskog forms from the supersymmetric grand unified scale, DHR obtained predictions for the quark masses and mixings. Using Monte Carlo methods we test these predictions against the latest determinations of the mixings, the CP-violating parameter κ and the B_d^0 - B_d^0 mixing parameter x_d . The acceptable solutions closely specify the quark masses and mixings, but lie at the edges of allowed regions at 90% confidence level.

One of the outstanding problems in particle physics is that of explaining the fermion masses and mixings. In the Standard Model (SM) the 6 quark masses, 3 charged lepton masses, the 3 quark mixings and the CP-violating phase of the Cabibbo-Kobayashi-Maskawa (CKM) matrix are introduced as phenomenological parameters. Over the years various models have been proposed to reduce the number of these free parameters [1], of which the best known is the Fritzsch model [2]. Recently Dimopoulos, Hall and Raby (DHR) have proposed an ansatz for fermion mass matrices [3] in the framework of minimal supersymmetric (SUSY) Grand Unified Theories (GUT). The DHR approach is based on the observation that some discrete symmetries present at the grand unification scale are broken in the low-energy theory. Thus some elements of the fermion mass matrices that vanish at the GUT scale are non-zero at the electroweak scale, and their low-energy values are calculable from the renormalization group equations. The fermion masses and mixings at the electroweak scale can thereby be expressed in terms of a smaller number of input parameters at the GUT scale. DHR work in the massless neutrino limit and relate the 13 SM parameters and a SUSY parameter $\tan\beta$ (discussed below) to 8 input parameters, leading to 6 predictions that include an allowed range of $147\{187 \text{ GeV}$ for the top quark mass (m_t). In comparison the Fritzsch approach gives $77 \leq m_t \leq 96 \text{ GeV}$ [1], which is nearly excluded in the SM by the CDF experiment [4] at a 90% confidence level (C.L.).

The DHR quark mass matrices at the scale m_t are

$$M_u = \begin{pmatrix} 0 & C & 0 \\ C & u & B \\ 0 & B & A \end{pmatrix} \begin{matrix} \text{B} \\ \text{C} \\ \text{A} \end{matrix} \begin{matrix} 0 \\ 1 \\ 0 \end{matrix} \frac{v \sin \beta}{\sqrt{2}} \quad M_d = \begin{pmatrix} 0 & F e^i & 0 \\ F e^{-i} & E & d \\ 0 & 0 & D \end{pmatrix} \begin{matrix} \text{B} \\ \text{C} \\ \text{A} \end{matrix} \begin{matrix} 0 \\ 1 \\ 0 \end{matrix} \frac{v \cos \beta}{\sqrt{2}} ; \quad (1)$$

where all the parameters are real, $\tan\beta = v_2/v_1$ in terms of the Higgs doublet vacuum expectation values, and $v = 246 \text{ GeV}$. The charged lepton mass matrix M_e is obtained from the above form of M_d by the substitutions $\beta = 0$; $d = 0$; $E \rightarrow 3E^0$; $D \rightarrow D^0$; $F \rightarrow F^0$. At the SUSY-GUT scale, the parameters u and d vanish and $D = D^0$; $E = E^0$; $F = F^0$, so the input mass matrix M_u is of the Fritzsch form [2] and M_d and M_e are of the Georgi-Jarlskog form [5], giving the GUT scale mass relations $m_b = m_s$; $m_s' = m_c$; $m_d' = 3m_e$

between quarks and leptons. The mass ratio prediction

$$(m_d = m_s)(1 - m_d = m_s)^2 = 9(m_e = m_\mu)(1 - m_e = m_\mu)^2 \quad (2)$$

holds at all scales.

The Wolfenstein parameterization [6] of the CKM matrix determined from the unitary matrices that diagonalize the DHR mass matrices can be expressed in terms of four angles (θ_i) and a complex phase (δ) as follows

$$V_{ud} = (s_1^2 + s_2^2 + 2s_1s_2 \cos \delta)^{\frac{1}{2}} = |V_{ud}| e^{i\phi_{ud}}; \quad (3a)$$

$$V_{us} = s_3 - s_4 = |V_{us}| e^{i\phi_{us}}; \quad (3b)$$

$$V_{ub} = s_2 = |V_{ub}| e^{i\phi_{ub}}; \quad (3c)$$

$$V_{cb} = s_1s_2 \sin \delta = |V_{cb}| e^{i\phi_{cb}}; \quad (3d)$$

with $s_i = \sin \theta_i$; $c_i = \cos \theta_i$ ($i = 1, 2, 3, 4$), where θ_2, θ_3 are the angles that diagonalize the matrix M_u , and θ_1, θ_4 are those for M_d [3]; only three of these angles are independent. These mixing angles are related to the quark masses and other parameters by

$$s_1 = \sqrt{\frac{m_b m_c}{m_s m_t}}; \quad s_2 = \sqrt{\frac{m_b m_c}{m_u m_t}}; \quad s_3 = \sqrt{\frac{m_b m_c}{m_s m_t}}; \quad s_4 = s_3 \sin \delta; \quad (4)$$

The evolution based on the SUSY renormalization group equations (RGE) from the GUT scale to the appropriate fermion mass scales, taking all SUSY particles and the second Higgs doublet degenerate at the scale of m_t [3], gives the following relations,

$$m_t = \frac{m_b m_c}{m_s} \frac{x}{V_{cb}^2} \frac{1}{b c^{1/2}}; \quad m_s = m_d = \frac{1}{3} m_{\bar{s}}^{1/2} x; \quad (5a)$$

$$\sin \delta = \frac{m_t}{V} \frac{3I}{2} \frac{1}{b c^{1/2}} y^{12}; \quad s_3 = \frac{V_{cb} m_b}{m_{\bar{s}}^{1/2}} x; \quad (5b)$$

where

$$x = (g_1)^{1/6} (g_2)^{3/2}; \quad y = x (m_b = m_c)^{1/2} b^{-1}; \quad (6a)$$

$$I = \prod_{i=1,2,3} (g_i)^{c_i = b_i}; \quad I(\delta) = \int_0^\delta d\ln \delta; \quad (6b)$$

The RGE parameters b_i ; c_i are given in Ref. [3]. In these equations the couplings α_1 and α_2 are evaluated at the scale m_t . The mass parameters are defined as m_q ($= m_q$) for quarks heavier than 1 GeV, and the lighter quark masses m_s ; m_d ; m_u are calculated at the scale $\mu = 1$ GeV.

Starting from the well-determined values [7], $\alpha_1(M_Z) = 0.016887$; $\alpha_2(M_Z) = 0.03322$, and evolving at one-loop level to their intersection determines the GUT scale $M_G = 1.1 \cdot 10^{16}$ GeV and the GUT coupling constant $\alpha_G = 1/25.4$. Evolving backwards, the strong coupling constant $\alpha_s(M_Z) = 0.106$ is obtained, consistent with the LEP result $\alpha_s = 0.118 \pm 0.008$ [8]. Also the values $\alpha_1(m_t) = 0.017$ and $\alpha_2(m_t) = 0.033$ are determined, as well as the factors $K_1(m_t) = 9.7$ and $I(m_t) = 110$. We have used a top-quark threshold of 180 GeV in the RGE, consistent with our output determination. In evolution below the electroweak scale we include 3-loop QCD and 1-loop QED effects in the running masses to obtain the evolution factors $b_q = 1.44$; $c_q = 1.80$ and $s_q = 1.95$ where $m_q = m_q(m_q) = m_q(m_t)$ for $q = b, c$ and $m_s = [m_s(1 \text{ GeV}) = m_s(1 \text{ GeV})] \cdot [m_s(m_t) = m_s(m_t)]$. Quark and lepton thresholds were handled by demanding that the couplings and running masses be continuous. The number of active flavors in the β -functions and in the anomalous dimensions was changed as each successive fermion was integrated out of the theory.

Following DHR, we take the following 8 relatively better-known parameters as inputs: m_e ; m_μ ; m_τ ; m_c ; m_b ; $m_u = m_d$; $|V_{cb}|$ and $|V_{cd}|$. We generate random values for all inputs within 90% C.L. (1.64 σ) ranges. The input mass values [9,10] are $m_e = 0.511 \text{ MeV}$; $m_\mu = 105.66 \text{ MeV}$; $m_\tau = 1784.1 \pm 2.7 \text{ MeV}$; $m_c(m_c) = 1.27 \pm 0.05 \text{ GeV}$; $m_b(m_b) = 4.25 \pm 0.1 \text{ GeV}$, where 1 errors are quoted. We also impose the theoretical constraint $0.2 \leq m_u = m_d \leq 0.7$ [11]. We next calculate m_d ; m_s from Eqs. (2) and (5a) (obtaining $m_d = 5.93 \text{ MeV}$, $m_s = 146.5 \text{ MeV}$), s_1 and s_2 from Eq. (4), s_3 from Eq. (5b) for the input of $|V_{cb}|$, s_4 from Eq. (4), and s_5 from $|V_{cd}|$ of Eq. (3a). Using these values we evaluate the magnitudes of all elements of the CKM matrix. We retain only those Monte Carlo events that satisfy the following ranges from the 1992 Review of Particle Properties [9],

$$V_{CKM} = \begin{pmatrix} 0.9747 \pm 0.0007 & 0.218 \pm 0.004 & 0.002 \pm 0.001 \\ 0.218 \pm 0.004 & 0.9735 \pm 0.0007 & 0.032 \pm 0.004 \\ 0.003 \pm 0.001 & 0.030 \pm 0.004 & 0.9985 \pm 0.0005 \end{pmatrix}; \quad (7)$$

as well as the ratio $V_{ub}/V_{cb} = 0.149$ [9]. Figure 1 shows a scatter plot of $\sin \theta_{13}$ versus m_t obtained from our Monte Carlo analysis; one sees that only a narrow wedge of the space is permissible.

From $\sin \theta_{13} = 1$ in Eq. (5), V_{cb} must satisfy the inequality

$$|V_{cb}| > \frac{x}{v} \sqrt{\frac{3I}{2(1-y^{12})}} \frac{m_b m_c}{m_{bc}} \sqrt{\frac{1}{2}} > 0.053; \quad (8)$$

which is just at the edge of the 90% C.L. allowed range. Calculating m_t from Eq. (5a) and $\sin \theta_{13}$ from Eq. (5b) and requiring that V_{cb} be within its allowed range, we find

$$174 < m_t < 183 \text{ GeV}; \quad \sin \theta_{13} > 0.954 \text{ (tan } \theta_{13} > 3.2); \quad (9)$$

This top quark mass determination is consistent with estimates from the electroweak radiative corrections [8] but is much more restrictive. The predicted value of $\tan \theta_{13}$ is large, which may have significant phenomenological implications for Higgs boson searches at colliders [12].

Next we include the constraints from the measured values

$$|J_K| = (2.259 \pm 0.018) \times 10^{-3} [9]; \quad r_d = 0.181 \pm 0.043 [13]; \quad (10)$$

of the CP-violating parameter J_K and the B_d^0 - \bar{B}_d^0 mixing parameter r_d . The theoretical formulas, including QCD corrections, can be found in Eqs. (2.1) and (2.10) of Ref. [14]. In our Monte Carlo analysis we allow variations of the bag-factors and B-decay constant over the following ranges [14]:

$$0.33 \leq B_K \leq 1.5; \quad 0.1 \text{ GeV} \leq \frac{q}{B_B f_B} < 0.2 \text{ GeV}; \quad (11)$$

taking $f_K = 160 \text{ MeV}$ and $M_K = 3.521 \times 10^{-15} \text{ GeV}$. The solutions so obtained closely specify the CKM matrix to be

$$V_{CKM} = \begin{pmatrix} 0 & & \\ & 0 & \\ & & 1 \end{pmatrix} \begin{pmatrix} 0.9749 \{ 0.9759 & 0.2185 \{ 0.2230 & 0.0027 \{ 0.0032 \\ 0.2185 \{ 0.2230 & 0.9735 \{ 0.9745 & 0.0530 \{ 0.0540 \\ 0.0106 \{ 0.0109 & 0.0518 \{ 0.0529 & 0.9985 \{ 0.9986 \end{pmatrix} ; \quad (12)$$

and predict the CP-violating phase to be in the range

$$70^\circ < \delta < 80^\circ ; \quad (13)$$

The inclusion of r_K and r_d almost uniquely determines the values of $|V_{td}|$ and $|V_{ts}|$. Since $|V_{cb}|$ is near its allowed upper limit, $|V_{ub}|$ is pushed to its lower end by the unitarity condition. The output value of the ratio

$$0.051 < |V_{ub}|/|V_{cb}| < 0.059 \quad (14)$$

is at the low end of the allowed range. Improved experimental determinations of $|V_{cb}|$ and $|V_{ub}|/|V_{cb}|$ will test the DHR ansatz. In terms of the Wolfenstein parameters [6], we find

$$\begin{aligned} 0.2185 < \lambda < 0.2230 ; \quad 1.07 < A < 1.13 ; \\ 0.195 < \bar{\lambda} < 0.243 ; \quad 0.105 < \bar{\lambda} < 0.129 ; \quad 0.222 < \sqrt{\lambda^2 + \bar{\lambda}^2} < 0.275 ; \end{aligned} \quad (15)$$

The output values of the mass ratios of the light quarks are

$$0.52 < m_u/m_d < 0.70 ; \quad m_d/m_s = 0.0405 ; \quad 0.021 < m_u/m_s < 0.028 ; \quad (16)$$

giving $3.08 < m_u < 4.15 \text{ MeV}$. These light quark masses and their ratios are consistent with those obtained in Refs. [10,11], but do not agree as well with some other recent studies [15], in which $m_u/m_d < 0.3$ was obtained.

The heavy quark masses are now constrained to the narrow ranges

$$1.19 < m_c < 1.23 ; \quad 4.09 < m_b < 4.20 ; \quad (17)$$

Another interesting result is restrictive ranges for the constants B_K and f_B

$$0.33 < B_K < 0.43 ; \quad 0.14 < \frac{q}{B_B f_B} < 0.17 \text{ GeV} ; \quad (18)$$

on which theoretical uncertainties have been problematic [16].

We conclude with some brief remarks. From Eq. (5a), m_t is inversely proportional to $|V_{cb}|^{1/2}$, and the theoretical uncertainty in this quantity could somewhat enlarge or close the window in m_t (and correspondingly the window in $|V_{cb}|$). The DHR analysis assumes dominance of the top quark Yukawa couplings in the RGE evolution. Since the output $\tan\beta$ may be large, the effects of fully including $|V_{cb}|$ and $|V_{ub}|$ in the evolution may not be negligible; this question deserves further study. Also two-loop renormalization group equations between M_Z and M_{GUT} should eventually be incorporated. We have studied the charged Higgs boson effects on K and r_d . With M_H degenerate with m_t , as assumed in the model, we found no significant changes in our results. This is due to the fact that Higgs effects are smaller at large $\tan\beta$ for the K and B systems. In summary, the DHR ansatz for fermion mass matrices is consistent with all current experimental constraints at 90% C.L.. It leads to almost unique values for m_t and quark mixings which make it an interesting target for future experiments.

ACKNOWLEDGMENTS

We thank M. Barnett and K. Hikasa for advance information from the Particle Data Group, G. Anderson and L. Hall for communications, and A. Manohar for comments on quark mass ratios. This work was supported in part by the U.S. Department of Energy under contract No. DE-AC02-76ER00881 and in part by the University of Wisconsin Research Committee with funds granted by the Wisconsin Alumni Research Foundation. T. Han was supported by an SSC Fellowship from the Texas National Research Laboratory Commission under Award No. FCFY 9116.

REFERENCES

- [1] See e.g., F. J. Gilman and Y. Nir, *Ann. Rev. Nucl. Part. Sci.* 40, 213 (1990); P. Kaus and S. Meshkov, *Phys. Rev. D* 42, 1863 (1990); X. G. He and W. S. Hou, *Phys. Rev. D* 41, 1517 (1990); C. H. Albright, *Phys. Lett. B* 246, 451 (1990).
- [2] H. Fritzsch, *Phys. Lett.* 70B, 436 (1977); 73B, 317 (1978).
- [3] S. Dimopoulos, L. J. Hall and S. Raby, OSU preprints DOE-ER-01545-566 (1991); DOE-ER-01545-567 (1991).
- [4] F. Abe et al., Central Detector Facility (CDF) Collaboration, *Phys. Rev. Lett.* 68, 447 (1992).
- [5] H. Georgi and C. Jarlskog, *Phys. Lett.* 86B, 297 (1979).
- [6] L. Wolfenstein, *Phys. Rev. Lett.* 51, 1945 (1983).
- [7] U. Amaldi, W. de Boer, and H. Furstenau, *Phys. Lett.* 260B, 447 (1991).
- [8] J. R. Carter, rapporteur talk at the 1991 Joint International Lepton-Photon Symposium & Europhysics Conference on High Energy Physics, Geneva, LP-HEP 91; P. Langacker, Univ. of Pennsylvania preprint UPR-0492T (1992), and references therein.
- [9] Particle Data Group, to appear in *Phys. Lett. B* (1992).
- [10] J. Gasser and H. Leutwyler, *Phys. Rep.* 87, 77 (1982).
- [11] D. Kaplan and A. Manohar, *Phys. Rev. Lett.* 56, 2004 (1986).
- [12] See e.g. V. Barger, M. S. Berger, A. L. Stange, and R. J. N. Phillips, Univ. of Wisconsin-Madison preprint MAD/PH/680 (1991), *Phys. Rev. D* (in press).
- [13] H. Schroder, DESY preprint DESY 91-139 (1991).
- [14] V. Barger, J. L. Hewett, and R. J. N. Phillips, *Phys. Rev. D* 41, 3421 (1990).
- [15] J. Donoghue and D. Wyler, *UM HEP-350* (1991); K. Choi, *UCSD/PTH 92/06* (1992).

[16] See e.g. C. Q. Geng and P. Turcotte, Univ. of Montreal Preprint, UdeM-LPN-TH-78 (1992), and references therein.

FIGURES

FIG .1. Scatterplot of $\sin \delta$ versus m_t from our Monte Carlo analysis of the DHR model, imposing the constraints of input masses and present values of CKM matrix elements.