

**CP Violation in top pair production
at an e^+e^- collider**

Darwin Chang^(1,2), Wai-Yee Keung^(3,4), Ivan Phillips⁽¹⁾

⁽¹⁾Department of Physics and Astronomy,

Northwestern University, Evanston, IL 60208, USA.

⁽²⁾Institute of Physics, Academia Sinica, Taipei, Taiwan, R.O.C.

⁽³⁾Physics Department, University of Illinois at Chicago, IL 60680, USA

⁽⁴⁾Theory Division, CERN, CH-1211, Geneva 23, Switzerland

Abstract

We investigate possible CP violating effects in e^+e^- annihilation into top quark pairs. One of the interesting observable effects is the difference in production rates between the two CP conjugate polarized states $t_L\bar{t}_L$ and $t_R\bar{t}_R$. The result is an asymmetry in the energy spectra of the lepton and the anti-lepton from the heavy quark decays. Another CP-odd observable is the up-down asymmetry of the leptons with respect to the reaction plane. These two asymmetries measure complementarily the absorptive and dispersive form factors of the electric dipole moment. Finally, as an illustration, we calculate the size of the CP violating form factors in a model where the CP nonconservation originates from the Yukawa couplings of a neutral Higgs boson.

I. Introduction.

Since the top quark is widely believed to be within the reach of the present collider machines, it is not unreasonable for theorists to imagine what we can learn from the top quark. The best place to study the top quark in detail is in an e^+e^- collider. One of the facts one would like to learn from the discovery of the top quark is the origin of the still mysterious CP violation. In this paper we investigate a way CP violation can manifest itself in the top pair production of an e^+e^- collider.

The top quark, due to its short lifetime, is believed to decay before it hadronizes[1]. Therefore the information about its polarization may be preserved in its decay products. If that is the case, then one can investigate the source of CP nonconservation by measuring the CP violating observable involving a polarized top pair in the final state. This idea of detecting the rate asymmetry between different polarized states was recently proposed by Schmidt and Peskin[2, 3]. For $t\bar{t}$ production through the virtual photonic or Z intermediate states, to the lowest order in the final state quark mass, the polarizations of the quarks are either $t_L\bar{t}_R$ or $t_R\bar{t}_L$. (Note that we have adopted the notation that \bar{t}_L is the antiparticle of t_R and should be left handed.) These two modes are CP self-conjugate. However since the top quark is heavy, there will also be large percentage of $t_L\bar{t}_L$ and $t_R\bar{t}_R$ modes which are CP conjugates of each other. Therefore one can consider a CP asymmetry in the event rate , $N(t_L\bar{t}_L) - N(t_R\bar{t}_R)$.

Schmidt and Peskin have shown how to detect the asymmetry $N(t_L\bar{t}_L) - N(t_R\bar{t}_R)$ through the energy spectra of prompt leptons[2, 3]. One assumes that the t quark decays semileptonically through the usual $V - A$ weak interaction. Knowing that the hadronization time is much longer than the decay time[1], one can analyze polarization dependence of its decay at the quark level. The top quark first decays into a b quark and a W^+ boson, which subsequently becomes $\ell^+\nu$. For heavy top quark, the W^+ boson produced in top decay is predominantly longitudinal. Due to the $V - A$ interaction, the b quark is preferentially produced with left-handed helicity. So the longitudinal W^+ boson is preferentially produced along the direction of the top quark polarization. Therefore the anti-lepton ℓ^+ produced in the W^+ decay is also preferentially in that direction. In the rest frame of the t , the angular distribution[4] of the produced ℓ^+ has the form $1 + \cos \psi$, with ψ as the angle between ℓ^+ and

the helicity axis of the t . Above the $t\bar{t}$ threshold, the top quark is produced with nonzero momentum. As a result of the Lorentz boost, the anti-lepton ℓ^+ produced in the decay of the right handed top quark t_R has a higher energy than that produced in the decay of the left handed top quark t_L . Similarly, the lepton l produced in the decay of \bar{t}_L has a higher energy than that produced in the decay of \bar{t}_R . Consequently, in the decay of the pair $t_L\bar{t}_L$ the lepton from \bar{t}_L has a higher energy than the anti-lepton from t_L ; while in the decay of $t_R\bar{t}_R$ the anti-lepton has a higher energy. Therefore one can observe $N(t_L\bar{t}_L) - N(t_R\bar{t}_R)$ by measuring the energy asymmetry in the leptons. It turns out that this energy asymmetry is sensitive only to the absorptive parts of CP violating form factors.

There is another equally interesting CP-odd effect in the azimuthal angular distribution, namely, the rate difference between the events with ℓ^\pm above the reaction plane and the events with ℓ^\pm below the reaction plane. Such an observable, like the previous one, is a direct measurement of CP violation and thus has no background from the CP conserving interactions. Unlike the previous case, this up-down asymmetry will probe the CP violating dispersive form factors. Though there have been many studies of CP violating observables in the literature[5, 6], we feel that the above two observables are simple and intuitive in nature and are easily implemented in future experiments.

Finally, we will use a generic neutral Higgs model with CP violation in the Yukawa couplings to illustrate how these observables can arise. Such mechanisms of CP violation are contained in many extensions of Standard Model including the simple two doublet model[7]. Throughout this paper, we focus our attention on CP non-conservation in the production mechanism only. There will be additional contributions if the usual $V - A$ decay amplitude is also modified by CP violating interactions[8].

II. CP violating form factors and amplitudes.

We start by writing down the general form factors of the t quark. The vertex amplitude $ie\Gamma^j$ for the virtual γ^* or Z^* turning into $t(p)$ and $\bar{t}(p')$ can be parametrized in the following expression:

$$\Gamma_\mu^j = c_v^j \gamma_\mu + c_a^j \gamma_\mu \gamma_5 + c_d^j i \gamma_5 \frac{p_\mu - p'_\mu}{2m_t} + \cdots, \quad j = \gamma, Z. \quad (1)$$

We use the tree-level values for c_v and c_a . They are

$$\begin{aligned} c_v^\gamma &= \frac{2}{3}, & c_a^\gamma &= 0, \\ c_v^Z &= (\frac{1}{4} - \frac{2}{3}x_W)/\sqrt{x_W(1-x_W)}, \\ c_a^Z &= -\frac{1}{4}/\sqrt{x_W(1-x_W)}. \end{aligned} \quad (2)$$

Here $x_W \simeq 0.23$ is the electroweak mixing angle in the Standard Model. The c_d terms are the electric dipole form factors. The spinor structure can be rewritten into another form using $i\gamma_5(p-p')_\mu = \sigma_{\mu\nu}(p+p')^\nu\gamma_5$. Other irrelevant terms, like the magnetic moments, are not listed in Eq.(1). It can also be easily shown that, in the limit $m_e = 0$, c_d is the only relevant form factor for the CP violating quantities we are interested in.

The helicity amplitudes $e^2 M(h_e, h_{\bar{e}}, h_t, h_{\bar{t}})$ for the process $e^-e^+ \rightarrow t\bar{t}$ at the scattering angle θ have been given in the literature[6]. For the initial configuration of $e_L\bar{e}_R$, we have

$$\begin{aligned} M(-+-+) &= [c_v^\gamma + r_L c_v^Z - \beta r_L c_a^Z](1 + \cos \theta) \\ M(-++-) &= [c_v^\gamma + r_L c_v^Z + \beta r_L c_a^Z](1 - \cos \theta) \\ M(-+--) &= [2t(c_v^\gamma + r_L c_v^Z) - \frac{i}{2}(c_a^\gamma + r_L c_a^Z)\beta/t] \sin \theta \\ M(-+++) &= [2t(c_v^\gamma + r_L c_v^Z) + \frac{i}{2}(c_a^\gamma + r_L c_a^Z)\beta/t] \sin \theta. \end{aligned} \quad (3)$$

Here we have used the convention [9] that CP invariance, when $c_d^{\gamma,Z}$ are turned off, is signified by the relation

$$M(\sigma, \bar{\sigma}; \lambda, \bar{\lambda}) = M(-\bar{\sigma}, -\sigma; -\bar{\lambda}, -\lambda). \quad (4)$$

The dimensionless variables are defined by, $t = m_t/\sqrt{s}$, $z = m_Z/\sqrt{s}$, $\beta^2 = 1 - 4t^2$. The Z -propagator and its coupling to the left-handed electron gives $-er_L/s$ with

$$r_L = (\frac{1}{2} - x_W)/[(1 - z^2)\sqrt{x_W(1 - x_W)}] \quad (5)$$

The cross section is

$$d\sigma(e_L\bar{e}_R \rightarrow t_{h_t}\bar{t}_{h_{\bar{t}}})/d(\cos \theta) = \frac{3}{2}\pi\alpha^2\beta|M(-+, h_t, h_{\bar{t}})|^2/s. \quad (6)$$

A color factor 3 has been included explicitly in the above formula. Similarly, we obtain formulas for the initial configuration $e_R\bar{e}_L$ with r_L replaced by r_R ,

$$r_R = -x_W/[(1 - z^2)\sqrt{x_W(1 - x_W)}], \quad (7)$$

and $\cos \theta$ by $-\cos \theta$, and $\sin \theta$ by $-\sin \theta$ in Eq.(3). In case of an unpolarized e^-e^+ machine, we must sum up initial configurations $e_L\bar{e}_R$ and $e_R\bar{e}_L$, and include the spin average factor $\frac{1}{4}$.

It is also interesting to note that, if the absorptive part of a scattering amplitude can be ignored (which is certainly true at the tree level), then the unitarity of the S matrix implies that the scattering matrix is hermitian. This hermiticity allows one to write down the constraint due to the CPT invariance as

$$M(\sigma, \bar{\sigma}; \lambda, \bar{\lambda}) = M^*(-\bar{\sigma}, -\sigma; -\bar{\lambda}, -\lambda). \quad (8)$$

We shall refered to this special case of CPT invariance as $\text{CPT}\hat{T}$ invariance. The $\text{CPT}\hat{T}$ invariance of course can be violated by the absorptive part of the scattering amplitude.

III. Leptonic Energy asymmetry

It is straightforward to see that the absorptive part $\text{Im } c_d^\gamma$ or $\text{Im } c_d^Z$ is required to produce the difference between configurations $t_L\bar{t}_L$ and $t_R\bar{t}_R$. This is expected because we need the absorptive part via the final state interactions to overcome the $\text{CPT}\hat{T}$ constraint. The asymmetry integrated over the angular distribution is

$$\delta \equiv \frac{[N(t_L\bar{t}_L) - N(t_R\bar{t}_R)]}{N(t\bar{t}; \text{all})} = \frac{\sum_{h=L,R} 2\beta(c_v^\gamma + r_h c_v^Z)(\text{Im } c_d^\gamma + r_h \text{Im } c_d^Z)}{\sum_{h=L,R} (3 - \beta^2)(c_v^\gamma + r_h c_v^Z)^2 + 2\beta^2 r_h^2 c_a^{Z^2}}. \quad (9)$$

Note that there is no CP violating contribution due to c_a^Z coupling in the numerator when the electron mass m_e is ignored.

We can make use of this asymmetry parameter δ to illustrate the the difference in the energy distributions of ℓ^+ or ℓ^- from the t or \bar{t} decays. The energy $E_0(\ell^+)$ distribution of a static t quark decay $t \rightarrow \ell^+ \nu b$ is very simple[4] in the narrow width Γ_W approximation when m_b is negligible. It can be represented as

$$f(x_0) = \begin{cases} x_0(1 - x_0)/D & \text{if } m_W^2/m_t^2 \leq x_0 \leq 1, \\ 0 & \text{otherwise.} \end{cases} \quad (10)$$

for on-shell W in the decay. Here we denote the scaling variable $x_0 = 2E_0(\ell^+)/m_t$ and the normalization factor $D = \frac{1}{6} - \frac{1}{2}(m_W/m_t)^4 + \frac{1}{3}(m_W/m_t)^6$. When the t quark is not static, but moves at a speed β with helicity L or R , the distribution expression becomes a convolution,

$$f_{R,L}(x, \beta) = \int_{x/(1+\beta)}^{x/(1-\beta)} f(x_0) \frac{\beta x_0 \pm (x - x_0)}{2x_0^2 \beta^2} dx_0. \quad (11)$$

Here $x = 2E(\ell^+)/E_t$. The kernel above is related to the $(1 \pm \cos \psi)$ distribution mentioned in the introduction. Similar distributions for the \bar{t} decay are related by CP conjugation at the tree-level. Using the polarization asymmetry formula in Eq.(9), we can derive an expression for the difference in the energy distributions of ℓ^- and ℓ^+ :

$$\frac{1}{N} \left[\frac{dN}{dx(\ell^+)} - \frac{dN}{dx(\ell^-)} \right] = \delta[f_L(x, \beta) - f_R(x, \beta)] . \quad (12)$$

Here distributions are compared at the same energy for the lepton and the anti-lepton, $x(\ell^-) = x(\ell^+) = x = 4E(\ell^\pm)/\sqrt{s}$. The count N includes events with prompt leptons or anti-leptons from the top pair production. It is useful to compare Eq.(12) with that of the overall energy distribution,

$$\begin{aligned} \frac{1}{N} \left[\frac{dN}{dx(\ell^+)} + \frac{dN}{dx(\ell^-)} \right] &= \frac{\sum_{h=L,R} 4\beta r_h c_a^Z (c_v^\gamma + r_h c_v^Z) [f_R(x, \beta) - f_L(x, \beta)]}{\sum_{h=L,R} (3 - \beta^2) (c_v^\gamma + r_h c_v^Z)^2 + 2\beta^2 r_h^2 c_a^{Z^2}} \\ &\quad + f_L(x, \beta) + f_R(x, \beta) . \end{aligned} \quad (13)$$

Here we only keep the dominant tree-level contribution. The first term of Eq.(13) is due to the two helicity modes that are CP self-conjugate.

Fig. 1




Fig. 1 shows the overall prompt lepton energy distribution of Eq.(13), and the ratio of the expressions in Eq.(12) and Eq.(13) per unit $\text{Im } c_d^\gamma$ or $\text{Im } c_d^Z$.

Note that our sample includes events with one or at least one prompt ℓ^\pm . However, if we are willing to use a smaller sample of prompt dilepton events (from the simultaneous semileptonic decays of both t and \bar{t}), a recent paper Ref. [10] has shown how one can use the kinematics of such events to analyze the helicities of t and \bar{t} . Therefore such knowledge can be used to enhance the detectability of the asymmetry by cutting away the CP self-conjugate modes in the denominator of δ in Eq.(9) [10]. However, the reduction in the size of event samples may be too high a price to pay.

IV. CP-odd up-down asymmetry.

It is known that explicit CP violation requires the CP nonconserving vertex as well as additional complex amplitudes. In the above case, this complex structure comes from the absorptive part due to the final state interactions. However, the complex structure can also come from other sources. One of these is the azimuthal phase $\exp(iL_z\phi)$ in the decay process. This will produce a CP-odd up-down asymmetry even with only the dispersive part of CP violating vertex, $\text{Re } c_d$.

The angular distribution of ℓ^+ from the t decay is specified by the the spin density matrix $\rho_{\lambda,\lambda'}$ of the top quark.

$$\rho(\theta)_{\lambda,\lambda'} = \mathcal{N}(\theta)^{-1} \sum_{h_e, h_{\bar{e}}, h_{\bar{t}}} M(h_e, h_{\bar{e}}, \lambda, h_{\bar{t}}) M^*(h_e, h_{\bar{e}}, \lambda', h_{\bar{t}}) . \quad (14)$$

Here \mathcal{N} is the normalization such that $\text{Tr}\rho=1$. ρ is hermitian by definition.

$$dN(\ell^+) = \left[(1 + \cos\psi)\rho_{++} + (1 - \cos\psi)\rho_{--} + 2\text{Re}(\rho_{+-}e^{i\varphi})\sin\psi \right] d\varphi d\cos\psi / (4\pi) . \quad (15)$$

The polar angles ψ and the azimuthal angle φ of ℓ^+ are defined in the t rest frame F_t , which is constructed by boosting the $t\bar{t}$ center of mass frame F_0 along the motion of the top quark. In F_0 , the z -axis is along the top momentum \mathbf{p}_t , the y axis is along $\vec{p}_{e^-} \times \vec{p}_t$, and the x -axis is given by the right-handed rule. The production plane is the $x - z$ plane.

Similarly, the angular distribution of ℓ^- from the \bar{t} decay is specified by the spin density matrix, $\bar{\rho}_{\bar{\lambda},\bar{\lambda}'}$, of the anti-top quark.

$$\bar{\rho}(\theta)_{\bar{\lambda},\bar{\lambda}'} = \mathcal{N}(\theta)^{-1} \sum_{h_e, h_{\bar{e}}, h_t} M(h_e, h_{\bar{e}}, h_t, \bar{\lambda}) M^*(h_e, h_{\bar{e}}, h_t, \bar{\lambda}') . \quad (16)$$

$$dN(\ell^-) = \left[(1 + \cos\bar{\psi})\bar{\rho}_{++} + (1 - \cos\bar{\psi})\bar{\rho}_{--} - 2\text{Re}(\bar{\rho}_{+-}e^{-i\bar{\varphi}})\sin\bar{\psi} \right] d\bar{\varphi} d\cos\bar{\psi} / (4\pi) . \quad (17)$$

The polar angles $\bar{\psi}$ and the azimuthal angle $\bar{\varphi}$ of ℓ^- are defined in the \bar{t} rest frame $F_{\bar{t}}$, which is similarly constructed by boosting the $t\bar{t}$ center of mass frame F_0 along the motion of the anti-top quark. It is important to keep in mind that the three coordinate systems F_0 , F_t and $F_{\bar{t}}$, have parallel directions of coordinate axes.

In the phase convention[9] such that Eqs.(4,8) are satisfied when CP is conserved, the following identities, first noticed by Gounaris et al.[11],

$$\rho(\theta)_{\lambda,\lambda'} = \bar{\rho}(\theta)_{-\lambda,-\lambda'} , \quad (18)$$

between the density matrix elements can be derived. Under CP conjugation, as we exchange ℓ^- and ℓ^+ , by definition one should make the angular substitutions $\bar{\psi} \rightarrow \pi - \psi$, $\bar{\varphi} \rightarrow \pi + \varphi$. In that case, one observes that the distribution in Eq.(17) is transformed into Eq.(15) provided Eq.(18) is satisfied. On the other hand, the $CPT^{\hat{}}$ invariance, when the absorptive amplitudes are ignored, implies

$$\rho(\theta)_{\lambda,\lambda'} = \bar{\rho}(\theta)_{-\lambda,-\lambda'}^* , \quad (19)$$

This is very similar to the CP and $CPT^{\hat{}}$ transformations in the case of $e^-e^+ \rightarrow W^-W^+$ as analyzed before[11, 12].

When the effect of the CP violating form factors are included in the analysis, one can form the following CP or $CPT^{\hat{}}$ odd combinations:

$$R(\pm)(\theta)_{\lambda,\lambda'} = \text{Re } \rho(\theta)_{\lambda,\lambda'} \pm \text{Re } \bar{\rho}(\theta)_{-\lambda,-\lambda'} ,$$

and

$$I(\pm)(\theta)_{\lambda,\lambda'} = \text{Im } \rho(\theta)_{\lambda,\lambda'} \pm \text{Im } \bar{\rho}(\theta)_{-\lambda,-\lambda'} .$$

Among them, $R(-)$ and $I(+)$ are $CPT^{\hat{}}$ odd; $R(-)$ and $I(-)$ are CP odd and all the others CP and $CPT^{\hat{}}$ even. The observation of $R(-)$ requires final state interactions due to $CPT^{\hat{}}$. Therefore it does not need to involve the the complex phase of the azimuthal dependence in Eqs.(15, 17). It can be decoded by analyzing the polar angular dependence of ψ or $\bar{\psi}$ in these equations. In the collider C. M. frame these dependence can be translated into the energy dependence of the corresponding lepton in the final state, which has already been studied in the previous section.

Here we will focus on $I(-)$ which does not require final state interactions. Since ρ is hermitian, the only nonzero component of $I(-)$ is $I(-)_{+-}$. It can be related to c_d as

$$\mathcal{N}(\theta)\text{Im} [\rho(\theta)_{+-} - \bar{\rho}(\theta)_{-+}] = 2 \sin \theta \sum_{h=R,L} (-h)(c_v^\gamma + r_h c_v^Z \pm \beta r_h c_a^Z \cos \theta) \text{Re} (c_d^\gamma + r_h c_d^Z) \beta / t, \quad (20)$$

where the contributions $h = R, L$ pick up the signs $+, -$ respectively. One can in principle make detailed angular analysis of the difference between Eq.(15) and its CP conjugate in Eq.(17) similar to what was done for the case of $e^- e^+ \rightarrow W^- W^+$ by Gounaris et al.[11] However, in an effort to find simpler observables which may be more intuitive and may be easier to detect, we shall consider the following partially integrated observable. Let $dN(\ell^+, \text{up})$ count events with ℓ^+ above the xz plane, *i.e.* $p_y(\ell^+) > 0$. Then, with other obvious notations, we define the following up-down asymmetry

$$\mathcal{A}^{u.d.}(\theta) = \frac{[dN(\ell^+, \text{up}) + dN(\ell^-, \text{up})] - [dN(\ell^+, \text{down}) + dN(\ell^-, \text{down})]}{[dN(\ell^+, \text{up}) + dN(\ell^-, \text{up})] + [dN(\ell^+, \text{down}) + dN(\ell^-, \text{down})]}. \quad (21)$$

It is evaluated for each scattering angle θ . The branching fraction of the t semileptonic decay cancels in the ratio. Integrating on ψ, φ or $\bar{\psi}, \bar{\varphi}$ over up or down hemispheres, we obtain $\mathcal{A}^{u.d.}(\theta)$ in a very simple form from Eqs.(15,17),

$$\mathcal{A}^{u.d.}(\theta) = \frac{1}{2}(\text{Im} \bar{\rho}(\theta)_{-+} - \text{Im} \rho(\theta)_{+-}) \quad (22)$$

As we sum up contributions from ℓ^\pm in each square bracket of Eq.(21), the asymmetry is insensitive to the sign of charge, it is obvious that a non-vanishing value of $\mathcal{A}^{u.d.}(\theta)$ is a genuine signal of CP violation. Although the angular distributions of the leptons derived from Eq.(3) will have corrections from the strong interaction, the corrections *cannot* fake the CP asymmetry as the effects due to the strong interaction cancel away in the differences.

To enhance statistics, it is useful to measure the integrated up-down asymmetry,


$$\mathcal{A}^{u.d.} = \frac{1}{\sigma} \int \mathcal{A}^{u.d.}(\theta) \frac{d\sigma}{d(\cos \theta)} d(\cos \theta). \quad (23)$$

From Eqs.(20-23), we obtain,

$$\mathcal{A}^{u.d.} = -\frac{3\pi}{16} \frac{\sum_{h=L,R} (-h)(c_v^\gamma + r_h c_v^Z)(\text{Re} c_d^\gamma + r_h \text{Re} c_d^Z) \beta / t}{\sum_{h=L,R} (3 - \beta^2)(c_v^\gamma + r_h c_v^Z)^2 + 2\beta^2 r_h^2 c_a^{Z^2}}. \quad (24)$$

Note that information on c_a^Z in Eq.(20) is lost when integrating the whole range of $\cos\theta$. However, one can easily find other convolution in Eq.(23) to recover the CP information due to c_a^Z .

Fig. 2



In Fig. 2, we plot the integrated up-down asymmetry $\mathcal{A}^{u.d.}$ per unit $\text{Re } c_d^\gamma$ or $\text{Re } c_d^Z$ for various parameters. It seems that $\mathcal{A}^{u.d.}$ per unit $\text{Re } c_d$ increases with \sqrt{s} , however, $\text{Re } c_d$ is usually energy dependent, characterized by the underlying scale of new physics. Above that scale, $\text{Re } c_d$ diminishes very fast. Therefore, the optimal choice of \sqrt{s} is about at the scale of new physics. Note that even if $\text{Re } c_d$ is relatively constant and the angular asymmetry is larger at higher energy, the event rate will become smaller because of the nature of the s -channel production.

To measure the up-down asymmetry, we need a good determination of the reaction plane. This is possible for those events in which one of the top quarks decays hadronically into jets. One may be concerned about the imperfect angular resolution of hadronic jets. We argue that even if the angles of jets are ambiguous at the level of 10 degrees or so, as long as some sort of orientation of the reaction plane can be defined, the asymmetry will not be smeared away by more than one order of magnitude. For example, one can simply discard events

that the definition of up or down is made ambiguous by this smearing. The details of such a smearing effect will strongly depend on future detectors.

V. Higgs Model.

Among various mechanisms for the CP violation, the one that may manifest itself most easily is the neutral Higgs mediated CP violation. Since the neutral Higgs couplings are typically proportional to the quark mass, the large mass of the top quark naturally gives large couplings to the neutral Higgs bosons. CP non-conservation occurs in the complex Yukawa coupling,

$$\mathcal{L}_{CPX} = -(m_t/v)\bar{t}(AP_L + A^*P_R)tH + (m_Z^2/v)BHZ^\nu Z_\nu. \quad (25)$$

Here $v = (\sqrt{2}G_F)^{-\frac{1}{2}} \simeq 246$ GeV. The complex coefficient A is a combination of model-dependent mixing angles. Simultaneous presence of both the real part $A_R = \text{Re } A$ and the imaginary part $A_I = \text{Im } A$ guarantees CP asymmetry. For example, in the low energy regime, it can give rise to the electric dipole moment of elementary particles [13, 14]. Here we will derive the CP violating form factors at high energy. They are induced at the one-loop level as shown in the Fig. 3. First, we calculate the absorptive parts according to the Cutkosky rules. The leading contribution to $\text{Im } c_d^\gamma$ comes from the rescattering of the top quark pair through the Higgs-boson exchange. Both CP violation and the final state effect are produced by the same one-loop graphs.

$$\text{Im } c_d^\gamma = c_v^\gamma \left(\frac{m_t}{v} \right)^2 \frac{A_R A_I t^2}{2\pi\beta} \left(1 - \frac{h^2}{\beta^2} \log\left(1 + \frac{\beta^2}{h^2}\right) \right). \quad (26)$$

The dimensionless variables are defined by, $t = m_t/\sqrt{s}$, $z = m_Z/\sqrt{s}$, $\beta^2 = 1 - 4t^2$ as before, as well as $h = m_H/\sqrt{s}$. For $\text{Im } c_d^Z$, there is a similar contribution. In addition, there could be a contribution due to the ZH intermediate state, Fig 1b, provided the kinematics is allowed.

$$\text{Im } c_d^Z = \frac{c_v^Z}{c_v^\gamma} \text{Im } c_d^\gamma - \frac{\alpha A_I B c_v^Z t^2}{2(1 - x_W) x_W \beta^2} [\beta_Z + (2t^2 + 2t^2 h^2 - 2t^2 z^2 - h^2) L]. \quad (27)$$

Here $\beta_Z^2 = 1 + h^4 + z^4 - 2z^2 - 2h^2 - 2h^2 z^2$, and the logarithmic factor

$$L = \frac{1}{\beta} \log \frac{1 - z^2 - h^2 - \beta\beta_Z}{1 - z^2 - h^2 + \beta\beta_Z}. \quad (28)$$

Our expression in Eq.(27) agrees with that in Ref.[15]. Note that, at the threshold ($\beta = 0$), $\text{Im } c_d^\gamma$ vanishes but $\text{Im } c_d^Z$ has a value

$$\text{Im } c_d^Z = -\frac{\alpha A_I B c_v^Z \beta_Z}{8(1-x_W)x_W} \left(\frac{1-z^2+h^2}{1-z^2-h^2} - \frac{\beta_Z^2}{3(1-z^2-h^2)^2} \right).$$

In case $m_Z + m_H < \sqrt{s} < 2m_t$, we still have the absorptive part $\text{Im } c_d^Z$, which is simply given by replacing the logarithm factor L in Eq. (27) by its continuation,

$$L = -\frac{2}{\sqrt{-\beta^2}} \arctan \frac{\beta_Z \sqrt{-\beta^2}}{1-z^2-h^2}. \quad (29)$$

The dispersive parts are obtained by the dispersion relation,

$$\text{Re } c_d^j(s) = \frac{1}{\pi} P \int_{s_0}^{\infty} \frac{\text{Im } c_d^j(s')}{s' - s} ds'. \quad (30)$$

The symbol P denotes the principal value of the singular integral. Note that the absorptive part is evaluated at s' above the threshold s_0 , which is $4m_t^2$ or $(m_Z + m_H)^2$ for $t\bar{t}$ or ZH intermediate states respectively. It is understood that the dimensionless variables, β , β_Z , t^2 , h^2 and z^2 , are defined with respect to s' .

Fig. 4





Fig. 4 shows typical sizes of c_d^j for various cases. It is of order of 10^{-2} to 10^{-3} in general. We must keep in mind that we only show the Higgs boson contribution in the perturbative

regime. In scenarios that the Higgs bosons interact strongly, the CP violating effect in the colliders could be much larger.

VI. Conclusion.

We have shown that the energy asymmetry or the up-down asymmetry of the prompt lepton events are sensitive to the CP violation in the top pair production in e^+e^- annihilation. These two asymmetries are complementary to each other for measuring the absorptive and the dispersive parts of the CP violating form factors $c_d^{\gamma,Z}$.

Fig. 5



The cross section for $e^-e^+ \rightarrow t\bar{t}$ at $\sqrt{s} = 400$ GeV, for $m_t = 170$ GeV, is ~ 1400 fb, among which about a quarter is due to the channels $t_L\bar{t}_L$ and $t_R\bar{t}_R$ (see Fig. 5). At an optimal luminosity of $10^{33} \text{ cm}^{-2}\cdot\text{s}^{-1}$, one year run of the futuristic NLC[16] will produce 20,000 prompt semi-leptonic events $b\bar{b}\ell^\pm X$. Note that we do not require both t and \bar{t} to decay semileptonically. Therefore, only one power of the branching fraction $2/9$ is involved. Such sample is good enough to analyze the CP violating form factor c_d at a level of ten percent. Although predictions from Higgs models is still smaller than this level of sensitivity, the measurement could provide directly the important, model-independent information about the form factors of the top quark interactions. For a heavier top quark, \sqrt{s} has to be raised

above the corresponding threshold, but not so far away as to incur the $1/s$ suppression in event rates.

D.C. wishes to thank the Theory Group at the Institute of Physics at Academia Sinica in Taipei, Taiwan and Department of Physics and Astronomy at University of Hawaii at Manoa for hospitality while this work was in progress. This work is supported by grants from Department of Energy and from National Science Council of Republic of China.

References


- [1] I. Bigi and H. Krasemann, Z. Phys. **C7**, 127 (1981); J. Kühn, Acta Phys. Austr. Suppl. **XXIV**, 203 (1982); I. Bigi, Y. Dokshitzer, V. Khoze, J. Kühn, and P. Zerwas, Phys. Lett. **181B**, 157 (1986).
- [2] C.R. Schmidt and M. E. Peskin, Phys. Rev. Lett. **69**, 410 (1992). C.R. Schmidt, Phys. Lett. **B293**, 111 (1992).
- [3] D. Chang and W.-Y. Keung, Phys. Lett. **B305**, 261 (1993) or hep-ph/9301265.
- [4] M. Jeżabek and J.H. Kühn, Nucl. Phys. **B320**, 20, (1989).
- [5] Additional CP-odd observables in collider physics have been studied by J.F. Donoghue and G. Valencia, Phys. Rev. Lett. **58**, 451 (1987); W. Bernreuther and O. Nachtmann, Phys. Rev. Lett. **63**, 2787 (1989); G. Valencia and A. Soni, Phys. Lett. **B263**, 517 (1991); A. Bilal, E. Massó, and A. De Rújula, Nucl. Phys. **355**, 549 (1991); D. Atwood and A. Soni, Phys. Rev. **D45**, 2405 (1991).
- [6] G. Kane, G.A. Ladinsky, and C.-P. Yuan, Phys. Rev. **D45**, 124 (1991).
- [7] A. Mendez and P. Pomarol, Phys. Lett. **B272**, 313 (1991); X. G. He, J. P. Ma and B. H. J. McKellar, Melbourne Preprint UM-P-92/75.
- [8] B. Grzadkowski and J.F. Gunion, Phys. Lett. **B287**, 237–246 (1992).
- [9] Note that we use a different phase convention from Ref.[6]. One can see sign flips between our Eq.(3) and Eq.(6.6) in Ref.[6].
- [10] G.A. Ladinsky, and C.-P. Yuan, Michigan State Preprint, MSUTH 92/07 (1992).

- [11] G. Gounaris, D. Schildknecht, F.M. Renard, Phys. Lett. **B263**, 291 (1991).
- [12] K. Hagiwara, R.D. Peccei, D. Zeppenfeld, and K. Hikasa, Nucl. Phys. **282** 253 (1987).
- [13] S. Weinberg, Phys. Rev. Lett. **63**, 2333 (1989); Phys. Rev. **D42**, 860 (1990).
- [14] S. Barr and A. Zee, Phys. Rev. Lett. **65**, 21 (1990); See also J. Gunion and R. Vega, Phys. Lett. **251B**, 21 (1990); D. Chang, W.-Y. Keung, and T.C. Yuan, Phys. Lett. **251B**, 608 (1990); D. Chang, W.-Y. Keung, and T.C. Yuan, Phys. Rev. **D43**, R14 (1991); R. Leigh, S. Paban, and R. Xu, Nucl. Phys. **B352**, 45 (1991).
- [15] W. Bernreuther, T. Schröder, and T.N. Pham, Phys. Lett. **B279**, 389 (1992).
- [16] See the proceeding of *Physics and Experiments with Linear Colliders*, ed. by R. Orava, P. Eerola, and M. Nordberg, 9–14 September 1991, at Saariselkä, pub. by World Scientific, Singapore (1992).


Figures

1. The energy distributions of prompt leptons, for the case that $m_t = 170$ GeV, $\sqrt{s} = 400$ GeV. Case (a) for $N^{-1}[dN/dx(\ell^+) + dN/dx(\ell^-)]$, and case (b) for $[dN/dx(\ell^+) - dN/dx(\ell^-)]/[dN/dx(\ell^+) + dN/dx(\ell^-)]$ per unit of $\text{Im}c_d^\gamma$ (solid) or $\text{Im}c_d^Z$ (dashed).
2. The weighted up–down asymmetry of prompt leptons, for the case that $m_t = 170$ GeV, versus \sqrt{s} , per unit of $\text{Re}c_d^\gamma$ (solid) or $\text{Re}c_d^Z$ (dashed).
3. Feynman diagrams for the process $e^+e^- \rightarrow t\bar{t}$. The tree amplitude interferes with those one–loop amplitudes with (a) the final state interactions due to the exchange of a Higgs boson, or (b) the intermediate state of the ZH bosons.
4. c_d^γ and c_d^Z versus \sqrt{s} for the case that $m_H = 100$ GeV and $m_t = 120$ GeV. The parameters are chosen to be $A_I = A_R = B = 1$.
5. The total cross sections are shown in the upper three curves for cases $m_t=120$ GeV (solid), 150 GeV (dashed), and 200 GeV (dash–dotted). The partial cross sections of the helicity configurations $t_L\bar{t}_L$ and $t_R\bar{t}_R$ are shown in the lower three curves correspondingly.

Figure 3



(a)



(b)