M A D /PH /777 D T P /93/58 R A L-93-054 July 1993

#### Event shape criteria for single-lepton top signals

V.Barger<sup>a</sup>, J.Ohnemus<sup>b</sup> and R.J.N.Phillips<sup>c</sup>

<sup>a</sup>Physics Dept., University of W isconsin, M adison W I 53706, USA <sup>b</sup>Physics Dept., University of Durham, Durham, DH1 3LE, England <sup>c</sup>Rutherford Appleton Laboratory, Chilton, OX11 0QX, England

#### Abstract

Single-lepton plus jets signals from tt production at hadron colliders generally give m ore spherically sym m etrical events than the principal backgrounds from W production. W e show that sphericity and aplanarity criteria, applied to lepton plus neutrino plus four jet nalstates at the Ferm ilab Tevatron pp collider, help to discrim inate; they can be used either to validate an eventual top signal or sim ply to reduce background. A n alternative circularity criterion in the transverse plane is less successful. The predicted top quark, an essential component of the Standard M odel (SM ), is apparently too heavy to have been discovered yet at high energy colliders; the present experimental limit is  $m_t > 113$  GeV from the CDF detector [1] and independently  $m_t > 103$  GeV from the D 0 detector [2]. The consistency of radiative corrections to all electroweak data however indicates that  $m_t = 150^{+19+15}_{24-20}$  GeV [3], and hence that the top quark can probably be discovered at the Fermilab Tevatron pp collider sconer or later [4]. Here the SM predicts m ainly tt pair production via QCD with t ! bW decays; the sm allest and cleanest signals are expected in two-lepton channels [5, 6], but larger single-lepton [7, 8] and all-jet [9] signals can also in principle be separated from backgrounds. Since the top signals will initially (and perhaps for a considerable tim e) be based on a sm all num ber of events, it will be in portant to suppress the backgrounds as far as possible in ways that do not seriously reduce the signals. In this note we propose the use of event-shape criteria to distinguish the top signal and reduce backgrounds in the relatively copious single-lepton-plus-4 jet channel.

The underlying idea is that Tevatron tt production will be mainly near threshold in the CM frame of the lowest-order QCD subprocesses qq ! tt or gg ! tt, and hence will m ostly lead to rather spherical event shapes in this fram e. P hase-space factors favor spherical con gurations; the tt spin correlations [10] and decay m atrix elem ents introduce substructure but do not drastically change this spherical tendency, which can help to distinguish the signal from backgrounds. This idea is very fam iliar in the context of heavy quark searches at e<sup>+</sup> e colliders and has recently been applied to 6-jet top signals at the Tevatron [9]; we propose here to apply it to lepton-neutrino-4 jet signals from tt ! bbW W ! bbqq<sup>0</sup>, decays (where the neutrino is partially measured via transverse energy balance). Backgrounds in this channel from b- and c-quark sem ileptonic decays can be removed by lepton  $p_{\rm T}$  and isolation cuts [7]. The major background then comes from W production plus 4 QCD jets, which remains comparable with the signal after all the usual cuts on transverse momenta  $p_T$ , pseudorapidities , and m issing transverse energy  $\mathbb{E}_{\mathbb{T}}$  [7,8]; we neglect contributions from W W or W Z plus 2 Q C D jets [11] that are an order of m agnitude sm aller. This background can be further suppressed by requiring a b-jet-tag, costing the signal a tagging-e ciency factor of order 0.3 at present [4]. A Itematively, we may expect the inherent collinear singularities to cause spatial correlations of QCD jets with each other and with the beam axis, giving less

2

spherical con gurations and hence a discrimination between background and signal via an appropriate choice of event-shape variable as follows.

F inst we restrict ourselves to m om enta in the transverse plane, in order to use the m ost direct neutrino information contained in its transverse m om entum de ned by  $p_T$  () =  $E_T$ . We consider the usual \circularity" event shape variable C de ned by

$$C = 2m \ln \frac{\frac{1}{F} (p_{T}^{i} \hat{n})^{2}}{\frac{1}{F} (p_{T}^{i})^{2}}; \qquad (1)$$

sum m ed over the lepton, neutrino, and 4 jet transverse m om enta and m in im ized w ith respect to the choice of the unit vector  $\hat{n}$ . This quadratic variable is unstable against the splitting or combining of jets; although C is well de ned for any given jet algorithm, it would be theoretically preferable to use a linearized variable such as

$$C^{0} = \frac{1}{2} \min \frac{\frac{1}{p_{T}^{i}} \hat{p}_{T}^{i}}{\frac{1}{p_{T}^{i}} \hat{p}_{T}^{i}} \hat{p}_{T}^{i}}; \qquad (2)$$

Both C and C  $^{0}$  are normalized to 1 for an ideal circular event and vanish for a linear con guration.

Next we consider event shapes in three dimensions. The longitudinal neutrino momentum  $p_L$  () is not measured; it can be inferred approximately by requiring that the lepton-neutrino invariant mass m (`) =  $M_W$ , but this leaves a two-fold ambiguity. We choose the solution that allows the most consistent the reconstruction (details below) and form the conventional normalized 3 3 momentum tensor

$$M_{ab} = \frac{P_{i} p_{a}^{i} p_{b}^{i}}{\frac{1}{i} (p^{i})^{2}}$$
(3)

from the lepton plus neutrino plus 4{jet m om enta  $p_a^i$  (i = 1;:::6;a = 1;2;3), in the CM fram e of these m om enta. M <sub>ab</sub> has three eigenvalues Q<sub>i</sub> with 0 Q<sub>1</sub> Q<sub>2</sub> Q<sub>3</sub> and Q<sub>1</sub> + Q<sub>2</sub> + Q<sub>3</sub> = 1; the sphericity S and aplanarity A are then de ned by S =  $\frac{3}{2}$ (Q<sub>1</sub> + Q<sub>2</sub>) and A =  $\frac{3}{2}$ Q<sub>1</sub>. (It would be theoretically preferable to use linearized variables such as thrust and acoplanarity, but for ease of computation we use S and A here). Thus S = 2A = 1 for ideal spherical events, S =  $\frac{3}{4}$  and A = 0 for plane circular events, and S = A = 0 for linear events. W e therefore expect the Tevatron top signal to give typically larger C; C<sup>0</sup>; S; and A than the W + jets background, and have m ade quantitative parton-level calculations at p = 1.8 TeV to con m this.

For illustration we compute the pp ! ttX production and decay distributions in lowest order with full spin correlation e ects [10], using the MRS set S<sub>0</sub> parton distributions of Ref. [12] evaluated at scale Q = m<sub>t</sub>, neglecting possible additional QCD jets and taking `+ + 4 jet nal states. The total tt production rate is norm alized conservatively to the central value of Ref. [13], although values about 20% higher have recently been proposed [14]. W e calculate the W + 4 jet background from the full tree-level matrix elements following Ref. [8], evaluating the parton distributions at scale Q = hp<sub>T</sub> i in accord with Tevatron results [15]. W e sum lepton avors and charges `= e ; . W e simulate calorim eter errors by gaussian uncertainties on jet and lepton energies, following the CDF values tabulated in Ref. [16], and calculate the apparent neutrino p<sub>T</sub> from overall E<sub>T</sub> in balance. W e in pose the following acceptance cuts on transverse momenta, pseudorapidities, and jet multiplicity N<sub>j</sub>, selecting regions where the top signal should be strong,

$$p_{T}(j) > 20 \text{ GeV}; \quad j(j)j < 2; \quad N_{j} = 4;$$
  
 $p_{T}(') > 20 \text{ GeV}; \quad j(')j < 2; \quad \mathbf{E}_{T} > 20 \text{ GeV}:$ 
(4)

We also set minimum jet-jet and lepton-jet separations

$$R(j; j) > 0.7;$$
  $R(j; j) > 0.7;$  (5)

where  $(R)^2 = ()^2 + ()^2$ , to m in ic some e ects of jet-nding algorithms and lepton isolation cuts [15]. We note that these R cuts already introduce some event-shape discrimination.

We rst compare the shapes of the C; C<sup>0</sup>; S; and A distributions for the signal and W + jets background, for the case  $m_t = 150 \text{ GeV}$ . The means and standard deviations for the signal (background) are

from which we see that the circularity variables o er little discrimination in this case, whereas the S and A variables show appreciable displacements between signal and background. The S and A distributions are shown in Fig.1. We see that the signal has signi cantly di erent distributions from the background, expecially for A; this could help to validate an eventual top signal (extracted by b-tagging say). A lternatively, cutting out events below a m in in um value  $S_{m in}$  or  $A_{m in}$  would reduce the background m ore than the signal; integrated cross sections for  $S > S_{m in}$  and  $A > A_{m in}$  are shown in Fig.2. We shall illustrate below the e ects of a choice  $S_{m in} = 0.20$ ;  $A_{m in} = 0.05$  (not speci cally optim ized), sim ilar to cuts suggested for all-jet signals [9]; this reduces the signal (background) by 15% (36%), giving a cleaner result and an increase in its statistical signi cance (signal)/ (background).

To extract a top quark m ass peak, we introduce a constrained event param eterization as follows. We rst compute the two solutions for  $p_L$  () by requiring m (') =  $M_W$ ; if they are form ally complex, we set the in aginary parts to zero, getting the single nearest-to-onshell solution. We then select the pair of iets with invariant m ass closest to M  $_{\rm W}$  , and identify them as the W ! jj decay candidates. The remaining two jets are candidates for b and b jets, if this is a tt event. From the four possible ways of pairing the latter two jets with our unique W ! jj choice and the two-fold W ! ' solution, we select whichever pairing gives the closest agreem ent between the resultant b+ W invariant m asses. The m ean of these two closest m (bW) values is dened as the reconstructed top m ass  $m_{t}$  (following the notation of Ref. [5]); taking the mean gives a narrower signal peak than either the sem ileptonic or the hadronicm (bW) value separately. Our m<sup>t</sup> variable di ers from those proposed in Refs. [7,8]. Figure 3 presents the calculated  $m_t$  distribution for signal and background, for  $m_t = 130$ , 150, and 170 G eV, rst with the standard acceptance cuts of Eqs. (4) { (5) (upper curves), and also with the additional event-shape cuts S > 0.20; A > 0.05 (low er curves). Hadronization e ects have not been included, except in their contributions to calorim eter resolution; we expect they will som ewhat further sm ear the  $m_t$  peak.

Our results show that

- (a) S and A distributions of top signals di er signi cantly from background in the lepton
   + neutrino + 4 jets channel;
- (b) these di erences could help to validate an eventual top signal or alternatively reduce background and improve signi cance;

(c) the signal would peak in our reconstructed top mass variable  $m_t$ .

E vent shape criteria have previously proved prom ising in the 6 jet channel [9]; they may also have value in lepton-plus-3 jet and dilepton channels.

### A cknow ledgm ents

We thank Walter Giele for providing a copy of the VECBOS program for W + 4 jet production and thank Tao Han for discussions. JO would like to thank the Rutherford Appleton Laboratory Theory G roup and the DESY Theory G roup for hospitality during the course of this work. JO is supported by the UK Science and Engineering Research Council. The work of VB is supported in part by the U.S.D epartment of Energy under Contract No.DE-AC02-76ER00881, in part by the Texas National Laboratory Research Commission under G rant No.RGFY 93-221, and in part by the University of W isconsin Research Committee with funds granted by the W isconsin A lum ni Research Foundation.

## References

- [1] CDF collaboration: A.Barbaro-Galtieri, report to EPS Conference, Marseilles, July 1993.
- [2] D 0 collaboration: M. Narain, XXV II<sup>th</sup> Recontres de Moriond, March 1993, to be published.
- [3] P. Langacker, Pennsylvania preprint UPR-0555-T (1993).
- [4] S.Protopopescu and G.P.Yeh, reports to Top Physics Sym posium, M adison, N ovember 1992.
- [5] H.Baer, V.Barger, J.Ohnemus, and R.J.N.Phillips, Phys.Rev.D 42, 54 (1990) and references therein.
- [6] T.Han and S.Parke, FERM ILAB-PUB-93/105-T.
- [7] H.Baer, V.Barger, and R.J.N. Phillips, Phys. Rev. D 39, 3310 (1989).
- [8] F.A. Berends, H. Kuif, B. Tausk, and W.T. Giele, Nucl. Phys. B 357, 32 (1991);F.A. Berends, J.B. Tausk, and W.T. Giele, Phys. Rev. D 47, 2746 (1993).
- [9] JM.Benlloch, N.Wainer, and W.T.Giele, FERM ILAB-PUB-93/060-T.
- [10] V.Barger, J.Ohnemus, and R.J.N.Phillips, Int.J.M od.Phys.A 4, 617 (1989).
- [11] V.Barger, T.Han, J.Ohnemus, and D.Zeppenfeld, Phys. Rev. D 41, 2782 (1990).
- [12] A D.Martin, R.G.Roberts, and W.J.Stirling, Phys. Rev. D 47, 867 (1993).
- [13] R K. Ellis, Phys. Lett. B 259, 492 (1991).
- [14] E.Laenen, FERM ILAB-PUB-93/155-T.
- [15] CDF collaboration: F.Abe et al, Phys. Rev. Lett. 70, 4042 (1993).
- [16] CDF collaboration: F.Abe et al., Phys. Rev. D 45, 3921 (1992).

# Figure Captions

- 1. Sphericity S and aplanarity A distributions for tt signal ( $m_t = 150 \text{ GeV}$ ) and W + jets background, for '+ + 4 jet events at the Tevatron.
- 2. Integrated signal and background cross sections for (a)  $S > S_{min}$  and (b)  $A > A_{min}$ , versus  $S_{min}$  and  $A_{min}$ , respectively.
- 3. Reconstructed top-quark m ass  $m_t$  distributions for Tevatron tt signals with  $m_t = 130$ , 150, 170 G eV, and W + 4 jets background. Upper curves are for S > 0; A > 0; low er curves are for S > 0.20; A > 0.05.



Figure 1



Figure 2



Figure 3