## QSSR ESTIMATE OF THE B<sub>B</sub> PARAMETER AT NEXT-TO-LEAD ING ORDER

S.Narison

Theoretical Physics Division, CERN CH - 1211 Geneva 23, Switzerland and Laboratoire de Physique M athem atique U niversite de M ontpellier II Place Eugene B ataillon 34095 M ontpellier Cedex 05, France and A A. Pivovarov<sup>1</sup> N ational Laboratory for High Energy Physics (KEK), 1-1 0 ho, T sukuba, Ibaraki 305, Japan

## ABSTRACT

We compute the leading  $_{\rm s}$  corrections to the two-point correlator of the O  $_{\rm B=2}$  operator which controls the B  $^0{\rm B}$   $^0$  m ixing. Using this result within the QCD spectral sum rules approach and some phenomenologically reasonable assumptions in the parametrization of the spectral function, we conclude that the vacuum saturation values B\_B ' B\_B ' 1 are satis ed within 15%.

CERN-TH.7140/94 PM 93/16 KEK Preprint 93-184 January 1994

 $<sup>^1{\</sup>rm O}\,n$  leave from Institute for Nuclear Research of the Russian A cademy of Sciences, M oscow , Russia

The violation of the CP symmetry is still one of the most intriguing phenomena in particle physics. The standard model (SM) or ers a pattern for an explanation of this violation through the complex phase of the unitary Cabibbo-Kobayashi-Maskawa quark mixing matrix for three generations. The rigorous experimental test of the pattern requires knowing the numerical values of some hadronic matrix elements at low energies with some non-perturbative QCD methods.

In this paper we discuss an estim ate of the  $B_d^0 - B_d^0$  m ixing parameter within the QCD spectral sum rules (QSSR) approach [1,2] along the lines of Ref. [3], but by including the new  $_s$  corrections and by taking care on the detailed contributions of di erent B -like states to the spectral function. In so doing, we estim ate the two-point correlator of the O  $_{B=2} = (b_L - d_L)^2$  operator, where  $q_L$  are left-handed quark elds

$$T(x) = h0 f O_{B=2}(x) O_{B=2}(0) f i:$$
 (1)

The leading term has a trivial expression in the con guration space. Indeed, to leading order in  $_{\rm s}$ , it simply reads

$$T_{0}(\mathbf{x}) = 2N_{c}^{2} + \frac{1}{N_{c}} + \frac$$

where S (x;m) is the free ferm ion propagator and N<sub>c</sub> stands for the number of quark colours. The primemeans taking only the part of the propagator that is proportional to a matrix. The function  $_5(x) = h0 \text{ Jr} j_5(x) j_5(0) \text{ Ji}$  is the two-point correlator associated to the current  $j_5 = bi _5 d$ .

One can also rewrite Eq. (2) in the form

$$T_0(x) = 2 1 + \frac{1}{N_c}$$
 (x) (x); (3)

where  $(x) = h0 j T j_L(x) j_L(0) j li and j_L = b_L d_L$ , which has the following Lorentz decomposition in x-space

$$(x) = (0 0 + g 0^2)_{T} (x^2)_{0} 0 0_{L} (x^2):$$

Form ulae (2) and (3) demonstrate an explicit factorization of the correlator (1) in the conguration space to leading order in  $_{\rm s}$ .

The dispersion representation in x-space for any two-point correlator  $_{j}(x)$  (j = T;L;5) has the form  $_{7}$ 

$$i_{j}(x^{2}) = \int_{s_{j}}^{2} r_{j}(s)D(x;s)ds;$$
 (4)

where D (x;s) is a free boson propagator with the m ass" p s. The spectral functions  $r_1$ , read to leading order in s:

$$r_{L}^{(0)}(s) = N_{c} \frac{1}{16^{-2}} z (1 - z)^{2}; r_{T}^{(0)}(s) = N_{c} \frac{1}{16^{-2}} \frac{1}{3} (1 - z)^{2} (2 + z);$$

$$r_5^{(0)}(s) = m_b^2 N_c \frac{1}{16^2} 2 \frac{(1-z)^2}{z};$$
 (5)

where  $z = m_b^2 = s_r m_b$  is the b-quark pole mass.

One can de ne the spectral function (s) of the full correlator T (x) in the same way as in Eq. (4) and express it, to rst order in  $_{\rm s}$ , in term softhe spectral functions  $r_{\rm i}$  (s) associated to the two-line correlators. Therefore

$$(s) = r_1 (s_1) r_2 (s_2) (s; s_1; s_2) ds_1 ds_2;$$
(6)

where

$$(s;s_1;s_2) = \frac{1}{16^2 s} q (s s_1 s_2)^2 4s_1s_2$$
 (7)

is the two-body phase-space factor, and the concrete form of the spectral functions  $r_j$  (s) entering Eq. (6) depends on the representation chosen for the correlator T (x) (as in Eqs. (2,3) for  $T_0(x)$ ). The integration region in Eq. (6) is determined by the properties of the phase-space factor (7) for corresponding representation of the whole correlator; (s;s<sub>1</sub>;s<sub>2</sub>) is supposed to be equal to zero within kinem atically forbidden regions.

To leading order in 1=N<sub>c</sub>, one can write the correlator T (x) as a product of two two-line correlators given in Eq. (4). It is worthwhile to notice that this decomposition is gauge-invariant and nite, i.e. it does not require any renormalization that is a relation of the vanishing of the operator O<sub>B=2</sub> anom alous dimension in leading order in 1=N<sub>c</sub>.

Including  $\ _{\rm s}$  corrections, the spectral function in the factorization approximation reads:

$$f_{act}(s) = {}_{0}(s)(1 + {}_{f}(s));$$
 (8)

where  $_0$  (s) generates the leading-order term and  $_f$  (s) is a (properly norm alized) factorized correction in the  $_s$  order. The colour structure is the following

$$_{0}(s) = N_{c}^{2} 1 + \frac{1}{N_{c}} \sim_{0} (s); \quad f(s) = C_{F} - \frac{s}{c} \sim_{f} (s);$$

where  $C_F = (N_c^2 - 1) = 2N_c$  for the SU  $(N_c)$  colour group and the quantities with the tilde contain no explicit  $N_c$  dependence.

The representation of the spectral density  $f_{act}$  (s) through those of two-line correlators for the vector-like decom position in Eq. (3) is:

$$\begin{aligned} & \sum_{\text{fact}} (\mathbf{s}) = \sum_{1}^{Z} ds_1 ds_2 \quad (\mathbf{s}; \mathbf{s}_1; \mathbf{s}_2) f \quad \frac{s_{12}^2}{4} + 2s_1 s_2 \quad \mathbf{r}_T \quad (\mathbf{s}_1) \mathbf{r}_T \quad (\mathbf{s}_2) \\ & + \frac{s_{12}^2}{4} \quad s_1 s_2 \quad (\mathbf{r}_T \quad (\mathbf{s}_1) \mathbf{r}_L \quad (\mathbf{s}_2) + \mathbf{r}_L \quad (\mathbf{s}_1) \mathbf{r}_T \quad (\mathbf{s}_2)) + \frac{s_{12}^2}{4} \mathbf{r}_L \quad (\mathbf{s}_1) \mathbf{r}_L \quad (\mathbf{s}_2) \end{aligned}$$

where  $s_{12} = s + s_1 + s_2$ . The non-factorizable corrections are of the 1=N<sub>c</sub> order and the full spectral density can be written in the form

$$(s) = _{0}(s) (1 + _{f}(s) + _{nf}(s))$$
(9a)

and

$$n_{\rm f}(s) = \frac{C_{\rm F}}{N_{\rm c}+1} \stackrel{s}{\longrightarrow} n_{\rm f}(s) = \frac{1}{2} \quad 1 \quad \frac{1}{N_{\rm c}} \stackrel{s}{\longrightarrow} n_{\rm f}(s):$$
 (9b)

O ne of the diagram s contributing to the non-factorizable part of the whole spectral density is given in Fig. 1.

Several comments are in order here. The decomposition (9) is gauge-invariant. The same is true for the anom alous dimension of the operator  $O_{B=2}$ . We use fourdimensional algebra of Dirac's matrices throughout the computation. This seems quite natural because it allows one to make Fierz rearrangements freely, which is crucial for establishing the validity of factorization. At the same time this approach simply implies the special choice of the renormalization scheme. So, this scheme is not the standard  $\overline{MS}$  one and it is not the real dimensional reduction as used in [4] either. To the considered order in s, however, the scheme dependence reduces to a certain choice of the normalization parameter . In principle, one can x the scheme by direct comparison of our results with the massless limit for corresponding correlator [5] or, to put it another way, by comparing the non-logarithm ic parts of corrections in the massless limit one can obtain the relation between our parameter

and the corresponding  $\frac{1}{M S}$  or  $D_{\text{im Red}}$ . Thus, the answer in any desired scheme can be easily recovered by considering the massless limit. In the present paper we do not dwell on this point.

We have computed numerically the spectral density (s) in the rst non-leading order in  $_{\rm s}$ . We analyse our results, paying special attention to the presentation of the entire spectral density (s) as a sum of factorizable and non-factorizable pieces, as in Eq. (9). This decomposition is useful from the theoretical point of view. It is also interesting from the 1=N  $_{\rm c}$  analysis, and it is quite convenient technically. As for the factorizable part of the spectral density (Eq. 8) the analytical expressions for two-line spectral functions  $r_{\rm T,L}$  (s) are well known to rst order in  $_{\rm s}$ , and we use them as they have been given in Refs. [6,7]. For the non-factorizable part of the spectral density, we have to compute the gluon-exchange diagram both for unequal mass lines and for equalm ass lines of a two-line correlator. We have done it using the REDUCE system of analytical computation [8] and the table of two-loop integrals given in [6,7]. A fler using a Fierz transform ation, these non-factorizable diagram s, i.e. the corresponding analytical expression is given by a product of two traces in D irac's indices. The transform ed diagram s are given in Fig. 2.

For diagram s with a gluon connecting b or d-quarks in the loop (Fig. 2(a,b)), the resulting representation can be rendered into a product of two scalar spectral densities of two-line correlators. The relevant part of the spectral density reads:

$$r_{m} (s) = \frac{1}{16^{-2}} s' (2(1 \quad 2z)v + C_{F} - \frac{s}{2} \quad 4v(1 \quad 2z)(\frac{1}{2} + 2\ln\frac{2}{m_{b}^{2}}) + 4(1 \quad 2z)(1 \quad v)\ln z + (1 + v)(4zv + (3 + v)(z \quad 1))\ln\frac{1 \quad v}{1 + v} + (1 \quad 2z)^{2} (u) + v(18z \quad 13) + 16z \quad 8 \overset{\circ}{;}$$

where = 2 D=2, D is the dimensional regularization parameter,  $v = \begin{bmatrix} 1 & \frac{4m_{\rm b}^2}{1} \\ 1 & \frac{4m_{\rm b}^2}{s} \end{bmatrix}$  $p = \frac{1}{1 - 4z}$ , (u) = 8 (Sp (u<sup>2</sup>) Sp (u)) + 4 (2 ln (1 u<sup>2</sup>) ln (1 + u)) ln u, u = (1 v) = (1 + v),

$$Sp(u) = \int_{0}^{2} \frac{\ln (1 - t)}{t} dt$$

For a zero m ass correlator (d-quark), the spectral density has the form

$$r_0(s) = \frac{1}{16^2}s(2 + C_F - \frac{s}{4}) + 4(\frac{1}{2} + 2\ln\frac{2}{s}) + 21$$

The corresponding representation for diagram s, with dierent quark avours tied with a gluon in the bop, is of the vector type. The nonfactorizable rst order corrections to the spectral densities are:

$$r_{L;T}(s) = r_{L;T}^{(0)}(s) + C_{F} - r_{L;T}^{(1)}(s);$$

$$\begin{aligned} r_{\rm L}^{(1)}({\rm s}) &= z \,(1-z)^2 \,(\frac{1}{+} + 2 \ln \frac{2}{m_{\rm b}^2}) & 2z \,(1-z)^2 \ln (1-z) \ln (\frac{z}{1-z}) \\ &+ z \,(1-z)^2 \,(4z^2 - 3z + \frac{1}{2}) \ln (\frac{z}{1-z}) + 4z \,(1-z)^2 {\rm Sp} \,(-\frac{z}{1-z}) + z \,(1-z) \,(2z-3) \ln (1-z) \\ &- z \,(4z^2 - 7z + \frac{3}{2}) \ln z + \frac{65}{12} z^3 - \frac{17}{2} z^2 + \frac{21}{4} z - \frac{2}{3}; \\ &3r_{\rm T}^{(1)}({\rm s}) = (z+2) \,(1-z)^2 \,(\frac{1}{-} + 2 \ln \frac{2}{m_{\rm b}^2}) & 2 \,(z+2) \,(1-z)^2 \ln (1-z) \ln (\frac{z}{1-z}) \\ &+ (1-z)^2 \,(2z^3 - 2z^2 + \frac{5}{2} z + 2) \ln (\frac{z}{1-z}) + 4 \,(z+2) \,(1-z)^2 {\rm Sp} \,(-\frac{z}{1-z}) \\ &+ z \,(1-z) \,(2z+1) \ln (1-z) + \,(3z^3 + 2z^2 - \frac{9}{2} z + 2) \ln z + \frac{41}{12} z^3 + \frac{3}{2} z^2 - \frac{75}{4} z + \frac{46}{3}; \end{aligned}$$

The whole computation can be subjected to a powerful test consisting in the cancellation of divergences by the one-loop renorm alization constant of the operator O<sub>B=2</sub>, which is known to be Z<sub>O<sub>B=2</sub> = 1 3(1 1=N<sub>c</sub>) $\frac{s}{4}$ . We checked this cancellation explicitly.</sub>

The QCD results for the spectral density are given in Table 1.

At larger s (say, s  $5m_b^2$ ), where the perturbative QCD result makes sense, one can notice that the non-factorizable correction dies out faster than the factorizable one, so that the full correction comes mainly from that of the two-line correlator. Therefore, at su ciently large s, the factorization (in the sense of Eqs. (2) and (3)) becomes exact and the 'form al' 1=N<sub>c</sub> suppression of non-factorizable corrections becomes num erically valid.

U sing the QSSR approach for estimating  $B_B$ , we can work with the moments:

$$M_{i}(s_{th}) = \frac{\sum_{s_{th}}}{4m_{b}^{2}} (s)s^{i}ds;$$
(10)

which can be decomposed as:

$$M_{i}(s_{th}) = M_{i}^{0}(s_{th}) 1 + M_{i}^{f}(s_{th}) + M_{i}^{nf}(s_{th});$$

according to the corresponding decom position of the spectral density, Eq. (9a).

We show in Table 2 the strength of the non-factorizable correction for s  $5 \text{m}_{\rm b}^2$ for the following set of input parameters:  $\frac{(5)}{MS} = 175 \text{ MeV}$ ,  $m_b = 4.6 \text{ GeV}$ , and with the one-loop expression for the strong coupling constant  $_{\rm s}$  () = 6 = 23 ln ( =  $\frac{(5)}{M_{\rm s}}$ ) at the point  $= m_b$ . For given regions of integration in Eq. (10), the m on ents of the factorizable spectral density are practically independent of the power i of the weight function s<sup>i</sup>. They change by less than 5% for i=1-10. The non-factorizable correction changes its sign within the integration region for s  $5 \pm 5 \text{ m}_{\text{b}}^2$ , and  $\pm 5 \text{ m}$  om ents are more sensitive to the power i. One can notice that the non-factorizable correction does not exceed a 15% level with respect to the full factorized spectral density at su ciently large s. The full order- s corrections (factorizable+ non-factorizable) can be quite large for both the spectral density itself and its m om ents depending on the energy s as they can reach a magnitude of 100% with respect to the leading term . Hopefully, the non-factorizable corrections, measured in term softhe fully factorized (lowest order + s term s) spectral density are, nevertheless, still m oderate. However, all the factorizable corrections to the correlator (1), no matter how large they are, can be absorbed into the calculation of the decay constant  $f_B$  from the two-point correlator with two quark lines, in such a way that the relevant corrections to the  $B_{\rm B}$  parameter are only due to the non-factorizable ones.

Contrary to the QCD part, the estim ate of the phenom enological contributions to the correlator is much more involved. For the analysis to be performed one might make the following phenom enological assumptions:

{ the m inim alchoice of operator: O  $_{B=2} = g _{B=2} @ B @ B$ , can give a good description of the spectral function.

{ the contributions of the B B and B B pairs to the spectral function are equal due to their approxim ate degeneracy and to the equal values of their decay constants [9] (this feature is fully satis ed in the large  $m_b \lim it$ );

{we do not have exotic contributions due to a singlet m ixture of colored states or due to som e four-quark-like states which m ay restore the N<sub>c</sub>-structure of the correlator in order to m atch the proper (1+ 1=N<sub>c</sub>) construction entering the standard de nition of the B<sub>B</sub> parameter.

W ithin these phenom enological assumptions, we can conclude that the e ect of inclusion of the perturbative corrections into the QCD part of the correlator on the B<sub>B</sub> parameter is reasonably small and the change is not larger than 15%. The absolute value of B<sub>B</sub> how ever cannot be unam biguously established using the perturbative part of the correlator only. A ssum ing that non-perturbative non-factorizable corrections are small and the factorizable corrections are properly taken into account through  $f_{\rm B}$  we and after correcting for exotic contributions in theoretical part of the correlator

B<sub>B</sub> ' B<sub>B</sub> ' (1:00 0:15)

where we have not explicitly written the smalle ect due to the anom alous dimension of the operator O  $_{\rm B=2}$ .

An improvement of our result needs an explicit quantitative check of the previous phenom enological assumptions. Most probably, the QSSR approach based on the three-point function [10] may be more appropriate, as it does not need some of the assumptions which we have used for the two-point function. Unfortunately, one has to face, in this case, a highly involved computation of the non-recursive three-bop non-factorizable diagrams in order to obtain the  $_{\rm s}$  corrections.

A cknow ledgem ent

SN.would like to thank A.Pich for discussions. A A P. acknow ledges the kind hospitality extended to him at the Laboratoire de Physique M athem atique of M ontpellier, where the main part of the present computation has been done and thanks the C entre International des E tudiants et Stagiaires (C E S) of the European C om - m unity for nancial support. The work of A A P. is partly supported by Japan Society for the P rom otion of Science (JSPS).

## References

- M A.Shifman, A.J.Vainshtein and V.J.Zakharov, Nucl. Phys. B147 (1979) 385, 448.
- [2] For a recent review, see e.g.: S.Narison, QCD spectral sum rules,
  W orld Scienti c Lecture Notes in Physics, vol. 26 (1989).
- [3] A.Pich, PhysLett. B 206 (1988) 322.
- [4] G.Altarelli, G.Curci, G.Martinelliand S.Petrarca, Nucl Phys. B187 (1981)461.
- [5] A.Pich and E. de Rafael, PhysLett. B158 (1985)477.
- [6] S.C. Generalis, PhD thesis, Open University preprint OUT-4102-13 (1984).
- [7] S.C. Generalis, J.P. hys. G 16 (1990) 785.
- [8] A.C. Hearn, REDUCE user's manual, RAND Publication CP78 (Rev 7/91), 1991.
- [9] S. Narison, PhysLett. B198 (1987) 104; B216 (1988)191, B308 (1993) 365, Z. Phys.C55 (1992) 55 and Talk given at the Third W orkshop on Charm Factory, 1-6th June 1993, Marbella, Spain, CERN preprint TH-7042/93 (1993).
- [10] A A Ovchinnikov and A A Pivovarov, PhysLett. B 207 (1988) 333.

s=m <sup>2</sup> <sub>b</sub>	f	nf	$_{nf} = (1 + _{f})$
5.5	1.03	0.02	0.01
6.0	0.95	0.21	0.11
6.5	0.89	0.29	0.15
7.0	0.84	0.32	0.17

Table 1: Norm alized spectral densities

i	$s_{th}=m_b^2$	M f	M <sub>nf</sub>	$M_{nf} = (1 + M_{f})$
0	5.5	1.07	-0.16	-0.08
	6.0	0.99	0.11	0.06
	6.5	0.93	0.23	0.12
	7.0	0.88	0.29	0.15
5	5.5	1.08	-0.21	-0.10
	6.0	1.00	80.0	0.04
	6.5	0.94	0.20	0.11
	7.0	0.89	0.27	0.14
10	5.5	1.09	-0.27	-0.13
	6.0	1.01	0.02	0.01
	6.5	0.96	0.16	80.0
	7.0	0.91	0.23	0.12

Table 2: Norm alized m om ents of spectral densities

This figure "fig1-1.png" is available in "png" format from:

http://arxiv.org/ps/hep-ph/9403225v1