

CP violation in photon-photon collisions.

Jose Wudka

Department of Physics
University of California, Riverside
Riverside, CA 92521

November 13, 2021

Abstract

The effective Lagrangian parametrization is used to determine the CP violating effects in $\gamma\gamma$ collisions. For the processes studied the effects are found to be very small, the one exception being scalar production.

1 Introduction

Probing physics beyond the Standard Model requires high precision experiments preferably, on quantities whose standard model values are suppressed. A well known example of this type of observable is the β parameter, whose value is very close to one due to the $SU(2)_R$ transformation properties of the Standard Model scalar doublet.

A particularly interesting set of processes for which the Standard Model contribution is very much suppressed consists of those for which CP is violated. In the Standard Model amplitudes for CP violating processes are associated with the phase of the Kobayashi-Maskawa matrix and are extremely small [1, 2]. In contrast, many kinds of new physics generate comparatively

Supported in part through funds provided by the Department of Energy and by the SSC fellowship FCFY 9211.

large amounts of CP violation [3]. CP violating observables are therefore very good candidates in which to look for new physics effects.

In this talk I will consider the possibility of observing CP violating processes within the gauge-boson and scalar sector of the Standard Model [4]; part of this work was done in collaboration with J. Gunion and B. Grzadkowski, a generalization is under investigation. With only Standard Model interactions these effects are negligible, but this need not be the case in general. The environment in which I will study these processes is the photon-photon collider. Though such a machine will probably be constructed using back-scattered laser radiation in an e^+e^- collider [5], in this talk I will consider, for clarity and brevity, an ideal monochromatic collider in which both photons can be given any desired polarization. As I will show, even in this utopian situation there are great difficulties in observing a clear signal for some of the processes considered.

The approach which I will follow in this talk is to parametrize the effects of new physics in a model and process independent way by using an effective lagrangian [6]. This is, by its very nature, a process and model independent approach which preserves all the successes of the Standard Model while incorporating new physics in a consistent manner. I will not describe the formalism in detail here but refer the reader to [6]. Briefly, what is required is to construct all dimension six operators containing Standard Model fields and respecting the symmetries of the Standard Model, which also violate CP. The effective lagrangian consists of the Standard Model lagrangian plus a linear combination of these operators with undetermined coefficients. The value of these coefficients cannot be determined without further knowledge of the physics underlying the Standard Model; nonetheless these couplings can be estimated using consistency conditions. These estimates are not numerically accurate, nonetheless they do provide reliable order of magnitude value, which is what is needed in order to determine the sensitivity of a given experiment to the scale of new physics.

In this talk I will consider the processes

$$! W^+W^- ; ZZ ; HH ; H^0 \tag{1}$$

Some comment on the fermion anti-fermion final state will be made at the end.

I will assume a "light" Higgs

$$m_H \ll 3 \text{ TeV} : \quad (2)$$

For the processes of interest the relevant dimension six operators are [7]

$$O_{W^2} = \text{Tr} [W^\mu W^\mu] \quad (3)$$

$$O_{B^2} = B^\mu B^\mu \quad (4)$$

$$O_{BW} = \text{Tr} [B^\mu W^\mu] \quad (5)$$

$$O_W = \text{Tr} [W^\mu W^\nu W^\rho] \quad (6)$$

So that the Lagrangian becomes

$$\begin{aligned} \mathcal{L} = \mathcal{L}_{SM} + \frac{1}{2} g^2 O_{BW} + g^2 O_W \\ + g^2 O_{W^2} + g^2 O_{B^2} : \end{aligned} \quad (7)$$

The scale Λ determines the limit of applicability of this parametrization of heavy physics effects: all processes studied using \mathcal{L} must have energies below Λ . Indeed, the assumption that heavy physics effects are summarized by a series of effective local operators can only be true if the energy scale of interest is significantly smaller than the scale of the heavy physics. Moreover if we are studying processes whose energies are such that the underlying physics is apparent, we would not bother to study their radiative effects in order to re-discover it. These remarks, though obvious, are often ignored in the literature.

The coefficients c_i will be chosen so that Λ corresponds to the scale where the heavy physics effects are observed directly. To estimate them suppose first that the heavy physics is weakly coupled; in this case one can verify that all the operators O_i are generated by loops by the underlying theory. We then expect

$$c_i \sim \frac{1}{16\pi^2} : \quad (8)$$

If the underlying theory is strongly interacting the argument required to estimate the coefficients c_i is the same as the one used in the so-called "naive dimensional analysis" [8]. The c_i are in fact running coupling constants defined by matching conditions at the scale Λ , at which the underlying

physics becomes apparent. Then consistency requires that a change in the renormalization mass μ with $c \approx O(1)$ should not change the order of magnitude of the μ . This gives $j_W j_{B,W} = 16^2$ and $j_{B,W} j_{B,W} = 1$.

For a strongly coupled theory, however, the Higgs mass is expected to receive large $O(\Lambda^2)$ corrections so that this scenario is in general inconsistent with the above assumption that the Higgs is light. The exception occurs when this mass is protected by a symmetry (such as supersymmetry). In this case, however, the low energy spectrum of the models are invariably richer than that of the Standard Model. I will therefore assume that a light Higgs is not viable for a natural strongly coupled heavy theory. In such a situation a different parametrization, the so called chiral representation, of the effective lagrangian is required and will not be considered here due to time limitations. Because of this I will adopt the estimates $j_{ij} = 16^2$.

2 Results

I will then consider an ideal photon collider where the photons have definite momentum and perfect polarizations. As mentioned above, I will not consider the realistic situation where the photons to be considered are produced by back-scattered laser light. This is done due to time limitations, and also to avoid complications which, though quantitatively very important, obscure to a certain degree the basic problems one has to deal with when trying to uncover new physics using the processes considered in this talk.

The photons' center of mass momenta are $k_{1,2} = \frac{1}{2} \sqrt{s} (1; \pm 1; 0; 0)$ with polarizations $\epsilon_{1,2} = \frac{1}{\sqrt{2}} (0; 0; \pm 1; \pm i)$, $j = 1, 2$. In all calculations I will choose $\epsilon_{1,2}$ so as to suppress (or in the optimal case to eliminate) the Standard Model contributions. When the final particles have the same mass the final particle momenta are $\frac{1}{2} \sqrt{s} (\cos \theta; \pm \sin \theta; 0)$ where θ is the velocity of the final particles and θ is the center of mass scattering angle. With these preliminaries I turn now to the various reactions.

2.1 $\gamma\gamma \rightarrow ZZ$:

To the order we are working in the effective lagrangian there are no contributions. The leading terms for this reaction come from dimension eight operators [9], and will not be considered further in this talk.

2.2 $! W^+ W^-$:

For this process the only contributing operators are O_W and O_{BW} ; if both W vector bosons are longitudinal only the second operator contributes. The relevant diagrams are

where the solid dot denotes an O_{BW} insertion.

Assuming $\sqrt{s} \gg m_W$ allows for the use of the equivalence theorem, so that the W vector bosons can be replaced by the corresponding Goldstone particles. For longitudinally polarized W bosons at these energies in the W state O_W does not contribute, so the W result will depend on O_{BW} only.

Choosing $\epsilon_{12} = \frac{1}{\sqrt{2}}(0; 0; 1; i)$ (i.e. $\epsilon = i$) yields the amplitude

$$A(! W^+ W^-) = \frac{g_W}{2} \frac{t}{s} \frac{1}{m_W^2 - t}; \quad (9)$$

where $g_W = 16\pi^2 g_{BW}$. Note that this amplitude is real. The total cross section corresponding to this amplitude is

$$\sigma(! W^+ W^-) = \frac{g_W^2 s}{192 \cdot 3 \cdot 4} \left[1 + O\left(\frac{m_W^2}{s}\right) \right]; \quad (10)$$

In the limit of large center of mass energy the Standard Model contribution vanishes; the amplitude for longitudinally polarized W vector bosons in the W state is [10]

$$A_{SM}(! W^+ W^-) = \frac{8i(1 - \frac{t}{s})}{1 - 2 \cos^2 \theta}; \quad (11)$$

where $\theta = \arcsin \frac{\sqrt{s}}{2m_W}$ (note that it is purely imaginary so it will not interfere with the O_{BW} contribution). The corresponding cross section is

$$\sigma_{SM} = \frac{2}{s} (1 - \frac{t}{s})^2$$

$$\sigma = \frac{1}{2} \ln \frac{1 + \sqrt{1 - 4m_W^2/s}}{1 - \sqrt{1 - 4m_W^2/s}}; \quad (12)$$

which indeed vanishes as $s \rightarrow 1: \sigma_{SM} \sim 2(2m_W/s)^2$ as $s \rightarrow 1$.

I will consider the observability of this process from three points of view

- i) Require $\sigma_{new} > \sigma_{SM}$; this implies $s^3 = (1536 m_W^2)^4$. Since we also want $\sigma > s$ (else the new physics can be probed directly), this implies $\sqrt{s} > 31 \text{TeV}$; for an accelerator of this energy scales of order 32TeV can be probed. This result is obtained by assuming that $\kappa_W = 1$.
- ii) Require $N_{new} > \frac{P}{N_{SM} + N_{new}}$, where N_{new} is the number of events generated by the new physics and N_{SM} Standard Model events. Using again $\sigma > s$ and $\kappa_W = 1$ this condition is equivalent to the requirement that the luminosity for the machine is greater than $2.4 \cdot 10^6/\text{fb}$.
- iii) Require that the forward backward asymmetry be greater than 0.1 and that there be more than 10 Standard Model events. This is equivalent to a luminosity above $2 \cdot 10^6/\text{fb}$.

This clearly illustrates the enormous problems one has to deal with: absurdly large luminosities have to be invoked in order to detect a signal.

This problem can be traced back to the estimate $\kappa_W = 1$. One might be tempted to relax this condition and assume, for example, $\kappa_W = 16^2$ in which case the required luminosities drop to $10/\text{fb}$. Unfortunately such large values for the coefficients are inconsistent with the whole approach. In other words, there is no consistent way of generating such large coefficients from the underlying dynamics without radically altering the Standard Model itself (for example, one would then expect $\kappa_W = 0(1)$).

2.3 σ_{HH} :

The contributing effective operators to this process are $O_{W, B, WB}$ appearing in the diagrams

where the heavy dot denotes an effective operator insertion.

The Standard Model contributions come from loops such as

whose evaluation is straightforward. For simplicity I will consider here only the expression for the Standard Model generated by a heavy top loop, this corresponds to the effective operator [11]

$$O_{\text{heavy top}} = \frac{35}{54} \frac{G_F}{8} \frac{1}{2} H^2 \frac{1}{4} F^2 \quad (13)$$

where G_F is the Fermi constant. Note that this operator vanishes for the choice of polarizations

$$1,2 = \frac{1}{2} (0;0;1; -1); \quad (\epsilon = 1) \quad (14)$$

this will be true for all the Standard Model contributions provided the KM mixing terms are ignored (which I will do in the following due to the smallness of these effects).

The cross section for this process and for the above choice of polarizations is

$$(\sigma_{\text{H H}}) = \frac{2^2 \frac{2}{3} \frac{S}{4}}{256}; \quad (15)$$

$$\sigma_{\text{H}} = 16^{-2} (\sigma_{W^+} + \sigma_{B^+} + \sigma_{B^+ W^+}); \quad (16)$$

Note that $\sigma_{\text{H}} \ll 1$.

Since the Standard Model contribution vanishes due to the choice of polarization vectors, the observability of this process is rate dominated. To estimate the observability of this process I require that 10 events be generated in one year. This corresponds to

$$\sqrt{s} < 0.2 \frac{100 \text{ fb}}{\sqrt{\mathcal{L}}} \text{ TeV} \quad (17)$$

where \mathcal{L} denotes the luminosity.

To determine the content of this result recall that, since we are assuming that the heavy physics is not directly observed, that $\sqrt{s} > 2m_H$. Assuming for example, $m_H = 250 \text{ GeV}$ requires $\mathcal{L} > 170 \text{ fb}$. On the other hand if $\sqrt{s} = 1 \text{ TeV}$ then $\mathcal{L} > 220 \text{ fb}$. For this process the required luminosities are not absurdly large as in the previous case, but they are still large requiring many years' integrated luminosity to observe even a marginal signal. As before this can be traced to the consistent estimates of the coefficients in the Lagrangian, if I had (incorrectly) taken $\kappa_H = 16^2$ the required luminosity would drop by two orders of magnitude.

2.4 Higgs production.

The same effective operators considered above can be used to study the production of single Higgs bosons in photon colliders. The graph is simply

where the heavy dot denotes an effective operator vertex.

Note that there is no Standard Model contribution (at tree level), and that the one loop contributions can be eliminated by the above choice of polarization vectors. The cross section is then

$$\sigma(\gamma\gamma \rightarrow H) = \frac{3g^2 m_H^2}{32 \pi^2 G_F^4} \left(\frac{\sqrt{s}}{m_H} \right) \quad (18)$$

so that, integrating this over the width of the Higgs and taking

$$\sigma_H = \frac{3g^2 m_H^3}{128 \pi m_W^2}; \quad (19)$$

corresponding to a mass above the $W W$ threshold, yields

$$\begin{aligned}
 &= \frac{Z_{m_H + m_H = 2}}{m_H m_H = 2} (\sigma_{! H}) \frac{d^4 p}{s} \\
 &= \frac{256}{3} \frac{m_W^2 s_W^2}{m_H^2} \frac{1}{m_H^2}
 \end{aligned} \tag{20}$$

so that,

$$\sigma = \frac{\sigma}{100 \text{ fb}} \frac{1}{m_H^2} \frac{16 \cdot 2^4}{m_H^2} \tag{21}$$

where σ and m_H are measured in TeV.

Taking, for example, $m_H = 1 \text{ TeV}$ and $\sigma = 10 \text{ fb}$ then values of σ below 5 TeV generate more than ten events. This is a non-trivial result: a 1 TeV accelerator can probe scales $\sqrt{2}$ times its energy using this reaction; for lower Higgs mass or higher luminosities the sensitivity to σ (as a multiple of $\frac{d^4 p}{s}$) improves.

2.5 fermion state

For the process $\sigma \rightarrow f f$ there is a forbidding zoo of operators that are CP violating and should be included in the effective lagrangian describing such processes. For example, denoting a left handed fermion doublet by F and a right handed singlet by f ,

$$\begin{aligned}
 &i \sigma^{YD} F F \quad \text{h.c.}; \\
 &i \sigma^{YID} F_I F \quad \text{h.c.}; \\
 &i \sigma^{Y'} F f' \quad \text{h.c.}; \\
 &F f' B;
 \end{aligned} \tag{22}$$

etc. The analysis of such contributions is under way.

3 Conclusions

A consistent application of the effective lagrangian method gives, for most processes considered in this talk, unobservable rates, even in the

optimal situation where the Standard Model contribution vanishes. This can be traced back to the consistent estimation of the coefficients of the effective operators.

An ad-hoc over-estimation of the coefficients of the operators can give very nice predictions which might be claimed to be observable in near future colliders. These results are, however, completely unreliable being based on an inconsistent model.

The best final state here considered is that of single Higgs production for which the accelerator becomes a very respectable probe into the physics underlying the Standard Model. If there is no Higgs, or if its mass lies beyond $4 \sqrt{3} \text{TeV}$, then new interactions can be expected at this energy. This scenario was not explored in this talk.

The experimentally benign case of a two fermion final state is currently being studied.

The author gratefully acknowledges the help of J. Gunion and B. Grzadkowski.

References

- [1] H.R. Quinn, these proceedings.
- [2] R. Pececi, these proceedings.
- [3] J.F. Gunion and B. Grzadkowski, Phys. Let. B 294 (1992) 361.
- [4] J.P. Ma and B.H.J. McKellar, Univ. of Melbourne report UM-P-93-16 (unpublished). Bulletin board reference hep-hp 9303203.
- [5] I.F. Ginzburg et. al., Nucl. Instrum. & Methods 205 (1983) 47; *ibid.* 219 (1984) 5. Nucl. Instrum. & Methods A 294 (1990) 72.
- [6] See, for example, J. Wudka, Univ. of Calif. Riverside report UCRHEP-T121 (unpublished) and references therein.

- [7] W .Buchm uller and D .W yler, Nucl. Phys. B 268 (1986) 621; see also W . Buchm uller et. al. , Phys. Let. B 197 (1987) 379.
- [8] H .G eorgi and A .M anohar, Nucl. Phys. B 234 (1984) 189.
- [9] U .Baur and E L .Berger, Phys. Rev. D 47 (1993) 4889.
- [10] H .Veltm an, Saclay report SA CLAY - SPHT - 93 - 111 (unpublished).
Bulletin Board reference hep-ph 9311261.
- [11] H . Steeger et. al. , Phys. Rev. Let. 59 (1987) 385. G .L. Lin et. al. ,
Phys. Rev. D 44 (1991) 2139; Univ. of M ichigan report UM -TH -93-05
(unpublished).