

Theoretical Physics Institute  
University of Minnesota

TPI-MINN-94/42-T  
UMN-TH-1323-94  
hep-ph/9501222  
31 December, 1994

**Determining  $\alpha_s$  from Measurements at  $Z$ :  
How Nature Prompts us about New Physics**

M. Shifman

*Theoretical Physics Institute, Univ. of Minnesota, Minneapolis, MN 55455*

E-mail address: SHIFMAN@VX.CIS.UMN.EDU

**Abstract**

The value of  $\alpha_s(M_Z)$  emerging from the so called global fits based mainly on the data at the  $Z$  peak (and assuming the standard model) is three standard deviations higher than the one stemming from the low-energy phenomenology. The corresponding value of  $\Lambda_{\text{QCD}}$  is very large,  $\sim 500$  MeV, and is incompatible with crucial features of QCD. If persists, the discrepancy should be interpreted as due to contributions to the  $Z$ -quark-antiquark vertices which go beyond the standard model.

The statement that precision measurements of electroweak physics at the  $Z$  peak give no evidence whatsoever of new physics lying beyond the standard model (SM) is becoming common place now. Moreover, the values of the SM parameters extracted from analysis of the LEP and SLD data are used as canonical. This refers, in particular, to the strong coupling constant. The so called global fits which assume validity of the standard model and are based on a large set of data (“high-energy data”) yield values of  $\alpha_s(M_Z)$  in the  $\overline{MS}$  scheme which cluster around 0.125 [1, 2], with the error bars 0.005 [3]. The corresponding value of  $\Lambda_{\text{QCD}}$  is about 500 MeV [5]. These numbers, accepted as the most exact results for the strong coupling constant existing at present, propagate further into a stream of papers, published in the last year or two, devoted to various aspects of QCD.

The question arises whether Quantum Chromodynamics can tolerate these numbers. I will argue below that the answer is negative. There are two reasons why I believe that  $\alpha_s(M_Z)$  must be close to 0.11 and the corresponding value of  $\Lambda_{\text{QCD}}$  close to 200 MeV (or even smaller). A more formal argument comes from consideration of traditional “low-energy” data, the cleanest of which is the evolution of moments of the structure functions in deep inelastic scattering (DIS). These measurements are abundant and have high statistics (for a review see [6]). Theoretical formulae for the moment evolution are known in the leading and next-to-leading logarithmic approximations; if one considers the *Euclidean* domain of  $Q^2$  above  $\sim 10 \text{ GeV}^2$  non-perturbative effects play no role in the  $Q^2$  evolution of the moments provided that the moments considered are not too high. A typical result for  $\alpha_s$  emerging in this way (scaled to the normalization point  $\mu = M_Z$ ) is  $0.113 \pm 0.005$  [7].

Other “low-energy” analyses – heavy quarkonia, jets at PEP and PETRA and so on – produce similar and even lower values of  $\alpha_s$ , with the only exception to be discussed in some detail below. However, in determining the value of  $\alpha_s$  from these data one encounters serious problems: due to the essentially Minkowskian nature of the jet and quarkonium calculations precise estimates of the role of non-perturbative effects are difficult and, hence, the corresponding results for  $\alpha_s$  are inconclusive. Moreover, in some instances above even the perturbative next-to-leading corrections are not calculated so far. It should be noted that lattice calculations (e.g. Ref. [8]) also produce  $\alpha_s(M_Z) = 0.115$  or less; being a lattice outsider I do not know how reliable these lattice results are.

The second, less formal, argument in favor of  $\Lambda_{\text{QCD}} \lesssim 200 \text{ MeV}$  is the success of the operator product expansion (OPE) [9] in an extremely wide range of applications to different QCD problems. Although less formal at the moment, this argument seems more convincing to me. The analysis is based on the *Euclidean* expansions (QCD sum rules [10]) ensuring, thus, the best possible control over non-perturbative contributions. Although Wilsonian OPE is valid in any consistent field theory, practically successful numerical predictions become possible only because of the existence of a *window* in QCD – a crucial phenomenon not completely understood theoretically [11]. The window is a Euclidean domain of momenta where in certain correlation functions perturbative corrections turn out to be numerically smaller

than non-perturbative ones. In typical instances after the Borel transformation the window stretches down to  $\sim 0.7$  GeV. This fact ensures a very fast transition from essentially perturbative regime to essentially non-perturbative one. An interpretation of the window phenomenon emerges if one assumes that  $b$ , the first coefficient in the Gell-Mann-Low function, is a large parameter (numerically large), all perturbative corrections are suppressed by powers of  $1/b$ , while the non-perturbative corrections do not contain this parameter. The window phenomenon would be impossible if  $\Lambda_{\text{QCD}} \gtrsim 500$  MeV. It should be stressed that the difference between  $\alpha_s(M_Z) = 0.125$  and  $\alpha_s(M_Z) = 0.11$  (i.e.  $\Lambda_{\text{QCD}} \approx 500$  MeV *versus* 200 MeV) is not merely quantitative but, rather, qualitative. In the first case one expects a rather slow and gradual transition from the perturbative regime to the non-perturbative one, while in the second case the non-perturbative (power) corrections blow up in the domain where perturbation theory still seems convergent.

Precise measurements of  $\Lambda_{\text{QCD}}$  from the QCD sum rules can be fully formalized. As a matter of fact, the old work [12] in this direction is seminal. It yields  $\Lambda_{\text{QCD}} < 210$  MeV at the one-loop level. Later analyses along these lines also exist. Further efforts, both theoretical and experimental, are needed in order to measure  $\Lambda_{\text{QCD}}$  in this way at the level of accuracy desirable today. But even the achieved level of accuracy rules out  $\Lambda_{\text{QCD}} \gtrsim 500$  MeV.

A discrepancy between the low-energy expectations for  $\alpha_s$  and the fits at  $Z$  alerted some theorists a few years ago (see below). Their arguments were largely overshadowed later by the assertion, worked out in a series of interesting and stimulating papers [13], that a precise low-energy determination of  $\alpha_s$  from  $\tau$  is possible,  $\alpha_s(M_\tau) = 0.33 \pm 0.03$ . This result implies, in turn, that  $\alpha_s(M_Z) = 0.120 \pm 0.003$ , in accord with the value measured at the  $Z$  peak. Ref. [13] presents the state of the art in estimating all known sources of non-perturbative corrections. The leading corrections come from the gluon and four-quark condensates and turn out to be negligibly small. This calculation, quite correct by itself, gives rise to a new doctrine – a very precise determination of  $\alpha_s(M_Z)$  from essentially perturbative formula for  $\Gamma(\tau \rightarrow \text{hadrons})$  is possible – and we are witnessing now how this doctrine is gradually becoming generally accepted in the community.

The problem with all approaches of this type is that it can not be formulated as a completely Euclidean analysis. One has to deal with Minkowskian spectral densities integrated with some weights over some *finite* energy range. If in the genuinely Euclidean calculation one can reliably judge the accuracy achieved by considering the retained correction terms, an estimate of the accuracy for Minkowskian averages (integrals over a finite energy range) based on individual condensate terms is grossly misleading. To see that this is indeed the case it is sufficient to consider a model spectral density suggested in the last section of Ref. [14]. This spectral density corresponds to an infinite series of equidistant poles, with one and the same residue, and it may be relevant in the large  $N_c$  limit in the channel with one heavy quark

and a massless antiquark,

$$\Pi_{\text{model}}(E) = -\psi(-E) + \frac{1}{E} = -\sum_{n=1}^{\infty} \frac{1}{E-n} + \text{Const},$$

where  $\psi$  is the logarithmic derivative of the  $\Gamma$  function. In order to mimic the standard QCD routine in the treatment of the spectral density we proceed as follows. One considers  $\Pi(E)$  at Euclidean (negative) values of  $E$ , expands in  $1/E$  and then takes the imaginary part of the expansion,

$$\Pi(E) = -\ln(-E) + \frac{1}{2E} + \sum_{n>0} \frac{(-1)^{n-1} B_n}{2n} \frac{1}{E^{2n}}, \quad (1)$$

$$\text{Im} \Pi(E) = \pi - \frac{\pi}{2} \delta(E) + \pi \sum_{n>0} \frac{(-1)^{n-1} B_n}{(2n)!} \delta^{(2n-1)}(E) \quad (2)$$

where  $B_n$  stand for the Bernoulli numbers. The constant term in  $\text{Im} \Pi(E)$  is an analog of the ‘‘perturbative’’ term; the rest is due to ‘‘non-perturbative’’ power corrections. Being extremely simple, the model contradicts no general requirements.

It is not difficult to check that the Borel transform of this function,  $\hat{B}\Pi(\epsilon)$ , considered at Euclidean (negative) values of the Borel parameter  $\epsilon$ , possesses the property we expect from the OPE-based analysis; namely, the exact result differs from the truncated series by a quantity of order of the last power term kept. At the same time the integrals

$$\int_0^{E_0} E^n dE \text{Im} \Pi$$

calculated from the expansion (2) differ from the exact values by a large amount,  $\sim 1/E_0$ , in spite of the fact that the power series for each given moment consists here of a finite number of terms. Thus, in the first moment ( $n = 0$ ), if the first power term is retained, the second one and all others are zero, and one would expect the absolute accuracy following the line of reasoning of Ref. [13].

Of course, the model considered is relevant only in the limit  $N_c = \infty$ . For  $N_c = 3$  a natural broadening of the resonances smears the spectral density and improves the accuracy of the Minkowskean calculations done with the truncated series. Still, the strength of the non-perturbative effects in the spectral density is represented not only by individual power terms of low dimensions but, also, by the asymptotic behavior of the power terms of high orders. The corresponding contribution is exponential in energy,

$$\Delta \text{Im} \Pi \propto e^{-CE} \sim \exp\{-C e^{C'/\alpha_s(E)}\} \quad (3)$$

and, therefore, is *not seen* in the truncated OPE expansions. Equation (3) gives an estimate of a deviation from duality which shows up when one descends from the asymptotically high to lower energies.  $C$  and  $C'$  in are constants, see Ref. [14] and the forthcoming publication [15] where the issue will be discussed in more detail. At the moment no reliable *purely theoretical* method exists that would allow one

to find the constant  $C$ . In other words, theoretical estimates of non-perturbative contributions in the Minkowskian quantities of the type of the total hadronic width of  $\tau$  or the  $e^+e^-$  annihilation cross section at a given energy are rather vague, and they definitely do not have such a great accuracy as is required today in the problem of  $\alpha_s$ . One has to invoke phenomenological information, and this reverses the problem. If we accept that  $\Lambda_{\text{QCD}}$  lies in the vicinity of 200 MeV we have to conclude that about 20% of the pre-asymptotic term in the hadronic  $\tau$  width comes from non-perturbative effects (violations of duality) so that actually  $\alpha_s(M_\tau) \approx 0.27$ . At the moment this conclusion must be considered as perfectly legitimate. (By the preasymptotic term I mean  $R_\tau - 3$  where  $R_\tau$  is defined in [13].) Similar and even larger violation of duality was shown [16] to take place in the inclusive semileptonic decays of the  $D$  mesons which are very close in mass to  $\tau$ 's.

What is usually done in the conference talks and review papers to lull the public opinion is averaging of two groups of data – low-energy and high-energy values of  $\alpha_s(M_Z)$ . Then the world average usually quoted is  $0.117 \pm 0.005$ ; it lies only  $\sim 1.5$  standard deviations from either of them, and the contradiction is hidden under the rug.

If the value of  $\alpha_s(M_Z) \approx 0.11$ , as it stems from the low-energy data, what is the way out? The most placid solution of the problem would be reversing the trend of the  $Z$  peak experiments. Only three years ago the corresponding global fits used to yield numbers for the strong coupling constant which did not contradict the above value. Since then the result was steadily increasing.

To bring the value of the strong coupling constant in line with the low-energy considerations one has to diminish the experimental number for the hadronic width of the  $Z$  by  $\sim 7$  MeV. Surprisingly, this 7 MeV is the excess of the hidden beauty produced at  $Z$ , compared to the SM expectations, detected recently [1, 2]. As was noted in Refs. [17, 18], if one allows for new physics in the  $Zb\bar{b}$  vertex to take care of this 7 MeV excess in  $\Gamma(Z \rightarrow b\bar{b})$  one then solves the  $\alpha_s$  puzzle too.

If no systematic bias is found (and experts say that this scenario is very unlikely) we are forced to look for physical explanations of the discrepancy. The SM global fits can be altered if there is a contribution due to new physics. As a matter of fact one of the explanations has been already proposed – light gluinos [19]. Gluinos with masses of order of a few GeV change the rate of running of the strong coupling (it becomes slower), so that both numbers, the “low-energy”  $\alpha_s$  and the “high-energy” one become compatible with each other and compatible with the estimate  $\Lambda_{\text{QCD}} \approx 200$  MeV. Many theorists, however, are reluctant to accept this scenario because of certain specific problems associated with the light gluinos. The niche for their existence is nearly closed experimentally.

It seems more appealing to assume that new heavy particles (with mass  $\gtrsim 100$  GeV) generate, through loops, a correction to the  $Z$ -quark-antiquark vertices, enhancing the hadronic decays of the  $Z$ . In order to convert  $\alpha_s(M_Z) = 0.125$  into  $\alpha_s(M_Z) = 0.11$  it is sufficient to ensure the enhancement of the hadronic width by  $\sim 0.4\%$ . The first idea that comes to one's mind is the fourth generation [20]. Then

the  $Z$  boson has a well-defined axial coupling to the doublet of quarks belonging to the fourth generation; let us denote them by  $T$  and  $B$ . Through the  $T$  and  $B$  loops the  $Z$  boson proceeds into a pair of gluons which are then coupled to light quarks,  $u, d, s, c, b$ . (One of the gluon propagators is contracted into a point due to the  $Z$ -boson quantum numbers). Interference of this graph with the tree  $Z$ -quark-antiquark vertices produces a correction to the hadronic decays of  $Z$  proportional to  $\ln m_T/m_B$ , so that the sign of the effect can be adjusted at will. The elegance of this mechanism becomes obvious if one takes into account the fact that the two-gluon intermediate state is weak isosinglet, so that the interference in the  $u\bar{u} + d\bar{d}$  channel cancels. The same happens in the  $s\bar{s} + c\bar{c}$  channel. The only surviving correction is in the  $b\bar{b}$  channel ( $t$  is too heavy to appear in the intermediate state and cancel it). Thus, the fourth generation can naturally enhance the  $Z$  decays into  $b\bar{b}$ , the mode where the current experimental data are known to disagree with theoretical expectations at the  $2\sigma$  level [1, 2]. If one could adjust the ratio  $\ln m_T/m_B$  in such a way as to totally erase the disagreement in the  $b\bar{b}$  then this  $b\bar{b}$  enhancement would be sufficient to simultaneously solve the  $\alpha_s$  problem. Experts say, however [21, 22], that a large ratio  $m_T/m_B$  is ruled out by consistency of the SM radiative corrections to the masses and polarization operators of the  $Z$  and  $W$  bosons. Stretching all numbers to their extremes I found that the fourth generation can be responsible for at most 1.5 MeV in  $\Gamma(Z \rightarrow b\bar{b})$ , instead of the desired 7.

Leaving the fourth generation aside we can turn to superpartners. The light gluino scenario has been already mentioned. The heavy (virtual) gluino effects have been also discussed in the literature. The correction is generated by the  $Z$  coupling to squarks which then exchange a gluino and convert into quarks. Both squarks and gluino are assumed to lie in the 100 GeV ballpark. This mechanism was studied in [23], and later, even in more detail, in [24]. The squarks/gluino effect in the  $b\bar{b}$  channel is specifically addressed in the works [25]. As it follows from these calculations, the sign of the gluino contribution is correct (i.e. it produces an enhancement) and, moreover, the effect can reach the desired 0.4% in the  $Z$  hadronic width provided that the gluino/squark masses are on the light side of the allowed mass domain. Thus, superpartners in loops can, in principle, solve both difficulties simultaneously.

It should be noted, though, that if the superpartners are responsible for bringing  $\alpha_s(M_Z)$  down to 0.11 the possibility of a straightforward Grand Unification within the Minimal Supersymmetric Standard Model (MSSM) is ruled out. Indeed, the simplest version of Grand Unification, with the squark and gluino masses in the 100 GeV ballpark, implies [26] that  $\alpha_s(M_Z) = 0.125$  or larger. This seems to be an exciting observation defying the standard boring great desert scenarios [27].

Of course, one can say that the  $\alpha_s(M_Z)$  problem – 0.125 versus 0.11 controversy – is only a  $3\sigma$  effect. Being translated in the language of  $\Lambda_{\text{QCD}}$  the difference becomes quite drastic. Moreover, having  $\Lambda_{\text{QCD}}$  in the ballpark of 200 MeV is crucial for consistency of a very large number of QCD-based calculations in the low-energy domain known to produce successful predictions which, seemingly, will not survive

if  $\Lambda_{\text{QCD}} \gtrsim 500$  MeV. In any case, the question is definitely ripe enough for an intensive public debate. The data on  $\alpha_s(M_Z)$  from the  $Z$  peak, as they exist now, at the very least must be taken as a clear hint that new physics is around the corner. Whether it is supersymmetry or something else [17] is hardly possible to decide at the moment. The supersymmetric explanation is good since (i) it goes through without twisting arms, i.e. it can naturally enhance the total hadronic and  $b\bar{b}$  widths of the  $Z$  by the desired amount without spoiling the rest of the well-fit picture with the electroweak radiative corrections (see e.g. [28]); (ii) supersymmetry is anyway a popular element of the present-day theory as the only available mechanism which might explain the lightness of the Higgs particles in a natural way (assuming, of course, that relatively light Higgs particles do exist). A minimal lesson one has to draw is important for QCD practitioners. At the moment it seems reasonable to abstain from using  $\alpha_s(M_Z) = 0.125$  (and the corresponding value of  $\Lambda_{\text{QCD}}$ ) as the best measurement of these key QCD parameters. If so, the problem of the semileptonic branching ratio deficit in the  $B$  mesons [29], which nearly disappeared [30] after the corresponding theoretical formulae were evaluated with the large  $\alpha_s$ , resurfaces again. It would be interesting to check whether penguins generated by superpartners can ensure sufficient enhancement of the non-leptonic modes of  $B$ . Work in this direction has already begun, with quite encouraging results [31]. The squark penguins can give rise to the chromomagnetic operators of the type  $\bar{s}\sigma_{\mu\nu}G_{\mu\nu}b$  with the coefficients less suppressed compared to the standard model expectations [32]. Then the  $b \rightarrow s + \text{gluon}$  transition can be responsible for, say, 20% of the hadronic width of the  $B$  meson, thus eliminating any difficulties with  $\text{Br}_{sl}(B)$  [31]. Simultaneously, the expectation value of the charm multiplicity in the  $B$  decays goes down, which is also welcome. Further analysis is needed to check the overall consistency. It is necessary to verify, for instance, that the  $b \rightarrow s\gamma$  rate is not enhanced beyond what is acceptable. Similar squark penguins can play a role in the  $\Delta I = 1/2$  rule in the strange particle decays.

Another question to be considered is as follows. If the violation of duality at  $\tau$  is at the level of 20% of the pre-asymptotic term, what is to be expected in the inclusive  $B$  decays?

All questions discussed above have been repeatedly considered in the literature previously – different aspects in separate publications. I combine them together. The only element which I add is my deep conviction that  $\Lambda_{\text{QCD}}$  can not be larger than  $\sim 200$  MeV. The measurements of  $\alpha_s$  at the  $Z$  pole must be interpreted as a direct indication on new physics. Since convictions are very hard to formalize this letter should be viewed as an open invitation to further discussions among experts.

Illuminating discussions with K. Hagiwara, H. Ohnishi, L. Roszkowski, M. Virchaux and especially P. Langacker, V. Novikov and M. Vysotsky are gratefully acknowledged. I would like to thank B. Holdom, P. Langacker and M. Vysotsky for pointing out to me Refs. [17, 18, 22, 25]. This work was supported in part by DOE under the grant number DE-FG02-94ER40823.

## References

- [1] M. Vysotsky, *Electroweak Theory at Z and above*, Plenary Talk at the Int. Conf. on High Energy Physics, Glasgow, 1994 [hep-ph/9412258].
- [2] P. Langacker, *Test of the Standard Model and Searches for New Physics*, to be published in *Precision Tests of the Standard Electroweak Model*, ed P. Langacker, World Scientific, Singapore, 1994 [hep-ph/9412361].
- [3] Ref. [2] quotes even a higher value,  $\alpha_s(M_Z) = 0.127 \pm 0.005$ . The errors are believed to be dominated by theoretical uncertainties. The most precise calculation is done for the total hadronic width  $\Gamma(Z \rightarrow \text{hadrons})$ . The error quoted above,  $\Delta\alpha_s(M_Z) = \pm 0.005$ , is essentially determined by analyzing  $\Gamma(Z \rightarrow \text{hadrons})$ . If these data are excluded, all other measurements at  $Z$  (event shape, jets,...) yield  $\alpha_s(M_Z) = 0.120 \pm 0.008$  where I quote the average value obtained in Ref. [4].
- [4] H. Ohnishi, *An experimental study of QCD and jets in  $e^+e^-$  annihilation*, PhD Thesis, Nagoya University, December 1994.
- [5] At the present level of accuracy it is important to specify what particular definition of the QCD scale parameter is accepted. Most commonly used is an artificial parameter called  $\Lambda_{\overline{\text{MS}}}$ , see Ref. [6] for an excellent review. The fact that  $\Lambda_{\overline{\text{MS}}}$  does not directly measure the scale where  $\alpha_s$  becomes unity and is, thus, quite inconvenient, is clearly seen from the  $N_f$  dependence of  $\Lambda_{\overline{\text{MS}}}$  ( $N_f$  is the number of flavors). To make contact with other papers easier I will continue using  $\Lambda_{\overline{\text{MS}}}$ , with a regret. Results extracted from deep inelastic scattering (DIS) are usually expressed in terms of  $\Lambda_{\overline{\text{MS}}}^{(4)}$  while those referring to the  $Z$  peak are expressed in terms of  $\Lambda_{\overline{\text{MS}}}^{(5)}$ . The numerical relation between these two parameters depends on where exactly one matches the running laws referring to  $N_f = 4$  and  $N_f = 5$ . The corresponding uncertainty in  $\alpha_s(M_Z)$  is  $\sim 0.001$ . The scale parameter which is most relevant for the genuine low-energy hadronic physics is  $\Lambda_{\overline{\text{MS}}}^{(3)}$ . The absolute majority of authors avoid giving this parameter in fearing of introducing further uncertainties associated with matching at the charmed quark threshold. Therefore, below I will quote the values of  $\Lambda_{\overline{\text{MS}}}^{(4)}$ , denoting this parameter, by definition, by  $\Lambda_{\text{QCD}}$ . The reader should remember that  $\Lambda_{\overline{\text{MS}}}^{(4)} \approx (3/2)\Lambda_{\overline{\text{MS}}}^{(5)}$  and, moreover,  $\Lambda_{\overline{\text{MS}}}^{(3)}$  is larger than  $\Lambda_{\overline{\text{MS}}}^{(4)}$ . The value of the ratio  $\Lambda_{\overline{\text{MS}}}^{(3)}/\Lambda_{\overline{\text{MS}}}^{(4)}$  is rather sensitive to the point of matching.
- [6] See e.g. G. Altarelli, *QCD and Experiment – Status of  $\alpha_s$* , in *QCD – 20 Years Later*, Proceedings of the 1992 Aachen Workshop, eds. P. Zerwas and H. Kastrup [World Scientific, Singapore, 1993], vol. 1, page 172.
- [7] An example of the DIS data analysis can be found e.g. in M. Virchaux and A. Milsztajn, *Phys. Lett.* **B274** (1992) 221. A nice compilation is given in Ref. [6].

The central average value of  $\alpha_s(M_Z)$  from DIS quoted in [6] is 0.112. The error bars given in this review are expanded, however, up to 0.007.

- [8] A.X. El-Khadra, G. Hockney, A. Kronfeld and P. Mackenzie, *Phys. Rev. Lett.* **69** (1992) 729; C. Davies, K. Hornbostel, G. Lepage, A. Lidsey, J. Shigemitsu and J. Sloan, *A Precise determination of  $\alpha_s$  from lattice QCD*, Preprint OHSTPY-HEP-T-94-013 [hep-ph/9408328].
- [9] K. Wilson, *Phys. Rev.* **179** (1969) 1499;  
K. Wilson and J. Kogut, *Phys. Reports* **12** (1974) 75.
- [10] M. Shifman, A. Vainshtein and V. Zakharov, *Nucl. Phys.* **B147** (1979) 385; for a recent review see *Vacuum Structure and QCD Sum Rules*, ed. M. Shifman, North-Holland, 1992.
- [11] V. Novikov, M. Shifman, A. Vainshtein and V. Zakharov, *Nucl. Phys.* **B249** (1985) 445.
- [12] S. Eidelman, L. Kurdadze and A. Vainshtein, *Phys. Lett.* **82B** (1979) 278.
- [13] S. Narison and A. Pich, *Phys. Lett.* **211B** (1988) 183; E. Braaten, S. Narison and A. Pich, *Nucl. Phys.* **B373** (1992) 581; F. Le Diberder and A. Pich, *Phys. Lett.*, **B286** (1992) 147;  
For a fresh review see A. Pich, *QCD Predictions for  $\tau$  Hadronic Width: Determination of  $\alpha_s(M_\tau)$* , Preprint FTUV/94-71 [hep-ph/9412273].
- [14] M. Shifman, *Theory of Preasymptotic Effects in Weak Inclusive Decays*, in Proc. of the Workshop *Continuous Advances in QCD*, ed. A. Smilga, [World Scientific, Singapore, 1994] [hep-ph/9405246].
- [15] M. Shifman, N. Uraltsev and A. Vainshtein, to be published.
- [16] B. Blok, R. Dikeman and M. Shifman, *Calculation of  $1/m_c^3$  terms in the total semileptonic width of the D mesons*, Preprint TPI-MINN-94/23-T [hep-ph/9410293].
- [17] B. Holdom, *Phys. Lett.*, **B339** (1994) 114.
- [18] J. Erler and P. Langacker, *Implications of High Precision Experiments and the CDF Top Quark Candidates*, Preprint UPR-0632T [hep-ph/9411203].
- [19] I. Antoniadis, J. Ellis and D. Nanopoulos, *Phys. Lett.* **B262** (1991) 109; M. Jezabek and J.H. Kühn, *Phys. Lett.* **B301** (1993) 121; L. Clavelli, *Phys. Rev.* **D46** (1992) 2112; L. Clavelli, P. Coulter and K. Yuan, *Phys. Rev.* **D47** (1993) 1973; J. Blümlein and J. Botts, *Phys. Lett.* **B325** (1994) 190.
- [20] The consideration of this point presented below was worked out in collaboration with V. Novikov and M. Vysotsky .

- [21] V. Novikov and M. Vysotsky, private communication.
- [22] V. Novikov, L. Okun, A. Rozanov, M. Vysotsky and V. Yurov, *Do the present electroweak precision measurements leave room for extra generations?*, Preprint CERN-TH-7252-94.
- [23] K. Hagiwara and H. Murayama, *Phys. Lett.* **B246** (1990), 533.
- [24] A. Djouadi, M. Drees and H. Konig, *Phys. Rev.* **D48** (1993) 308.
- [25] M. Bouwens and D. Finnel, *Phys. Rev.* **D44** (1991) 2054; G. Kane, C. Kolda and J. Wells, *Phys. Lett.*, **B338** (1994) 219.
- [26] G. Kane, C. Kolda, L. Roszkowski and J. Wells, *Phys. Rev.* **D49** (1994) 6173; P. Langacker and N. Polonsky, *Phys. Rev.* **D47** (1993) 4028 [for an update see N. Polonsky, *Unification and Low-Energy Supersymmetry at One and Two-Loop Orders*, PhD Thesis, University of Pennsylvania, 1994].
- [27] By the simplest version of Grand Unification I mean the one which ignores subtleties due to possible GUT-scale thresholds and high-dimension (nonrenormalizable) operators. Including these effects one can, perhaps, descend somewhat below  $\alpha_s(M_Z)=0.125$ . How much below – this question requires further investigation.
- [28] G. Altarelli, *Precision Tests of the Standard Model: Theory*, in Proc. Int. Europhysics Conference on High-Energy Physics, Marseille, 1993, eds. J. Carr and M. Perrottet, Editions Frontieres, Gif-sur-Yvette, 1994, page 689.
- [29] I. Bigi, B. Blok, M. Shifman and A. Vainshtein, *Phys. Lett.* **B323** (1994) 408.
- [30] E. Bagan, P. Ball and V. Braun, *Charm quark mass corrections to non-leptonic inclusive B decays*, Preprint TUM-T31-67-94 [hep-ph/9408306]; E. Bagan, P. Ball, V. Braun and P. Gosdzinsky, *Theoretical update of the semileptonic branching ratio of B mesons*, Preprint DESY-94-172 [hep-ph/9409440].
- [31] A. Kagan, *Implications of TeV Flavor Physics for the  $\Delta I = 1/2$  Rule and  $Br_{sl}(B)$* , Preprint SLAC-PUB-6626/94 [hep-ph/9409215].
- [32] A. Vainshtein, V. Zakharov and M. Shifman, *ZhETF* **72** (1977) 1275 [*JETP* **45** (1977) 670]; M. Shifman, A. Vainshtein and V. Zakharov, *Phys. Rev.* **D18** (1978) 2583; (E) *Phys. Rev.* **D19** (1979) 2815.