

Thermal Two Point Function of a Heavy Muon in hot QED plasma within Bloch-Nordsieck Approximation

K. Takashiba

Department of Physics, Faculty of Science,
Osaka City University, Sugimoto, Sumiyoshi-ku,
Osaka, 558, Japan
(February 3, 2022)

Abstract

The thermal propagator of a heavy muon propagating in a hot QED plasma is examined within the Bloch-Nordsieck approximation, which is valid in the infrared region. It is shown that the muon damping rate is finite, in contrast to the lower-order calculation with hard thermal loop resummations taken into account.

11.10W x, 12.20.m, 12.38.Lg, 13.35.Bv

Typeset using REVTeX

Electronic address: niegawa@ocux.physics.osaka-cu.ac.jp

I. INTRODUCTION

Because of much interest in the early Universe as well as in the quark-gluon plasma to be produced in heavy-ion collisions, many authors have studied gauge theories at high temperature. Thermal propagators of bosons in real-time thermal field theory carry the Boltzmann factor $1/(e^{E/T} + 1)$. For a massless boson like gauge boson, this factor behaves as $1/T$ at the infrared region, $k \rightarrow 0$, which leads to stronger infrared divergences in thermal amplitudes than in vacuum theory. In vacuum theory, it is well known [13] that all the physical quantities are free from infrared singularities, provided one sums over all degenerate final states (Bloch-Nordsieck mechanism). An important exception to this general statement is the so-called Coulomb singularity. For thermal reaction rates, due to the stronger infrared singularities mentioned above, it is not known if they do not possess infrared singularities, except again for Coulomb singularities. Up until now, a proof is available of absence of leading infrared singularities in a generic thermal reaction rate [4] and of all infrared singularities in a bremsstrahlung process within some specific approximation [5].

It has been established [6,7,14] that, to obtain a consistent perturbation expansion, resummations of hard-thermal-loop (HTL) contributions are necessary. According to this HTL resummation scheme, infrared behaviors of thermal amplitudes are softened or screened compared with those in naive perturbative calculation. This softening renders some otherwise divergent physical quantities finite [7,8]. There are, however, some other physical quantities which are still infrared divergent to leading order in HTL resummation scheme. A well-known example is the quark (muon) damping rate in a hot QCD (QED) plasma [9]. As a matter of fact, the quark (muon) damping rate diverges due to the Coulomb singularity. In naive perturbation calculation, the divergence is of power type. In HTL resummation scheme, as mentioned above, the infrared behaviour is softened and the infrared divergence becomes to be of logarithmic type. It is expected, at least for quark case, that the infrared singularity is screened when still higher order resummation is performed. As far as the infrared behaviour of a spinor field in QED is concerned, Bloch-Nordsieck approximation is a good approximation [10,11]. In this paper, the thermal two-point function of a heavy muon in a QED plasma is computed within this approximation. Various properties of the result are discussed, among those is an especially important conclusion: The decaying behavior of a heavy muon does not suffer from divergence.

II. THERMAL TWO POINT FUNCTION OF A HEAVY MUON

By the term "heavy muon" we mean so heavy a muon that it is not thermalized, i.e., $e^{E/T} \gg 1$. The hot QED plasma means that the temperature T of the plasma is much higher than the electron mass, $T \gg m_e$. As discussed in Sec. I, the infrared behavior of the theory is responsible for the divergent damping rate. Thus, we employ the Bloch-Nordsieck approximation to the muon sector of the QED Lagrangian. The Lagrangian density is obtained [10,11] by substituting a constant vector u ($u^2 = 1$) for \not{p} in the muon sector of the standard QED Lagrangian density:

$$L^{\text{total}} = L + L_{\mu} + L_A ; \quad (2.1)$$

$$L = \overline{(iu \quad @ + eu \quad A \quad M)} ; \quad (2.2)$$

$$L = \overline{(i \quad @ + e \quad A \quad m)} ; \quad (2.3)$$

$$L_A = \frac{1}{4} F \quad F ; \quad (2.4)$$

$$F = @ \quad A \quad @ \quad A ; \quad (2.5)$$

where $;$ $;$ $M ; m$, and A stand, respectively, for heavy muon eld, electron eld, muon mass, electron mass and photon eld. It is known that, in the infrared region, the spinor structure of the heavy muon does not play any significant roles [10,11]. Then the above Lagrangian density well approximates the exact QED Lagrangian in the infrared region.

To obtain the thermal two-point function of the heavy muon, we start from the Euclidean or imaginary-time formalism of thermal eld theory, initiated by Matsubara [12], in which a time arrow flows from zero to $i=T$ in a complex time plane. In this formalism, the energy is discrete; $p_4 = 2nT$ ($n = 0; 1; 2; \dots$) for a boson eld and $p(2n+1)T$ for a fermion eld. At the final stage, we analytically continue the two-point function, thus obtained, to real time (or real energy). A four vector $a = (a; a_4)$ in the imaginary-time formalism is obtained from the counterpart $(a_0; a)$ in Minkowski space by an analytic continuation, $a_0 \rightarrow i a_4$. Similar continuation is made for tensors.

The generating functional is introduced as usual,

$$Z = \int D \overline{D} \quad D \quad \overline{D} \quad D A \exp S ; \quad (2.6)$$

$$S = \int_0^T d^4x \quad L^{\text{total}} - \frac{1}{2} (@ \quad A^2) + \overline{+} + \overline{+} + \overline{+} + \overline{+} + j \quad A ; \quad (2.7)$$

where the covariant gauge is employed. In Eq. (2.7), $;$ $;$ $j;$ are sources being conjugate to the elds with which they couple. After integrating over $;$ $;$ and $;$ in Eq. (2.6), the two point function of the heavy muon is obtained as,

$$\begin{aligned} G_E &= \frac{1}{Z} \frac{\overline{Z}(\overline{x}) \quad \overline{Z}(y)}{\overline{Z}} [; ; ; j] ; \quad = 0 \\ &= \frac{1}{Z} \int D A \det(iu_E \quad @ \quad eu \quad A \quad M) \\ &\quad \det(i \quad @ \quad e \quad A \quad m) G(x \quad y; eA; M) \\ &\quad \exp \frac{1}{4} \int_0^T d^4x F \quad F - \frac{1}{2} (@ \quad A^2) ; \end{aligned} \quad (2.8)$$

where

$$(iu_E \quad @ \quad eu \quad A \quad M) G(x \quad y; eA; M) = E(x \quad y) ; \quad (2.9)$$

$$E(x \quad y) = T \sum_{k_4 = \text{odd}}^X \frac{d^3k}{(2\pi)^3} \exp[ik \quad (x \quad y)] ; \quad (2.10)$$

$$k_4 = (2n+1)T \quad (n = 0; 1; 2; \dots) ; \quad (2.11)$$

In Eq. (2.10), " $k_4 = \text{odd}$ " indicates that the summation runs over n as defined in Eq. (2.11).

Since the heavy muon is not thermalized, the muon determinant has been reduced to $\det(iu_E - M)$ as in vacuum theory [10]. Using the formula $\det a = \exp(\text{Tr} \ln a)$ and Furry's theorem [13], we have

$$\det(i\partial_\mu - eA - M) = \det(i\partial_\mu - M) \exp \left[\frac{e^2}{2} \int d^4 z_1 d^4 z_2 (A(z_1) G(z_1, z_2; m) (A(z_2) G(z_2, z_1; m) + O(e^4)) \right]; \quad (2.12)$$

$$G_0(x, y; m) = T \sum_{k_4 = \text{even}}^X \int^Z \frac{d^3 k}{(2\pi)^3} \frac{\exp[ik \cdot (x - y)]}{k + m}; \quad (2.13)$$

Here $G_0(x, y; m)$ denotes a bare electron propagator. As seen from Eq. (2.8) with Eq. (2.12), in the heavy muon two-point function, the contributions of electrons enter through thermal electron loops. Here we employ the HTL resummation scheme for the electron sector: To leading order, among such electron loops, only HTLs that yield corrections to soft-photon external lines are important. In hot QED considered here, the $O(e^4)$ term in Eq. (2.12) vanishes [14] to leading order in the HTL resummation scheme. We then ignore it in the following.

Following [10], we introduce the integral representation,

$$G(x, y; eA; M) = \int_0^Z d\tau \exp i (iu_E - \partial_\mu eA - M + i\tau) G(x, y); \quad (2.14)$$

where τ is a "convergence" factor. Defining the function $U(\tau)$ as,

$$U(\tau) = \exp i (iu_E - \partial_\mu eA - M + i\tau) G(x, y); \quad (2.15)$$

$$G(x, y; eA; M) = \int_0^Z d\tau U(\tau); \quad (2.16)$$

we have

$$\frac{\partial U}{\partial \tau} = (iu_E - \partial_\mu eA - M + i\tau) U(\tau); \quad (2.17)$$

$$U(0) = G(x, y); \quad (2.18)$$

Now we assume the following form for the solution to Eq. (2.17) with Eq. (2.18),

$$U(\tau) = T \sum_{q_4 = \text{odd}}^X \int^Z \frac{d^3 q}{(2\pi)^3} \exp iK \cdot (x - y) \tilde{A}; \quad (2.19)$$

$$+ iq \cdot (x - y) - i(M + e u q \cdot \tau);$$

Eq. (2.17) with Eq. (2.19) leads to

$$\frac{\partial K}{\partial} = u_E \frac{\partial K}{\partial x} + e u_E A ; \quad (2.20)$$

$$K(x=0) = 0 ; \quad (2.21)$$

where $u_E = (u; u_4)$. It is straightforward to solve this equation in the form :

$$K(x, y;) = eT \sum_{k_4 = \text{even}}^x \frac{d^3 k}{(2)^3} \exp[ik \cdot (x - y)] A(k)_E u d^0 \exp[iu \cdot k^0] ; \quad (2.22)$$

$$k_4 = 2nT \quad (n = 0; 1; 2; \dots) ; \quad (2.23)$$

where " $k_4 = \text{even}$ " indicates that the summation runs over n as defined in Eq. (2.23).

Using all the formulae displayed above, and carrying out the integration by part, we obtain for G_E in Eq. (2.8):

$$G_E = i D A_0 \sum_{q_4 = \text{odd}}^x \frac{d^3 q}{(2)^3} \exp[iq \cdot (x - y)] i (M + u \cdot q) \cdot$$

$$\frac{e^2}{2} \sum_z d^4 z_1 d^4 z_2 \text{Tr} (A) G(z - z_2; e = 0; m) (A) G(z - z_4; e = 0; m)$$

$$+ d^4 z \frac{1}{2} A \cdot \partial^2 \cdot 1 - \partial \cdot \partial A$$

$$ieT \sum_{k_4 = \text{even}}^x \frac{d^3 k}{(2)^3} A(k)_E \exp[ik \cdot (x - y)] d^0 \exp[iu \cdot k^0] ; \quad (2.24)$$

Going to the momentum space,

$$G_E = \int d^4(x - y) \exp[i p \cdot (x - y)] G \quad (2.25)$$

we have,

$$G_E = i D A_0 \sum_{l_4 = \text{even}}^x \frac{d^3 l}{(2)^3} A(l) \cdot l^2 \cdot 1 - \frac{1}{l^2} \cdot l \cdot l + (l) \cdot A(l)$$

$$ieT \sum_{l_4 = \text{even}}^x \frac{d^3 l}{(2)^3} \exp[i l \cdot (x - y)] A(l)_E d^0 \exp[i l \cdot u^0] ; \quad (2.26)$$

where ϵ is the thermal vacuum polarization tensor of the photon:

$$(l) = e^2 T \sum_{k_4 = \text{odd}}^x \frac{d^3 k}{(2)^3} \text{Tr} \frac{1}{(k - l) + M} \frac{1}{k + M}$$

$$= \epsilon^{\mu\nu} \frac{1}{2} T + \epsilon^{\mu\nu} + \frac{1}{2} \frac{l^\mu l^\nu}{l^2} ; \quad (2.27)$$

The exponent in Eq. (2.26) is quadratic with respect to A , and we can perform the functional integration over A :

$$G_E = i \int_0^Z d\exp i(M + u_E p_c) + \frac{1}{2^2} D(\cdot; u_E); \quad (2.28)$$

$$D(\cdot; u_E) = (2)^{3D} T^X d^D \frac{1}{(1-u)^2} \frac{1}{\cos(1-u)} \\ + \frac{1}{l^2 + T} : u^2 : \frac{(1-u)^2}{l^2}; + \frac{1}{l^2 + L} : u_E^2 : \frac{(1-u)^2}{l^2} u^2 + \frac{(1-u)^2}{l^2}; \\ + \frac{1}{l^2} ; \quad (2.29)$$

where α is the fine structure constant. Here, in order to regulate the ultraviolet divergence, we have employed the dimensional regularization, $D = 3 - \epsilon$ with $\epsilon > 0$. (A dimensionful parameter is introduced, as usual, so that e remains dimensionless.) In the HTL approximation, which we employ for the electron sector, the explicit form of Π_T and Π_L are known [6,9,15]:

$$\tau = \frac{1}{2} m_{sc} \frac{l_4^2}{r^2} : 1 \quad @ \quad 1 + \frac{r^2}{l_4^2} A \frac{i l_4}{2 j j} L(i l_4; j j); \quad ; \quad (2.30)$$

$$L = m_{sc} \frac{r^2}{R^2} \left(1 - \frac{i l_4}{2 j l_j} L(i l_4; j l_j) \right); \quad (2.31)$$

where

$$m_{sc} = \frac{eT}{3}; \quad (2.32)$$

$$L(i_4; j) = \ln \frac{i_4 + j}{i_4} : \quad (2.33)$$

m_{sc} in Eq. (2.32) is thermal Debye mass of the thermal photon. In obtaining Eqs. (2.29)–(2.33), the electron mass m_e has been ignored (cf. the first paragraph).

Let us turn back to $D(\cdot; u_E)$ in Eq. (2.29). After a straightforward but tedious calculation (cf. Appendix A), we have

$$D(\mathbf{u}_E) = \frac{1}{2} \left(\frac{u_4^2}{\mathbf{u}_E} \left(\frac{1}{2} + \ln \frac{2}{p - m_{sc}} \right) - \frac{1}{2} \frac{e^{m_{sc} \mathbf{u}_E}}{\mathbf{u}_E m_{sc}} + E_i(\mathbf{u}_E m_{sc}) \right) \\ + \frac{1}{2} \left(\frac{1}{\mathbf{u}_E} + \frac{1}{2} \left(1 + \ln \frac{2}{p - m_{sc}} \right) \right) (1)^2 \mathbf{u}_E \\ + \frac{1}{2} \left(\frac{u_E^2}{\mathbf{u}_E} \left(\frac{1}{2} - \ln \frac{p}{2} \right) + \frac{2}{2} + \frac{2 \ln \frac{4}{2}}{2} \frac{2 u_4}{\mathbf{u}_E} \tan^{-1} \frac{u_4}{\mathbf{u}_E} \right)$$

$$\begin{aligned}
& 2^2 T \frac{u_E^2}{j^2} \ln \frac{T}{2} + \ln \frac{u_E}{j} + 4^2 T \frac{u_E^2}{j^2} \int_0^1 dx \frac{1}{x} \frac{1}{e^{2T(j \cdot u_4 x)} - 1} \\
& + \frac{u_E^2}{j(j \cdot u_4)} \ln(1 - e^{2T(j \cdot u_4)}) + c.c.: \\
& + 2^2 T \frac{u_4 u_E^2}{j^2} \int_0^1 dx \frac{1}{j \cdot u_4 x} \frac{1}{e^{2T(j \cdot u_4 x)} - 1} + c.c.: \\
& + \left(\frac{1}{2} - \ln 2 \right) + \left(\frac{1}{2} + \ln 4 T \right) - 2 \ln(1 - e^{2T(j \cdot u_4)}) \\
& 2 j \cdot T \frac{1}{e^{2T(j \cdot u_4)} - 1} + \frac{1}{2} \ln(1 - e^{2T(j \cdot u_4)}) + c.c.: \\
& + T (j \cdot u_4) \frac{1}{e^{2T(j \cdot u_4)} - 1} + c.c.: \\
& + 3 i u_4 T \int_0^1 dx \frac{1}{e^{2T(j \cdot u_4)} - 1} + c.c.: \\
& i 2^2 T^2 u_4 \int_0^1 dx (j \cdot u_4 x) \frac{e^{2T(j \cdot u_4 x)}}{(e^{2T(j \cdot u_4 x)} - 1)^2} + c.c. ; \quad (2.34)
\end{aligned}$$

where γ_E and Ei are, respectively, the Euler constant and the exponential integral function and $u_E = \sqrt{u_4^2 + j^2}$. The details are given in Appendix A.

An analytic continuation of G_E in Eq. (2.28) with Eq. (2.34) back to Minkowski space, which is obtained through $p_4 \rightarrow i(p_4 + i)$; $u_4 \rightarrow i(u_4 + i)$, yields the retarded function. We normalize it through MS scheme [3], which amounts to choose the renormalization factor Z_2 as

$$Z_2 = e^{\frac{3}{2} - \frac{1}{2}} ; \quad (2.35)$$

Then we obtain,

$$G_R = i \int_0^1 d \exp \left(i (M^2 - u \cdot p) \right) \frac{1}{2^2} D_R ; \quad (2.36)$$

where $u \cdot p = u_\mu p^\mu$ is the scalar product in Minkowski space.

III. PHYSICAL CONTENTS OF THE TWO POINT FUNCTION

We study the behaviors of D_R in Eq. (2.36) in various limits of parameters involved. We divide the region $0 < 1$ into several regions, which are discriminated by the two parameters $A = \frac{1}{j^2}$, $B = \frac{1}{m_{sc} j^2}$ (see Appendix A).

a) A

$$D_R \sim 2 \left(1 + \frac{3}{2} \frac{1}{2} + \ln 2 \right) \frac{i}{2} \quad (3.1)$$

b) A B

$$D_R = 2^2 T \left(\frac{1}{j} \ln T \frac{i}{2} + 1 - \frac{j}{2} \right) + 2 \frac{3}{2} \frac{1}{2} \ln \frac{2^{p-T}}{j} \frac{u_0}{j} \ln \frac{u_0 + j}{u_0 + j} - i^2 \frac{u_0}{j} \quad (3.2)$$

c) B

$$D_R = 2^2 T \frac{1}{j} \ln T \frac{i}{2} + 1 - \frac{u_0^2}{j^2} \ln \frac{T}{j n_{sc}} \frac{i}{2} + \frac{1}{2} 2^2 T \frac{u_0^2}{j^2 m_{sc}} + 2 \frac{3}{2} \frac{1}{2} \ln \frac{2^{p-T}}{j} \frac{u_0}{j} \ln \frac{u_0 + j}{u_0 + j} - i^2 \frac{u_0}{j} : \quad (3.3)$$

In the above equations, use has been made of $u^2 = 1$.

When the muon is far from the mass shell $M \approx p \approx u \approx M$, in Eq. (2.36), the region A gives the dominant contribution:

$$G_R = \frac{A (1 + \epsilon)}{M u} \frac{1}{\pi (M u - p)} ; \quad (3.4)$$

$$= \frac{3}{2} \left(\epsilon \right) ; \quad (3.5)$$

$$A = \exp - \left(1 + \frac{3}{4} \left(\ln \epsilon \right) \right) ; \quad (3.6)$$

where ϵ is the gamma function. When $\epsilon > 3$, $\epsilon < 0$ in Eq. (3.4). This is just as in vacuum theory [10], which we do not discuss any further. When $\epsilon = 3$, G_E in Eq. (3.4) behaves as the bare propagator.

We are now in a position to discuss the decaying behavior of the heavy muon. For this purpose, we should study G_R near on the muon mass shell $M \approx p \approx u \approx 0$. The time dependence of the heavy-muon two-point function is obtained through the Fourier transformation;

$$G_R = \int \frac{dp_0}{2} G_R \exp[i p_0 t] \\ = \frac{M}{E} \exp[- i E t] \exp \frac{1}{2} D_R (= M t = E ; u) : \quad (3.7)$$

In obtaining Eq. (3.7), use has been made of the on-shell approximation $(M + u - p) \approx u' - E$. From Eq. (3.7) and Eqs. (3.1)-(3.3), it is straightforward to obtain the time dependence of G_R :

a) A

$$G_R = \frac{M}{E} e^{-i E t} \frac{M t}{E} \exp - \left(1 + \frac{3}{4} \left(\ln \epsilon \right) \right) \epsilon \quad (3.8)$$

b) A B

$$G_R = \frac{M}{E} e^{iEt} \left(\frac{M T t}{E} \exp \left(\frac{M T t}{E} \left(\frac{1}{j} - \frac{i}{2} + \frac{1}{2} \frac{j}{j^2} \right) \right) + \left(\frac{3}{2} - \frac{1}{2} \ln \frac{2^{p-T}}{u_0} \right) \frac{u_0}{2j} \ln \frac{u_0}{u_0 + j} + i \frac{u_0}{2j} \right) \quad (3.9)$$

c) B

$$G_R = \frac{M}{E} e^{iEt + F(t)}; \quad (3.10)$$

$$F(t) = \frac{M T t}{E} \left(\frac{M T t}{E} \ln \frac{M T t}{E} \left(\frac{1}{j} - \frac{i}{2} + \frac{1}{2} \frac{u_0^2}{j^2} \ln \frac{T}{j m_{sc}} \left(\frac{i}{2} + \frac{1}{2} \right) \right) - \left(\frac{u_0^2}{j^2} \frac{1}{m_{sc}} \left(\frac{3}{2} - \frac{1}{2} \ln \frac{2^{p-T}}{u_0} \right) + \frac{u_0}{2j} \ln \frac{u_0}{u_0 + j} + i \frac{u_0}{2j} \right) \right); \quad (3.11)$$

We discuss two cases separately. It is to be noted that $j = \frac{1}{M} p = \frac{v}{1 - v^2}$ with v being the external muon velocity:

$$0 < v < 1$$

In the region a) above, the behavior of G_R is gauge-parameter dependent, and we cannot draw any physically sensible conclusion. In the region b), the factor $\frac{M T t}{E} \exp \left(\frac{M T t}{E} \left(\frac{1}{j} - \frac{i}{2} + \frac{1}{2} \frac{j}{j^2} \right) \right)$ in G_R , Eq. (3.9), dominates over others. Then, the leading part of G_R is gauge-parameter independent and increases as t increases. In the region c) the dominant part of G_R is gauge-parameter independent, which first increases and then decreases. The turning point in t is approximately given by $\frac{E}{M T} \left(\frac{T}{m_{sc} j} \right)^{1=v^2}$.

$$v \rightarrow 1$$

In the region a) and b) the situation is the same as the first sentence in the above case. In the region c) G_R decreases monotonously.

Let us discuss the large t region c) more elaborately. The leading contribution comes from logarithmic parts of $F(t)$ in Eq. (3.11) (cf. Eq. (3.3)). It is to be noted that the gauge-parameter dependent part in Eq. (3.11) may be neglected and Eq. (3.11) is gauge independent:

$$F(t) = \frac{M T t}{E} \left(\frac{M T t}{E} \ln \frac{M T t}{E} \left(\frac{u_0^2}{j^2} \ln \frac{T}{m_{sc} j} \right) \right); \quad (3.12)$$

A striking feature is that Eq. (3.12) is divergence free or finite. The damping rate γ_d is usually defined through the behavior of G_R at large t ; $G_R \sim e^{iEt - \gamma_d t}$ ($t \rightarrow \infty$). Were it not for the factor $\ln t$ in Eq. (3.12), the damping rate would be identified from Eq. (3.10) with Eq. (3.12). Eq. (3.12) indicates that, at large t , the heavy muon in a QED plasma decays much faster than the constant damping rate. In any case, the quantity, $F(t)$, in Eq. (3.12) governs the damping behavior of the heavy muon. The behavior of $F(t)$ in Eq. (3.12) is as follows:

$$0 < v < 1$$

$F(t)$ decreases after $t = t_d \left(\frac{E}{M T} \left(\frac{1}{1 - v^2} \right)^{1-v^2} \right)$. This means that after t_d , the heavy muon begins to decay.

$$v \rightarrow 1$$

In this case,

$$F(t)' = -\frac{M T t}{E \frac{1}{1 - v^2}} \ln \frac{M m_{sc} t}{E \frac{1}{1 - v^2}} : \quad (3.13)$$

This is always negative and the heavy muon decays.

IV . S U M M A R Y A N D D I S C U S S I O N

In this paper we have derived and discussed the two point function of a heavy muon passing through a QED plasma within the Bloch Nordsieck approximation. The assumption of heavy muon means that the muon line is hard. The electron sector only comes in as thermal radiative corrections to photon lines. Among those, we only take into account the HTL corrections to soft photon lines. HTL corrections to vertices are not necessary.

It is well known that, to leading order of HTL resummation scheme, the damping rate of a quark (muon) diverges due to the Coulomb-b-type infrared singularity even to the lowest non-trivial order of resummed perturbation theory. It is because only the electric sector of soft gluon (photon) line is screened by the thermal mass. As far as QCD is concerned, the infrared singularity is screened by the magnetic mass which is expected to arise through still higher order resummations. But in QED, no magnetic mass is induced.

In this paper, within the Bloch Nordsieck approximation, it is shown that the "damping rate" of a heavy muon in a hot QED plasma is finite. More precisely, the heavy muon decays as $t^{\frac{M T}{E} + \frac{1}{2}}$ at large t .

Finally it is worth mentioning the case where the HTL corrections to the soft photon line are ignored. (Explicit computation is given in Appendix B). For $M \gg p \gg M$, the gauge parameter dependent parts of G_R (cf. Eq. (2.36)) is the same as Eq. (3.5). Also for $\epsilon = 3$, G_R reduces essentially to the bare propagator. In the nearly on the mass shell case the time dependence of G_R is quite different from that of Eq. (3.10) with Eq. (3.12):

$$G_R' = \frac{M}{E} e^{-iEt} \exp \left[-\frac{M T}{E} t^{\frac{M T}{E} + \frac{1}{2}} \right]; \quad (4.1)$$

the anti-damping behavior! From this fact, we learn that the HTL resummation for soft photon lines plays an crucial role for obtaining a physically sensible damping behavior.

A C K N O W L E D G E M E N T

I gratefully thank Prof. A. Niegawa for his kind help and thank many volunteers who develop public domain softwares.

APPENDIX A : CALCULATION OF $D(\mathbf{k}; U_E)$

According to the HTL resummation scheme, when the external photon lines are soft $l_4; l_5 \ll O(eT)$, the HTL corrections are necessary. Since l_4 is $2\pi nT$ ($n = 0; 1; 2; \dots$), l_4 with $n \neq 0$ is no longer in the soft region. Then we should take care of the $n = 0$ mode only.

The explicit expression for $T_{\mu L}$ in Eq. (2.27), within the HTL approximation, are given in [7,15]. From these expression, we have

$$T_{\mu L}(n=0) = 0; \quad L_{\mu}(n=0) = m_{sc}^2 = \frac{e^2 T^2}{3} : \quad (A1)$$

We now decompose $D(\mathbf{k}; U_E)$ in Eq. (2.29) into two parts:

$$\begin{aligned} D(\mathbf{k}; U_E) &= T(2) \sum_{l_4=even}^Z d^D l D(\mathbf{k}; U_E; l_4; l) \\ &+ T(2) \sum_{l_4=even}^Z d^D l D(\mathbf{k}; U_E; l_4; l) \\ &+ T(2) \sum_{l_4=even}^Z d^D l D(\mathbf{k}; U_E; l_4; l); \end{aligned} \quad (A2)$$

where the prime on the summation symbol in Eq. (A2) means to take summation over l_4 with non-zero modes. We further decompose Eq. (A2) as

$$T(2) \sum_{l_4=even}^Z d^D l D(\mathbf{k}; U_E; l_4; l) = \sum_{m=1}^3 D_m; \quad (A3)$$

$$T(2) \sum_{l_4=even}^Z d^D l D(\mathbf{k}; U_E; l_4; l) = \sum_{m=1}^2 D_m^0; \quad (A4)$$

where

$$D_1 = T(2) \sum_{l_4=even}^Z d^D l \frac{1}{(l^2 - u^2)} \frac{(u_4)^2}{l^2 + m_{sc}^2}; \quad (A5)$$

$$D_2 = T(2) \sum_{l_4=even}^Z d^D l \frac{1}{(l^2 - u^2)} \frac{u^2}{l^2}; \quad (A6)$$

$$D_3 = T(1)(2) \sum_{l_4=even}^Z d^D l \frac{1}{(l^2 - u^2)} \frac{e^0}{l^2} \frac{1}{A}; \quad (A7)$$

$$D_1^0 = T(2) \sum_{l_4=even}^Z d^D l \frac{1}{(l^2 - u^2)} \frac{u_E^2}{l^2}; \quad (A8)$$

$$\begin{aligned} D_2^0 &= T(1)(2) \sum_{l_4=even}^Z d^D l \frac{1}{(l^2 - u^2)} \frac{\cos(l \cdot u)}{(l \cdot u)^2} \\ &\quad \frac{1}{l^2} : \end{aligned} \quad (A9)$$

It is to be noted that the integrand of D_m ; ($m = 1; 2; 3$) and D_m^0 ; ($m = 1; 2$) are regular at $l = 0$.

We first evaluate D_m ; ($m = 1; 2; 3$). Carried out the integration over the polar angle $\cos^{-1}(\frac{l}{l_4})$, Eq. (A 5) becomes

$$D_1 = T(2) \frac{u_4^2}{j^2} 2^{1-2} (=2) \int_0^Z dl (l^2 + m_{sc}^2)^{-1} \left(\frac{1}{l^2} e^{ijl} + \text{c.c.} \right) : \quad (\text{A } 10)$$

In order to perform the integration over l , it is convenient to analytically continue the integrand into a complex l plane. By deforming the integration contour (as shown in Fig. 1), we obtain,

$$\begin{aligned} D_1 &= T(2) \frac{u_4^2}{j^2} 2^{1-2} (=2) \frac{j}{2} m_{sc} \int_0^Z \frac{dl}{m_{sc}} \frac{(l^2 - m_{sc}^2)^{-2}}{l^2} e^{i\frac{Z-l}{2}} (1 - e^{ijl}) + \text{c.c.} \\ &\quad + i \lim_{\epsilon \rightarrow 0} \int_0^Z \frac{dl}{m_{sc}} \frac{(l^2 - m_{sc}^2)^{-2}}{l^2} (1 - e^{ijl}) + \text{c.c.} \\ &= 2^2 T \frac{u_4^2}{j^2} \left[\frac{1}{m_{sc}} + \ln \frac{2}{m_{sc}} - \frac{1}{2} \right] \frac{1}{j m_{sc}} e^{ij m_{sc}} + E_i(j m_{sc}) \\ &= 2^2 T \frac{u_4^2}{j^2} \left[\frac{1}{m_{sc}} + \ln \frac{2}{m_{sc}} - \frac{1}{2} \right] \\ &\quad + 2^2 T \frac{u_4^2}{j^2} \left[\ln(j m_{sc}) + O(j^2 m_{sc}) \right] (j m_{sc} - 1) \\ &\quad + 2^2 T \frac{u_4^2}{j^2} \left[\frac{1}{m_{sc}} + O\left(\frac{e^{-j m_{sc}}}{m_{sc}}\right) \right] (j m_{sc} - 1) : \quad (\text{A } 11) \end{aligned}$$

Through similar procedures, D_2 and D_3 may be evaluated:

$$D_2 = 2^2 T j \left[\frac{1}{m_{sc}} + \ln \frac{2}{m_{sc}} + \frac{1}{2} \right] : \quad (\text{A } 12)$$

$$D_3 = (-1)^2 T j : \quad (\text{A } 13)$$

The terms proportional to $1/m_{sc}$, which correspond to UV divergent piece at $l = 0$ or $D = 3$, is to be eliminated later by renormalization.

Let us evaluate D_m^0 ($m = 1; 2$) in Eq. (A 4). To evaluate both Eq. (A 8) and (A 9), it is convenient to introduce

$$F(l_4) = T(2) \int_0^Z d^D l \frac{1}{(l^2 - l_4^2)^2} \frac{1}{l^2 + m^2} : \quad (\text{A } 14)$$

The integral in Eq. (A 8) is obtained from $F(l_4)$ with $m = 0$, while the integral in Eq. (A 9) is obtained from $\partial F(l_4) = \partial$ with $m = 0$. Carrying out the integrations over the polar angle and using the technique of deforming the integration contour in the l -plane, we obtain

$$F(l_4) = T(2) \int_0^Z dl \frac{1}{j^2} \left[\frac{l_4^2}{l^2} + \frac{u_4^2}{j^2} + \dots \right] = 2$$

$$\begin{aligned}
& + \sin \frac{Z_1}{2} p \frac{1}{l_4^2 + i\epsilon} dl \frac{1}{l^2} (l_4^2 + i\epsilon)^{-2} \\
& \left(\frac{1}{(u_4 l_4 + i\epsilon)^2} + \text{cc} - \frac{e^{iu_4 l_4} \epsilon}{(u_4 l_4 + i\epsilon)^2} + \text{cc} \right) \# \\
& : \tag{A 15}
\end{aligned}$$

As to the integration of the third term in last brackets in Eq. (A 15), we first carry out the integration by parts and the integration by deforming the contour C_1 to $C_2 \cup C_3$ in Fig. 2. Then we shift the integration variable to get

$$\begin{aligned}
F(l_4) &= T(2)^{-1} = 2 \left(= 2 \right) \frac{1}{i\epsilon} \frac{l_4^2}{1 + \frac{u_4^2}{i\epsilon}} + \\
& + \sin \frac{Z_1}{2} (l_4^2 + i\epsilon)^{-\frac{1}{2}} \int_0^{Z_1} dl \frac{(l^2 - 1)^{-2}}{il\epsilon + p \frac{u_4}{l_4^2 + i\epsilon}} + \text{cc} \\
& + \frac{Z_1}{i\epsilon} p \frac{1}{l_4^2 + i\epsilon} dl \frac{e^{il_4 l}}{1} + \frac{1}{i\epsilon} \frac{e^{il_4 u_4}}{i\epsilon} \frac{p}{l_4^2 + i\epsilon} + \text{cc} : \\
& + \frac{i}{i\epsilon} \int_0^{Z_1} \frac{u_4}{i\epsilon} dl \frac{e^{il_4 l} (p \frac{u_4}{l_4^2 + i\epsilon} - il)}{l_4^2 + i\epsilon} + \text{cc} : \tag{A 16}
\end{aligned}$$

Using Eq. (A 16) with $\epsilon = 0$ and carrying out the summation over l_4 , we obtain

$$\begin{aligned}
D_1^0 &= \sum_{l_4 = \text{even}}^X u_E^2 F(l_4) \\
&= 2^{-2} T \frac{u_E^2}{i\epsilon} \left[\frac{1}{2} \ln^2 - \frac{2}{\epsilon} + 2 \ln \frac{4}{2} \frac{3=2T}{2} \frac{2 u_4}{i\epsilon} \tan^{-1} \frac{u_4}{i\epsilon} \right. \\
& \quad \left. + 2^{-2} T \frac{u_E^2}{i\epsilon} \ln \frac{T}{2} + \ln \frac{u_E}{i\epsilon} \right] + 4^{-2} T \frac{u_E^2}{i\epsilon} \int_0^{Z_1} dx \frac{1}{x e^{2^{-T} i\epsilon x} - 1} \\
& + \frac{u_E^2}{i\epsilon (i\epsilon - u_4)} \ln(1 - e^{2^{-T} (i\epsilon - u_4)}) + \text{cc} : \\
& + 2^{-2} T \frac{u_4 u_E^2}{i\epsilon} \int_0^{Z_1} dx \frac{1}{i\epsilon - u_4 x} \frac{1}{e^{2^{-T} (i\epsilon - u_4 x)} - 1} + \text{cc} : \tag{A 17}
\end{aligned}$$

D_2^0 in Eq. (A 9) may be evaluated as:

$$D_2^0 = \sum_{l_4 = \text{even}}^X T(2)^{-1} d^D l \frac{1}{l^4} \frac{\partial^3}{\partial \epsilon^3} F(l_4) \# ; \tag{A 18}$$

and we obtain,

$$D_2^0 = \left(-1 \right) + \frac{1}{\epsilon} + \ln \frac{2}{\epsilon} + \left(-\ln 4 T \right) + 2 \ln(1 - e^{2^{-T} i\epsilon}) + 2 i\epsilon T \frac{1}{e^{2^{-T} i\epsilon} - 1}$$

$$\begin{aligned}
& \frac{1}{2} \ln(1 - e^{2T(\hbar j \hbar u_4)}) \quad \text{cc:} \\
& T(\hbar j \hbar u_4) \frac{1}{e^{2T(\hbar j \hbar u_4)} - 1} \quad \text{cc:} \\
& 3\hbar u_4 T \int_0^1 dx \frac{1}{e^{2T(\hbar j \hbar u_4)} - 1} + \text{cc:} \\
& + i2^{-2} T^2 u_4 \int_0^1 dx (\hbar j \hbar u_4 x) \frac{e^{2T(\hbar j \hbar u_4 x)}}{(e^{2T(\hbar j \hbar u_4 x)} - 1)^2} + \text{cc:} \quad \# \quad (A19)
\end{aligned}$$

Now let us study the behaviors of D_m^0 ($m = 1, 2$) in the following two limits (\hbar_A is as in the text, $\hbar_A = \frac{1}{\hbar j T}$):

a)

$$D_1^0 \sim \frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2u_E}{2\hbar j} + 2^{-2} T \frac{u_E^2}{\hbar j} \left(\frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2\hbar j}{2\hbar j} \right); \quad (A20)$$

$$D_2^0 \sim \left(\frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2u_E}{2\hbar j} \right) + \left(\frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2\hbar j}{2\hbar j} \right) 2T \hbar j; \quad (A21)$$

and,

b)

$$D_1^0 \sim \frac{1}{2} + \frac{1}{2} - \ln \frac{4^{-3=2} T}{2} \frac{u_4}{\hbar j} \tan^{-1} \frac{u_4}{\hbar j} + 2^{-2} T \frac{u_E^2}{\hbar j} \left(\frac{1}{2} + \frac{1}{2} + \ln \frac{2\hbar j}{u_E T} \right); \quad (A22)$$

$$D_2^0 \sim \left(\frac{1}{2} + \frac{1}{2} - \ln \frac{4^{-3=2} T}{2} \right); \quad (A23)$$

Substituting all the results obtained above into Eq. (A2), we deduce the following behaviors of $D(\hbar; u_E)$ in various limits (\hbar_B is as in the text, $\hbar_B = \frac{1}{m_{sc} \hbar j}$):

a)

$$D(\hbar; u_E) \sim \frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2u_E}{2\hbar j} \quad \left(\frac{1}{2} + \frac{1}{2} + \ln \frac{2u_E}{2\hbar j} \right); \quad (A24)$$

b)

$$\begin{aligned}
D(\hbar; u_E) \sim & 2^{-2} T \frac{u_E^2}{\hbar j} (\ln T u_E + \frac{1}{2}) + \frac{1}{2} + \frac{1}{2} - \ln \frac{4^{-3=2} T}{2} \frac{u_4}{\hbar j} \tan^{-1} \frac{u_4}{\hbar j} \\
& \left(\frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2\hbar j}{2\hbar j} \right) \left(\frac{1}{2} + \frac{1}{2} - \ln \frac{4^{-3=2} T}{2} \right); \quad (A25)
\end{aligned}$$

c)

$$\begin{aligned}
D(\mu; u_E)' &= 2^2 T \frac{u_E^2}{j^2} (\ln T u_E + 1) + 2 \left(\frac{1}{2} + \frac{1}{2} \ln \frac{4^{3=2} T}{2} \right) \frac{u_4}{j^2} \tan^{-1} \frac{u_4}{j^2} \\
&\quad + (1)^2 T j^2 \left(\frac{1}{2} + \frac{1}{2} \ln \frac{4^{3=2} T}{2} \right) \\
&\quad + 2^2 T j^2 \frac{u_4^2}{j^2} \left(\frac{1}{2} + 1 \ln j^2 m_{sc} \right) + 2^2 T \frac{u_4^2}{j^2 m_{sc}} : \quad (A26)
\end{aligned}$$

The remaining tasks are to eliminate UV divergence through renormalization with MS scheme and to perform the integration (see text).

APPENDIX B: NO HTL CORRECTION TO THE PHOTON PROPAGATOR

We set $T_{\mu L} = 0$ in Eq. (2.29). In this case, Eq. (2.29) reduces to

$$D(\mu; u_E) = D_{1+2} + D_3 + \sum_{m=1}^{\infty} D_m^0 \quad (B1)$$

$$\begin{aligned}
&= T(2) \int d^D l \frac{1}{(l^2 - u^2)} \frac{1}{(l^2 - u^2)} \\
&\approx \frac{u_E^2}{l^2} + (1) \frac{1}{l^2} : \quad (B2)
\end{aligned}$$

Here

$$D_{1+2} = T(2) \int d^D l \frac{1}{(l^2 - u^2)} \frac{u_E^2}{l^2} ; \quad (B3)$$

$$D_3 = (1) T(2) \int d^D l \frac{1}{(l^2 - u^2)} \frac{1}{l^2} : \quad (B4)$$

Similar procedures as in Appendix A leads to

$$D_{1+2} = 2^2 T \frac{u_E^2}{j^2} \left(\frac{1}{2} + \frac{1}{2} \ln \frac{2 j^2}{p} \right) ; \quad (B5)$$

$$D_3 = (1)^2 T j^2 : \quad (B6)$$

Substituting the above Eqs. (B5) and (B6) and Eqs. (A20)-(A23) to Eq. (B1), we have

a) Λ

$$D(\mu; u_E) = 2 \left(\frac{1}{2} + \frac{1}{2} + 1 + \ln \frac{2 u_E}{2} \right) + (1) \left(\frac{1}{2} + \frac{1}{2} + \ln \frac{2 u_E}{2} \right) : \quad (B7)$$

b) Λ

$$D(\cdot; u_E)_{m_{sc}=0} = 2^{-2} T \frac{u_E^2}{|j|} (\ln T u_E + (1) + 2 \frac{1}{-} + \frac{1}{2} \ln \frac{4^{-3=2} T}{2} \frac{u_4}{|j|} \tan^{-1} \frac{u_4}{|j|} \\ + 2 (1) T |j| (1) \frac{1}{-} + \frac{1}{2} \ln \frac{2 u_E}{2} : \quad (B 8)$$

REFERENCES

- [1] F. Bloch, A. Nordsieck, Phys. Rev. 52, 54 (1937).
- [2] D. R. Yennie, S. C. Frautschi, and H. Suura, Ann. Phys. (N.Y.) 13, 379 (1961); T. D. Lee and M. Nauenberg, Phys. Rev. 133, B1549 (1964); J. M. Jauch and F. Rohrlich, The Theory of Photons and Electrons (Springer-Verlag, 1980).
- [3] T. Muta, Foundations of Quantum Chromodynamics (World Scientific, Singapore, 1987).
- [4] A. Niegawa, K. Takashiba, Nucl. Phys. B 370, 335 (1992); See also M. Le Bellac and P. Reynaud, in: Banff/CAP workshop on thermal field theory, edited by F. C. Khanna, R. Kobes, G. Kunstatter, and H. Umezawa, (World Scientific, Singapore, 1994) p.440.
- [5] H. A. Weldon, Phys. Rev. D 44, 3955 (1991); 49, 1579 (1994).
- [6] R. D. Pisarski, Phys. Rev. Lett. 63, 1129 (1989).
- [7] E. Braaten and R. D. Pisarski, Nucl. Phys. B 337, 569 (1990).
- [8] E. Braaten and M. H. Thoma, Phys. Rev. D 44, 1298, R2625 (1991); J. I. Kapusta, P. Lichard, and D. Seibert, ibid. D 44, 2774 (1991); R. Baier, H. Nakkagawa, A. Niegawa, and K. Redlich. Z. Phys. C 53, 433 (1992); ibid. 45, 4323 (1992).
- [9] C. P. Burgess and A. L. Marini, Phys. Rev. D 45, R17 (1992); A. Rebhan, ibid. D 46, 482 (1992). R. Baier, H. Nakkagawa, A. Niegawa, in Proceedings of the Workshop on High Temperature Field Theory, Winnipeg, July 1992 Can. J. Phys. (to be published); R. D. Pisarski, Phys. Rev. D 47, 5589 (1993); M. E. Carrington, University of Winnipeg Report No. WIN-93-xx, 1993 (unpublished); T. Altherr and D. Seibert, CERN Report No. CERN-TH.6882/93, 1993 (unpublished). See, also, V. V. Lebedev and A. V. Smilga, Ann. Phys. (N.Y.) 202, 229 (1990); Physica A 181, 187 (1992); T. Altherr, E. Petitgirard, and T. del Rio Gaztelumutia, Phys. Rev. D 47, 703 (1993); A. V. Smilga, Bern University Report No. BUTP-92/39, 1992 (unpublished); S. Peigne, E. Pilon, and D. Schi, Orsay Report No. LP THE Orsay 93-13, 1993 (unpublished); R. Baier and R. Kobes, Phys. Rev. D 50, 5944 (1994); A. Niegawa, Phys. Rev. Lett 73, 2023 (1994).
- [10] N. N. Bogoliubov, D. V. Shirkov, INTRODUCTION TO THE THEORY OF QUANTIZED FIELDS (Interscience Publishers Ltd., London, 1959)
- [11] A. Kerner and N. G. Stefanis, Phys. Rev. D 40, 2103 (1989).
- [12] T. Matsubara, Prog. Theor. Phys. 14, 351 (1955); See also J. I. Kapusta, Finite-temperature field theory (Cambridge University Press, 1989); N. P. Landsman and Ch. G. van Weert, Phys. Rep. 145, 141 (1987).
- [13] See e.g. C. Itzykson and J.-B. Zuber, Quantum Field Theory (McGraw-Hill, New York, 1980); J. D. Bjorken and S. D. Drell, Relativistic Quantum Fields (McGraw-Hill, New York, 1965).
- [14] E. Braaten and R. D. Pisarski, Nucl. Phys. 339, 310 (1990); See also, J. Frenkel and J. C. Taylor, Nucl. Phys. B 334, 199 (1990).
- [15] V. P. Silin, Zh. Eksp. Teor. Fiz. 38, 1577 (1960) [Sov. Phys. JETP 11, 1136 (1960)]; V. V. Klimov, Yad. Fiz. 33, 1734 (1981) [Sov. J. Nucl. Phys. 33, 934 (1981)]; Zh. Eksp. Teor. Fiz. 82, 336 (1982) [Sov. Phys. JETP 55, 199 (1982)]; H. A. Weldon, Phys. Rev. D 26, 1394 (1982); R. D. Pisarski, Physica 158A, 146 (1989).

FIGURES

FIG .1. Integration contour in the complex l plane

FIG .2. The integration along the contour C_1 may be evaluated along the contour $C_2 \stackrel{L}{=} C_3$.




Fig. 1

This figure "fig1-1.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9501223v1>




Fig. 2

This figure "fig1-2.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9501223v1>