hep-ph/9501240 CERN-TH.7537/94 GeF-TH-1/95 IFUM 489/FT

# A Study of U ltraviolet R enorm alon A m biguities in the D eterm ination of $_{\rm s}$ from D ecay

G.Altarelli, P.N ason<sup>1</sup> and G.R idol <sup>2</sup>

CERN TH-D ivision, CH-1211 Geneva 23, Switzerland

#### A bstract

The divergent large-order behaviour of the perturbative series relevant for the determ ination of s from decay is controlled by the leading ultraviolet (UV) renorm alon. Even in the absence of the rst infrared (IR) renorm alon, an ambiguity of order  $2 = m^2$  is introduced. We make a quantitative study of the practical implications of this ambiguity. We discuss the magnitude of UV renorm alon corrections obtained in the large N f lim it, which, although unrealistic, is nevertheless interesting to some extent. We then study a number of improved approxim ants for the perturbative series, based on a change of variable in the B orel representation, such as to displace the leading UV renorm alon singularity at a larger distance from the origin than the str IR renorm alon. The spread of the resulting values of  $_{s}$  (m<sup>2</sup>) obtained by di erent approximants, at di erent renorm alization scales, is exhibited as a measure of the underlying ambiguities. F inally, on the basis of m athem atical m odels, we discuss the prospects of an actual improvement, given the signs and magnitudes of the computed coe cients, the size of  $_{\rm s}$  (m  $^2$ ) and what is known of the asymptotic properties of the series. Our conclusion is that a realistic estimate of the theoretical error  $_{\rm s}$  (m<sup>2</sup>) 0:060, or  $_{\rm s}$  (m<sup>2</sup><sub>z</sub>) cannot go below 0:006.

CERN-TH.7537/94

December 1994

<sup>&</sup>lt;sup>1</sup>On leave of absence from INFN, Sezione diM ilano, M ilan, Italy.

<sup>&</sup>lt;sup>2</sup>On leave of absence from INFN, Sezione diGenova, Genoa, Italy.

#### 1. Introduction

The possibility of measuring  $_{s}$  from -decay has been extensively studied in a series of interesting papers, in particular by Braaten, Narison and Pich [1-4]. The relevant quantity is R = (! + hadrons)= (! + 1), with l = e; . At present [5] the ALEPH collaboration nds

$$(\mathbf{R})_{\rm exp} = 3.645 \quad 0.024: \tag{1.1}$$

Based on this result, it is argued, if the form alism of QCD sum rules is assumed, according to SVZ [6], that

$$_{\rm s}$$
 (m<sup>2</sup>) = 0:355 0:021 (1.2)

and nally

$$_{\rm s}$$
 (m $_{\rm z}^2$ ) = 0:121 0:0016 (exp) 0:0018 (th) = 0:121 0:0024: (1.3)

Given that m is so small, this determination of  $_{s}$  (m $_{z}^{2}$ ) appears [7–10] a bit too precise!

In defence of this method [11] one can certainly point out that R has several combined advantages. D ropping som e inessential complications, R is an integral in s of a spectral function R (s) which is the analogue of  $R_{e^+e^-}$  (s) but for the case of charged weak currents. Thus, rst, R is even more inclusive than  $R_{e^+e}$  (s) and one expects that the asymptotic regime is more precocious for more inclusive quantities. Second, one can use analyticity in order to transform the relevant integral into an integral over the circle  $i_{5j} = m^2$  [4]. This not only gives some con dence that the appropriate scale of energy for the evaluation of R is of order m, but also shows that the integration over the low-energy dom ain helps very much in smearing out the complicated behaviour in the resonance region. A lso important is the presence of a phase space factor that kills the sensitivity of the spectral function near Res = m<sup>2</sup>, where there is a gap of validity of the asymptotic approximations due to the vicinity of the cut singularities and also to the nearby charm threshold. On the circle jsj= m<sup>2</sup>, asymptotic form ulae should be approximately valid for the correlator. The perturbative component of R (s) is known up to term s of order  $(m^2)^3$  [12]. One can hope to get som e control of the non-perturbative corrections by using the operator

product expansion and som e estim ate (either experim ental or by som e m odel) of the dom inant condensates, in the spirit of the QCD sum rules [6].

This series of virtues of R is indeed real but would not be su cient in itself to justify the precision on  $_{s}$  (m<sup>2</sup>) which is claim ed. The real point is that no corrections of order  $1=m^2$  are assumed to exist. The fact that there is no operator with the corresponding dimension in the short distance expansion is not su cient, because there could be non-leading corrections in the coe cient function of the leading operator. We think it is a fair statem ent that there is no theorem that quarantees the absence of  $^{2}$  =m<sup>2</sup> term s in R in the massless limit; no theorem that proves that term s of order  $^{2}$  = m<sup>2</sup> cannot arise from the mechanism that generates con nement. But even if in principle the above theorem would exist, still, in practice, there would be ambiguities on the leading-term perturbative expansion of order  $^{2}$ =m<sup>2</sup> from the ultraviolet renorm alon sequence associated to the divergence of the perturbative series for the spectral function [13-21]. The present note is mainly devoted to a quantitative discussion of the impact of UV renorm alon ambiguities on the determ ination of  $_{s}$  (m<sup>2</sup>). On the basis of the accumulated knowledge on renormalon behaviour, we address the question of what is the theoretical error on  $_{s}$  (m<sup>2</sup>) and examine possible ways to decrease it. We discuss the magnitude of UV renormalon corrections obtained from explicit calculations [18-21], which although based on unrealistic simpli ed schemes, are nevertheless interesting to some extent. We then study a number of improved approximants for the perturbative series, based on a change of variable in the Borel representation [14,17], such as to displace the leading UV renorm alon singularity at a larger distance from the origin than the stin frared (IR) renorm alon. The spread of the resulting values of  $_{s}$  (m<sup>2</sup>) obtained by di erent approximants, at di erent renorm alisation scales is exhibited as a measure of the underlying am biguities. Finally, on the basis of m athem atical m odels, we discuss the prospects of an actual im provem ent, given the signs and the magnitudes of the computed coe cients, the size of  $_{s}$  (m<sup>2</sup>) and what is known of the asymptotic properties of the series. Our conclusion is that a realistic extinate of the theoretical error cannot go below  $\int_{S} (m^2)$ 0:060, or

 $_{\rm s}$  (m  $_{\rm z}^2$ ) 0:006.

The organisation of this article is as follows. In sect 2 we summarise the basic formulae and discuss di erent procedures to do the integration over the circle that di er by resumming or not an in nite series of  $\large^2$  term s". We discuss the

relative merits of the various procedures and their scale dependence. In sect.3 we introduce the problem s related to the divergence of the perturbative series, we review the Borel transform method and the renorm alon singularities. In sect.4 we derive som e useful form ulae obtained in the Borel space after integration on the circle. In sect 5 we consider the explicit form for the leading UV renorm alon singularity derived in perturbation theory in the large  $N_f$  limit,  $N_f$  being the number of avours. This lim it is not meant to be realistic, but, for orientation, we evaluate the quantitative in pact that such an UV renorm alon would have on the determ ination of  $_{\rm s}$  (m  $^2$ ). We nd that this e ect is rather sm all. In sects.6,7, which contain the main original results of this work, we introduce and study a num ber of im proved approxim ants that could in principle suppress the ambiguity from the leading UV renorm alon. We study the combined e ects of di erent, a priori equivalent, procedures, di erent accelerators of convergence and di erent choices of the renorm alisation scale. We also study in a simple mathematical model under which conditions for the known coe cients of the series the accelerator method leads to a better approximation of the true result. Finally, in sect. 8 we present our conclusion.

# 2. Basic Form ulae and Truncation Ambiguities

The quantity of interest is the integral over the hadronic squared mass s in decay of a function R (s) analogous to  $R_{e^+e^-}$  (s), weighted by a phase-space factor. In the lim it of massless u;d;s quarks we have [1-3]:

$$R = \int_{0}^{Z_{m^{2}}} \frac{ds}{m^{2}} 2 \, 1 \, \frac{s}{m^{2}} \, \frac{1}{1 + \frac{2s}{m^{2}}} \, R \, (s): \qquad (2.1)$$

R (s) is proportional to the imaginary part of a current-current correlator:

$$R(s) = \frac{N}{2} Im(s) = \frac{N}{2i} [(s+i)(s-i)];$$
 (2.2)

The norm alization factor N is de ned in such a way that, in zeroth order in perturbation theory, R(s) = 3. In turn, the correlator (s) is related to the Adler function D (s), de ned in such a way as to remove a constant:

D (s) = 
$$s \frac{d}{ds} N$$
 (s): (2.3)

By rst integrating by parts and then using the Cauchy theorem one obtains for R the result

$$R = \frac{1}{2 i} \prod_{j:j=m^2}^{I} \frac{ds}{s} = 1 + \frac{s}{m^2} + \frac{s}{m^$$

The Adler function D has a perturbative expansion of the form :

D (s) = 
$$D_{n=0}^{0} D_{n}a(s)^{n} D_{1}a(s) + D_{2}a^{2}(s) + D_{3}a^{3}(s) + \dots$$
  
(2.5)

where  $a = {}_{s} = , D^{0} = 3(1 + )$  where is a known smallelectroweak correction, and, for N  $_{f} = 3$  in the  $\overline{MS}$  scheme,

$$D_{1} = 1$$

$$D_{2} = \frac{11}{2} \quad 4 \quad (3) \quad + \frac{C_{A}}{12} \quad \frac{C_{F}}{8} = 1:640$$

$$D_{3} = \frac{151}{18} \quad \frac{19}{3} \quad (3) \quad 4^{-2} + 2C_{A} \quad \frac{31}{6} \quad \frac{5}{3} ( (3) + (5))$$

$$+ 2C_{F} \quad \frac{29}{32} \quad \frac{19}{2} \quad (3) + 10 \quad (5) \quad + C_{A}^{2} \quad \frac{799}{288} \quad (3)$$

$$+ C_{A}C_{F} \quad \frac{827}{192} + \frac{11}{2} \quad (3) \quad + C_{F}^{2} \quad \frac{23}{32} \quad = 6:371; \quad (2.6)$$

where (3) = 1:20206 and (5) = 1:03693,  $C_A = N_C = 3$  and  $C_F = (N_C^2 - 1) = (2N_C) = 4=3$ . The quantity =  $(11C_A - 2N_f) = 12$  is the rst beta function coe cient [22]:

$${}^{2}\frac{\mathrm{da}\left({}^{2}\right)}{\mathrm{d}^{2}} = \mathrm{a}\left({}^{2}\right) ; \quad (\mathrm{a}) = \mathrm{a}^{2}\left(1 + {}^{0}\mathrm{a} + :::\right) \quad (2.7)$$

 $(=9=4, ^{0}=16=9 \text{ for } N_{f}=3).$ 

The expansion in eq. (2.5) de nes the Adler function at all complex s with a cut for s > 0. In the spacelike region, where s < 0, a (s) is real and given asymptotically by ( $^2 > 0$ ):

$$\frac{1}{a(s)} = \frac{1}{a(2)} + \log \frac{s}{2} = \log \frac{s}{2};$$
 (2.8)

If we want a (s) at some complex value of the argument, e.g. s = jsjexp(i), we can use the formula

$$a(s) = \frac{a(s)}{1 + a(s)i}$$
(2.9)

where the angle is on the upper tip of the cut for s real and positive, + on the lower tip and zero on the negative real axis. The more accurate two-loop expression is given by

$$a(s) = \frac{a(js)}{1 + a(js)ji + {}^{0}a(js)jlog(1 + a(js)ji)};$$
 (2.10)

The expansion for R (s) (for s real and positive) can be obtained from that of D by the relation T

R (s) = 
$$\frac{1}{2 \text{ i}} \int_{s^0 \neq s}^{1} \frac{ds^0}{s^0} D$$
 (s<sup>0</sup>): (2.11)

Perform ing the integration by using the expansion for D in eq. (2.5) and the expression in eq. (2.10) for a complex argument, one obtains

$$R (s) = D^{0} 1 + F_{1}a(s) + F_{2}a^{2}(s) + F_{3}a^{3}(s) + :::;$$
 (2.12)

and

$$F_1 = D_1; F_2 = D_2; F_3 = D_3 \frac{2}{3}:$$
 (2.13)

The origin of the  $^{2}$  <sup>2</sup>=3 term is easily understood. By using the one-loop expansion for a (s), eq. (2.9), one gets

$$R (s) = D^{0} 1 + \frac{1}{2 i} \log \frac{1 + i a(s)}{1 i a(s)} + \dots = D^{0} 1 + a(s) \frac{2^{2} a^{3}(s)}{3} + \dots$$
(2.14)

We have the following observations on this result. First, for  $N_f = 3$ , the coe cients  $F_{1;2;3}$  in the expansion of R (s) for decay coincide with those of  $R_{e^+e^-}$  because the potentially dimension of R (s) for decay coincide with those of  $R_{e^+e^-}$  because the potentially dimension of R (s) for decay coincide with Q<sub>1</sub> being the quark charges, vanish in this case. Second, we observe that eq. (2.13) is obtained by a truncation of higher-order terms in the quantity  $2a^2 2' 0.7$  with  $a = a (m^2)$ . Note that a similar problem of truncation arises when the integration over the circle in eq. (2.4) is performed. In the early treatments of this problem (e.g. in ref. [2]) the expression of a (s), which appears on the circle, is taken from eq. (2.10) and expanded in a consistently to the order  $a^3$ . W ith this procedure, one obtains

$$R^{(BNP)} = D^{0} 1 + H_{1}a(m^{2}) + H_{2}a^{2}(m^{2}) + H_{3}a^{3}(m^{2}) + :::;$$
(2.15)

with

$$H_1 = 1; H_2 = 5.2023; H_3 = 26.3666:$$
 (2.16)

W e will refer to this result as BNP form ula (for the authors of ref. [2]). M ore recently in ref. [4] it was advocated that a better procedure for performing the integration on the circle is to keep the full three-loop expression for a (  $jsje^{i}$ ), according to the form ula

$$R^{(LP)} = \frac{D^{0}}{2i} \int_{\substack{z \neq 1 \\ z \neq 1}}^{I} \frac{dz}{z} (1 \ z)^{3} (1+z)^{h} + D_{1}a(zm^{2}) + D_{2}a^{2}(zm^{2}) + D_{3}a^{3}(zm^{2}) + \dots$$
(2.17)

It is this procedure which is currently adopted (LeD iberder-Pich, or LP method). At xed experimental value of R, the two procedures lead to values of  $f_{\rm c}$  (m<sup>2</sup>) that dier by terms of order  $(2a^2)(2a^2)a^2 = 0.01$  or  $(2a^2)^2a^2$ 0.05. These  $\large$  $^2$  term s" always arise when one goes from the spacelike to the timelike region (e.g. sim ilar term s arise [22] when one relates D rell-Y an processes to electroproduction or fragm entation functions to structure functions). There have been m any discussions in the past on the opportunity of resumming these terms [23]. If there was a good argum ent to consider the expansion for D (s) in some respect superior to that for R (s) it could be worthwhile at low energies to keep the expression in eq. (2.14) in its resummed form rather than to expand in  ${}^{2}a^{2}$ . A glance at eqs. (2.6) shows 3, there are no explicit \large  $^2$  term s" in the coe cients D<sub>n</sub> of the that, for n expansion for D . Sim ilarly when the integration on the circle is performed with the complete formula for a  $(\dot{p}\dot{p}^{i})$  all terms are kept up to order  $a^{3}(2a^{2})^{n}$ , i.e. up to order  $a^3$  one expands in a but keeps  $2a^2 a^2$  unexpanded. Sure enough this sequel of term s exists in reality, so why not take them into account? However, the counterargum ent is that there are in perturbation theory terms involving<sup>2</sup> that arise from origins other than the spacelike-tim elike connection and, in any case, there are many terms of the same general magnitude (for example, the term proportional to  $^2$  in D<sub>3</sub>, eq. (2.6)), so that the advantage of keeping this particular class of term s is likely to be completely illusory.

One can further consider the scale dependence of the di erent procedures. One can write

$$R^{(BNP)} = D^{0} 1 + H_{1}a(^{2}) + H_{2}a^{2}(^{2}) + H_{3}a^{3}(^{2}) + \dots ; \qquad (2.18)$$

where

$$\text{H}_{1}(^{2}) = \text{H}_{1}$$

$$H_{2}^{2}(^{2}) = H_{2} \quad H_{1} \quad \log \frac{m^{2}}{2}$$

$$H_{3}^{2}(^{2}) = H_{3} \quad 2H_{2} \quad \log \frac{m^{2}}{2} + H_{1} \quad ^{2} \log^{2} \frac{m^{2}}{2} \quad ^{0} \log \frac{m^{2}}{2}^{\#}: \quad (2.19)$$

A nalogously, we can study the scale dependence in the LP method. In this case we have

$$R^{(LP)} = \frac{D^{0}}{2i} \int_{\dot{z}\neq 1}^{I} \frac{dz}{z} (1 z)^{3} (1+z)^{h} + D_{1}a(z^{2}) + D_{2}a^{2}(z^{2}) + D_{3}a^{3}(z^{2}) + \dots$$
(2.20)

where the  $D_n$  coe cients are related to the  $D_n$  as the  $H_n$  to the  $H_n$  in eqs. (2.19).

The results of the LP and BNP methods are shown in g.1, where, assuming a measured value of 3.6 for R, we show the corresponding determination of  $_{\rm s}$  (m<sup>2</sup>) as a function of the renormalization scale. For comparison, we also show (dashed curve) the determination obtained with a simplified LP method, in which we use the one-loop expression eq. (2.9) for a. For > 1 G eV there is less dependence if the resummed expressions are used. This stability is often taken as a possible indication that resumming is better.

In conclusion, we agree that the resummed formulae provide a less am biguous result with respect to a change of scale than the unresummed expression. However, it is true that a priori it is not possible to guarantee that a more accurate result is obtained in one way or the other. As a consequence the spread shown in g.1 for di erent choices of and of procedure is to be taken as a realam biguity. In particular the large discrepancy at 1 G eV is a genuine signal of trouble, especially in view of the fact that several proposed scale — xing procedures lead to sm all values of , e.g. m inim al sensitivity, BLM scheme etc. [24]. A sm all value of is also suggested by physical considerations, because the average hadronic m ass is well below m . So, on the one hand, one cannot sensibly reject the option of sm all values of . On the other hand, the corresponding value of a (m<sup>2</sup>) becomes very am biguous at sm all .

## 3. Renorm alons and Borel Transform ation

In the following discussion we rst study the properties of D (s) in itself, and only later we consider the integration over the circle. The problem that we now consider




Figure 1: The value of  $_{\rm s}$  (m  $^2$ ) obtained from R  $\,=\,3$  :6, as a function of the renorm alization scale  $\,$  .

is the well known fact that the series for D is divergent. Indeed one can identify sequences of diagram s, depicted in g. 2, called \renorm alon" term s [13-21], that provide the leading behaviour at large n for the n-th coe cient of the expansion for D. The renorm alon contribution is of the form :




Figure 2: Renorm alon diagram s for a two-point correlator.

$$P_n = C_k n \ln^k - [1 + O(1=n)]$$
 (n large): (3.1)

Note the n! behaviour which implies that the series is divergent. Here,  $C_k$  and  $_k = 0$  are not known for the real theory but only, to some extent, in the large-N<sub>f</sub> expansion [18-21], and the index k runs over a discrete set of values:

$$k = 1; 2; 3; ::: Ultraviolet (UV) Renormalons$$
  
 $k = +1(?); +2; +3; ::: Infrared (IR) Renormalons (3.2)$ 

In the above list we om it the contribution from instantons, which appear at rather large values of k. They have been computed in ref. [25] and shown to be small. The UV or IR renorm alons arise from the lim its of large or small virtuality, respectively, for the exchanged gluon (s). A shinted by the question mark, the k = +1 IR renorm alon is probably absent in perturbation theory, but the issue is not really settled [16-21]. The absence of this term is necessary for the consistency of the assumption that all non-perturbative e ects can be absorbed in the condensates. In the following we will assume that the k = 1 IR renorm alon is indeed absent. In view of the divergence of the perturbative expansion, one can possibly give a m caning to the quantity

$$d(a) = \frac{D}{D^{0}} \quad 1 = D_{1}a + D_{2}a^{2} + D_{3}a^{3} + \dots$$
(3.3)

by the Borel transform method [27]. One de nes the perturbative expansion of the Borel transform B (b) of d (a) by removing the n! factors:

$$B (b) = \sum_{n=0}^{M} D_{n+1} \frac{n^n}{n!} = D_1 + D_2 b + D_3 \frac{b^2}{2} + \dots$$
(3.4)

Then, form ally

$$d(a) = \int_{0}^{2} db e^{b=a} B(b)$$
(3.5)

in the sense that the expansion for B (b) reproduces the expansion for d(a) term by term . W hat is needed for d(a) to be well de ned is that the integral converges (this cannot be true at all s [13,17] because of the singularities of d(a) in the s plane, but this problem can be neglected in our context) and that B (b) has no singularities in the integration range. But, as already mentioned, the large-n expansion of B (b) leads to singularities on the real axis. In fact, at large n, B (b) is essentially given by a geometric series:

B (b) 
$$C_k^{X} n^{k} \frac{b^{n}}{k} C_k (k+1) 1 \frac{b^{k}}{k} + \text{less singular terms (3.6)}$$

so that it is singular at b = k = . Thus the UV renorm alons correspond to singularities at b = 1 = ; 2 = ;::: and the IR renorm alons at +2 = ;+3 = ;::: A sa consequence, the convergence radius of the expansion for B (b) near the b-origin is determined by the UV renorm alon at b = 1 = , independent of the existence of the IR renorm alon at b = +1 = .

Thus, the perturbative expansion for B (b) can be directly used only to perform the integration up to b = +1 = . The contribution from b = +1 = . up to b = 1, where the expansion is not valid, could typically lead to term s of order  $^{2}$ =s (or even worse). For example, if B (b) is su ciently well behaved at b = +1 = . and at b = 1,

$$d(a) = \int_{1=}^{2} db e^{b=a} B(b) \quad aB(1=) \exp(1=a) \quad aB(1=)^{2} = s; \quad (3.7)$$

where we used a <sup>1</sup> ' log (s= <sup>2</sup>) and the fact that the exponential cuts away all large-b contributions so that B (b) was approximated by its value near b = +1=. From a dimension point of view, at large n, the series for d (a) is dominated by the UV renormalon behaviour with D<sub>n</sub> n n!()<sup>n</sup>. At xed small a, the individual terms  $\mathfrak{P}_n \mathbf{j} \mathbf{a}^n$  rst decrease with n, then atten out and eventually increase because of the n! factor. The best estimate of the sum is obtained by stopping at the minimum, for n  $n_{opt}$ , given by  $\mathfrak{P}_n \mathbf{j} \mathbf{a}^n$   $\mathfrak{P}_{n-1} \mathbf{j} \mathbf{a}^{n-1}$ , or  $n_{opt}$  1= a. From the theory of asymptotic series [27], the corresponding uncertainty d (a) is of order  $\mathfrak{P}_{n_{opt}} \mathbf{j}^{n_{opt}}$ :

(where the Stirling approximation was used:  $n!' n^n e^{-n^p 2 n}$ ). Thus one could improve the accuracy of the perturbative expansion by computing more subleading term suntil  $n_{opt}$  is reached and then add a residual term of order <sup>2</sup>=s. Note that the estimate  $n_{opt}$  1= a 4 indicates a rather small value. However this estimate is obtained from the behaviour of the leading UV renormalon series, while there is no alternation of signs and in general no evidence of renormalon behaviour in the few known terms of the series.

W hile an accuracy of order <sup>2</sup>=s is what one gets in practice from the threeloop expression of d(a), it is true that, in principle, if there is no IR renormalon at b = +1=, d(a) can be better de ned. In fact, as the location of the leading UV renormalon at b = 1= is not in the integration range, there is the possibility of de ning B (b) by analytic continuation up to b = +2=. If this is realised then the remaining am biguity, of order ( $^{2}=s$ )<sup>2</sup>, is unavoidable because the corresponding singularity at b = +2= is on the realaxis, so that an arbitrary procedure to go around it must be de ned and the di erence between two such procedures would be of that order. However, since operators of dimension 4 do exist in the operator expansion, this am biguity can be reabsorbed in the non-perturbative condensate term s [13-15]. W e also understand that the absence of the IR renormalon at b = +1= is necessary for the consistency of the SVZ [6] approach because operators of dimension 2 are absent in this channel. If there would be a singularity at b = +1= the corresponding am biguity could not be absorbed in a condensate. There are indeed indications in perturbation theory that the st R renorm alon does not appear [19,21]. However there could be non-perturbative sources of breaking of the operator expansion at nonleading level. A fiter all no theory of connement could be built up from perturbation theory and renorm alons. But, in practice, independent of the existence of the IR renorm alon at b = +1=, the accuracy to be expected from the st three terms in the expansion for d(a), as they have been used so far in the actual determ ination of s, is of order <sup>2</sup>=s.

# 4. The Integration over the C ircle in the Borel Transform Form alism

In this section we show that the integration over the circle in eq. (2.4) for R is particularly simple in the Borel representation. Starting from eqs. (2.9) and (3.5) we have

$$r = \frac{R}{D^{0}} \quad 1 = \frac{1}{2i} \prod_{j=m^{2}}^{I} \frac{ds}{s} \quad 1 \quad \frac{s}{m^{2}} \prod_{j=m^{2}}^{j} \frac{ds}{s} \quad 1 \quad \frac{s}{m^{2}} \prod_{j=m^{2}}^{j} \frac{ds}{s} \prod_$$

We work in the approximation where the two-loop coe cient <sup>0</sup> in the beta function is neglected. Then, according to eq. (2.9), we can replace 1=a (s) by 1=a+i, where a is a ( $j_{2}$ ) = a ( $m^{2}$ ), invert the integration order and write  $s=m^{2} = exp(i)$ :

$$r = \int_{0}^{Z_{1}} db e^{b=a} B (b) \frac{1}{2i}^{Z} \quad id (1 + e^{i})^{3} (1 e^{i}) e^{ib} : \qquad (4.2)$$

The integration is easily perform ed, with the result

$$r = \int_{0}^{Z_{1}} db e^{b=a} B(b) \frac{12 \sin(b)}{b(b \ 1)(b \ 3)(b \ 4)}$$
$$= \int_{0}^{Z_{1}} db e^{b=a} B(b) F(b): \qquad (4.3)$$

We see that, in rst approximation, the e ect of going from a (s) to a (js) by integrating over the circle is to multiply the Borel transform B (b) by the factor F (b). For real x, the function F (x) is shown in g.3. It is an entire function in the whole complex plane with a good behaviour at in nity on the real axis. The factor F (x) has simple zeros at the location of all UV renorm alons and also of all IR renorm alons with




Figure 3: P bt of the function F(x).

the exception of those at b = 1 (if any), 3 and 4. Since, in general, the corresponding singularities are not simple poles, they are not eliminated, but their strength is attenuated (this point will be discussed in more detail in sect. 5.). Equation (4.3) obviously coincides with the LP approach [4] in the limit  $^{0} = ^{00} = 0.0$  ne can also repeat the procedure by expanding in a ( $^{2}$ ) instead of a (m $^{2}$ ), according to eq. (2.20). The resulting dependence is shown in g.1 together with the analogous results for the BNP [2] and the LP form ulae.

# 5. Large-N $_{\rm f}$ Evaluation of R enorm alons and their R esum – m ation

As well known, the typical renorm alon diagrams of QED and QCD can be evaluated in the large-N<sub>f</sub> lim it and their structure is simple in this lim it [18-21]. In the abelian case the large-N<sub>f</sub> lim it corresponds to the large-lim it, and this is believed to be true also in the non-abelian gauge theory in spite of the fact that the beta function in this case cannot be evaluated only in term s of vacuum polarisation diagram s. The sequence of dom inant term s in <sup>n</sup> generalises the term s in and <sup>2</sup> that appear in D'<sub>2</sub> and D'<sub>3</sub> respectively. As argued in a recent paper, ref. [20], the determ ination of the exact behaviour of the UV renorm alon series m ay be very di erent from the one indicated in the large-lim it. W hile we do not know the complete form of the leading UV renorm alon at b = 1=, we can nevertheless compute, for orientation, the quantitative in pact of its approximate form at large on the determ ination of the leading UV renorm alon at b = 1= to the Borel transform B (b):

$$B(b) = \frac{2}{9}e^{5=3} \left[7 + 2n\right](x)^{n} = \frac{2}{9}e^{5=3} 7 \frac{2x}{1+x} \frac{1}{1+x}; \quad (5.1)$$

where x = b and the factor  $e^{5=3}$  transforms the result from the MOM into the  $\overline{MS}$  scheme. We observe that the leading UV renormalon is a double pole. Since in the large-limit <sup>0</sup> can be neglected, the corresponding expression for R in the approximation of eq. (4.3) is appropriate, and it turns the double pole into a simple pole.

We now study the num erical e ect of including the whole renorm alon series with

respect to a truncated result up to the order  $b^2$ . We is consider the impact on d(a), i.e. before the integration on the circle. In order to get the correction to d(a) from the higher-order terms in the UV renorm alon we must subtract from B (b) its expansion up to 0 ( $b^2$ ) and perform the inverse B orel transform, eq. (3.5):

$$d(a) = \frac{1}{2} \int_{0}^{Z_{1}} dx e^{x=a} B(x) \frac{2}{9} e^{5=3} (7 \quad 9x + 11x^{2}) :$$
 (5.2)

For = 27=12 = 2.25 and a = 0.12 (or a = 0.27) one nds d 3.7 10<sup>-3</sup> which corresponds to a 2% increase in the value of  $_{s}$  (m<sup>2</sup>) at xed d(0.12) = 0.155 ( $_{s}$  (m<sup>2</sup>) 0.007).

W e now repeat the same exercise for the function r, given in eq. (4.3), obtained after integration over the circle. W e compute the variation

$$r(a) = \frac{1}{2} \int_{0}^{Z_{1}} dx e^{x=a} B(x) \frac{2}{9} e^{5=3} (7 \quad 9x + 11x^{2}) F(x): (5.3)$$

Numerically we nd r(0.12) 6.0  $10^{3}$  which, at xed r(0.12) = 0.220, again corresponds to  $_{s}$  (m<sup>2</sup>) 0.007. Thus the extra factor F (b) has practically no in uence on the e ect on  $_{s}$  (m<sup>2</sup>) of the nearest UV singularity.

In conclusion the overall e ect of the UV renorm alon singularity in this model is sm all and not much changed by the integration over the circle.

# 6. Search for M ore Convergent Approxim ants

A ssum ing that indeed there is no IR renorm alon at b = +1 = one can in principle try to obtain by analytic continuation a de nition of the Borel transform, valid on the positive real baxis up to b = +2 =, outside the radius of convergence of its expansion. We now discuss how the analytic continuation could be implemented in practice.

Starting from eq. (3.4) we can make a change of variable [14,17] z = z (b) with inverse b = b(z), z(0) = 0 and z(1) = 1 (so that the interval from 0 to 1 in b is mapped into the 0 to 1 range in z), such that the IR singularities are mapped onto the interval between  $z_0 = z(2=)$  and 1 and the UV singularities are pushed away at

 $jzj z_0$ . Changing variable one obtains

$$d(a) = \int_{0}^{Z_{1}} db e^{b=a} B(b) \int_{0}^{Z_{1}} dz \frac{db}{dz} e^{b(z)=a} B(b(z)):$$
(6.1)

U sing the expansion

$$b(z) = c_1 z + c_2 z^2 + \dots$$
 (6.2)

the series

B (b) = D<sub>1</sub> + D<sub>2</sub>b + D<sub>3</sub>
$$\frac{b^2}{2}$$
 + ::: (6.3)

goes into

$$B(p(z)) = D_{1} + D_{2}c_{1}z + (D_{2}c_{2} + D_{3}\frac{c_{1}^{2}}{2})z^{2} + \dots$$
(6.4)

which is convergent up to  $z = z_0$ , while the original between expansion was convergent only up to b = 1 =, corresponding to  $z(1 = ) < z_0$ . The improved approximation for d(a) is therefore given by

$$d(a) \quad ' \quad \sum_{z}^{Z_{20}} dz \quad \frac{db}{dz} = b(z)=a \quad D_{1} + D_{2}c_{1}z + (D_{2}c_{2} + D_{3}\frac{c_{1}^{2}}{2})z^{2} + \cdots + \frac{d}{2}$$

$$= \int_{0}^{Z_{20}} dbe^{b=a} \quad D_{1} + D_{2}c_{1}z(b) + (D_{2}c_{2} + D_{3}\frac{c_{1}^{2}}{2})z(b)^{2} + \cdots ; \quad (6.5)$$

where the full expression of z as function of b is inserted in the integral. In this way, an in nite sequence of terms is added to the bexpansion. For a small, the upper limit of integration can be replaced with in nity without signi cante ect.

O ne possible example is given by [17]:

n

$$z(b) = \frac{p}{p} \frac{1+b}{1+b+1} + b(z) = \frac{4z}{(1-z)^2}$$
: (6.6)

In this case the stUV singularity is at z = 1, and all higher UV renorm alons are on the unit circle jzj = 1. IR renorm alons are between  $z_0 = \begin{pmatrix} p \\ 3 \end{pmatrix} = \begin{pmatrix} p \\ 3 \end{pmatrix} = \begin{pmatrix} p \\ 3 \end{pmatrix} + 1$  and z = 1. In this example,  $c_1 = 4 = , c_2 = 8 = .0$  ther examples are

$$z(b) = \frac{b}{k+b}$$
 !  $b(z) = \frac{kz}{(1-z)}$ ; (6.7)

with k = 1;2 or 3. Also in these cases the set IR renormalon at b = 2 becomes the closest singularity to z = 0, while the UV are pushed further away. Here we have  $c_1 = c_2 = k = .$ 

Before discussing numerical applications, we observe that the present method relies simply on the position of the IR and UV renorm alon singularities in the Borel plane and not on the nature and the strength of the singularities. We have seen that the integration over the circle in eq. (4.3) does not change the position of the singularities in the b plane, but simply a lects their strength. Thus, we can as well consider the elect of the accelerators on the expansions for  $R^{(BNP)}$  given in eq. (2.18) or on the LP expression of eq. (2.20). For example, the improved version of eq. (4.3) simply becomes

$$r' \int_{0}^{Z_{1}} dbe^{b=a} D_{1} + D_{2}c_{1}z(b) + (D_{2}c_{2} + D_{3}\frac{c_{1}^{2}}{2})z(b)^{2} + \dots F(b):$$
(6.8)

We now consider the following numerical exercise. We assume that experiments have measured R = 3.6. We then compute  $a(m^2)$  with the LP formula as a function of the scale , and we perform the same calculation applying our acceleration procedures to the LP method [4]. The results are shown in g. 4. We see that relatively large dimensions in the tted value of  $s(m^2)$  are obtained, especially at large for dimensions and in comparison to the non-accelerated formulae. We do not see a priori compelling reasons to prefer one or the other procedure. The fact that a priori equivalent methods lead to results with a sizeable spread must be considered as an indication of a real ambiguity. Even if we only consider the method of ref. [4] for the integration over the circle, it is impossible to go below an uncertainty of the order  $s(m^2) = 0.050$  for in the range from 1 to 3 GeV. The ambiguity becomes even larger if we extend the comparison to the formulae with truncation in  $2a^2/2$  a  $\{la BNP (g. 5)$ .

# 7. Study of M ore Convergent Approxim ants in a M odel

The method for accelerating the convergence discussed in the previous section only relies on the position of the singularities of the Borel transform and not on their nature and strength. It looks rather surprising that one can compensate for the e ect of renorm alons without actually knowing their form in detail. In this section we study a simple mathematical model to clarify under which conditions the method can be successful, in the sense that it provides a better approximation to the true result.




Figure 4: E ect of the accelerators on the determ ination of  $_{s}$  (m<sup>2</sup>) with the LP m ethod, for R = 3:6. The curves b, c and d refer to the change of variable of eq. (6.7), while e refers to eq. (6.6).




Figure 5: As in Fig. 4, for the BNP method.

W e consider as a model the case where the Borel transform is exactly specied by

$$B_{\text{true}}(b) = 1 + D_{2}b + D_{3}\frac{b^{2}}{2} + \frac{x^{4}}{a} (b)^{n} + \frac{1}{a} (b)^{n}$$
$$= 1 + D_{2}b + D_{3}\frac{b^{2}}{2} + \frac{1}{(1+b)} + \frac{(+1)}{2} (b)^{2} (b)^{2} (7.1)$$

The added sum stands for the higher-order contribution that could arise from a leading UV renorm alon at b = 1 = w ith a degree of singularity specified by and a xed overall strength given by . We will take = 1 in the following discussion. It is convenient to re-express eq. (7.1) in term s of x = b:

$$B_{\text{true}}(\mathbf{x}) = 1 + \overline{D}_{2}\mathbf{x} + \overline{D}_{3}\frac{\mathbf{x}^{2}}{2} + \frac{1}{(1+\mathbf{x})} + 1 + \mathbf{x} + \frac{(+1)}{2}\mathbf{x}^{2}$$
(7.2)

where  $\overline{D}_2 = D_2 =$  and  $\overline{D}_3 = D_3 = 2^\circ$ . Similarly we can introduce  $B_{pert}(x)$  and  $B_{accel}(x)$ , the perturbative Borel functions without and with acceleration, respectively:

$$B_{pert}(x) = 1 + \overline{D}_2 x + \overline{D}_3 \frac{x^2}{2}$$
 (7.3)

$$B_{accel}(x) = 1 + \overline{D}_{2}\overline{c}_{1}z(x) + (\overline{D}_{2}\overline{c}_{2} + \overline{D}_{3}\frac{\overline{c}_{1}^{2}}{2})z(x)^{2}; \qquad (7.4)$$

where  $\overline{c}_{1;2} = c_{1;2}$ . In all cases the corresponding d function,  $d_{true}$ ,  $d_{pert}$  and  $d_{accel}$  is given by  $Z_1$ 

$$d(a) = \int_{0}^{2} dx e^{-x^{2} a} B(x):$$
 (7.5)

W e consider the ratio

$$H = \frac{d_{\text{true}}}{d_{\text{true}}} \frac{d_{\text{accel}}}{d_{\text{pert}}} = 1 \quad \overline{D}_2 I_2 (a) \quad \overline{D}_3 I_3 (a);$$
(7.6)

where the quantities  $I_{2,3}$  are given by

$$I_{2}(a) = \frac{1}{I_{0}(a)} \int_{0}^{Z_{1}} dx e^{x = a} \overline{c}_{1} z(x) + \overline{c}_{2} z(x)^{2} x$$
(7.7)

$$I_{3}(a) = \frac{1}{2} \frac{1}{I_{0}(a)} \int_{0}^{Z_{1}} dx e^{x=a} \overline{C}_{1}^{2} z(x)^{2} x^{2}$$
(7.8)

and

$$I_{0}(a) = \int_{0}^{Z_{1}} dx e^{x=a} \frac{1}{(1+x)} + x \frac{(+1)}{2} x^{2}$$
(7.9)

C learly,  $\frac{1}{H}$  j < 1 is the condition for the acceleration method to be successful. In particular for H = 0 d<sub>true</sub> and d<sub>accel</sub> coincide. For each value of a and , in a given model speci ed by z(x) and the corresponding coe cients  $\overline{q}_{i2}$ , the condition H = 0is satisfied on a straight line in the plane  $\overline{D}_2; \overline{D}_3$ , while the inequality  $\frac{1}{H}$  j < 1 is satisfied in a band defined by two straight lines parallel to the H = 0 line. In gs.6-8 we plot the lines H = 0 for a = 0.27 for for and z(x) given by eq. (6.6) (case labelled by 0), or by eq. (6.7) with k = 1 or 2 (cases 1 and 2). The values of in




Figure 6: Lines corresponding to H = 0 for the method  $0^{"}$  (solid),  $1^{"}$  (dashed) and  $2^{"}$  (dotted) for = 0.5. The circle corresponds to expansion of 1=(1 + x).

gs. 6-8 are = 0:5;1;2. We see that for each choice of z(x) the lines have dimension erent negative slopes. The lines tend to cross each other in a region of the plane not far from the point  $\overline{D}_2 = -$ ,  $\overline{D}_3 = -$  ( + 1), i.e. the values that correspond to the rst few terms of the expansion of the asymptotic function 1=(1 + x). The region where




Figure 7: As in Fig. 6, for = 1.




Figure 8: As in Fig. 6, for = 2.

the lines cross is more sharply de ned if is small, i.e. if the asymptotic function is not too singular. In gs. 9{10 we show the bands  $\frac{1}{10}$  j< 1 for = 0.5;1 in cases 0 and 1.




Figure 9: Bands corresponding to  $\frac{1}{2}$  j < 1 for the m ethod \0" (solid), \1" (dashed) for = 0.5.

The conclusion is that the m ethod for accelerating the convergence works well only if the coe cients  $\overline{D}_2; \overline{D}_3$  resemble those of the asymptotic series, in other words if the known terms in the expansion are su ciently representative of the asymptotic series. In particular we see that it is very unlikely to get an improvement if the coe cients  $\overline{D}_2$  and  $\overline{D}_3$  are of the same sign, as is unfortunately the case for the series of interest for us (see eq.6).

From a di erent point of view we now consider the simple function B (b) given by

B (b) = 
$$\frac{1}{(1+b)}$$
 (7.10)




Figure 10: As in Fig. 9, for = 1.

and we plot the relation of the exact result

$$d(a) = \int_{0}^{Z_{1}} dx e^{x = a} B(x)$$
(7.11)

with its accelerated or non-accelerated series approximants, as a function of a and of the order of the expansion. The results obtained for the accelerating function z (b) given in eq. (6.6) (the case labeled by 0) and = 1;2 are shown in gs. 11,12. We




Figure 11: E ect of the resummation technique described in the text, for a function with Borel transform B (b) = 1=(1 + b); = 1

see that when, as in this case, the coe cients of the expansion coincide with their asymptotic form the accelerated formulae provide a much better approximation to the true result, more so if the singularity is weaker (i.e. is smaller). The non-accelerated formulae are only good for small enough a. The physically interesting case of a 0.27 appears to be at the limit of the range where the non-accelerated formulae are acceptable.




Figure 12: E ect of the resummation technique described in the text, for a function with Borel transform B (b) = 1=(1 + b); = 2

### 8. Conclusion

The determ ination of  $_{s}$  (m<sup>2</sup>) or  $_{s}$  (m<sup>2</sup><sub>z</sub>) from decay is nom inally very precise and the experimental errors are extremely small at LEP. Certainly the dominant ambiguity is at present the theoretical error. The nom inal precision is large because in the massless limit no explicit 1=m<sup>2</sup> corrective terms are present in the operator expansion. But it has become clear by now that one cannot sensibly talk of powersuppressed corrections if the ambiguities in the leading term are not under control [28]. We think that there is no real theorem that prevents non-perturbative corrections in the coe cient function of the leading term in the operator expansion at the level of  $^{2}$ =m<sup>2</sup>. Equivalently, there could be an IR renorm alon singularity at b = +1=, which

would create an irreducible (being located on the integration path) am biguity of order  ${}^2 = m^2$ . In all-order perturbative evaluations of the singularity pattern in the B orel plane the IR singularity at b = +1 = is probably absent, a result consistent with the idea that all irreducible am biguities can be reabsorbed in condensates. Even if the IR renorm alon singularity at b = +1 = is indeed absent, the radius of convergence of the expansion is limited by the leading UV renorm alon singularity at b = -1 =. If this disease is not cured or cannot be cured the resulting am biguity is still of order

 $^{2}$ =m $^{2}$ . In principle the problem could be solved if the exact nature and strength of the singularity was known, by simply taking its e ect into account in the evaluation of  $_{\rm s}$  (m<sup>2</sup>), along the way indicated in section 5 in the case of the estimate of the singularity in the unrealistic limit of large . But the exact determ ination of the singularity appears to be beyond the scope of presently known methods. In the actual case where the UV renorm alon singularity at b = 1 = 1 is not specied, one can still try, in principle, to bypass the problem by a transform ation of variables that pushes the leading UV singularity to a larger distance from the origin than the st  $\mathbb{R}$  renorm alon singularity at b = +2 = . Expanding in the new variable is equivalent to add a speci ed in nite sequel of terms to the original expansion. The convergence of the series should be improved by these accelerators of convergence and the ambiguity decreased. We have studied the quantitative e ect of implementing a number of such accelerators with di erent choices of the renorm alisation scale . An indication of the size of the ambiguities on  $_{s}$  (m<sup>2</sup>) is obtained from the spread of the results for di erent starting form ulae (e.g. with a ( s) taken in the integration over the circle in its renorm alisation group improved form or in a xed order truncated

expansion), di erent accelerator functions and di erent choices of the renorm alisation scale. The di erence between the resum m ed or truncated expression for a (s) on the circle are expecially large at small values of , while the variations induced by the di erent accelerators are expecially pronounced at large values of . The relative stability of the unaccelareted result of ref. [4] versus changes of appears as largely accidental in that the accelerated form ulae based on it are much less stable at large . It was argued in ref. [19] that if one expands in  $_{\rm s}$  (<sup>2</sup>) instead of expanding in  $_{\rm s}$  (m  $^2$ ) the scale dependence of the UV renorm alon correction becomes of order  $(^{2}=m^{2})(m^{2}=^{2})^{2}$ . Can then one be safe if is chosen su ciently large? Clearly in the true result the sum of the perturbative terms plus the remainder must be scale independent. When is changed, the number of terms to be added before the series becom es asymptotic changes and must compensate for the di erence. The increased sensitivity of the accelerated formulae at large is not encouraging for invoking that large is safer. All together, from g.4 we nd it di cult to im agine that the theoretical error on the strong coupling can be taken smaller then, say,  $_{\rm s}$  (m<sup>2</sup>) 0.050 (which approximately corresponds to  $_{\rm s}$  (m<sup>2</sup>) 0:005).

The accelerator method is based on the mere knowledge of the position of the singularity and not on its precise form . C learly such a method can only work if the known terms of the expansion carry enough information on the asymptotic form of the series. We have quantitatively con med this statement by studying the problem on a simple mathematical model where the true result is known. The performance of di erent accelerators is studied as a function of the coe cients of the rst few term s. These results indicate that there is little hope of improving the ambiguity from the leading UV renorm alon because the stafew coe cients of the actual expansion show no evidence for the asymptotic behaviour, in particular no sign alternance. This last argument (as well as the one on the dependence of the UV renorm alon) can be interpreted in di erent ways. If one is a great optim ist, he can argue that the series does not resemble at all to the renorm alon asymptotics, hence the norm alisation of the renorm alon term is very small (as is the case for the explicit form of the singularity obtained in the large N  $_{\rm f}$  lim it). Or, if one is more cautious, as one should be in estimating errors, he can say that since the known terms do not show sign of asymptotia, they are dominated by subasymptotic e ects and cannot be used to estim ate the rem ainder. In this spirit we do not propose the accelerators as a better way to determ ine the true result but simply as a criterium to evaluate the theoretical

error. In fact, while all accelerators tend to increase the resulting value of  $_{\rm s}$ , the amount of the upward shift is sizeably di erent for di erent accelerators.

In order to by pass all possible objections one should be able to t at the same time  $_{\rm s}$  (m  $^2$ ) and C  $_2$ , the coe cient of 1=m  $^2$  corrections to R . Note that in the A LEPH m om ent analysis [5]C<sub>2</sub> is xed to zero while the coe cients of som e higher-dim ension operators are tted. This is not very relevant to the main issue. If  $C_2$  is not xed it is found that the sensitivity to  $_{\rm s}$  (m  $^2$ ) is much reduced. In an interesting paper N arison [29] attempted to put an upper bound on  $C_2$  from the data on  $e^+e_-$ ! hadrons. This is an important issue that would deserve further study. Our interpretation of the analysis of ref. [29] is that values of  $C_2$  of order (500 M eV)<sup>2</sup> are not at all excluded. Narison [29] derives a more stringent lim it  $jc_2 j < (374 \text{ MeV})^2$  but we feel he relies too much on the so called optim isation procedure. Indeed, som ething that should be a constant in a dummy variable turns out to be a steep parabola. The value at the tip is taken, with a small error, as the best estimate because of the vanishing of the derivative at that point, instead of considering the span of the results in a priori reasonable range for the irrelevant parameter. In a recent paper [30] an estim ate of C<sub>2</sub> from Argus data on hadronic decay was obtained and the results are compatible with  $jC_2 j < (500 \text{ M eV})^2$ .

We ignored here other possible sources of error beyond those arising from higher orders in perturbation theory. These include errors from the freezing mechanism for s, errors from the translation of s (m<sup>2</sup>) in term s of s (m<sup>2</sup><sub>z</sub>), from the region of the circle integration near the positive real axis and so on. These errors are presumably smaller [9] than our current estimate of the error from higher order terms in the perturbative espansion. Taking all the other uncertainties into account we end up with a total theoretical error around s (m<sup>2</sup><sub>z</sub>) 0:006. As a result, in spite of the fact that our estimate of the error is larger than usually quoted, the determ ination of s (m<sup>2</sup><sub>z</sub>) from remains one of the best determ inations of the strong coupling constant.

# References

- [1] K.Schilcher and M.D.Tran, Phys. Rev. D 29 (1984)570.
- [2] E.Braaten, Phys. Rev. Lett. 60 (1988)1606;
   S.Narison and A.Pich, Phys. Lett. B 211 (1988)183;

E.Braaten, Phys. Rev. D 39 (1989)1458;

E.Braaten, S.Narison and A.Pich, Nucl. Phys. B 373 (1992)581.

- [3] M.Luo and W.J.Marciano, preprint BNL-47187 (1992).
- [4] F.LeD iberder and A.Pich, Phys. Lett. B 286 (1992)147.
- [5] LDu ot, Proceedings of the 3rd W orkshop on Tau Lepton Physics, M ontreux, 1994.
- [6] M A.Shifman, A.J.Vainshtein and V.J.Zakharov, Nucl. Phys. B 147 (1979)385, Nucl. Phys. B 147 (1979)448, Nucl. Phys. B 147 (1979)519.
- [7] J.Pumplin, Phys. Rev. D 41 (1990)900.
- [8] G.Altarelli, in \QCD-Twenty Years Later", Aachen 1992, eds. P.Zerwas and H A.Kastrup, World Scientic, Singapore (1992).
- [9] G A ltarelli, Proceedings of the 3rd W orkshop on Tau Lepton Physics, M ontreux, 1994.
- [10] T N. Truong, Preprint Ecole Polytechnique, EP-CPth A 266.1093 (1993); Phys. Rev. D 47 (1993)3999.
- [11] S.Narison, Proceedings of the 3rd W orkshop on Tau Lepton Physics, M ontreux, 1994.
- S.G. Gorishny, A.L. Kataev and S.A. Larin, Phys. Lett. B 259 (1991)144;
   L.R. Surguladze and M.A. Samuel, Phys. Rev. Lett. 66 (1991)560.
- [13] G.'tHooft, in \TheW hysofSubnuclearPhysics", Erice 1977, ed. by A Zichichi, Plenum, New York.
- [14] B. Lautnup, Phys. Lett. B 69 (1977)109;
   G. Parisi, Phys. Lett. B 76 (1978)65, Nucl. Phys. B 150 (1979)163.
- [15] F D avid, Nucl. Phys. B 234 (1984)237;A H. M ueller, Nucl. Phys. B 250 (1985)327.

- [16] G B.W est, Phys. Rev. Lett. 67 (1991)1388,67913732 (E);
   L S.Brown and L G.Ya e, Phys. Rev. D 45 (1992)398;
   L S.Brown, L G.Ya e and C.Zhai, Phys. Rev. D 46 (1992)4712.
- [17] A.H.Mueller, in \QCD-Twenty Years Later", Aachen 1992, eds. P Zerwas and H.A.Kastrup, World Scientic, Singapore.
- [18] V.J.Zakharov, Nucl. Phys. B 385 (1992)452.
- [19] M.Beneke and V.J.Zakharov, Phys. Rev. Lett. 69 (1992)2472;
   M.Beneke, Nucl. Phys. B 385 (1992)452.
- [20] A J.Vainshtein and V J.Zakharov, University of Minnesota preprint TPI-M INN 94/9T (1994).
- [21] C.N.Lovett-Tumer and C.J.M axwell, University of Durham Preprint DPT/94/58.
- [22] See, for example, G. Altarelli, Phys. Rep.81 (1982)1.
- [23] M.R.Pennington and G.G.Ross, Phys. Lett. B 102 (198)167;
  G.Parisi, Phys. Lett. B 90 (1980)295;
  G.Curci and M.Greco, Phys. Lett. B 92 (1980)175;
  A.P.Contogourisetal, Phys. Rev. D 25 (1982)1280; Phys. Rev. D 28 (1983)1644;
  Int. J.M od. Phys. A 5 (1990)1951.
- [24] P.M. Stevenson, Phys. Rev. D 23 (1981)2916;
  G.G nunberg, Phys. Lett. B 221 (1980)70; Phys. Rev. D 29 (1984)2315;
  S.Brodsky, G.P. Lepage and P.B. Mackenzie, Phys. Rev. D 28 (1983)228;
  H.J. Lu, Phys. Rev. D 45 (1992)1217;
  M. Neubert, preprint CERN-TH .7487/94;
  A.L.K ataev and V.V. Starshenko, preprint CERN-TH .7198/94; CERN-TH .7400/95;
  P.A.Raczka and A. Szym acha, W arsaw University preprint IFT /13/94, hep-ph 9412236.
- [25] P.Nason and M.Porrati, Nucl. Phys. B 421 (1994)518;
   I.I.Balitsky, M.Beneke and V.M.Braun, Phys. Lett. B 318 (1993)371;
   P.Nason and M.Palassini, CERN-TH.7483/94, hep-ph/9411246.
- [26] O.V. Tarasov, A.A. Vladim irov and A.Yu. Zharkov, Phys. Lett. B 93 (1980)429.

- [27] GN.Hardy, \D ivergent Series", Oxford University Press, 1949.
- [28] A.H.Mueller, Phys. Lett. B 308 (1993)355.
- [29] S.Narison, Phys. Lett. B 300 (1993)293.
- [30] CA.Dom inguez, University of Cape Town Preprint, UCT-TP-221/94.