

A Model Independent Approach to Future Solar Neutrino Experiments?

S.M. Bilenky^{(a,b)Y} and C. Giunti^{(a)Z}

^(a) INFN, Sezione di Torino, and Dipartimento di Fisica Teorica,
Università di Torino, Via P. Giuria 1, I-10125 Torino, Italy

^(b) Joint Institute for Nuclear Research, Dubna, Russia

Abstract

It is shown that from the data of future solar neutrino experiments (SNO and Super-Kamiokande), in which high-energy ^8B neutrinos will be detected, it will be possible in a model independent way: 1) To reveal the presence of sterile neutrinos in the flux of solar neutrinos on the earth; 2) To obtain the initial flux of ^8B ν_e 's; 3) To determine the probability of ν_e 's to survive as a function of neutrino energy.

[?] Talk presented by S.M. Bilenky at the Conference Beyond the Standard Model IV, Lake Tahoe, California, December 13-18, 1994.

^Y E-mail address: BILENKY@TO.INFN.IT

^Z E-mail address: GIUNTI@TO.INFN.IT

Solar neutrino experiments are a powerful tool for the investigation of the problem of neutrino mixing and for the search of new physics beyond the Standard Model. It is well known that from the analysis of the existing data follow some indications in favor of non-zero neutrino masses and mixing [1]. This analysis is based, however, on the assumption that the Standard Solar Model [2] (SSM) correctly predicts the solar neutrino fluxes from the different reactions of the solar thermonuclear cycles.

In this report we will consider the SNO [3] and Super-Kamiokande [4] (S-K) experiments, scheduled to start in 1995-96. We will show that these experiments will allow to investigate neutrino mixing in a model independent way and to determine the initial ^8B neutrino flux directly from the data.

In the SNO experiment solar neutrinos will be detected through the observation of the CC, NC and CC+NC elastic scattering (ES) processes:

$$\nu_e + d \rightarrow e + p + p; \quad (1)$$

$$\bar{\nu}_e + d \rightarrow e^+ + p + n; \quad (2)$$

$$\nu_e + e \rightarrow \nu_e + e; \quad (3)$$

In the S-K experiment solar neutrinos will be detected through the observation of the process (3), with an event rate that is expected to be about 50 times larger than in the current Kamiokande experiment. Due to the high energy thresholds, only high energy ^8B neutrinos will be detected in both experiments. The initial flux of ^8B ν_e 's is given by

$$\Phi_e^0(E) = X(E); \quad (4)$$

where $X(E)$ is a known function (determined by the phase space factor of the decay $^8\text{B} \rightarrow ^8\text{Be} + e^+ + \nu_e$) and Φ is the total flux.

As it is well known (see, for example, Ref.[5]), a discovery of transitions of active neutrinos into sterile states would be a discovery of new physics beyond the Standard Model. We will show first that the SNO and S-K experiments could allow to reveal the presence of such transitions in a model independent way.

Let us consider the NC process (2). For the average total probability of transitions of $\bar{\nu}_e$ into all active neutrinos we have

$$\sum_{\nu_e}^* P_{\bar{\nu}_e \rightarrow \nu_e} = \frac{N^{NC}}{X_d}; \quad (5)$$

where N^{NC} is the NC event rate and

$$X_d = \int_{E_{th}^{NC}}^Z d(E) X(E) dE \approx 4.72 \cdot 10^{43} \text{ cm}^2; \quad (6)$$

In the general case of transitions of $\bar{\nu}_e$'s into active as well as into sterile neutrinos

$$\sum_{\nu_e}^* P_{\bar{\nu}_e \rightarrow \nu_e}(E) = 1 - P_{\bar{\nu}_e \rightarrow \nu_s}(E); \quad (7)$$

where $P_{e \rightarrow s}(E)$ is the total probability of transition of e 's into all possible sterile states. It is clear from Eqs.(5) and (7) that from the measurement of N^{NC} it is impossible to reach any conclusions about $e \rightarrow s$ transitions without an assumption on the value of the total flux.

Let us take into account, however, that in the future SNO and S-K experiments solar neutrinos will be detected also through the observation of CC and CC+NC reactions. We have

$$\sum_{e \rightarrow s} P_{e \rightarrow s} = \frac{N^{ES}}{N^{NC}} : \quad (8)$$

Here

$$N^{ES} = \int_{E_{th}^{ES}}^Z \Phi_e(E) X_e(E) dE \quad 2.08 \cdot 10^{45} \text{ cm}^2 dE \quad (9)$$

and

$$N^{NC} = \int_{E_{th}^{NC}}^Z \Phi_e(E) X_e(E) dE ; \quad (10)$$

where N^{ES} is the ES event rate and $\Phi_e(E)$ is the flux of e on the earth, which will be determined in the SNO experiment from the measurement of the electron spectrum in the CC reaction (1). From Eqs.(5) and (8) we obtain the following lower bounds:

$$hP_{e \rightarrow s} \geq \frac{N^{ES}}{N^{NC}} ; \quad (11)$$

$$hP_{e \rightarrow s} \geq \frac{N^{ES}}{N^{NC}} ; \quad (12)$$

where

$$R_{NC}^{ES} = \frac{N^{ES}}{N^{NC}} : \quad (13)$$

is a measurable quantity (that does not depend on θ). Thus, if $R_{NC}^{ES} \neq 1$ it will mean that sterile neutrinos exist and Eq.(11) or Eq.(12) will give a lower bound for the average $e \rightarrow s$ transition probability. Let us notice that in the case $R_{NC}^{ES} = 1$ no conclusion about sterile neutrinos can be reached.

Several relations analogous to (11) and (12), which will allow to test in a model independent way whether there are transitions of solar e 's into sterile states, were derived in Ref.[6]. If all of them will not reveal the presence of $e \rightarrow s$ transitions, it will be natural to assume that $P_{e \rightarrow s}(E) = 1$. In this case the total ^8B neutrino flux can be determined directly from the experimental data. From Eqs.(5) and (8) we obtain

$$\frac{N^{NC}}{X_d} \quad \text{and} \quad \frac{N^{ES}}{X_e} : \quad (14)$$

A comparison of the flux obtained from Eq.(14) with the ^8B neutrino flux predicted by the SSM will be an important test of the model.

The SNO and S-K experiments will allow to determine the ν_e survival probability directly from measurable quantities:

$$P_{\nu_e \rightarrow \nu_e}(E) = \frac{\Phi_{\nu_e}(E)}{\Phi_{\nu_e}^{\text{total}}(E)} ; \quad (15)$$

$$hP_{\nu_e \rightarrow \nu_e}^{\text{ES}} = \frac{1}{\Phi_{\nu_e}^{\text{total}} \Phi_{\nu_e}^{\text{ES}}} \frac{N^{\text{ES}}}{N^{\text{total}}} \Phi_{\nu_e}^{\text{total}} :$$

Here $\Phi_{\nu_e}(E)$ is the flux of solar ν_e 's on the earth that will be measured by SNO and the total flux is given by Eq.(14).

In conclusion, we have shown that future solar neutrino experiments will allow to obtain model independent informations about neutrino mixing and the initial solar ν_e flux. These experiments have a good potential for revealing new physics beyond the Standard Model.

References

- [1] GALLEX Coll., Phys. Lett. B 285, 390 (1992).
- [2] J.N. Bahcall, Neutrino Physics and Astrophysics, Cambridge University Press, 1989; J.N. Bahcall and M.H. Pinsonneault, Rev. Mod. Phys. 64, 885 (1992); S. Turck-Chieze et al., Phys. Rep. 230, 57 (1993).
- [3] The Sudbury Neutrino Observatory Collaboration, Phys. Lett. B 194 (1987) 321; H.H. Chen, Nucl. Instr. Meth. A 264 (1988) 48.
- [4] A. Suzuki, Proc. of the Workshop on Elementary Particle Picture of the Universe, KEK, Japan, 1987; C.B. Bratton et al., Proposal to Participate in the Super-Kamiokande Project, Dec. 1992.
- [5] S.M. Bilenky and S.T. Petcov, Rev. Mod. Phys. 59, 671 (1987).
- [6] S.M. Bilenky and C. Giunti, DFTT 63/94 (hep-ph@xxx.janlgov/9411373), to be published in the Proc. of The First Arctic Workshop on Future Physics and Accelerators, Saariselka, Finland, August 1994.