In the m inim al supersymmetric standard model, the three gauge couplings appear to unify at a mass scale near 2 10^{16} GeV.We investigate the possibility that intermediate scale particle thresholds modify the running couplings so as to increase the unication scale. By requiring consistency of this scenario, we derive some constraints on the particle content and locations of the intermediate thresholds. There are remarkably few acceptable solutions with a single cleanly de ned intermediate scale far below the unication scale. UM -TH -95-01 UF IFT +HEP -95-1

Raising the unication scale in supersymmetry

Stephen P.Martin

Randall Physics Laboratory, University of Michigan, Ann Arbor M I 48109

and

Pierre Ramond

Institute for Fundam ental Theory Departm ent of Physics, University of Florida Gainesville FL 32611

ABSTRACT: In the minimal supersymmetric standard model, the three gauge couplings appear to unify at a mass scale near 2 10^{16} GeV.We investigate the possibility that intermediate scale particle thresholds modify the running couplings so as to increase the unication scale. By requiring consistency of this scenario, we derive some constraints on the particle content and locations of the intermediate thresholds. There are remarkably few acceptable solutions with a single cleanly de ned intermediate scale far below the unication scale.

1. Introduction

Data from LEP suggests that with N = 1 supersymmetry[1] at low energy (1 TeV), the three gauge couplings of the standard model converge to unify[2] at one scale M_X 2 10^{16} GeV. This apparent unication is predicated on two assumptions. One is that the weak hypercharge coupling is normalized to its unication into a higher rank Lie group, such as SU (5), SO (10) or E₆. The second is the absence of intermediate thresholds between 1 TeV and M_X. This apparent unication of couplings may be regarded as a \prediction" of the low energy value of sin² _W given the measured value of the strong coupling constant, and is a tantalizing hint of a unifying structure, such as superstring theory or a supersymmetric G rand Unied Theory.

W hile it is clear that the three gauge couplings have a much better chance to unify with low energy supersymmetry than without, it may be premature to unequivocably announce their unication, and this simple picture may have to be modied. The main reasons are the large experimental uncertainties in the value of the QCD coupling constant and ignorance of the detailed structure of the supersymmetric thresholds.

Thus it may be that the gauge couplings do not exactly unify at M $_X$. In that case, we may want to alter this simple picture by adding at least one intermediate threshold between the SUSY scale and the \uni cation" scale at M $_X$. The question of interest is whether the couplings can then be made to unify at a larger scale after introduction of the new intermediate threshold (s), caused by particles with vector-like electroweak quantum numbers. These modify the running of the gauge couplings above the intermediate thresholds to achieve true uni cation at the scale M $_U$, which we take to be larger than M $_X$. By requiring consistency of this scenario, we can derive constraints on the particles at the intermediate thresholds and relations between the intermediate scales M $_X$ and M $_U$.

There are several reasons to pursue this line of inquiry. One is that interm ediate m ass scales appear in m any extensions of the m inim al supersymmetric standard m odel (M SSM), such as those which incorporate a light invisible axion [3] or m assive neutrinos through the see-saw mechanism [4]. A nother is to explain the near zero values of m any of the Yukawa matrix elements through mixing the known particles with vector-like particles. These particles m ay appear at interm ediate thresholds.

Our prim ary motivation, however, is superstring theory which indicates that the uni-

cation energy should be more than one order of magnitude above M_X. The elective low energy theories generated by superstrings contain, in addition to the three chiral families, many vector-like particles, incomplete remnants of 27 and $\overline{27}$ representations of E₆. These vector-like particles have electroweak singlet masses, assumed to be, in the absence of any special mechanism, of the order of the highest scale around, in this case the P lanck mass. However, these theories have a larger invariance group than that of the MSSM, and must develop intermediate thresholds below the string scale to break the invariance group to that of the MSSM. This is typically achieved by at directions in the potential.

If the true scale of gauge coupling uni cation is higher than the apparent uni cation scale because of interm ediate scale thresholds as assumed here, one may view the \success" of gauge coupling uni cation as just an accident. We are implicitly taking the point of view that it is not completely accidental, and that it is still possible to understand gauge coupling uni cation through calculable perturbative means. We therefore assume that the three gauge couplings remain perturbative up to the uni cation scale M_U, and that the reason behind the raising of the uni cation scale is not some artifact of e.g. stringy threshold elects, but is really due to the presence of intermediate scale thresholds. We also assume that the normalization of weak hypercharge is indeed the standard one appropriate for uni cation with SU (2)_L and SU (3)^c into a simple gauge group. (Ref. [5] explores the possibility of di erent normalizations of the hypercharge as a means of raising the uni cation scale.)

This paper is organized as follows. In section 2 we develop the form alism for uni cation of couplings with one interm ediate scale threshold, and then for several interm ediate thresholds. In section 3 we discuss the e ects of various possibilities for the new particles at the interm ediate scale(s), including both new chiral super elds and new gauge vector super elds. In section 4 we discuss the results for one interm ediate scale with raised uni cation. Here we not tight constraints on the particle content and location of the interm ediate scale. Section 5 deals with results for more than one interm ediate scale, using as an example a particular three-fam ily superstring model.

2.0 ne-Loop Equations W ith New Thresholds

Let us begin by recalling som e salient facts about the running of the gauge couplings.

Since we will be comparing the running of gauge couplings with an interm ediate scale to the \tem plate" case of the MSSM, it will be su cient to use one-loop renormalization group equations only. The three gauge couplings run with scale according to

$$i^{1}(t) = i^{1}(t_{X}) + \frac{b_{i}}{2}(t - t_{X});$$
 (2:1)

where

$$_{i}(t) = \frac{g_{i}^{2}(t)}{4}$$
;

are the couplings for the three gauge groups, i = 1;2;3 for U $(1)_Y$, SU $(2)_L$, and SU $(3)^c$, respectively. The scale is given by

$$t = \ln (=_0);$$

where 0 is an arbitrary reference energy, and

$$t_X = \ln (M_X = 0)$$
;

is the uni cation scale. For N = 1 supersymmetry we have

$$b_{i} = 3c_{adjoint} \qquad c_{r}; \qquad (2.2)$$

where the c_r 's are the Dynkin indices of the representations, and the sum is over the left-handed chiralmultiplets. The hypercharge is normalized so that

$$b_1 = \frac{3}{20} X_r Y_r^2;$$

corresponding to the electric charge

$$Q = I_3 + \frac{Y}{2} :$$

For the three fam ilies and two H iggs doublets of chiral super elds in the M inim al Supersym m etric Standard M odel (M SSM), we have

$$b_1 = \frac{33}{5}$$
; $b_2 = 1$; $b_3 = 3$:

We start with the trajectories for $_1$ and $_2$ since their values at low energies are known with the greatest accuracy. We denet_X as the scale at which these two appear to meet in the MSSM:

$$x^{1}$$
 1^{1} (t_X) = 2^{1} (t_X) :

The extrapolated data, with N = 1 supersymmetry around 1 TeV, show that $_X^{1}$ 24:5, with M $_X^{2}$ 10¹⁶ G eV. However we do not assume precisely the same value for $_3(t_X)$ at that scale, since we are assuming that the \uni cation" at M $_X$ is only apparent; rather we set

$$_{\rm X}^{1} = _{3}^{1} (t_{\rm X}) + ;$$

introducing the parameter which parameterizes our ignorance about $_{3}$ (M $_{Z}$), our ignorance about the precise location of the SUSY thresholds, and our negligence of two-loop e ects. The present uncertainties indicate that

using the most conservative estimate. We contrast this situation by noting that without low energy supersymmetry, the same parameters have the values $_X^{1}$ 42, M $_X$ 10¹³ GeV, and 5.

Case of One Interm ediate Threshold

A ssum e rst only one interm ediate threshold above the supersymmetric thresholds, at the scale

$$t_{I} = \ln (M_{I} = 0); \quad t_{I} < t_{X}:$$

The previous equations are still valid as long as we are below the interm ediate threshold, that is

for t_I . At the interm ediate threshold $t = t_I$, new vector-like particles with electroweak singlet m asses at M_I, alter the b_i coe cients to new values

$$b_i! b_i i; i = 1;2;3;$$

with all $_{i}$ positive as long as the matter is made up of chiral super elds. We assume that their elect is to push the true unication scale to the new value t_{U} with $t_{U} > t_{X}$. Thus, above the intermediate threshold, all three gauge couplings must satisfy

$$_{i}^{1}(t) = _{U}^{1} + \frac{1}{2} (b_{i} _{i})(t t_{U}); t_{I} t t_{U};$$

where there is only one coupling at uni cation, $_{\rm II}$.

We have thus two ways of writing the equations for the gauge couplings below the interm ediate threshold; one is given by (2.4), the other by

$$_{i}^{1}(t) = _{U}^{1} + \frac{b_{i}}{2}(t t_{I}) + \frac{1}{2}(b_{i} i)(t_{I} t_{U}); \quad i = 1;2;3:$$
 (2:5)

C om parison of the two yields the three consistency equations

$$b_{i} \quad i \quad \frac{t_{U} \quad t_{I}}{t_{U} \quad t_{X}} = \frac{2}{t_{U} \quad t_{X}} \quad U^{1} \quad x^{1} \quad i = 1;2;$$

$$b_{3} \quad 3 \quad \frac{t_{U} \quad t_{I}}{t_{U} \quad t_{X}} = \frac{2}{t_{U} \quad t_{X}} \quad U^{1} \quad x^{1} + :$$

$$(2:6)$$

By subtracting the rst two, we obtain the constraint

$$\frac{28}{5} (2 1) \frac{t_{U} t_{I}}{t_{U} t_{X}} = 0; \qquad (2:7)$$

which indicates that $_{2}$ 1 must be positive. The di erence between the second and the third equations in (2.6) yields

$$4 \quad \frac{2}{t_{U} \quad t_{X}} \quad (3 \quad 2) \quad \frac{t_{U} \quad t_{I}}{t_{U} \quad t_{X}} = 0 : \qquad (2.8)$$

The remaining equation yields the value of the gauge coupling at uni cation

$$U^{1} = X^{1} \frac{1}{2} [2(t_{U} t_{I}) + t_{U} t_{X}] :$$
 (2:9)

W ith only non-exotic matter at the intermediate threshold, the combinations

$$q_{32}$$
 and $\frac{2}{5}r_{21}$;

are integers. Then (2.7) and (2.8) can be rewritten as

$$\frac{r}{14} = \frac{t_U + t_X}{t_U + t_I};$$
(2:10)

and

$$\frac{q}{4} = \frac{t_{\rm U} + t_{\rm X}}{t_{\rm U} + t_{\rm I}} = 2$$
(2:11)

It may be protable to consider an elementary geometric derivation of (2.10) and (2.11). Consider the evolution of two inverse gauge couplings, which meet at a scale t_X , and assume that they both change directions at a lower scale t_I , to meet at the larger scale t_U , as shown in Figure 1.

Figure 1

The ratios of the slopes of the lines above $t_{\rm I}$ satisfy, for ${\ }_1{\ }^1$ and ${\ }_2{\ }^1$

$$\frac{b_2}{b_2^0} \frac{b_1}{b_1^0} = \frac{OB}{OB^0} = \frac{t_X}{t_U} \frac{t_I}{t_I} ; \qquad (2:12)$$

from which (2.10) follows. We can apply the same technique to the evolution of $_2$ and $_3$ (including the nearm iss at M $_X$ parametrized by) to obtain (2.11).

W em ay think of the interm ediate threshold as a \lens" which refocuses the lines ${}_{1}^{1}$ (t), ${}_{2}^{1}$ (t), and ${}_{3}^{1}$ (t) so that they meet at t_{U} rather than t_{X} . If q > 4, the interm ediate threshold acts as a divergent lens, and the two lines for ${}_{2}^{1}$ and ${}_{3}^{1}$ never intersect. If q = 4, the same two lines are parallel and again never meet. Thus we must have q < 4 for the two curves to intersect beyond t_{I} . In addition, q cannot be negative or would be too large. This is easy to understand, since q < 0 corresponds to a strongly focusing lens which would make ${}_{2}$ and ${}_{3}$ meet at a lower scale than they would in the M SSM. To avoid having ${}_{2}$ and ${}_{3}$ meet prematurely, would have to be large and positive when q < 0. To see this, note that we can write

$$= \frac{1}{2} [4 (t_U \quad t_X) \quad q(t_U \quad t_I)]: \qquad (2:13)$$

So, for instance if q = 1, we not that even in the case of small hierarchies M_X = M_I = 10 and M_U = M_X = 10, one has = 2.2, which corresponds to a larger error than the experimental uncertainties on ₃ warrant. For more substantial hierarchies, or for more negative values of q, the situation becomes rapidly even worse. Thus it is su cient to consider only the four cases, q = 0;1;2;3. Sim ilarly, from (2.10) we not that if r 14, the $\frac{1}{1}$ and $\frac{2}{2}^{1}$ lines will never meet, while if r < 0, they will meet prematurely, implying a lowered scale of unication. If r = 0, the unication scale is not raised and M_U = M_X. Thus we have 0 < r < 14.

The scale of true uni cation can be extracted from (2.11) and (2.10) in term s of M $_X$, M $_I$, and the parameters q; and r respectively:

$$M_{U} = M_{X} \frac{M_{X}}{M_{I}} e^{2} = (4 \ q)$$
 (2:14)

$$M_{\rm U} = M_{\rm X} - \frac{M_{\rm X}}{M_{\rm I}} \stackrel{\rm r=(14 \ r)}{=} :$$
 (2:15)

Taken together, these im ply

$$q = \frac{2}{7}r + \frac{2}{t_U + t_I}$$
; (2:16)

or equivalently

$$= \frac{1}{r} (t_U t_I) \frac{r}{7} \frac{q}{2} :$$
 (2:17)

If > 0, then r m ust be a positive integer in the range $\frac{7}{2}q < r < 14$. On the other hand if is negative, we have $0 < r < \frac{7q}{2}$. The special case = 0 yields a non-trivial result only when 2r = 7q. In that case, for non-exotic m atter, the only solution is q = 2; r = 7, and from (2.14) or (2.15), M_X is the geometric mean between M_I and M_U. This corresponds to the seem ingly perverse case of the gauge couplings unifying both with and without the intermediate threshold! For non-zero , the hierarchies of scales are summarized by the two equations

$$\frac{M_{X}}{M_{I}} = \exp \left(\frac{(14 \ r)}{2r \ 7q} \right); \qquad (2:18)$$

$$\frac{M_{U}}{M_{X}} = \exp \frac{r}{2r \quad 7q} \quad : \tag{2:19}$$

A nother constraint which should be taken into account is that our equations are m eaningless if the gauge couplings become too large. It is dicult to say exactly how large is too large, but if we arbitrarily require that U^{1}_{U} 2; then given the num erical value X^{1}_{X} 25, from (2.9) we obtain (safely neglecting t_{U} t_{X}):

$$_{2}(t_{U} t_{I}) < 145:$$
 (2.20)

Multiple Intermediate Thresholds

So far we have assumed only one intermediate threshold between 1 TeV and M_X, but as previously discussed, this may not be a realistic assumption. More generally, suppose there are N distinct intermediate mass scales M_{Ia} (a = 1 ::: N) between 1 TeV and the uni cation scale. At each of these N thresholds, _{1a}, _{2a}, and _{3a} are the decreases in slope of the running inverse gauge couplings. One may then use the master form ula (2.12) iteratively to build the corresponding equations. The results are

$$t_{U} \quad t_{X} = \frac{1}{4} \frac{X^{N}}{a=1} q_{a} (t_{U} \quad t_{Ia}) + \frac{1}{2};$$

and

$$t_U = t_X = \frac{1}{14} \sum_{a=1}^{X^N} r_a (t_U = t_{Ia})$$
:

where $q_a = {}_{3a} {}_{2a}$ and $r_a = 5({}_{2a} {}_{1a})=2$ for each of the N thresholds. Now requiring $t_U t_X > 0$ constrains the particle content. One may view this case as one of multiple lenses, some divergent, some convergent.

These multiple thresholds act like one e ective lens, which leads us to recast these equations by choosing a single e ective intermediate scale \bar{t}_I which should reject the \average" of the individual thresholds in some sense. The choice of \bar{t}_I is to some extent arbitrary, as long as $\bar{t}_I < t_X < t_U$, and indeed the appropriate choice for a de nition of \bar{t}_I depends on the particular example being studied. Then one de nes:

$$- \sum_{i=1}^{N} \frac{t_{U} t_{Ia}}{t_{U} \overline{t}_{I}}; \qquad (2.21)$$

in which each $_{ia}$ is weighted more (less) when the corresponding intermediate scale t_{Ia} is lower (higher) than \overline{t}_I . In terms of

$$\overline{q} \quad \begin{array}{c} X^{N} \\ q \\ a=1 \end{array} \quad \begin{array}{c} t_{U} \quad t_{Ia} \\ t_{U} \quad \overline{t}_{I} \end{array} ; \quad \overline{r} \quad \begin{array}{c} X^{N} \\ r_{a} \\ a=1 \end{array} \quad \begin{array}{c} t_{U} \quad t_{Ia} \\ t_{U} \quad \overline{t}_{I} \end{array} ; \quad (2:22)$$

we obtain

$$\frac{\overline{q}}{4} = \frac{t_U \quad t_X \quad =2}{t_U \quad \overline{t}_I} ; \qquad (2.23)$$

and

$$\frac{\overline{r}}{14} = \frac{t_U \quad t_X}{t_U \quad \overline{t}_I} :$$
 (2.24)

The gauge coupling at uni cation is

$$U^{1} = X^{1} \frac{1}{2} - (t_{U} - t_{I}) + t_{U} - t_{X} : \qquad (2.25)$$

Note that the above equations have the same form as in the case of a single intermediate threshold, but with \averaged" quantities $\overline{t}_I, \overline{q}, \overline{r}$, etc. In fact, one still has the constraints

$$0 \quad \overline{q} < 4 ; \tag{2.26}$$

$$0 < \overline{r} < 14$$
; (2.27)

from requiring that the coupling constants unify, but not too early. The main di erence is that \overline{q} , \overline{r} , and $\overline{2}$ need not be integers. Each of the equations (2.8)-(2.20) derived in the case of a single interm ediate threshold now hold with t_{I} , i_{I} , q, r replaced by \overline{t}_{I} , \overline{i}_{I} , \overline{q} , \overline{r} .

3. Particles at the Interm ediate Threshold (s)

In order to analyze each case in detail, it is convenient to list the possible representations of the new particles that generate the interm ediate thresholds, and compute their $_{\rm i}$ coe cients.

For the purposes of this paper, we focus on low energy theories that could have originated from superstring theories. Thus we restrict ourselves to representations contained in $27;\overline{27}$ and 78 representations of E₆, under the decomposition

$$\rm E_{6}$$
 SU(2)_L SU(3)^C U(1)_Y :

In som e string com pacti cations, speci cally with higher levelK ac-M oody, chiralmultiplets transform ing as the adjoint can survive[6], with their rem nants appearing in the low energy theory. The results are summarized in Table 1.

CH IRAL SUPERMULT IPLETS						
R epresentation	1	2	З	q	r	#
(2;1 ^c) ₁ + c:	<u>м</u> Б	1	0	1	1	n ₁
(1;1 ^c) ₂ + c:	<u>6</u> 5	0	0	0	3	n ₂
(1;3 ^c) $\frac{4}{3}$ + c:	യഥ	0	1	1	4	n ₃
(1;3 ^c) _{2/3} + c:	<u>2</u> 5	0	1	1	1	n ₄
$(2;3^{C})_{\frac{1}{3}} + c:$	<u>1</u> 5	3	2	1	7	n ₅
(2;3 ^c) $\frac{5}{3}$ + c:	5	3	2	1	5	n ₆
(3;1 ^c) ₀	0	2	0	2	5	n ₇
(1;8 ^c) ₀	0	0	3	3	0	n ₈

Table 1

The last three representations in Table 1 appear only in the adjoint of E₆. We note that

for all these representations, 5(2 1) is even, and r is an integer. M ore generally, it can be shown that all representations for which 5(2 1) is odd necessarily describe leptons with half-integer electric charges, or quarks which yield bound states with non-integer charges. It follows from Table 1 that

> $q = n_1 + n_3 + n_4 \quad n_5 \quad n_6 \quad 2n_7 + 3n_8;$ $r = n_1 \quad 3n_2 \quad 4n_3 \quad n_4 + 7n_5 \quad 5n_6 + 5n_7;$

where there are n_i vector-like representations at the interm ediate threshold. In the superstring compactication scenario, the vector-like representations come from the fundamental of E₆. If there are no chiral super eld rem nants of the adjoint, $n_6 = n_7 = n_8 = 0$. We see from the above that the quantity q + rm ust be a multiple of 3:

 $q + r = 6n_5 \quad 3n_2 \quad 3n_3 \quad 6n_6 + 3n_7 + 3n_8$:

We should also take into account the possibility that the gauge group is enlarged above the interm ediate scale(s). In such a case, it is possible to identify the coupling constants for the enlarged gauge group and run the new gauge couplings up to the high scale. How ever, it is not really necessary to do so. Instead, one can simply follow the running of the three low energy gauge couplings even though they are embedded within the larger gauge group at high scales. Because of the assumption that the gauge couplings are properly normalized for unication into a simple gauge group, one can take into account the elects of gauge bosons and gauginos living at the intermediate scales by simple step functions in the beta functions. Therefore we generalize our analysis to include possible vector-like rem nants of a single vector supermultiplet adjoint of E $_6$. Table 2 is exactly the same as the rst 6 rows of Table 1, except that the entries now appearmultiplied by the factor 3, in accordance with the form ula (2.2) for the b_i , since they belong to the vector supermultiplet.

VECTOR SUPERMULT PLETS							
R epresentation	1	2	3	q	r	#	
(2;1 ^c) ₁ + c:	9 <u> </u> 5	3	0	3	3	N ₁ (1)
(1;1 ^c) ₂ + c:	<u>18</u> 5	0	0	0	9	N ₂ (2)
$(1;3^{\rm C}) = \frac{4}{3} + {\rm C}:$	<u>24</u> 5	0	3	3	12	N ₃ (2)
(1;3 ^c) _{2/3} + c:	6 <mark>1</mark> 5	0	3	3	З	N 4 (1)
(2;3 ^c) _{1/3} + c:	<u>3</u> 5	9	6	3	21	N 5 (2)
(2;3 ^c) $\frac{5}{3}$ + c:	15	9	6	3	15	N ₆ (1)

Table 2

The numbers in parentheses relect the multiplicity of the representation in a single adjoint of E_6 . The adjoint contains also ve singlets with no hypercharge, as well as the triplet which contains the SU (2) gauge and gaugino elds and the color octet of gluons and gluinos, which are already contained in the M SSM.

In order to account for the representations already present in the W ess-Zum ino multiplets, we simply have to replace n_i by $n_i^0 = n_i$ 3N i Thus, the previous formulae still apply, with the di erence that the n_i^0 can now be negative.

For each choice of possible subgroups of E₆ as gauge group above M_I, we can write down (up to several inequivalent embeddings) the non-zero N_i's corresponding to the gauge bosons which get m ass at M_I. Note that we must only consider gauge groups with N₅ = N₆ = 0, because otherwise SU (2)_L and SU (3)^c would necessarily be uni ed at M_I, which is in con ict with the fact that they have di erent couplings at that scale. (The corresponding gauge bosons surely could not have interm ediate scale m asses in any case, because of proton decay bounds.) So, we list the possibilities according to rank:

Rank 4:

 $Case 0: SU(2)_{L}$ SU(3)^C U(1); AllN_i = 0.

Rank 5:

Case 1: SU (2)_L SU (3)^C SU (2) U (1); (a) AllN_i = 0 or (b) N₂ = 1.

Case 10: SU (2)_L SU (6)^C; N₂ = 2, N₃ = 2, N₄ = 1

In each of cases 1, 3, 6, and 9, there are inequivalent embeddings of the standard m odel gauge group, resulting in two di erent possibilities for the N_i. There are also acceptable subgroups of E₆ obtained by adding U (1) factors to the rank 4 and 5 possibilities listed above. The extra U (1) factors do not contribute to the N_i, and do not a ect the one loop renorm alization group equations for the gauge couplings.

If the gauge group above the interm ediate scale is larger than the standard model's, there must appear at the same interm ediate threshold chiral super elds containing standard model singlets, to break the gauge group. In particular models, one must check for their presence and that the order parameters do not produce unacceptable R-parity violation (or baryon number violation in models which have alternative discrete symmetries).

4. Results: One Interm ediate Threshold

In the case of only one threshold, one can combine the results of the previous two sections to enumerate the possibilities for raising the uni cation scale. In section 2 we found that q = 0;1;2;3 and that r is an integer between 0 and 14, and from section 3 we found that q + r is a multiple of 3. Of special interest, perhaps, are the cases for which $M_X = M_I$ is large, so that the di erent scales are cleanly separated and m ay be de nitely associated with di erent physics. For example, if M_I is to be associated with an invisible axion scale, we expect $M_I = 10^{10-2}$ GeV, so that $M_X = M_I = 10^{6-2}$. If we want the

hierarchy M $_X = M_I$ to be large, without having j jbe too large or giving M $_U$ outside of the correct range between M $_X$ and the P lanck scale, there are tight restrictions which we now discuss, classified in terms of the value of q.

W hen q = 0, the unit cation scale does not depend on r. It is given by

$$\frac{M_{U}}{M_{X}} = \exp \frac{1}{2} ; \qquad (4:1)$$

while the allowed values of r are multiples of three, r = 3;6;9;12, corresponding to

$$\frac{M_{X}}{M_{T}} = \exp \left(\frac{11}{2} \left(\frac{11}{3}; \frac{4}{3}; \frac{5}{9}; \frac{1}{6}\right)\right) ; \qquad (4.2)$$

respectively. Clearly, must be positive in order to raise the uni cation scale in this case, with larger values of corresponding to more substantial hierarchies in M $_{\rm U}$ =M $_{\rm X}$ and M $_{\rm X}$ =M $_{\rm I}$. However, note that the hierarchy M $_{\rm X}$ =M $_{\rm I}$ is severely limited unless r = 3, and even then M $_{\rm X}$ =M $_{\rm I}$ cannot exceed 6 10³ for < 1:5. In the cases r = 6;9;12, M $_{\rm X}$ =M $_{\rm I}$ cannot be large.

For q = 1, the uni cation scale is given by

$$M_{\rm U} = \frac{M_{\rm X}^{4=3}}{M_{\rm I}^{1=3}} e^2 \quad ^{=3}; \qquad (4:3)$$

independent of r. The possible values of r are 2, 5, 8, and 11, and the results for in terms of the hierarchies $t_U = t_X$ and $t_X = t_I$ from (2.18) and (2.19) are given in Table 3.

Table 3

r	2	5	8	11
$=(t_U t_X)$: 48	:19	36	: 43
$=(t_X t_I)$: 08	:11	: 48	1 : 6

From Table 3 and (4.3), we can see that the hierarchy M_X =M_I can be very large if r = 2 or 5. The case r = 2 can accomedate intermediate scales as low as 10^8 GeV for negative, and r = 5 can give M_I as low as 10^{10} GeV , for positive. The case r = 8 does not allow M_X =M_I to be larger than about 20, because otherwise we see from Table 3 that

would be larger than allowed by the experimental constraint (2.3). The case r = 11 does not allow M $_X = M_I$ to be large enough to be meaningful at all.

For q = 2, the uni cation scale is given by

$$M_{\rm U} = \frac{M_{\rm X}^2}{M_{\rm I}} e$$
 : (4:4)

The possible values of r are 1, 4, 7, 10, and 13, and the results for are given in Table 4.

r	1	4	7	10	13
$=(t_U t_X)$	3 : 8	: 48	0	:19	:29
$=(t_X t_I)$	29	:19	0	: 48	3:8

Table 4

C learly, in the case r = 13 there can be no appreciable hierarchy in M_X =M_I, because of the constraint (2.3) on . In the case r = 10, the constraint on implies that M_X =M_I can be at most 20 or so. The case r = 1 can give M_X =M_I as large as 200, but then does not allow M_U to be signi cantly larger than M_X. In the case r = 7, must be zero, as we have already noted, and from (4.4), the hierarchy M_X =M_I must be less than 10^3 in order that M_U not exceed the P lanck scale. The remaining case r = 4 can allow M_X =M_I to be as large as about 3 10^3 , but no larger, because otherwise we see from Table 4 that would be too negative.

For q = 3, the uni cation scale is given by

$$M_{\rm U} = \frac{M_{\rm X}^{4}}{M_{\rm T}^{3}} e^2 \qquad : \qquad (4.5)$$

The possible values of r are 3, 6, 9, and 12, and the results for are given in Table 5.

1 able 5					
r	3	6	9	12	
=($t_U t_X$)	1 : 6	: 48	:11	: 08	
$=(t_X t_I)$: 43	: 36	:19	: 48	

Table 5

C learly there is no way to get even an order of magnitude hierarchy in M $_X = M_I$ in the case r = 12, because otherwise from (4.5), M $_U$ would exceed the P lanck scale since is positive. The other cases have negative , and therefore can accom odate a slightly larger hierarchy; for p = 3; 6; 9, one can have M $_X = M_I$ as large as 30, 70, and 50 respectively, without having be too negative or exceeding the P lanck-scale bound on M $_U$.

To sum marize the preceding results, there are remarkably few cases in which one can have a large hierarchy of scales $M_X = M_I$. Only in the cases q = 1; r = 2 and q = 1; r = 5can one hope to have $M_X = M_I$ 10⁴. These appear to be the only acceptable cases if one w ishes to associate M_I with an invisible axion scale (or anything else below 10^{12} GeV). The cases q = 0; r = 3 and q = 2; r = 4 and q = 2; r = 7 can give hierarchies which are roughly in the range $M_X = M_I$ 10³. All of the other cases give sm aller upper limits for $M_X = M_I$.

In the superstring scenario, an estimate of string e ects indicates that the scale of string uni cation should be related to the gauge coupling through the form ula [7]

$$M_{\rm U} = 2.5^{\rm P} \frac{10^{18} \, \text{GeV}}{10^{18} \, \text{GeV}}$$
 (4:6)

Taking M_X = 10^{16} GeV, and $_{U}^{1} < _{X}^{1}$ 25, eq. (4.6) in plies that contact with the superstring can be made provided that M_U=M_X > 50.

As an example, suppose we take = 0.82 with r = 5, q = 1. Then eq. (4.8) can be satisfied together with the other constraints by $n_1 = 4$, $n_2 = n_3 = 0$, $n_4 = 6$ and $n_5 = 1$. We not that

$$M_{\rm U} = 7.5 \ 10^{17} {\rm GeV}$$
; $M_{\rm I} = 4.4 \ 10^{12} {\rm GeV}$; ${}_{\rm U}^{1} = 11$:

This is one of the solutions with low r for which there is only one interm ediate threshold well separated from M_X. It is interesting that most of the solutions with just one interm ediate scale threshold do not allow M_X = M_I to be very large.

5. Results: Several Interm ediate Thresholds

In most superstring theories, the elective low energy gauge group at the string scale is larger than the standard model gauge group, and it is necessary to have several intermediate scale thresholds. Even if there is only one order parameter associated with the intermediate

scale, the m asses of the vector-like particles are related to that order parameter by various dimensionless couplings which are certainly not always close to unity. This will result in som e \sm earing" of the threshold associated with each order parameter. Thus in a realistic model, the assumption of just one intermediate scale is probably not justified. However, we can still protably analyze the situation in terms of the averaged quantities $\overline{t}_{I}; \overline{q}; \overline{r}$, etc. which were introduced in section 2. These quantities summarize the elects of the intermediate scale mass thresholds in terms of a single elective intermediate scale, with the main difference being that \overline{q} and \overline{r} need not be integers.

Let us apply our analysis to the interesting example of the 3-family G epner-Schimmrigk superstring model [8,9]. Below the string scale, the surviving gauge group is the

subgroup of E $_6$, corresponding to our case 7 (of section 3) with N $_1 = 1$ and N $_2 = 2$. This gauge group is subsequently broken to

corresponding to our case 1 (b) with N₂ = 1, and then to the standard m odel gauge group. There are thus at least two a priori distinct interm ediate scale order parameters associated with each reduction in rank. The chiral super elds which survive below the string scale are classified under the gauge group SU (3)_L SU (3)^C SU (3)_R as:

9 leptons	(3;1; <u>3</u>)
6 m irror leptons	(3;1;3)
3 quarks	(3;3;1)
3 antiquarks	(1;3;3)

and, unlike most other string models, no mirror quarks $(3;\overline{3};1)$ or mirror antiquarks $(1;3;\overline{3})$.

This particle content includes, besides the chiral super elds for the three fam ilies of quarks and leptons and two H iggs doublets of the m in in alsupersym m etric standard m odel, chiral super elds corresponding to

$$n_1 = 20;$$
 $n_2 = 6;$ $n_3 = 0;$ $n_4 = 3;$ $n_5 = 0:$

C om bining these with the vector super elds, we have a total vector-like particle content yielding

$$n_1^0 = 17;$$
 $n_2^0 = 0;$ $n_3^0 = 0;$ $n_4^0 = 3;$ $n_5^0 = 0:$

Thus if all of these particles were concentrated at just one intermediate mass scale, we would have

$$_1 = 57=5;$$
 $_2 = 17;$ $_3 = 3$

giving

 $q_{total} = 14$ and $r_{total} = 14$: (5:1)

These values lie outside the range established by (2.26) and (2.27). If the particle thresholds a ect the gauge coupling uni cation in a perturbative and meaningful way below M_X , there must be some smearing, with the \averaged" quantity \overline{q} higher than q_{total} and \overline{r} lower than r_{total} . O therwise, from the discussion in section 2, 3 and 2 would meet too early (just above the intermediate scale) and 2 and 1 would never meet. It is clear that to move things in the right direction, the contributions of the 2a to each of \overline{q} and \overline{r} should be weighted less heavily than those of 1a and 3a. This can only occur if the masses of the electrosinglet down quark vector-like chiral super elds corresponding to n_4 are smaller than the average elds of the other particles. (Note that in this example, N $_3 = N_4 = 0$.)

Let us denote by \bar{t}_{n_1} , \bar{t}_{n_2} , and \bar{t}_{n_4} the arithmetic means of the scales associated with the chiral super elds corresponding to the weak doublet vector-like leptons, n_1 , the weak singlet charged leptons, n_2 , and the down-like electroweak singlet quarks, n_4 , respectively. Sim ilarly, the arithmetic means of the scales associated with the vector supermultiplets corresponding to N_1 and N_2 are denoted by \bar{t}_{N_1} and \bar{t}_{N_2} . Then it is convenient to choose for the electric intermediate scale $\bar{t}_I = \bar{t}_{n_4}$, which is just the scale associated with the electric threshold for $_3$. With this choice, one nds:

$$\overline{q} = 20 \quad \frac{t_{U} \quad \overline{t}_{n_{1}}}{t_{U} \quad \overline{t}_{I}} + 3 \quad \frac{t_{U} \quad \overline{t}_{N_{1}}}{t_{U} \quad \overline{t}_{I}} + 3; \quad (5.2)$$

$$\overline{\mathbf{r}} = \overline{\mathbf{q}} + 18 \quad \frac{\overline{\mathbf{t}}_{n_2} \quad \overline{\mathbf{t}}_{N_2}}{\mathbf{t}_{T} \quad \overline{\mathbf{t}}_{T}} \quad ; \tag{5:3}$$

$$_{2} = \overline{q} + 3$$
: (5:4)

Note that \overline{t}_I cannot be larger than \overline{t}_{N_1} or \overline{t}_{N_2} , because the vectorlike color triplets can only obtain their masses at or below the scale at which the gauge group is broken down

to that of the standard m odel. A lso, two of the vectorlike pairs corresponding to n_1 m ust have m asses at scales below $\bar{t}_{N_{1,2}}$, for the same reason. Since these contribute negatively to the RHS of (5.2), the net positive contributions to \bar{q} are quite limited. So we see that the only way to obtain $0 < \bar{q} < 4$ is for the vectorlike weak doublet leptons, corresponding to n_1 to be located (on average) well above \bar{t}_I . From (5.3), one can also see that the scale \bar{t}_{n_2} associated with the charged lepton chiral super elds must also be located above \bar{t}_{N_2} . Finally, we see from (5.4) that if the thresholds are arranged appropriately for gauge coupling unication, then \bar{t}_2 is automatically not larger than 3, so that the constraint (2.20) from perturbativity of the couplings does not limit the elective intermediate scale \bar{t}_I at all. Another way to see this is to note that the slope of 3^1 can never be negative with this particle content. (O fecure, in models with a larger sector of strongly interacting chiral super elds, the requirement of perturbativity can be quite in portant.)

If some of the chiral super elds have masses located far below M $_{\rm X}$, we have seen that some of these must include the color triplet elds corresponding to n_4 . This can be understood from the fact that only these color triplets give a positive contribution to \overline{q} among the chiral super elds of the model. One should note, however, that there is a potential embarrassment associated with such light color triplets; they can easily lead to proton decay at unacceptable rates if their masses are below M $_{\rm X}$, depending on their couplings to the quark and lepton super elds of the M SSM . This can be avoided if e.g. one assumes the existence of a discrete symmetry [10] prohibiting some or all of the baryon num ber and lepton num ber violating couplings. A ctually, the presence of vectorlike down-type quarks below M_X seems to be a fairly general feature of string-type models in which interm ediate scale thresholds are used to raise the uni cation scale; see for example [11,12]. One can understand this sem i-quantitatively by examining the values of q and r for the chiral supermultiplets in Table 1. Only the chiral super elds corresponding to n_3 and n_4 can give a positive contribution to \overline{q} . However, the super elds for n_3 (which are innocuous for proton decay) also give a relatively large negative contribution to \overline{r} . Since \overline{q} and \overline{r} both must be positive to raise the uni cation scale, it seems that the color triplet with electric charge 1=3 corresponding to n_4 must be weighted relatively heavily in the averaged quantities. This is another way of saying that they are relatively light compared to the other chiral super elds which are important in redirecting the running gauge couplings to their new meeting point. Of course, one can always achieve a raised

uni cation scale fairly safely by employing only thresholds which are close to M $_X$. In most superstring models[12], this is almost required, since the large number of strongly interacting chiral super elds would cause the gauge couplings to be non-perturbative if the elective intermediate scale were much lower than about 10^{15} GeV.

6.Conclusion

In this paper, we have exam ined the possibility that the true uni cation scale can be raised above its apparent value of $2 \ 10^{16}$ G eV by calculable perturbative m eans. It m ight seem rather surprising that in the MSSM the gauge couplings should appear to be nicely headed for uni cation at M $_{\rm X}$, only to be redirected to a new m eeting place at M $_{\rm U}$. Indeed, the apparent perverseness of this situation allow s us to put som e non-trivial constraints on the scenario. In the simplest case of just one cleanly de ned intermediate scale, it is striking that the hierarchy M $_{\rm U}$ = M $_{\rm I}$ is generally quite limited. In the probably more realistic case of a \sm eared" interm ediate scale or several interm ediate scales, one cannot be as precise because of the vastly increased number of unknown parameters. However, one can still put useful constraints on the placem ent of the interm ediate scales and particles, by writing things in terms of a single e ective intermediate threshold. Here too, in most realistic m odels based on superstrings, there is a tendency form any of the vector-like particles to be very heavy, based simply on the requirem ent that the gauge coupling rem ain perturbative and thus calculable in principle at high energies. Even in models like the one considered in section 5, in which the absence of a large number of vector-like strongly interacting particles causes perturbativity to be easily maintained, one nds that it is di cult to raise the uni cation scale consistently with interm ediate scales much below M $_{\rm X}$. If one insists on having some chiral super elds at relatively low intermediate scales, we nd that generally these chiral super elds include color triplets with electric charge 1=3, which m ay be dangerous for proton decay without assuming some extra symmetry.

The di culty in obtaining examples in which a raised uni cation scale is achieved due to a relatively low intermediate scale corresponds to our intuition that it would be surprising if the uni cation of gauge couplings were totally accidental. The lower the intermediate scale(s) are, the more we must regard the apparent success of the uni cation of gauge couplings as just a perverse accident. On the other hand, if there are intermediate scale thresholds which are only slightly below the uni cation scale, then the near

20

perfect uni cation of couplings should be regarded as partly, but certainly not com pletely, accidental. This scenario seem s to be the one preferred by superstring m odels.

The work of SPM. was supported in part by the United States D epartm ent of Energy. The work of PR. was supported in part by the United States D epartm ent of Energy under grant D = FG 05-86 = R 40272.

References

- 1) For reviews, see H.P.Nilles, Phys.Rep.110 (1984) 1 and H.E.Haber and G.L.Kane, Phys.Rep.117 (1985) 75.
- 2) P.Langacker, in Proceedings of the PASCOS90 Sym posium, Eds.P.N ath and S.R eucroft, (W orld Scientic, Singapore 1990) J. Ellis, S. Kelley, and D. Nanopoulos, Phys. Lett. 260B (1991) 131; U.Am aldi, W. de Boer, and H. Furstenau, Phys. Lett. 260B (1991) 447; P.Langacker and M. Luo, Phys. Rev. D 44 (1991) 817.
- 3) Jihn E.Kim, Phys.Rep.149 (1987) 1.
- 4) M .G ell-M ann, P.R am ond, and R .Slansky, in Sanibel Talk, CALT -68-709, Feb 1979, and in Supergravity, (N orth H olland, Am sterdam, 1979; T .Yanagida, in Proc. of the W orkshop on Uni ed Theories and Baryon N um ber in the Universe, T sukuba, Japan, 1979, edited by A .Saw ada and A .Sugam oto (KEK Report N 0.79-18, T sukuba, 1979).
- 5) L.E. Ibanez, Phys. Lett. B 318 (1993) 73.
- 6) S. Chaudhuri, S.W. Chung, and J. Lykken, FERM ILAB PUB-94-137-T, Talk given at the 2nd FT W orkshop on Yukawa Couplings and The O rigins of M ass, G ainesville, February 1994; S. Chaudhuri, S.W. Chung, G. Hockney, and J. Lykken, Ferm ilab preprint PR IN T-94-0170, August 1994.
- 7) V.S.Kaplunovsky, Nucl.Phys.B 307 (1988) 145 and Erratum, preprint Stanford-ITP-838 (1992).
- 8) R. Schimmrigk, Phys. Lett. B 193 (1987) 175; D. Gepner, Princeton University report, 1987.
- 9) S.Cordes and Y.Kikuchi, Int.J.M od. Phys. A 6 (1991) 5017; J.W u, R.A mow itt

and P.Nath, Int.J.Mod.Phys.A 6 (1991) 381; J.W u and R.Amowitt, Phys. Rev.D 49 (1994) 4931.

- 10) D.J.Castano and S.P.Martin, Phys.Lett.B 340 (1994) 67, and references therein.
- 11) J.Ellis, C.Kounnas and D.V.Nanopoulos; Nucl.Phys.B247 (1984) 373; S.Kelley, J.L.Lopez and D.V.Nanopoulos, Phys.Lett.B278 (1992) 140.
- 12) B.R.Greene, K.H.Kirklin, P.J.Miron, and G.G.Ross, Phys.Lett.B180 (1986) 69; Nucl.Phys.B278 (1986) 667; Nucl.Phys.B292 (1987) 606.