Z_2 Z_2 orbifold compactication {

the origin of realistic free ferm ionic models

A lon E. Faraggi
School of Natural Sciences, Institute for Advanced Study
O lden Lane, Princeton, NJ 08540
E-mail:faraggi@sns.ias.edu

ABSTRACT

A llthe realistic free ferm ionic m odels utilize a set of basis vectors, the NAHE set, that correspond to Z $_2$ – Z $_2$ orbifold compactication with nontrivial background elds. I argue that the realistic features of free ferm ionic models, like the number of generations and the ferm ion mass spectrum are due to the underlying Z $_2$ – Z $_2$ orbifold compactication.

As a uni ed theory of gravity and the gauge interactions, heterotic string theory should reproduce the matter and symmetry content of the Standard Model and determine the fermion mass spectrum. Presently we do not know what is the dynamical mechanism that selects the unique string vacuum, and, a priori, there is a large number of potentially viable superstring models.

The notion, however, that there is a huge number of string models is somewhat misleading. By just imposing one or two phenomenological criteria, like three generations and a gauge group that can be reduced at low energies to the standard model gauge group, already one nds that the number of possibilities is substantially reduced. Imposing further phenomenological constraints may indeed single out a unique superstring model. If such a model is constructed, it will certainly be of use in trying to learn about the dynamical mechanism that chooses the string vacuum.

The task of constructing phenom enologically viable string models seem shopeless. While in tendimensions the string vacuum is more or less unique, in four dimensions there is a huge number of equivalent candidates. The string consistency constraints impose a number of degrees of freedom and those degrees of freedom produce a symmetry that is larger than the observed symmetry at low energies. Furthermore the number of chiral generations is also determined in the four dimensional vacuum and is correlated with the gauge degrees of freedom. A bottom (up approach, in which dierent blocks of the standard model are assembled together piece by piece, is not adequate. Rather, what is required is a top (bottom approach in which the features of the standard model are carved out of the more symmetric string vacua.

To appear in Proceedings of the Beyond the Standard Model IV Conference, Lake Tahoe, California, December 13 (18, 1994.

Is there a guiding principle that may distinguish among the equivalent string vacuum? String vacua exhibits a new kind of sym metry, usually referred to as target { space duality², which is a generalization of the R! 1=R duality in the case of S^1 . At the self{dualpoint, $R_j = 1=R_j$, space{time sym metries are enhanced. For appropriate choices of the background elds the space{time sym metries are maximally enhanced³. At the maximally sym metric point the internal degrees of freedom that are needed to cancel the conformal anomaly may be represented in terms of internal free fermions propagating on the string world { sheet. It is plausible that if string theory is realized in nature then the true string vacuum is in the vicinity of the highly sym metric self{ dual point. It may turn out that near that point the free fermionic formulation⁴ provides a good approximation to the true string vacuum. However, the number of consistent free fermionic models is still enormous.

As is well known in (2,2) string models that adm it a geometrical interpretation the number of chiral generations is half the Euler number of the six dimensional compactied manifold. Following LEP data it is plausible to assume that only three complete generations exist in nature. How can three generations arise from a six dimensional compactied space. The answer may be simple. The six dimensional compactied space is divided into three factors of two. In the orbifold language⁵, divide the six dimensional space, which is compactied on a attorus, by a Z_2 Z_2 discrete symmetry. In that case the Z_2 Z_2 orbifold model produces exactly three twisted sectors. In the Z_2 Z_2 orbifold on a six dimensional space, the number three is deeply rooted in the structure of the models. Thus, the Z_2 Z_2 can very naturally lead to models with three generations. Namely, each light generation comes from a different twisted sector of the Z_2 Z_2 orbifold model.

It appears that Z_2 Z_2 orbifold on the attorus of the six dimensional compactied space, can very naturally lead to three generations. However, in general, Z_2 Z_2 orbifold on generic lattices do not lead to three generation models. For example the Z_2 Z_2 orbifold on SO $(4)^3$ lattice did not yield three generation models. In contrast, the Z_2 Z_2 models at the free fermionic point in toroidal compactication space, realized by the NAHE set^{6;7}, do produce three generation models. The dierence is seen by examining the number of xed points in the two compactications with (2,2) world sheet supersymmetry. On the SO $(4)^3$ lattice the Z_2 Z_2 produces sixteen generations, from each twisted sector. On the SO (12) lattice, which corresponds to the free fermionic point in the toroidal compactication space, it produces eight chiral generations, from each twisted sector. In the fermionic three generation constructions each one of the complex planes of the Z_2 Z_2 orbifold is modded out by additional Z_2^3 symmetries, thus reducing the number of generations to one generation from each twisted sector.

In the (2,2) ferm ionic constructions one starts from a set of boundary condition vectors that produces an N = 4 supersymmetric model with SO (12) E₈ E₈ gauge group³. One then adds two boundary condition vectors that correspond to the

 Z_2 Z_2 tw isting. The resulting gauge group is SO $(4)^3$ E_6 U $(1)^2$ E_8 with N=1 space{time supersymmetry. In this model there are twenty four chiral generations from the boundary condition vectors that correspond to twisted sectors and three pairs of chiral and antichiral generations from the untwisted sector. The number of twisted and untwisted moduli is equal to the number of generations. In addition the untwisted and twisted sectors produce E_6 E_8 singlets that are obtained by acting on the vacuum with oscillators that arise from the fermionic degrees of freedom that correspond to the six internal compaction dimensions.

In the orbifold formulation the same model is obtained by applying a Z₂ twist to a torodialy compactifed SO (12) lattice and E_8 E_8 gauge symmetry. The 36 free param eters of the six dimensional metric and the antisymmetric tensor eld param eterize the six dimensional compactied space. For generic values of these param eters the gauge symmetry that arises from the six dimensional compactied torus is U (1)6. For speci c choices of the background parameters the U (1)6 of the com pacti ed torus is enlarged. To reproduce the SO (12) E₈ E₈ gauge group of the free ferm ionic model, the metric G $_{\rm ij}$ is the C artan matrix of SO (12) and the antisym m etric tensor eld is given by, $B_{ij} = G_{ij}$ for i > j; $B_{ij} = 0$ for i = j and $B_{ij} = G_{ij}$ for i < j. For $R_I = \frac{\pi}{2}$ and with the chosen background elds, the right{m oving m om enta produce the root vectors of SO (12), thus reproducing the sam e gauge group as in the free ferm ionic model. The orbifold model is obtained by m oding out the six dim ensional torus by a discrete symmetry group. The massless spectrum contains states from the untwisted and twisted sectors. In the case of \standard em bedding" the num ber of chiral fam ilies is given by one half of the Euler characteristic. To translate the ferm ionic boundary conditions to twists and shifts in the bosonic formulation the real ferm ionic degrees of freedom that correspond to the com pacti ed dim ensions are bosonized. The ferm ionic boundary condition vectors, b. and b_2 , then translate to Z_2 Z_2 twist on the compactified coordinates and to shifts on the gauge degrees of freedom. It is then seen that symmetries and spectrum of the orbifold model coincide with those of the corresponding ferm ionic model.

The realistic free ferm ionic models correspond to models with (2,0), rather than (2,2), world sheet supersymmetry. The transition from the (2,2) models to the (2,0) models can be regarded as choosing a GSO phase between the two boundary condition vectors that produce the spinorial of SO (16). The GSO projection projects out the massless states from these sectors and the resulting gauge group is SO (12) SO (16) SO (16), with N = 4 space (time supersymmetry. Alternatively, one of the spinorial vectors may be enlarged with additional four periodic complex fermions in the hidden sector. The E_8 E_8 gauge group is modiled to SO (16) SO (16), as in the rst construction. The analysis with respect to the number of xed points is identical to the case with (2,2) world (sheet supersymmetry. However, in this case the observable gauge group after applying the E_2 E_3 is SO (10) U (1) rather than E_6 , and the U (1) is \anomalous". The twisted sectors produce spinorial and vectorial sixteen of

the observable SO (10) and hidden SO (16) gauge groups, respectively.

The structure of the Z_2 Z_2 orbifold with (2,0) world sheet supersymmetry and standard embedding, is common to all the realistic free fermionic models. Three generation models are obtained by adding three additional boundary condition basis vectors, beyond the NAHE set. The additional boundary condition vectors models of the three complex planes of the Z_2 Z_2 orbifold by a Z_2^3 symmetry and break the observable SO (10) symmetry to one of its maximal subgroups SU (5) U (1), SO (6) SO (4) or SU (3) SU (2) U (1)².

The ferm ion mass spectrum is also seen to originate from the Z_2 structure, realized by the NAHE set. The untwisted sector produces three pairs of Higgs doublets and a combination of the vectors that break the SO (10) symmetry produces one or two additional pairs. Due to the horizontal symmetries in the Z_2 Z_2 orbifold models, each pair of Higgs doublets couples only to states from one of the twisted sectors, producing couplings $16_{1}10_{1}$ j = 1;2;3. The cancelation of the anom alous U (1) D {term equation by singlet VEVs, gives P lanck scale mass to several Higgs doublets. As a result, there exist models in which only one mass term, namely the top quark mass term, exist at the cubic level of the superpotential. The mass term s for the lighter quarks and leptons are obtained from nonrenormalizable term s. The nonrenormalizable terms contain SO (10) singlets with nonvanishing VEVs, that are required to cancel the anom alous U (1) D (term equation. Thus, the nonrenorm alizable term s become e ective renormalizable terms that are suppressed relative to the leading cubic level term s. Due to the horizontal symmetries and the singlet VEVs one generation is necessarily light⁸. Sim ilarly, the mixing terms arise generically from nonrenom alizable terms of the form $16_{i}16_{i}1016_{i}16_{i}$ ", where the set two 16 are in the spinorial representation of the observable SO (10), the 10 is in the vector representation of the observable SO (10), the last two 16 are in the vector representation of the hidden SO (16) and n is a combination of SO (10) SO (16) scalar singlets^{9,3}. The \mathbb{Z}_2 \mathbb{Z}_2 orbifold structure gives rise to the horizontal sym m etries that m ay be needed to understand the matter mass spectrum. Requiring adequate generation mixing and the form of the mixing terms necessitates that we give nonvanishing VEVs to some of the hidden sector 16 representations. In Ref. [10] it was shown that this is possibly the source of supersym metry breaking in these models.

A cknow ledgem ents

This work is supported by DOE grant DE-FG02-90ER40542.

References

- 1. D J. G ross et. al., NuclPhys. B 256 (1985) 253.
- 2. A. Giveon et. al., Phys Rep. 244 (1994) 77, and references therein.
- 3. A E. Faraggi, PhysLett. B326 (1994) 62, hep-ph/9311312...
- 4. I. Antoniadis, C. Bachas, and C. Kounnas, NuclPhys. B289 (1987) 87; H. Kawai, D.C. Lewellen, and S.H.-H. Tye, NuclPhys. B288 (1987) 1.
- 5. L.Dixon, JA. Harvey, C. Vafa and E.Witten, Nucl. Phys. B274 (1986) 285.
- 6. I. Antoniadis et. al., Phys.Lett. B231 (1989) 65.
- 7. A E. Faraggi, NuclPhys. B 387 (1992) 239, hep-th/9208024.

- 8. A E. Faraggi, NuclPhys. B403 (1993) 101, hep-th/9208023; NuclPhys. B407 (1993) 57, hep-ph/9210256.
- 9. A E. Faraggi and E Halyo, PhysLett. B307 (1993) 305, hep-ph/9301261; NuclPhys. B416 (1994) 63, hep-ph/9306235.
- 10. A E. Faraggi and E Halyo, IASSNS {HEP $\{94/17, hep-ph/9405223.$