QCD Radiative Corrections to the Leptonic Decay Rate of the B_c M eson

Eric Braaten and Sean Fleming

Department of Physics and Astronomy, Northwestern University, Evanston, IL 60208

A bstract

The QCD radiative corrections to the leptonic decay rate of the B $_{\rm C}$ m eson are calculated using the form alism of nonrelativistic QCD (NRQCD) to separate short-distance and long-distance e ects. The B $_{\rm C}$ decay constant is factored into a sum of NRQCD matrix elements each multiplied by a short-distance coe cient. The short-distance coe cient for the leading matrix element is calculated to order $_{\rm S}$ by matching a perturbative calculation in full QCD with the corresponding perturbative calculation in NRQCD. This short-distance correction decreases the leptonic decay rate by approximately 15%.

The study of heavy quarkonium systems has played an important role in the developm ent of quantum chrom odynam ics (QCD). Som e of the earliest applications of perturbative QCD were calculations of the decay rates of charm onium [1]. These calculations were based on the assumption that, in the nonrelativistic limit, the decay rate factors into a short-distance perturbative part associated with the annihilation of the heavy quark and antiquark and a long-distance part associated with the quarkonium wavefunction. This simple factorization assumption fails for P-wave states [2], which satisfy a more general factorization formula containing two nonperturbative factors [3]. Calculations of the annihilation decay rates of heavy quarkonium have recently been placed on a solid theoretical foundation by Bodwin, Braaten, and Lepage [4]. Their approach is based on nonrelativistic QCD (NRQCD), an elective eld theory that is equivalent to QCD to any given order in the relative velocity v of the heavy quark and antiquark [5]. Using NRQCD to separate the short-distance and long-distance e ects, Bodwin, Braaten, and Lepage derived a general factorization formula for the inclusive annihilation decay rates of heavy quarkonium. The short-distance factors in the factorization formula can be calculated using perturbative QCD, and the long-distance factors are de ned rigorously in terms of matrix elements of NRQCD that can be evaluated using lattice calculations. The general factorization formula applies equally well to S-waves, P-waves, and higher orbital-angular-momentum states, and it can be used to system atically incorporate relativistic corrections to the decay rates.

Since the top quark decays too quickly to produce narrow resonances, the only heavy quark-antiquark bound states that remain to be discovered are the bcm esons and their antiparticles. The possibility that bcm esons may be discovered at existing accelerators has stimulated much recent work on the properties of these mesons [6] and on their production cross sections at high energy colliders [7]. Once produced, a bcm eson will cascade down through lower energy bc states via hadronic or electrom agnetic transitions to the pseudoscalar ground state Bc which decays weakly. The discovery of the Bcm eson will require a detailed understanding of its decay modes. In this paper, we compute the short-distance QCD radiative correction to the leptonic decay rate of the Bc. We use the formalism of NRQCD to factor the amplitude for the decay into short-distance one cients multiplied by NRQCD matrix elements. The short-distance one cient for the leading matrix element is calculated

to next-to-leading order in $\ _{\rm s}$ by m atching a perturbative calculation in full QCD with the corresponding perturbative calculation in NRQCD .

The leptonic decay of the B $_{\rm c}$ proceeds through a virtual W $^+$ as in Figure 1. The W $^+$ couples to the B $_{\rm c}$ through the axial-vector part of the charged weak current. A llQCD e ects, both perturbative and nonperturbative, enter into the decay rate through the decay constant $f_{\rm B}_{\rm c}$, de ned by the m atrix element

$$h0 \ b \quad {}_{5}c \ B_{c}(P) \ i = i f_{B_{c}}P ; \tag{1}$$

where $\beta_c(P)$ is the state consisting of a B_c with four-momentum P. It has the standard covariant norm alization $hB_c(P)$ $\beta_c(P)$ i = $(2)^32P^{0-3}(P^0-P)$ and its phase has been chosen so that f_{B_c} is real and positive. In terms of the decay constant f_{B_c} , the leptonic decay rate is

$$(B_{c}! \quad ^{+} \quad) = \frac{1}{8} \mathcal{Y}_{bc} \mathcal{J}^{2} G_{F}^{2} M_{B_{c}} f_{B_{c}}^{2} m^{2} \quad 1 \quad \frac{m^{2}}{M_{B_{c}}^{2}} ; \qquad (2)$$

where V_{bc} is the appropriate K obayashi-M askawa m atrix element, G_F is the Fermi constant, M_{B_c} is the m ass of the B_c m eson, and m \cdot is the m ass of the charged lepton.

The formula (1) provides a nonperturbative de nition of the decay constant $f_{\rm B_{\, c}}$, so that it can be calculated using lattice QCD simulations. One of the di-culties with such a calculation is that it requires a lattice with large volume and ne lattice spacing, since the strong interactions must be accurately simulated over many distance scales. The long-distance scales range from $1=_{\rm QCD}$, the scale of nonperturbative elects associated with gluons and light quarks, to the scale $1=(m_{\rm c}v)$ of the meson structure, where v is the typical relative velocity of the charm quark. The short-distance scales include the Compton wavelengths $1=m_{\rm c}$ and $1=m_{\rm b}$ of the heavy quark and antiquark. A more elective strategy for calculating $f_{\rm B_c}$ is to separate short-distance elects from long-distance elects, to calculate the short distance elects analytically using perturbation theory in $_{\rm S}$, and to use lattice simulations only for calculating the long-distance elects. Having already taken into account the short-distance elects, one can use a much coarser lattice which provides enorm ous savings in computer resources.

An elegant way to separate short-distance and long-distance e ects is to use NRQCD,

an e ective eld theory in which heavy quarks are described by a nonrelativistic Schroedinger eld theory of 2-com ponent Pauli spinors. The lagrangian is

$$L_{NROCD} = L_{light} + {}^{y}_{c} iD_{0} + D^{2} = (2m_{c})_{c} + {}^{y}_{b} iD_{0} D^{2} = (2m_{b})_{b} + :::;$$
 (3)

where L $_{\rm light}$ is the usual relativistic lagrangian for gluons and light quarks. The 2-com ponent eld $_{\rm c}$ annihilates charm quarks, while $_{\rm b}$ creates bottom antiquarks. The typical velocity v of the charm quark in the meson provides a small parameter that can be used as a non-perturbative expansion parameter. Lattice simulations using the terms given explicitly in (3) can be used to calculate matrix elements for the B $_{\rm c}$ with errors of order v^2 . To obtain higher accuracy, additional terms, represented by ::: in (3), must be included. There are 8 terms that must be added to decrease the errors to order v^4 .

To express the decay constant f_{B_c} in terms of NRQCD matrix elements, we must express the axial-vector current b $_5c$ in terms of NRQCD elds. Only the = 0 component contributes to the matrix element (1) in the rest frame of the B_c . This component of the current has an operator expansion in terms of NRQCD elds:

$$b^{0}_{5}c = C_{0} (m_{b}; m_{c})^{y}_{b} + C_{2} (m_{b}; m_{c}) (D_{b})^{y}_{c} + :::;$$
 (4)

The short-distance coe cient C $_0$ and C $_2$ can be determined by matching perturbative calculations of the matrix elements in full QCD and NRQCD. A convenient choice for matching is the matrix element between the vacuum and the state jobi consisting of a c and a b on their perturbative mass shells with nonrelativistic four-momenta p and p^0 in the center of momentum frame: $p + p^0 = 0$. The matching condition is

$$\text{hOjb} \,\, ^0 \,\, _5 \text{cjdbi} \,\, = \, \text{C}_0 \,\, \text{hOj} \,\, ^{\text{Y}}_{\text{b}} \,\, _{\text{c}} \text{jdbi} \,\, + \,\, \text{C}_2 \,\, \text{hOj(D} \,\, _{\text{b}})^{\text{Y}} \,\, \, \text{D}_{\text{c}} \text{jdbi} \,\, + \,\, :::; \quad \text{(5)}$$

where P-QCD and P-NRQCD represent perturbative QCD and perturbative NRQCD, respectively. At leading order in $_{\rm s}$, them atrix element on the left side of (5) is $v_{\rm b}$ ($_{\rm p}$) $_{\rm 5}u_{\rm c}$ (p). The D irac spinors are

$$u_{c}(p) = \frac{s}{\frac{E_{c} + m_{c}}{2E_{c}}} = \frac{!}{\frac{p}{E_{c} + m_{c}}};$$
 (6)

$$v(p) = \frac{s}{\frac{E_b + m_b}{2E_b}} = \frac{(p)}{\frac{E_b + m_b}{E_b + m_b}} ; \qquad (7)$$

where and are 2-component spinors and $E_Q = m_Q^2 + p^2$. Making a nonrelativistic expansion of the spinors to second order in $p=m_Q$, we nd

$$v_b(p)^0 = v_b(p) = v_b^y = 1 = v_b^y = v_b^$$

At leading order in $_{\rm s}$, the m atrix elements on the right side of (5) are $_{\rm b}^{\rm y}$ c and p² $_{\rm b}^{\rm y}$ c. The short distance one cients are therefore C $_{\rm 0}$ = 1 and

$$C_2 = \frac{1}{8 \operatorname{m}_{\text{red}}^2}; \tag{9}$$

where m $_{red}$ = m $_{b}$ m $_{c}$ = (m $_{b}$ + m $_{c}$) is the reduced m ass.

To determ ine the short distance one cients to order $_{\rm S}$, we must calculate the matrix elements on both sides of (5) to order $_{\rm S}$. We will calculate the order- $_{\rm S}$ correction only for the one cient C $_{\rm O}$, since the contribution proportional to C $_{\rm C}$ is suppressed by v². The one cient C $_{\rm O}$ can be isolated by taking the limit p! 0, in which case the matrix element of (D $_{\rm D}$) D $_{\rm C}$ vanishes. We rst calculate the matrix element on the left side of (5) to order $_{\rm S}$. The relevant diagrams are shown in Fig. 2. The tree diagram in Fig. 2a gives a product of D irac spinors $v_{\rm D}$ $_{\rm O}$ $_{\rm S}$ uc. We calculate the loop diagrams in Fig. 2b-2d in Feynman gauge, using dimensional regularization to regularize both infrared and ultraviolet divergences. More enturn integrals are analytically continued to D = 4 2 spacetime dimensions, requiring the introduction of a regularization scale . The elects of the quark self-energy diagrams in Fig. 2b and 2c is to multiply the matrix element by the renormalization constants $^{\rm P}$ $_{\rm Z}$, where

$$q \frac{1}{Z_Q} = 1 + \frac{2}{3} + \frac{1}{4} \frac{1}{IIV} + \frac{1}{2} \frac{1}{IR} + \frac{3}{4} (\log 4) + \frac{3}{2} \log \frac{m_Q}{m_Q} + 1 :$$
 (10)

The subcript UV or IR on the 's indicates whether the divergence is of ultraviolet or infrared origin. Combining the propagators using the Feynman parameter trick and using the fact that the external quarks band care on-shell, the vertex correction from Fig. 2d can be reduced to the tree-level diagram in Fig. 2a multiplied by the factor

$$= \frac{64 \text{ i.s.}}{3} \int_{0}^{Z_{1}} dx \int_{0}^{Z_{1}x} dy^{2} \frac{d^{D}k}{(2)^{D}} \frac{1}{[k^{2} (xp^{0} yp)^{2} + i.]^{3}}$$

$$= 2(1 \text{ x y})p^{-0} + p \text{ xm}_{c}^{2} + y \text{ m}_{b}^{2} + (x + y) \text{ m}_{b} \text{ m}_{c}$$

$$+ 2(1 \text{) } (xm_{b} ym_{b})^{2} (1 \text{) } (xp^{0} yp)^{2} \frac{(1 - \frac{3}{2})}{(2 \text{)}} k^{2} \text{ : (11)}$$

A fter integrating over k, it is convenient to change variables from the Feynm an parameters to s=x+y and t=x=s. The s-integrals are trivial, but the t-integrals must be evaluated with care using the imprescription in the denominator. Setting $p^{-0}p^{-1}m_bm_c(1+v^2=2)$, where v is the relative velocity of the p and p and taking the p in it p is the integral reduces to

$$= \frac{2}{3} \cdot \frac{1}{2} \cdot \frac{1}{uv} + \frac{1}{IR} + \frac{3}{2} (\log 4) + \frac{2}{v} \cdot \frac{3 \cdot \frac{m_b}{m_b + m_c}}{3 \cdot \frac{m_b}{m_b + m_c}} \log \frac{m_c}{m_b + m_c} \log \frac{m_b}{m_b} + \frac{1}{v} \cdot \frac{1}{IR} + \frac{1}{2} \log \frac{2m_{red}v}{v} + \log 4$$
; (12)

where m $_{\rm red}$ = m $_{\rm b}$ m $_{\rm c}$ = (m $_{\rm b}$ + m $_{\rm c}$). Multiplying the tree-level matrix element by the vertex factor 1+ and by the renormalization constants $^{\rm p}\overline{\rm Z}_{\rm b}$ and $^{\rm p}\overline{\rm Z}_{\rm c}$, we obtain the nalanswer for the matrix element to order $_{\rm s}$:

h0 to
0
 5 cycloi $_{P-QCD} = v_{b} ^{0}$ $^{5}u_{c} + \frac{2}{3} \frac{^{u}}{v} + \frac{3}{2} \frac{m_{b}}{m_{b} + m_{c}} \log \frac{m_{b}}{m_{c}} = 3$

$$\frac{i}{v} \frac{1}{I_{R}} = 2 \log \frac{2m_{red}v}{v} + \log 4$$
:(13)

Note that the ultraviolet divergences have cancelled. The imaginary part of (13) arises because it is possible for the quark and antiquark created by the current to scatter on-shell.

We next compute the NRQCD matrix element on the right side of (5) to order $_{\rm s}$. The relevant Feynman diagrams are again shown in Fig. 2. The tree diagram in Fig. 2a gives

a product of Pauli spinors $\frac{y}{b}$ c. It is convenient to calculate the loop diagram s in Coulom b gauge. In this gauge, the coupling of transverse gluons to heavy quarks is proportional to the heavy quark velocity, and therefore does not contribute in the lim it v! 0. We therefore need only calculate the contribution from Coulomb exchange. The wavefunction renormalization factors associated with the diagram s in Fig. 2b and Fig. 2c are trivial: $\frac{p}{Z_b} = \frac{p}{Z_c} = 1$. The diagram in Fig. 2d reduces to a multiplicative correction to the tree-level matrix element from Fig. 2a:

$$= \frac{16 \text{ i s}}{3} 2^{\frac{Z}{2}} \frac{d^{D}q}{(2)^{D}} \frac{1}{q^{2}} \frac{1}{E_{b} + q_{b}} \frac{1}{(p+q)^{2} = 2m_{b} + i} \frac{1}{E_{c} q} \frac{1}{(p+q)^{2} = 2m_{c} + i};$$
(14)

where $E_Q = p^2 = 2m_Q$. The infrared-divergent integral has been dimensionally regularized. A fler using contour integration to integrate over the energy q_0 of the exchanged gluon, we not that (14) reduces to an integral over the gluon's three-m omentum:

$$= \frac{32 \quad sm_{\text{red}}}{3} \quad {}^{2} \quad \frac{d^{3} \quad q}{(2 \quad)^{3} \quad q} \quad \frac{1}{q^{2}} \frac{1}{q^{2} + 2p \quad q} \quad i$$
 (15)

Evaluating the regularized integral in (15), we obtain

$$h0j_{b}^{y}_{c}jcbi_{p-NRQCD} = y_{b}^{c} 1 + \frac{2}{3} \frac{z}{v} \frac{i}{v} \frac{1}{IR} 2 log \frac{2m_{red}v}{v} + log 4$$
; (16)

where v is the relative velocity of the b and c: $\dot{p} j = m_{red} v$. Note that the infrared divergences and the factors of 1=v are identical to those in the matrix element (13) for full QCD.

Comparing the matrix elements (13) and (16), we can read of the short-distance coefcient C $_{0}$ from the matching condition (5):

$$C_0 = 1 + \frac{s (m_{red})}{m_b + m_c} \frac{m_b m_c}{m_b + m_c} \log \frac{m_b}{m_c} 2:$$
 (17)

We have chosen m $_{\rm red}$ as the scale of the running coupling constant. To the accuracy of this calculation, any scale of order m $_{\rm b}$ or m $_{\rm c}$ would be equally correct. Setting m $_{\rm b}$ = 4.5 GeV and m $_{\rm c}$ = 1.5 GeV, we not that $_{\rm s}$ (m $_{\rm red}$) 0.34, and C $_{\rm 0}$ 0.85.

Our nalresult for the decay constant of the B c is

$$if_{B_c}M_{B_c} = C_0 h0j_{b_c}^{y_c} + C_2 h0j(D_b)^{y_c} D_c B_c i$$
: (18)

The short-distance coe cient C $_0$ is given to next-to-leading order in $_s$ in (17), while C $_2$ is given to leading order in (9). The uncertainties consist of the perturbative errors in the short distance coe cients and an error of relative order v^4 from the neglect of matrix elements that are higher order in v^2 . To achieve an error of order v^4 , the matrix element h0 j(D $_b$) y D $_c$ y y can be calculated by lattice simulations using those terms in the NRQCD lagrangian that are given explicitly in (3), but h0 j y $_b$ $_c$ y $_c$ in ust be calculated using an improved action. If an error of order v^2 is suicient accuracy, than the matrix element h0 j(D $_b$) y D $_c$ y $_c$ i can be dropped, and h0 j y $_b$ $_c$ y $_c$ i can be calculated using only those terms in the NRQCD lagrangian that are given explicitly in (3). The parameters in this lagrangian are $_s$ and the quark masses m $_b$ and m $_c$, all of which can be tuned so as to reproduce the spectroscopy of charmonium and bottom onium. These simulations should be very accurate for the intermediate case of the bc system. In the absence of any lattice calculations of the bc system, the matrix element h0 j y $_c$ y $_c$ ican be estimated using wavefunctions at the origin from nonrelativistic potential models:

$$\mathfrak{J}0j_{b}^{y} \circ \mathfrak{B} \circ i \mathfrak{f} \qquad 2M_{B} \circ \frac{3}{2} \mathfrak{R} (0) \mathfrak{f} :$$
 (19)

The factor of 2M $_{\rm B_{c}}$ takes into account the relativistic norm alization of the state $\beta_{\rm c}i$.

Although it may not be relevant to the B_c meson, it is interesting to consider the $\lim_b m_c$. In this $\lim_b m_c$. In this $\lim_b m_c$. In this $\lim_b m_c$, the perturbative correction in (17) contains a large logarithm of $\lim_b m_c$. Heavy Quark E ective Theory can be used to sum up the leading logarithms of $\lim_b m_c$. B]:

$$C(s)$$
! $\frac{s(s)}{s(m_b)}$! $\frac{s(s)}{s(m_b)}$! $\frac{s(s)}{s(s)}$ $1 + \frac{s(m_c)}{s(s)}$ $\log \frac{s}{m_c}$ 2; (20)

where n_f is the number of light avors, including the charm quark, and where is an arbitrary factorization scale.

We have calculated the short-distance QCD radiative correction to the leptonic decay rate of the B $_{\rm C}$ m eson. This calculation provides an illustration of the factorization methods

based on NRQCD that were developed in Ref. [4]. The decay constant f_{B_c} of the B_c was factored into short-distance coe cients multiplied by NRQCD matrix elements. The radiative correction to the coe cient of the leading matrix element decreases the leptonic decay rate of the B_c m eson by about 15%.

This work was supported in part by the U.S.D epartment of Energy, D ivision of High Energy Physics, under Grant DE-FG 02-91-ER 40684.

R eferences

- [1] T. Applequist and H. D. Politzer, Phys. Rev. Lett. 34, 43 (1975); A. de Rujula and S.L. Glashow, Phys. Rev. Lett. 34, 46 (1975); R. Barbieri, R. Gatto, and R. Kogerler, Phys. Lett. B 60, 183 (1976); V. A. Novikov, L. B. Okun, M. A. Shifman, A. J. Vainshtein, M. B. Voloshin, and V. J. Zakharov, Phys. Rep. C 41 (1978).
- [2] R. Barbieri, R. Gatto, and E. Rem iddi, Phys. Lett. B 61, 465 (1976); R. Barbieri, M. Cao, and E. Rem iddi, Nucl. Phys. B 162, 220 (1980); R. Barbieri, M. Cao, R. Gatto, and E. Rem iddi, Phys. Lett. B 95, 93 (1980); Nucl. Phys. B 192, 61 (1981).
- [3] G. T. Bodwin, E. Braaten, and G. P. Lepage, Phys. Rev. D 46, R 1914 (1992).
- [4] G.T. Bodwin, E. Braaten, and G.P. Lepage, Argonne preprint ANL-HEP-PR-94-24 (hep-ph 9407339).
- [5] W E. Caswell and G. P. Lepage, Phys. Lett. B 61, 465 (1976).
- [6] Y.-Q. Chen, A cadem ia Sinica preprint (Ph.D. thesis); C.-H. Chang and Y.-Q. Chen, Phys.Rev.D 49, 3349 (1994); E. Eichten and C. Quigg, Ferm ilab preprint FERM ILAB-PUB-94/032-T (hep-ph 9402210); V.V. Kiselev, A.K. Likhoded, and A.V. Tkabladze, Serpukhov preprint IFVE-94-51 (hep-ph 9406339).
- [7] C.H. Chang and Y.Q. Chen, Phys. Lett. B 284, 127 (1992); Phys. Rev. D 46, 3845 (1992); Phys. Rev. D 48, 4086 (1993); E. Braaten, K. Cheung, and T.C. Yuan, Phys. Rev. D 48, 5049 (1993); K. Cheung, Phys. Rev. Lett. 71, 3413 (1993); K. Cheung and

- T.C. Yuan, Phys. Lett. B 325, 481 (1994); V.V. K iselev, A.K. Likhoded, and M.V. Shevlyagin, Z. Phys. C 63, 77 (1994); S.R. Slabospitsky, Serpukhov preprint IFVE-94-53 (hep-ph 9404346); C.H. Chang, Y.Q. Chen, G.P. Han, and H.-T. Jiang, Beijing preprint A S-ITP-94-24 (hep-ph 9408242); A.V. Berezhnoi, A.K. Likhoded, and M.V. Shevlyagin, Serpukhov preprint IFVE-94-48 (hep-ph 9408284).
- [8] See, for example, B.Grinstein, Ann.Rev.Nucl.Part.Sci. 42, 101 (1992); M.Neubert, Phys.Rep. 245, 259 (1994).

Figure Captions

- 1. Diagram for the annihilation of B $_{\rm c}$ into lepton pairs via a virtual W $^+$. The shaded oval represents the wavefunction of the B $_{\rm c}$.
- 2. D iagram s for the m atrix elements h0 jb 0 $_5$ cjbbi in perturbative QCD and h0 j y_b $_c$ jbbi in perturbative NRQCD.

This figure "fig1-1.png" is available in "png" format from:

http://arxiv.org/ps/hep-ph/9501296v2