# Self-energy of Heavy Quark

M.A. Ivanov and T. Mizutani

## D epartm ent of P hysics V irginia Polytechnic Institute and State U niversity B lacksburg, VA 24061

M arch 25, 2024

#### Abstract

W e dem onstrate that to calculate the self-energy of a heavy quark in the heavy quark lim it (or the heavy ferm ion lim it in what is called the Baryon Chiral Perturbation Theory), the use of standard dimensional regularization provides only the quantum lim it: opposite to the heavy quark (or classical) lim it that one wishes to obtain. W e thus devised a standard ultraviolet/infrared regularization procedure in calculating the one- and two-loop contributions to the heavy quark self-energy in this limit. Then the one-loop result is shown to provide the standard classical Coulomb self-energy of a static colour source that is linearly proportional to the untraviolet cuto . All the two-loop contributions are found to be proportional  $\ln(=)$  where is the infrared cuto . Often only the contribution from the to bubble (light quarks, gluon and ghost) insertion to the gluon propagator has been considered as the O (  $_{\rm s}$ ) correction to the C oulom b energy to this order. O ur result shows that other contributions are of the same magnitude, thus have to be taken into account.

Perm anent address: Bogoliubov Laboratory of Theoretical Physics, Joint Institute for Nuclear Research, 141980 Dubna (Moscow region), Russia

### 1 Introduction

Recently, many discussions have been devoted to the de nition of the of heavy quark mass. This quantity is the basic parameter in the heavy quark physics. Generally (not only for the heavy quarks in the heavy quark lim it in which we are interested, in the follow ing) one of the most popular choices within the perturbative scheme is the pole mass. In reference [1] it has been shown that the pole mass is gauge independent and infrared nite at the two-bop level. It was checked at the three-bop level in [2] where the relation between the mass of the modied minimal subtraction scheme and the scheme-independent pole m as was also obtained. Later authors of work  $\beta$  argue that precise de nition of the pole mass in the heavy quark limit may not be given once nonperturbative e ects are included (a more analytical approach leading to the identical conclusion has been given independently in [4]). To demonstrate that, they adopted the standard e ective running constant (as a typical non-perturbative quantity) obtained from the sum m ation of leading logarithms to calculate the C oulomb energy of a static (or in nitely heavy) colour source. This has produced an auxiliary singularity for small momentum (the so-called, infrared renorm alon) which, they claim ed, leads to the uncertainty of order  $o_{CD} = m_0$  in the de nition of the pole mass. An important observation which served as one of the bases to this dem onstrative calculation is that the widely used dimensional regularization may not be used to analize self-energy diagrams in the heavy quark limit, in particular, for separating the infrared and ultraviolet domains. The best example, as we shall also present in the following section, is that this regularization method leaves out the linear divergence that de nes the self-energy of a static quark in the lowest order: the Coulom b energy. A loo it should be emphasized here that the Heavy Q uark L in it does make sense only after introducing a scale parameter to the theory with which the quark mass may be com pared, and there is som ething aw kward about it in the dimensional regularization method.

It may be useful to point out here that a quite analogous situation to the heavy quark lim it can be found in the problem of the classical lim it h ! 0 for the self-energy of an electron in QED [5]. It was found that in the lim it h ! 0, orm ore precisely for the electron C om pton wave length h=m c  $r_0$ , where parameter  $r_0$  regularizes sm all distances, the energy of the C oulom b eld by a change of radius  $r_0$  is reproduced in the second order of

1

perturabative theory whereas the higher order contributions were shown to vanish in this lim it.

Here we rst study the quantum and classical limits for the lowest order contributions to the quark self-energy and quark-gluon vertex, and show that the standard dimensional regularization method only reproduces the quantum limit. Since one may clearly identify the classical limit to be equal to the currently fashonable heavy quark limit, we could conclude that the dimensional regularization method is not suited for studying this case. Then in the following section we calculate the two-loop contributions to the quark self-energy in the heavy quark limit, exploiting the conventional ultraviolet and infrared regularization method. We have found that each contribution to this order is all proportional to  $\ln(=)$  where and are the ultraviolet and infrared cuto s, respectively and that, unlike in QED, these contributions do not mutually cancel out. We point out that the gluonic bubble diagram s (Fig.4d) (im plicitly) taken often as the majour contribution in some processes is of the same magnitude as those from vertex and quark self-energy diagram s (Figs. 4a-4c) for the self-energy in the heavy quark limit. In the last section some comments will be given regarding the alleged infrared renorm alon and the Baryon (or heavy ferm ion) Chiral Perturbation Theory.

### 2 One-loop diagram

The self-energy of quark with mass m in the second order of perturbative theory is de ned by the diagram in Fig.1,

$$m_2 = {}_2(m) = C_F g_S^2 \frac{d^4k}{(2)^4 i} \frac{1}{(k^2)} \frac{1}{m} \frac{1}{k} g_S^{-\frac{1}{2}} \frac{1}{m}$$
 (1)

where  $C_F = (N_c^2 - 1)=2N_c$ , for a gauge group SU ( $N_c$ ). First, we note that this expression coincides with the electrom agnetic self-energy except for the trivial factor  $C_F g_s^2$ . W hile the integral has no infrared divergence, it develops the linear ultraviolet divergence. W e are interested in this quantity in the heavy quark lim it: m ! 1 . A swasm entioned in [3], the dimensional regularization cannot serve this purpose. To make this point more clear, we calculate the quark self-energy by way of introducing an ultraviolet regularization,

$$\frac{1}{(k^2)} ! \frac{F(k^2)}{(k^2)} :$$
 (2)

Here, the function F ( $k^2 = {}^2$ ) is assumed to decrease rapidly in the Euclidean region  $k_E^2 = k^2 ! + 1$  providing the ultraviolet convergence of the Feynman integrals. Now, the heavy quark limit (or classical limit) means the case where m in comparison with the quantum limit: m . The both limits may be realized using the Mellin representation for the form factor F ( $k^2 = {}^2$ ) under calculation of the integral in eq. (1). We can write in the Euclidean region

$$F(k_{E}^{2}) = \sum_{n=0}^{X^{1}} c_{n} (n)^{n} (k_{E}^{2})^{n} = \frac{1}{2i} \sum_{+i1}^{Z^{1}} \frac{d}{\sin n} c(n) (k_{E}^{2});$$
(3)

where  $c(n) = c_n$  for n being positive integers, and 0 < < 0.5.

Finally, we have

$$m_{2} = C_{F} g_{s}^{2} \frac{1}{2i} \int_{+i1}^{Z_{i1}} \frac{d}{\sin} c(\frac{1}{2}) \frac{1}{2} \frac{d^{4}k}{(2)^{4}i} \frac{4m}{(k^{2})^{1}} \frac{2(k+6)}{(m^{2}-(k+p)^{2})}$$

$$= C_{F} g_{s}^{2} m \frac{1}{2i} \int_{+i1}^{Z_{i1}} \frac{d}{\sin} c(\frac{1}{2}) \frac{1}{2} \frac{d^{2}k}{(1-2)^{2}} \frac{d^{2}k}{(1-2)^{2}} \frac{d^{2}k}{(1-2)^{2}} \frac{d^{4}k}{(2-1)^{4}i} \frac{4}{(2-1)^{2}} \frac{2(1-2)}{(1-2)^{2}} \frac{d^{4}k}{(2-1)^{4}i} \frac{4}{(2-1)^{2}} \frac{2(1-2)}{(1-2)^{2}} \frac{d^{4}k}{(2-1)^{4}i} \frac{4}{(2-1)^{2}} \frac{2(1-2)}{(1-2)^{2}} \frac{d^{4}k}{(1-2)^{2}} \frac{d^{4}k}$$

To get the leading term in m = in the quantum limit, we move the integration counter to the right and take into account of the rst double pole at = 0. Assuming that F (0) = 1, we have

$$m_2^{qu} = \frac{3}{4} C_F m \ln \frac{2}{m^2} + 0$$
 (1) : (5)

To get the leading term in =m in the classical lim it, we move the integration counter to the left and take into account the rst pole at = 0.5. From the standard relation between the M ellin and its inverse transform ations

c( 0:5) = 
$$2 dt f(t^2);$$

wehave

$$m_2^{cl} = \frac{s}{2} C_F c(0.5) = \frac{s}{2} C_F dtF(t^2)$$
: (6)

If we compare the above result with the calculation of the expression in Eq.(1) devising the dimensional regularization,

$$m_2 = \frac{3}{4} C_F m N + \frac{5}{3} ln \frac{m^2}{2}$$
 (7)

where  $N = 1 = + \ln 4$ , (: the Euler-M ascheroni constant) and is the subtraction point, one can see that it only reproduces the quantum limit Eq.(5) for the quark self-energy by identifying as the ultraviolet cuto. Thus, the naive dimensional regularization does not allow to see the heavy quark limit.

Ordinarily, studies of heavy quark lim it may be carried out by using the expansion of quark propagator in the inverse quark mass:

$$S (f_{0} + f_{0}) j_{f_{0} = m} = i \frac{M_{0} + f_{0} = 2m}{[1 + i = 2m + f_{0}(k^{2} = m^{2})][k_{4} + 2m if_{0}(k^{2} = m^{2})]}$$
$$= \frac{i M_{0}}{k_{4} + i} + O(\frac{1}{m})$$
(8)

Here,  $k_4 = ik_0$ ,  $M_0 = (1=2)(I + {}^0)$ , and  $f(x) = (1=2)({}^p \overline{1+x} 1)$ . It is readily seen that the D irac spinor u (p) for quark m ay be chosen as

so that  $u(p)u(p) = u(p)^{-0}u(p) = 1$ , and  $u(p) \sim u(p) = 0$ .

In the second order: Eq.(1), upon introducing the ultraviolet regulator to the otherw ise linearly divergent expression, the leading contribution reads

$$m_{2}^{cl} = _{2}(m) = 2 _{s}C_{F} - \frac{Z}{(2)^{3}} \frac{d^{3}\tilde{\kappa}}{\tilde{\kappa}^{2}} - \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}}$$
 (9)

which obviously coincides with the Eq.(6). One can see that this expression is the classical C oulomb energy of a static color source. To give more physical meaning to Eq.(9), one may introduce the charge density de ned as

$$(\mathbf{r}^{2}) = \frac{Z}{(2)^{3}} e^{i\mathbf{\tilde{k}}\mathbf{r}^{q}} \overline{\mathbf{F}}(\mathbf{\tilde{k}}^{2})$$
 (10)

supposing that F ( $k^2$ ) > 0, and the electron (quark) "radius"  $r_0 = 1 = .$ 

Then, we have

$$m_{2}^{cl} = \frac{g_{s}^{2}}{2r_{0}}C_{F} \int_{F}^{ZZ} dr_{1} dr_{2} (r_{1}^{2}) \frac{1}{4} \frac{1}{\frac{1}{r_{1}} r_{2}} (r_{2}^{2})$$
(11)

in accordance with the classical result.

As was mentioned above, there is no infrared divergence in the second order, but it plays the crucial role in higher orders in perturbative expansion.

To understand its role, we compute the quark-gluon vertex induced by both quarkgluon and self-gluon interactions: Figs 2-3. For simplicity we om it all the common coefcients and color factors.

(a) Quark-gluon interaction (diagram in Fig.2):

By introducing the gluon mass serving to regularize the infrared singularity we get

$${}_{1}(p;p) = \frac{Z}{(2)^{4}i} \frac{d^{4}k}{2k^{2}} \frac{1}{k^{2}} \frac{(m + k + j_{0})}{m^{2}} \frac{(m + k + j_{0})}{(k + p)^{2}} \dot{j}_{p=m}$$
(12)

Using the form factor as in Eq. (2) one obtains

$$\frac{8}{1} (p;p) = \begin{cases} \frac{1}{(4-)^2} & 2 \ln \frac{2}{2} & \ln \frac{2}{m^2} + O(1) & \text{if } m \\ \frac{1}{(4-)^2} & 2 \ln \frac{2}{2} + O(1) & \text{if } m \end{cases}$$

O ne can see that the infrared singularity de nes the behavior of the vertex in the heavy quark limit. Again result obtained from the dimensional regularization for ultraviolet divergence corresponds only to the quantum case:

$$_{1}$$
 (p;p) =  $\frac{1}{(4)^{2}} 2 \ln \frac{2}{2} N + \ln \frac{m^{2}}{2} + O(1)$  : (13)

Note that if we do not introduce the infrared cuto within the dimensional regularization scheme, we not the auxiliary 1= pole in the expression for the vertex in Eq.(12) leading to

$$_{1}$$
 (p;p) =  $\frac{1}{(4)^{2}}$  2N<sup>IR</sup> N<sup>UV</sup> + 3 ln  $\frac{m^{2}}{2}$  + 0 (1) :

(b) Self-gluon interaction (diagram in Fig.3):

This contribution may be written as

$${}_{2}(p;p) = \frac{Z}{(2)^{4}i} \frac{d^{4}k}{(2)^{4}i} \frac{1}{(2 + k^{2})^{2}} \frac{2k^{2}}{2k^{2}} \frac{4k (k + 2m k) (k + p)^{2}}{m^{2} (k + p)^{2}} \frac{p (k + p)^{2}}{p}$$
(14)

This integral turns out to be free from infrared singularity. Using the ultraviolet form factor gives

$${}_{2} (p;p) = \begin{cases} 8 \\ \frac{1}{(4)^{2}} & 3 \ln \frac{2}{m^{2}} + O(1) & \text{if } m \\ \\ \vdots & O(\frac{1}{m}) & \text{if } m \end{cases}$$

It is readily seen that this diagram gives no contribution in the heavy quark lim it. A gain result obtained from the dimensional regularization for ultraviolet divergence corresponds only to the quantum case:

$$_{2}(p;p) = \frac{1}{(4)^{2}} 3N \qquad 3\ln\frac{m^{2}}{2} + O(1) :$$
 (15)

## 3 Two-loop D iagram s in the Heavy Quark Lim it

The contributions to the quark self-energy from the two-loops (or fourth order in coupling constant) are dened by the diagrams in Fig.4. As a consequence of our discussion in the previous section, we shall not adopt the dimensional regularization, but merely use the ultraviolet form factor Eq.(2) together with the infrared regularization of the gluon propagator in the following.

### (a) Diagram in Fig. 4a.

With and being the ultraviolet and infrared cuto masses, as in the previous section, one nds

$$m_4^{(a)} = {}_4^{(a)} (m) = C_4^{(a)} g_s^4 I_4^{(a)} ({}^2)$$
 (16)

where  $^{2}$   $^{2}=^{2}$ ;  $C_{4}^{(a)}=1$  (for QED), and  $t^{a}t^{b}t^{a}t^{b}=C_{F}=2N_{c}$  (for QCD). The integral  $I_{4}^{(a)}$  ( $^{2}$ ) is equal to

$$I_{4}^{(a)}(^{2}) = i \frac{\sum_{(E)}^{Z} \frac{d^{4}k_{1}}{(2)^{4}}}{(2)^{4}} \frac{d^{4}k_{2}}{(2)^{4}} \frac{F(k_{1}^{2})}{k_{1}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{2}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k_{1}^{2})}} \frac{1}{(k_{14} + k_{24} + \frac{E(k_{1}^{2})}{k_{24}^{2} + \frac{E(k_{1}^{2})}{k_{24}^{2} + \frac{E(k_{1}^{2})}{k_{2}^{2} + \frac{E(k$$

Here, to be de nite, the integration is over the dimensionless Euclidean variables denoted as (E) under the integration sign. The calculation of this integral is given in Appendix. We have

$$m_4^{(a)} = 2 \ _{s}C_F \ \frac{Z}{(2)^3} \frac{d^3 \kappa}{\kappa^2} \frac{F(\kappa^2)}{\kappa^2} \ \frac{s}{4} \frac{1}{N_c} \ln(1 = 2)$$
 (18)

(b) Diagram in Fig. 4b.

Here one obtains

$$m_4^{(b)} = 4^{(b)} (m) = C_4^{(b)} g_s^4 I_4^{(b)} (2^2)$$
 (19)

where  $C_4^{(b)} = if^{abc}t^at^bt^c = (N_c=2)C_F$  (of course there is no QED counterpart to this diagram). The integral  $I_4^{(b)}$  (<sup>2</sup>) is given as

$$I_{4}^{(b)}({}^{2}) = \frac{\sum_{(E)}^{Z} \frac{d^{4}k_{1}}{(2)^{4}} \frac{d^{4}k_{2}}{(2)^{4}} \frac{d^{4}k_{3}}{(2)^{4}} \frac{F(k_{1}^{2})}{(2)^{4}} \frac{F(k_{1}^{2})}{k_{1}^{2} + \frac{1}{2}} \frac{F(k_{2}^{2})}{k_{2}^{2} + \frac{1}{2}} \frac{F((k_{1} - k_{2})^{2})}{(k_{1} - k_{2})^{2} + \frac{1}{2}}}{\frac{N_{b}}{(k_{14} + i)(k_{24} + i)}}$$

$$(20)$$

with the num erator N  $_{\rm b}$  equal to

 $N_{b} = (f_{x_{1}} \ 2 \ f_{y_{2}})M_{0} \ M_{0} + M_{0} (f_{x_{1}} + f_{x_{2}})M_{0} + M_{0} \ M_{0} (f_{x_{2}} \ 2 \ f_{y_{1}}):$ 

Since the integration is over Euclidean variables here,  $\& = {}^{0}ik_{4} ~ \&$ . With this it is easy to check that u (p)N<sub>b</sub>u (p) = 0, which means that the contribution from the diagram in the Fig.4b behaves as =m for large m, thus vanishes in the heavy quark limit.

(c) Diagram in Fig. 4c.

This contribution may be written as

$$m_4^{(c)} = 4^{(c)} m = C_4^{(c)} g_s^4 I_4^{(c)} (2^2)$$
 (21)

where  $C_4^{(c)} = 1$  (for QED), and  $t^a t^a t^b t^b = C_F^2$  (for QCD). The integral  $I_4^{(c)}$  (<sup>2</sup>) is equal to

$$I_{4}^{(c)}(^{2}) = i \frac{Z}{(c)} \frac{d^{4}k_{1}}{(2)^{4}} \frac{d^{4}k_{2}}{(2)^{4}} \frac{F(k_{1}^{2})}{(2)^{4}} \frac{F(k_{1}^{2})}{k_{1}^{2} + 2} \frac{F(k_{2}^{2})}{k_{2}^{2} + 2} \frac{1}{(k_{14} + i)} \frac{1}{(k_{14} + k_{24} + i)} \frac{1}{(k_{14} + i)}$$

$$\frac{1}{(k_{14} + i)} \frac{1}{(k_{24} + i)} \frac{1}{(k_{14} + i)} = I_{4}^{(a)} (2)$$
(22)

Here, we have taken into account the mass renorm alization in the second order (the dark disk in Fig. 4c). From the above result, it is easy to see that in QED the contributions coming from Fig. 4a and Fig. 4c cancel. This is a crucial point in solving the problem of classical limit h ! 0 in QED [5] since the contribution to the self-energy of a massive ferm ion from the photon vacuum polarization is proportional to

<sub>m</sub> (k) / (g k<sup>2</sup> k k ) ln 1 + 
$$\frac{k^2}{m^2}$$

and vanishes in the in nite mass limit: m  $!\ 1$  .

ForQCD one nds

$$m_4^{(c)} = 2 \ _{s}C_F \ \frac{Z}{(2)^3} \frac{d^3\tilde{\kappa}}{\tilde{\kappa}^2} \frac{F(\tilde{\kappa}^2)}{4} \ \frac{s}{4} 2C_F \ln(1 = 2)$$
 (23)

(d) Diagram in Fig. 4d.

The contribution from the diagram s in F ig. 4d, involving the vacuum polarization by m assless quarks, gluons, and ghost is written as

$$m_{4}^{(d)} = {}_{4}^{(d)} (m) = 2 {}_{s}C_{F} \frac{Z}{(2)^{3}} \frac{d^{3}\tilde{\kappa}}{\tilde{\kappa}^{2}} \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}} ren (\tilde{\kappa}^{2})$$
(24)

In the above expression, the renormalized vacuum polarization reads (see Appendix for a som ewhat detailed calculation)

$$ren (\tilde{\kappa}^{2}) = \frac{s}{4} \int_{0}^{Z^{1}} d \ln 1 + (1 ) \frac{\tilde{\kappa}^{2}}{2} 2N_{c} 1 + 4 (1 ) 4n_{f} (1 )$$

$$= \frac{s}{4} \tilde{b} \ln \frac{\tilde{\kappa}^{2}}{2} + O (1)$$
(25)

upon expanding in small parameter  $^2$ . Here the appearance of the coe cient  $\tilde{D} = (10N_c \quad 2n_f)=3$  is due to the use of the Feynman gauge.

Summing up all the contributions from the one- and two-bops, one nds

$$m = 2 {}_{s}C_{F} {}^{Z} \frac{d^{3}\tilde{\kappa}}{(2)^{3}} \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}} 1 {}_{s} \frac{s}{4} N_{c} \ln(1 = {}^{2}) + \frac{b}{N_{c}} \ln(\tilde{\kappa}^{2} = {}^{2}) :$$

$$(26)$$

$$! 2 {}_{s}C_{F} {}^{Z} \frac{d^{3}\tilde{\kappa}}{(2)^{3}} \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}} 1 {}_{s} \frac{s}{4} \ln({}^{2} = {}^{2})N_{c} \frac{2}{3} + \frac{b}{N_{c}} :$$

Here, we express the nal result in term of the rst coe cient in the Gell-M ann-Low function b =  $(11N_c 2n_f)=3$ . By adopting the number (avor) of light quarks to be three:  $n_f = 3$ , we can see for  $N_c = 3$  that the O ( $_s$ ) corrections to the leading static C oulom b self-energy coming from the bubble insertions to the gluon propagator is of the sam e order of magnitude as those from other diagram s: Figs.4a-c, so the latter contribution should be included.

In ref. [3] the self-energy of the heavy quark in the heavy quark lim it was calculated by introducing the running constant  $_{\rm s}$  ( $k^2$ ) in the Coulomb energy expression: Eq.(9), and by introducing the explicit integration cuto in place of the ultraviolet form factor F ( $k^2 = {}^2$ ) (Fig.5),

$$m = \frac{8}{3} \frac{\frac{Z}{(2)^{3}}}{\frac{R^{2}}{5}} \frac{\frac{d^{3}\tilde{K}}{(2)^{3}}}{\frac{R^{2}}{5}}$$
(27)

with

$${}_{s}(\mathbf{\tilde{k}}^{2}) = \frac{{}_{s}({}^{2}_{0})}{1 ({}_{s}({}^{2}_{0})\mathbf{b}=4)\ln({}^{2}_{0}=\mathbf{\tilde{k}}^{2})}:$$
(28)

According to Ref.[3], the zero of the denom inator in the above expression de nes the position of the infrared renorm alon, leading to the uncertainty in the pole mass. Here, we are interested in the O  $\begin{pmatrix} 2 \\ 5 \end{pmatrix}$  contribution from Eq.(27),

$$m_{4} = -\frac{s}{2} \frac{b}{3} \int_{0}^{2^{1}} dt \ln \frac{1}{t^{2}} = -\frac{s}{2} \frac{b}{3} \int_{0}^{2^{1}} dt \ln \frac{1}{t^{2}} dt \ln \frac{1}{t^{2}}$$

Here one might regard  $_0$  as corresponding to our (in this respect one could argue that the value of the integration cuto in Eq.(27) should be di erent from the scale  $_0$  introduced in Eq.(28). This would introduce an additional contribution of a logarithm ic type to Eq.(29), which would however not prohibit the appearance of the alleged renormalon). This result may be compared with the part of the expression in Eq.(26) which is proportional to b=N c. Since we have not calculated the higher loop contributions, we cannot draw any de nite conclusion. However, in view of the di erence in our result: Eq.(26) and that of ref.[3], it might be possible that due to the contributions other than the bubble insertions in the gluon propagator the characteristics (position, etc.) of the alleged infrared renorm alon in the heavy quark self-energy would be di erent from the original one, or even the renorm alon might not be there.

### 4 D iscussion

To sum marize our present note, rst we have demonstrated that the popular dimensional regularization method yeilds only the quantum limit, thus unsuited for the calculation of the self-energy of the heavy quark in the heavy quark (or the classical) limit which is just the opposite of the quantum limit. Thus we employed next the conventional method for ultraviolet and infrared regularizations to calculate the one- and two-loop contributions to the self-energy of the heavy quark in the heavy quark limit. The one-loop result is found as the standard C oulom b-like self-energy of a static colour souce whose radius is characterized by the ultraviolet regularization mass of the theory. Then all the two-loop contributions have turned out to be proportional to  $\ln(=)$  with the infrared cuto

, while the corresponding contributions in QED internally cancel out to vanish [5]. Of those two-loop contributions, only the bubble insertion to the gluon propagator in the low rst order (C oulom b) contribution (Fig.4d) has been discussed offen. We have found that other contributions are of the same size and have to be taken into account.

From our study we have reached a couple of observations (possibly, speculations). The rst is that since the contributions depicted in Figs.4a-c are not very di erent in magnitude from the bubble insertions to the gluon propagator at the two-loop level, the infrared renorm alon in the heavy quark selfenergy resulting from the simple bubble chain summation, due to the adoption of the running coupling constant, m ight be modied regarding its position, etc. or m ight even be absent once a due consideration of the remaining higher-loop e ects are incorporated.

The second observation is concerned with the dimensional regularization. Our assertion regarding the inappropriateness of in this method in the context of the heavy quark limit should be valid also in what is called the heavy baryon chiral perturbation theory (HBCHPT), (see for example a review [6] and references therein) which combines the ordinary Chiral Perturbation theory (CHPT) for the octet peudoscalar mesons coupled with baryonic (nucleons, deltas, etc.) degrees of freedom. Here very offen the Lagrangian (or the S-matrix) for the system is expanded in the inverse power of the baryon mass(es), exactly like in the heavy quark expansion realized in the heavy quark limit. The the meson loop integrals arising from the meson-baryon interactions are conventionally carried out using the dimensional regularization. Now according to our study in this note, this method is only compatible with the opposite quantum limit ! Thus it is our feeling that the existing results in HBCHPT should be examined again carefully from the present context.

#### A cknow legem ents

The authors would like to acknow ledge a helpful discussion with A rkady Vainshtein. This work was supported in part by the grant  $D \to FG$  05-84ER 40143 provided by the U.S. Department of Energy.

## Appendix

(a) The calculation of the integral 
$$I_4^{(a)}$$
 ( <sup>2</sup>).

U sing the obvious equality in Eq. (17)

$$\frac{1}{k_{14} + k_{24} + i} \frac{1}{k_{24} + i} = \frac{1}{k_{24} + i} - \frac{1}{k_{14} + k_{24} + i} \frac{1}{k_{14}}$$

and the well-known form ula

$$\frac{1}{k_4 + i} = \frac{P}{k_4} \quad i \quad (k_4)$$

one nds

$$I_{4}^{(a)}({}^{2}) = \frac{1}{4} \frac{Z}{(2)^{3}} \frac{d^{3}\tilde{\kappa}}{(2)^{3}} \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}} J({}^{2})$$

where

$$J(^{2}) = 2 \int_{0}^{\frac{\pi}{2}} dt \frac{f(0)}{t^{2}} = 4 \int_{0}^{\frac{\pi}{2}} dt f^{0}(t^{2}):$$

W e de ne the function f ( $t^2$ ) as

$$f(t^{2}) = \frac{Z}{(2)^{3}} \frac{d^{3}\tilde{k}}{t^{2} + \tilde{k}^{2} + z^{2}}$$

U sing the M ellin representation: Eq.(3) for the form factor F  $(k^2)$  and integrating over  $\tilde{k}$ , one can get

$$J(^{2}) = \frac{p}{2} \frac{1}{2i} \sum_{i=1}^{Z-i} \frac{d}{\sin} c() \frac{(1-2)}{(i-1)} \sum_{j=0}^{Z^{1}} d(1) \sum_{j=0}^{Z^{1}} dtt^{2} + \sum_{j=1}^{Z^{1}} dtt^{2} + \sum_{j=1}^{Z^{2}} \sum_{j=1}^{Z^{2}} \frac{d}{\sin} c() (2) \frac{(1+1)}{2i} \frac{(1+1)}{2i} \frac{1}{4} \ln \frac{1}{2i};$$

for small . Finally, we have

$$I_{4}^{(a)}({}^{2}) = \frac{1}{(4)^{2}} \frac{d^{3} \tilde{\kappa}}{(2)^{3}} \frac{F(\tilde{\kappa}^{2})}{\tilde{\kappa}^{2}} \ln \frac{1}{2} + O(1):$$

(a) The calculation of  $ren (q^2)$ .

As an example of the calculation of diagrams in Fig.4, we here present the details of the calculation of the vacuum polarization  $ren (q^2)$  by massless gluons, ghosts, and quarks (see, insertions to the gluon propagator in Fig.4d). We proceed in adopting the Feynman gauge. We shall use the dimensional regularization in the intermediate calculations, but the nal result will not depend on the choice of the regularization.

We de ne the S-m atrix element describing the vacuum polarization as

$$S = \frac{i}{2} \sum_{n=1}^{Z Z} dx_1 dx_2 A_{1}^{c_1} (x_1) \sum_{n=1}^{c_1 c_2} (x_1 - x_2) A_{2}^{c_2} (x_2)$$

with its Fourier transform de ned as

$$\sum_{\substack{c_1 c_2 \\ 1 2}}^{c_1 c_2} (q) = dx e^{iqx} \sum_{\substack{c_1 c_2 \\ 1 2}}^{c_1 c_2} (x):$$

For simplicity we do not introduce new notation for this value.

The contribution coming from the gluons and ghosts may be written in the form

$$\sum_{1=2}^{C_1 C_2} (q) = g_s^2 f^{abc_1} f^{abc_2} d^{2} \frac{Z^1}{d} + \frac{Z^2}{(2)^{D} i [R^2 (1)^2 + N_{ghost}^{1-2})^2}$$

where

$$N_{ghon}^{12} = g^{12} (D \quad 1) \frac{6}{D} k^{2} + [1 \quad 4 \quad (1 \quad )]_{q}^{2}$$
$$+ [q^{2}g^{12} \quad q^{1}q^{2}] 6 \quad D + (4D \quad 6) \quad (1 \quad );$$

$$N_{ghost}^{12} = g^{12} \frac{2}{D}k^2 + 2(1)q^2 2(1)[q^2g^{12} q^1q^2]$$
:

First, we show that the apparent non-gauge invariant term proportional to  $g^{1/2}$  does vanish. For this objective we use the identity:

<sup>Z</sup> 
$$d^{D} k \frac{ak^{2} + bq^{2}}{[k^{2} q^{2}]^{2}} = \frac{1}{2} d^{D} k \frac{D a + (2 D)b}{[k^{2} q^{2}]}$$

which may be proved by using the following expression,

$$\frac{k^2}{[k^2 - q^2]^2} = k^2 \frac{d}{dk^2} \frac{1}{[k^2 - q^2]} = \frac{1}{2}k \frac{d}{dk} \frac{1}{[k^2 - q^2]};$$

and integrating it by parts. Then the coe cient of the apparently non-gauge invariant contribution becomes

$$\begin{array}{c} {}^{Z^{1}} \\ {}^{0} \\ {}^{0} \end{array} \overset{d}{(1 \ )} \overset{Z}{=} \frac{d^{D} k}{\left[ \begin{array}{c} k^{2} \\ k^{2} \end{array} \right] (1 \ ) d^{2} \right]} \\ = {}^{Z} \\ {}^{D} k \frac{1}{\left[ \begin{array}{c} k^{2} \\ k^{2} \end{array} \right] } \overset{Z^{1}}{0} d \left[ \begin{array}{c} (1 \ ) \right]^{D \ = 2 \ 2} \\ {}^{D \ = 2 \ 2} \end{array} \begin{array}{c} 1 + 4 \frac{(D \ 1)}{(D \ 2)} (1 \ ) \\ (D \ 2) \end{array} \begin{array}{c} (1 \ ) \end{array} \begin{array}{c} ) \\ = {}^{Z} \\ {}^{D} k \frac{1}{\left[ \begin{array}{c} k^{2} \\ k^{2} \end{array} \right] } \\ \frac{1}{\left[ \begin{array}{c} k^{2} \\ k^{2} \end{array} \right]} \end{array} \begin{array}{c} \frac{2 \left( D \ = 2 \ 1 \right)}{(D \ 2)} + 4 \frac{(D \ 1)}{(D \ 2)} \frac{2 \left( D \ = 2 \right)}{(D \ 2)} \end{array} \begin{array}{c} 0 : \end{array}$$

Finally one nds the gauge invariant form for the vacuum polarization including both gluons and ghosts

$$\sum_{\substack{c_1 c_2 \\ c_1 c_2}} (q) = f^{abc_1} f^{abc_2} [q^2 g^{1/2} - q^1 q^2] \quad gluon+ghost (q^2)$$

with gluon+ ghost being equal to

$$g_{\text{luon+ghost}}(q^2) = \int_{0}^{Z^1} d^2 \frac{Z}{(2)^{\text{D}}} \frac{d^{\text{D}}k}{(2)^{\text{D}}i} \frac{[\text{D} \quad 6 \quad 4(\text{D} \quad 2) \quad (1)}{[k^2 \quad (1 \quad )q_1^2]^2})$$

Note that this expression has both infrared and ultraviolet logarithm ic divergences in four dimension. To regularize the infrared divergence we introduce the gluon (ghost) mass to the denominator. Then the renormalized vacuum polarization is written as

$$\begin{array}{rcl} & \operatorname{ren}_{ghon+ghost}(q^2) & = & \operatorname{ghon+ghost}(q^2) & \operatorname{ghon+ghost}(0) \\ \\ & = & \frac{s}{4} \int_{0}^{Z^1} d \ln [1 & (1 & )\frac{q^2}{2}] 2N_c [1 + 4 & (1 & )] \\ \\ & = & \frac{s}{4} q^2 \int_{4^{-2}}^{Z^1} \frac{dm^2}{m^2} \frac{q^2}{1 + 2m^2} \frac{1}{m^2} \frac{q^2}{q^2} \frac{1}{m^2} \frac{1}{m^2} \frac{q^2}{q^2} \frac{1}{m^2} \frac{1}{m^2} \frac{q^2}{q^2} \frac{1}{m^2} \frac{1}{m^2} \frac{q^2}{q^2} \frac{1}{m^2} \frac{1}{m^2} \frac{1}{q^2} \frac{1}{q^2}$$

The calculation of the vacuum polarization by quarks turns out to go along the sim ilar line with the result identical to the above one up to a constant factor. The combined result is the one in Eq.(25). Note that in this equation the momentum is measured in units of

### References

- [1] R.Tarrach, Nucl. Phys. B 183, 384 (1981).
- [2] N Gray, D J Broadhurst, W Grafe. K Schilcher, Z Phys. C 48, 673 (1990).
- [3] I.I.Bigi, M A Shifman, N G Uraltsev, A I.Vainshtein, Phys. Rev. D 50, 2234 (1994).
- [4] M Beneke and V M Braun, Nucl. Phys. B 426, 301 (1994); hep-ph/9411229.
- [5] G.V.E. mov, M.A.Ivanov, O.A.M.ogilevsky Ann.ofPhys. 103, 169 (1977).
- [6] U.-G. Meissner, Lecture given at Indian Summer School on Electron Scattering o Nucleons and Nuclei, Prague, Czech, September, (1994);hep-ph/9411300.

### List of Figures


Fig. 1. The lowest order (one loop) quark self-energy diagram .


Fig. 2. An O ( $_{\rm s}$ ) correction to the quark-gluon vertex by quark-gluon interaction.

Fig. 3. an O (  $_{\rm s}$ ) correction to the quark-gluon vertex by gluon-gluon interaction.

Fig. 4. Quark self-energy diagrams, to two loops. The mass counter term from the lowest order contribution is indicated as a blob in D iagram c. In D iagram d inserted in the gluon propagator are the gluon, ghost and light quark loops, respectively.

Fig. 5. Bubble chain in the gluon propagator causing the infrared renorm alon, according to Ref. 3.










Fig.4a

Fig.4b




Fig.4d




Fig.5