CHIRAL DYNAM ICS IN NUCLEONS AND NUCLEI

 $V \cdot B emard^{1}$

Centre de Recherche Nucleaire et Universite Louis Pasteur de Strasbourg Physique Theorique, BP 28, F {67037 Strasbourg Cedex 2, France

N.Kaiser²⁾

Technische Universitat Munchen, Physik Department T 30 James-Franck-Strae, D {85747 Garching, Germany

Ulf-G.Meiner³⁾

Universitat Bonn, Institut fur Theoretische Kemphysik, Nussallee 14{16, D {53115 Bonn, Germany

em ail: ¹⁾bernard@cmhp4.in2p3.fr, ²⁾nkaiser@physik.tu-muenchen.de ³⁾m eissner@pythia.itkp.uni-bonn.de

ABSTRACT:

W e review the implications of the spontaneous chiral symmetry breaking in QCD for processes involving one, two or more nucleons.

Commissioned article for Int. J.M od. Phys. E

CRN 95/3 TK 951

February 1995

CONTENTS

1. INTRODUCTION				
2. CHIRAL SYMMETRY IN QCD	8			
 2.1. E lem entary introduction to chiral sym m etry 2.2. Three{ avor QCD 2.3. Chiral perturbation theory 2.4. M odelling the pion 	8 9 11 21			
3. THE PION {NUCLEON SYSTEM	26			
 3.1. E ective Lagrangian 3.2. Extrem e non {relativistic lim it 3.3. Renorm alization 3.4. Low {energy constants and the role of the (1232) 3.5. A spects of pion {nucleon scattering 3.6. The reaction N ! N 3.7. The pion {nucleon vertex 	26 33 38 42 46 50 54			
4.NUCLEON STRUCTURE FROM ELECTROW EAK PROBES	58			
 4.1. E lectrom agnetic form factors of the nucleon 4.2. Nucleon C om pton scattering 4.3. A xial properties of the nucleon 4.4. Threshold pion photo{ and electroproduction 4.5. Two{pion production 4.6. W eak pion production 	58 63 76 79 88 94			
 5. THE NUCLEON {NUCLEON INTERACTION 5.1. General considerations 5.2. Nucleon {nucleon potential 5.3. M ore than two nucleons 5.4. Three{body interactions between nucleons, pions and photons 5.5. Exchange currents 	102 102 106 112 114 118			
 6. THREE FLAVORS, DENSE MATTER AND ALL THAT 6.1. Flavor SU (3), baryon m asses and {term s 6.2. K aon nucleon scattering 6.3. The pion in m atter 6.4. M iscellaneous om issions 	122 122 130 134 138			
APPENDICES	142			

I. IN TRODUCTION

E ective eld theories (EFTs) have becom e a popular tool in particle and nuclear physics. An elective eld theory di ers from a conventional renormalizable ("fundamental") quantum eld theory in the following respect. In EFT, one only works at low energies (where "low" is de ned with respect to som e scale speci ed later) and expands the theory in powers of the energy/characteristic scale. In that case, renorm alizability at all scales is not an issue and one has to handle strings of non { renorm alizable interactions. Therefore, at a given order in the energy expansion, the theory is specified by a nite num ber of coupling (low {energy) constants (this allow seg. for an order{by {order renorm alization). All observables are param etrized in terms of these few constants and thus there is a host of predictions for m any di erent processes. O by jourly, at som e high energy this e ective theory fails and one has to go over to a better high energy theory (which again m ight be an EFT of som e fundam ental theory). The trace of this underlying high energy theory are the particular values of the low energy constants. The EFT presum ably studied in most detail is chiral perturbation theory (CHPT). The central topic of this review will be the application of this fram ework when nucleons (baryons) are present, with particular emphasis on processes with exactly one nucleon in the initial and one nucleon in the nal state. Before elaborating on these particular aspects of CHPT, it is useful to make some general comments concerning the applications of EFTs.

EFT scome into play when the underlying fundam ental theory containsm assless (or very light) particles. These induce poles and cuts and conventional Taylor expansions in powers of momenta fail. A typical example is QED where gauge invariance protects the photon from acquiring a mass. One photon exchange involves a propagator 1=t, with t the invariant four{momentum transfer squared. Such a potential can not be Taylor expanded. A classic example to deal with such e ects is the work of Euler and H eisenberg [1.1] who considered the scattering of light by light at very low energies, ! m_e , with ! the photon energy and m_e the electron mass. To calculate the scattering am plitude, one does not need fullQED but rather integrates out the electron from the theory. This leads to an elective Lagrangian of the form

$$L_{e} = \frac{1}{2} (\vec{E}^{2} - \vec{B}^{2}) + \frac{e^{4}}{360^{-2} m_{e}^{4}} (\vec{E}^{2} - \vec{B}^{2})^{2} + 7 (\vec{E} - \vec{B}^{2})^{2} + \dots$$
(1.1)

which is nothing but a derivative expansion since E and B contain derivatives of the gauge potential. Stated di erently, since the photon energy is small, the electrom agnetic

elds are slow ly varying. From eq.(1.1) one reads o that corrections to the leading term are suppressed by powers of $(!=m_e)^4$. Straightforward calculation leads to the cross section (!) $!^6=m_e^8$. This can, of course, also be done using fullQED [1.2], but the EFT caculation is much simpler. The results of [1.2] nicely agree with the ones making use of the EFT for ! m_e .

A similar situation arises in QCD which is a non{abelian gauge theory of colored quarks and gluons,

$$L_{QCD} = \frac{1}{4g^2} G^a G^{,a} + qi D q qM q$$

$$= L_{QCD}^0 + L_{QCD}^I$$
(12)

with $M = \text{diag}(m_u; m_d; m_s; :::)$ the quark mass matrix. For the full theory, there is a conserved charge for every quark avor separately since the quark masses are all di erent. However, for the rst three avors (u;d;s) it is legitim ate to set the quark masses to zero since they are small on a typical hadronic scale like e.g. the {meson mass. The absolute values of the running quark masses at 1 GeV are m_u ' 5 MeV, m_d ' 9 MeV, m_s ' 175 MeV, i.e. $m_u = M$ ' 0:006, $m_d = M$ ' 0:012 and $m_s = M$ ' 0:23 [1.3]. If one sets the quark masses to zero, the left{ and right{handed quarks de ned by}}

$$q_{\rm L} = \frac{1}{2} (1 \ _5) q; \quad q_{\rm R} = \frac{1}{2} (1 + _5) q$$
 (1:3)

do not interact with each other and the whole theory adm its an U (3) U (3) symmetry. This is further reduced by the axial anomaly, so that the actual symmetry group of three avormassless QCD is

$$G = SU(3)_L \quad SU(3)_R \quad U(1)_{L+R}$$
 (1:4)

The U (1) sym m etry related to baryon num ber conservation will not be discussed in any further detail. The conserved charges which come along with the chiral SU (3) SU (3) symm etry generate the corresponding Lie algebra. In the sixties and seventies, manipulations of the commutation relations between the conserved vector (L+R) and axial (vector (L-R) charges were called "PCAC relations" or "current algebra calculations" and lead to a host of low energy theorem s and predictions [1.4]. These rather tedious manipulations have now adays been replaced by EFT methods, in particular by CHPT (as will be discussed later on). Let us com e back to QCD. O ne quickly realizes that the ground state does not have the full symmetry G, eq.(1.4). If that were the case, every known hadron would have a partner of the sam e m ass but with opposite parity. C learly, this is in contradiction with the observed particle spectrum. Further argum ents that the chiral symmetry is not realized in the W igner{W eylmode are given in section 2. The physical ground state must therefore be asymmetric under the chiral SU $(3)_{\rm L}$ SU (3), [1.5]. In fact, the chiral symmetry is spontaneously broken down (hidden) to the vectorial subgroup of isospin and hypercharge, generated by the vector currents,

$$H = SU(3)_{L+R} \quad U(1)_{L+R}$$
 (1:5)

As mandated by Goldstone's theorem [1.6], the spectrum of massless QCD must therefore contain N $_{\rm f}^2$ 1 = 9 1 = 8 massless bosons with quantum numbers $\frac{3}{5}$ = 0 (pseudoscalars) since the axial charges do not annihilate the vacuum. Reality is a bit more complex. The quark masses are not exactly zero which gives rise to an explicit chiral symmetry breaking (as indicated by the term L_{QCD}^{I} in eq.(1.2)). This is in agreement with the observed particle spectrum { there are no massless strongly interacting particles. However, the eight lightest hadrons are indeed pseudoscalar mesons. These are the pions ($;^{0}$), the kaons (K $; K^{0}; K^{0}$) and the eta (). One observes that M M_K M which indicates that the masses of the quarks in the SU (2) subgroup (of isospin) should be considerably smaller than the strange quark mass. This expectation is borne out by actual calculation of quark mass ratios. A lso, from the relative size of the quark masses m_{u;d} m_s one expects the chiral expansion to converge much more rapidly in the two{ avor case than for SU (3)_f. These basic features of QCD can now be explored in a similar fashion as outlined before for the case of QED.

As already noted, the use of EFTs in the context of strong interactions preceeds QCD. The W and identities related to the spontaneously broken chiral symmetry were explored in great detail in the sixties in the context of current algebra and pion pole dom inance [1.4,1.7]. The work of Dashen and Weinstein [1.8], Weinberg [1.9] and Callan, Colem an, Wess and Zum ino [1.10] clari ed the relation between current algebra calculations and the use of e ective Lagrangians (at tree level). However, only with W einberg's [1.11] sem inalpaper in 1979 it became clear how one could system atically generate loop corrections to the tree level (current algebra) results. In fact, he showed that these loop connections are suppressed by powers of $(E =)^2$, with E a typical energy (four{ the scale below which the EFT can be applied (typically the mass m om entum) and of the rst non {G oldstone resonance, in QCD 'M). The method was system at ized by Gasser and Leutwyler for SU $(2)_f$ in Ref.[1.12] and for SU $(3)_f$ in Ref.[1.13] and has become increasingly popular ever since. The basic idea of using an elective Lagrangian instead of the full theory is based on a universality theorem for low energy properties of eld theories containing massless (or very light) particles. Consider a theory (like QCD)

at low energies. It exhibits the following properties:

L is symmetric under some Lie group G (in QCD: $G = SU(3_k)$ SU(3), SU(3),

The ground state j > is symmetric under H G (in QCD: H = SU (3)). To any broken generator of G there appears a massless Goldstone boson (called "pion") with the corresponding quantum numbers ($J^P = 0$ in QCD).

The Goldstone bosons have a nite transition amplitude to decay into the vacuum (via the current associated with the broken generators). Thism atrix element carries a scale F, which is of fundamental importance for the low energy sector of the theory (in QCD: < $0 A^{a} j^{b} > =$ ip ${}^{ab}F$, with F the pion decay constant in the chiral limit).

There exists no other massless (strongly interacting) particles.

 $E\,xp\,licit\,$ sym m etry breaking (like the quark m ass term in QCD) can be treated in a perturbative fashion.

M atter elds (such as the spin $\{1/2 \text{ baryons}\}$ can be incorporated in the EFT according to the strictures of non $\{1/2 \text{ baryons}\}$ realized chiral sym m etry. How ever, special care has to be taken about their m ass terms (see below).

Now any theory with these properties books the same (in more than two space-time dimensions). This means that to leading order the solution to the W ard identities connected to the broken symmetry is unique and only contains the scale F. Thus, the EFT to lowest order is uniquely xed and it is most econom ical to form ulate it in terms of the G oldstone elds [1.14]. In fact, one collects the pions in a matrix (valued function (generally denoted 'U') which transforms linearly under the full action of G. In QCD, a popular choice is U (x) = exp[i ^a ^a (x)=F] with ^a (a = 1;:::;8) the G ell{M ann matrices and U⁰(x) = RU (x)L^y under chiral SU (3)_L SU (3)_k (with L;R an element of SU (3)_{L;R}). A coordingly, the pion elds transform in a highly non{linear fashion. This is a characteristic feature of EFT s.

The inclusion of the lowest lying baryon octet in the EFT of the strong interactions again preceds QCD, see e.g.[1.4,1.7[1.11]. However, the rst systematic analysis of QCD G reen functions and current m atrix {elements due to G asser, Sainio and Svarc is much more recent [1.15]. They showed that the fully relativistic treatment of the spin { 1/2 m atter elds (the nucleons) spoils the exact one {to {one correspondence between the loop expansion and the expansion in sm allm om enta and quark m asses. This can simply be understood from the fact that the nucleon massm does not vanish in the chiral lim it and thus an extra scale is introduced into the problem . Stated di erently, nucleon four{ m om enta can never be small. This problem was overcom e by Jenkins and Manohar [1.16] who used methods borrowed from heavy quark EFT to eliminate the troublesom e baryon m ass term . This am ounts to considering the baryons as very heavy, static sources. Consequently, all the mass dependence is shu ed into a string of interaction vertices with increasing powers of 1=m and a consistent power counting scheme emerges. In this review, we wish to sum marize the developments which have taken place over the last few years, with particular emphasis on the two{ avor sector and processes with one nucleon line running through the pertinent Feynm an diagram s. To our opinion, these are the best studied processes from the theoretical as well as from the experimental side. However, there has also been considerable activity concerning processes involving two (or more) nucleons starting from the work of W einberg [1.17] plus extensions to the three{ avor case, dense m atter and m uch m ore. To sum m arize the present state of the art, we believe that to rigorously test the consequences of the spontaneous chiral sym m etry breaking of QCD in nucleon and nuclear studies, calculations to order 0 (E⁴ = 4) are m and atory in m any cases. On the experim ental side, the advances in m achine and detector technology have lead, are leading and will lead to m any m ore data of unprecedented accuracy. These will serve as a good testing ground of the chiral structure of QCD.

In ref.[1.15], the two- avor case was considered. However, the problem s related to the non (vanishing mass in the chiral lim it generalize straightforwardly to avor SU (3). In this introduction, we therefore casually switch between the term s 'nucleon' and 'baryon'.

A nother non {perturbative m ethod which is much used in studying baryon properties at low energies is lattice gauge theory (LGT). To our opinion, LGT has not yet reached a su cient accuracy to describe dynam ical processes such as pion production or C om pton scattering in the non {perturbative regim e. H ow ever, we would like to stress that one should consider these m ethods as com plem entary. For exam ple, one hopes that in the not too distant future LGT will signi cantly contribute by supplying e.g. num erical values for the pertinent low {energy constants.

The material is organized as follows. In section 2, we give an elementary introduction to chiral symmetry, discuss three{ avor QCD and give a brief account of CHPT for the meson sector. We also show how one can model the Goldstone pion in a quark m odel language. Section 3 contains the basic discussion of the pion { nucleon Lagrangian, its construction, the extrem e non { relativistic lim it and the renorm alization procedure to order E³. We give a complete list of the num erical values of the low {energy constants for the next {to { leading order e ective Lagrangian $L_{N}^{(2)}$ and sum m arize to what extent these values can be understood from a resonance exchange picture. The inclusion of the spin {3/2 decuplet, i.e. the (1232), as an active degree of freedom in the EFT is critically examined. Applications to pion { nucleon scattering and the reaction N! Ν are also discussed. Section 4 is devoted to the nucleon as probed by electroweak currents. We discuss in detail such topics as the electrom agnetic form factors, C om pton scattering, axial properties and, furtherm ore, single and double pion production with real and virtual photons as well as W {bosons. Together with section 3, this is the main body of the work presented in this review. Section 5 contains the extensions to system s with two and more nucleons. Here, a complication arises due to the appearance of IR divergences in reducible diagram swhich leads to a modi cation of the power counting scheme. This is discussed in some detail and the pertinent method of applying the chiral power counting only to the irreducible diagram s together with the solution of a Schrodinger or Lippm ann {Schwinger equation to generate the S{m atrix is then applied to the potential between two, three and four nucleons. Since the construction of the NN { potential from the chiral Lagrangian involves a large number of low {energy constants, it appears to be favorable for certain applications to supply as much phenom enological input as possible, i.e. by taking the two{body pion {nucleon and the nucleon {nucleon interaction suitably parametrized from phenomenology. The chiralmachinery is then used to provide the remaining three{body forces. A s an example pion{deuteron scattering is discussed. Sim ilarly, in the description of the meson exchange currents it is argued that the nuclear short{range correlations indeed suppress the badly known contact terms thus leading to a more predictive scheme than for the NN {potential. Section 6 contains extension to the three{ avor sector, kaon {nucleon scattering, the density {dependence of pion properties in m atter and gives a sum m ary of topics not treated in detail. M any of these developm ents are only in their infancy and we therefore have decided m ore to highlight the weak points than to give any details. However, the reader is supplied with su ciently many references on these topics to get a more detailed (and eventually less biased) picture. The appendices contain various technicalities such as a sum m ary of the

pertinent Feynm an rules or the de nition of loop functions which are needed for actual calculations. To keep the sections self{contained, the relevant references are given at the end of each section.

REFERENCES

- 1.1 H.Euler, Ann. Phys. (Leipzig) 26 (1936) 398;
 H.Euler and W.Heisenberg, Z.Phys. 98 (1936) 714.
- 1.2 R.Karplus and M.Neum an, Phys. Rev. 83 (1951) 776.
- 1.3 J.G asser and H. Leutwyler, Phys. Reports C 87 (1982) 77.
- 1.4 S. L. Adler and R. F. Dashen, "Current Algebras and applications to particle physics", Benjam in, New York, 1968.
- 1.5 Y.Nam bu and G.Jona (Lasinio, Phys. Rev. 122 (1961) 345; 124 (1961) 246.
- J.Goldstone, Nuovo Cim. 19 (1961) 154;
 J.Goldstone, A.Salam and S.W einberg, Phys. Rev. 127 (1962) 965.
- 1.7 H. Pagels, Phys. Reports 16 (1975) 219.
- 1.8 R.Dashen and M.Weinstein, Phys. Rev. 183 (1969) 1261.
- 1.9 S.W einberg, Phys. Rev. 166 (1968) 1568.
- 1.10 S.Coleman, J.W ess and B.Zumino, Phys. Rev. 177 (1969) 2239; C.G.Callan, S.Coleman, J.W ess and B.Zumino, Phys. Rev. 177 (1969) 2247.
- 1.11 S.W einberg, Physica 96A (1979) 327.
- 1.12 J.G asser and H. Leutwyler, Ann. Phys. (N.Y.) 158 (1984) 142.
- 1.13 J.G asser and H. Leutwyler, Nucl. Phys. B 250 (1985) 465.
- 1.14 H. Leutwyler, Ann. Phys. (N.Y.) 235 (1994) 165.
- 1.15 J.Gasser, M.E.Sainio and A.Svarc, Nucl. Phys. B 307 (1988) 779.
- 1.16 E. Jenkins and A.V. Manohar, Phys. Lett. 255 (1991) 558.
- 1.17 S.W einberg, Phys. Lett. 251 (1990) 288.

In this section, we rst discuss chiral sym m etry on an elementary level. We extend these considerations to three{ avor QCD and the form ulation of its elective low {energy eld theory in term s of the G oldstone bosons related to the spontaneous chiral sym m etry breaking. We also outline brie y how the structure of the pion can be m odeled in a four{ quark interaction cut{old theory of the Nam bu{Jona-Lasinio type.

II.1. ELEM ENTARY INTRODUCTION TO CHIRAL SYMMETRY

Before discussing full QCD, let us give a few very introductory remarks about chiral symmetry. The reader familiar with this concept is invited to skip this section. To be speci c, consider a free and massless spin $\{1/2 \text{ (Dirac)} \text{ eld,} \}$

$$L[] = i \quad 0 \quad (2:1)$$

The state of a free relativistic ferm ion (of arbitrary mass) is completely characterized by its energy E, its momentum p and its helicity $\hat{h} = \sim$ p=pj. For massless ferm ions helicity is identical to chirality with $_5$ the chirality operator (one speaks of chirality related to the greek word for "hand"). Let us decompose the spinor into a right{ and a left{handed component,}}

$$= \frac{1}{2} (1 _{5}) + \frac{1}{2} (1 + _{5})$$

= P_L + P_R
= _L + _R
(2:2)

O by iously, the operators $P_{L,R}$ are projectors,

$$P_{\rm L}^{2} = P_{\rm L}; P_{\rm R}^{2} = P_{\rm R}; P_{\rm L} \quad P_{\rm R} = 0; P_{\rm L} + P_{\rm R} = 1$$
 (2.3)

with the property*

$$\frac{1}{2}\hat{h}_{L;R} = \frac{1}{2}_{L;R}$$
(2:3)

This shows that the states $_{L;R}$ are helicity eigenstates. In terms of these elds, the Lagrangian (2.1) takes the form

$$L[_{L};_{R}] = i_{L} @_{L} + i_{R} @_{R}$$

$$(2:4)$$

O ne notices that the left { and right { handed ferm ion m odes do not com m unicate. Stated di erently, one can apply separate U $(1)_{L,R}$ transform ations which leave the Lagrangian invariant,

$$L ! e^{i_{L}} L; R ! e^{i_{R}}$$

$$(2:5)$$

^{*} For a massive ferm ion, the P_{L;R} are still projectors but do not yield exactly the helicity.

leading to conserved left{ and right{handed currents,

$$J^{I} = I \qquad I; \quad I = L; R \tag{2.6}$$

with

$$(0 J)^{i} = 0; I = L; R$$
 (2:7)

Equivalently, one can construct conserved vector and axial{vector currents,

$$V = ; @ V = 0$$

A = 5; @ A = 0 (2:8)

since $J_{L;R} = (V \land A)=2$. To reiterate, chiral sym m etry m eans that for m assless ferm ions chirality is a constant of m otion. A ferm ion m ass term explicitly breaks this sym m etry since it m ixes the left{ and right{handed com ponents,

$$M = {}_{L}M {}_{R} + {}_{R}M {}_{L}$$
(2:9)

To make chiral symmetry a viable concept for massive fermions, the corresponding eigenvalues of the mass matrix have to be small compared to a typical energy scale of the system under consideration. As an example, we will now consider the case of three{ avor Quantum chrom odynamics (QCD).

II.2. THREE {FLAVOR QCD

The standard m odel of the strong, electrom agnetic and weak interactions involves three generations of ferm ion doublets, alas six di erent quark avors. From these six quark types, three are labelled 'light' (u;d;s) and the other three 'heavy' (c;b;t). Here light and heavy refers to a typical hadronic scale M_H 1 G eV. In fact, $m_c > M_H$ and $m_{b,t} >> M_H$ whereas typical values of the light quark m asses at a renorm alization point of 1 G eV are [2.1]

$$m_d = 5$$
 2M eV; $m_d = 9$ 3M eV; $m_s = 175$ 55M eV (2:10)

Note that there exist some controversy about these values, for reviews with detailed references see e.g. [2.2,2.3,2.4]. In the three{ avor sector, the QCD Lagrangian takes the form

$$L_{QCD} = \frac{1}{2g^2}G^a G^{,a} + qi$$
 (@ iG)q qMq $\frac{1}{16^2}G^a G^{,a}$ (2:11)

with $q^{T}(x) = (u(x); d(x); s(x)), G$ the gluon eld, G the corresponding eld strength tensor and $G_{ia} = \frac{1}{2}$ G_{a} its dual. The last term in (2.11) is related to the strong

CP {problem. In what follows, we will set = 0. The quark mass matrix can be chosen to be diagonal,

$$M = diag(m_u; m_d; m_s)$$
(2:12)

In (2.11), we have not made explicit the generators related to the local SU $(3)_{colour}$ transformations. From the chiral symmetry point of view we rewrite (2.11) as

$$L_{QCD} = L_{QCD}^{0} \quad qM q \qquad (2:13)$$

and L_{OCD}^0 is invariant under the global transform ations of the group

$$G = SU(3)_L \quad SU(3)_R \quad U(1)_V \quad U(1)_R$$
 (2:14)

P rojecting onto left{ and right{handed quark elds, $q_{L;R} = (1=2)(1_{5})q$, these transform under the chiral group SU (3)_L SU (3)_k as

$$q_{\rm L} ! e^{iT^{a} \frac{a}{L}} q_{\rm L}; q_{\rm R} ! e^{iT^{a} \frac{a}{R}} q_{\rm R}; a = 1; ...; 8$$
 (2:15)

with the generators T^a (a=1,:::,8) given in terms of the G ell-M ann SU (3) m atrices via $T^{a} = {}^{a}=2$ with Tr (T^aT^b) = ${}^{ab}=2$. In what follows, we will not be concerned with the vectorial U (1) symmetry related to the baryon current q q and the anom alous U (1)_A current. It is believed that the axial U (1) is broken by instanton e ects [2.5]. To the global SU (3)_L SU (3)_k symmetry of L^{0}_{QCD} one associates $16 = 2 (N_{f}^{2} 1)$ conserved currents,

$$V^{a} = q T^{a}q; \quad (0 V)^{a} = 0$$

 $A^{a} = q 5T^{a}q; \quad (0 A)^{a} = 0$
(2:16)

with the corresponding conserved charges

$$Q_{V}^{a} = \begin{cases} Z \\ d^{3}xV_{0}^{i}(x); & \frac{dQ_{V}^{a}}{dt} = 0 \\ Z \\ Q_{A}^{a} = d^{3}xA_{0}^{i}(x); & \frac{dQ_{A}^{a}}{dt} = 0 \end{cases}$$
(2:17)

Of course, in the presence of quark mass terms, this symmetry is explicitly broken.

O ne m ight now ask the question whether this chiral symmetry is also manifest in the ground state or the particle spectrum of QCD? In fact, there are numerous indications that this is not the case. The realization of the chiral symmetry in the W igner mode (i.e. all generators de ned in (2.17) annihilate the vacuum) would lead to degenerate hadron doublets of opposite parity in plain contradiction to the observed spectrum. Furtherm ore, in the W igner phase the vector{vector and axial-vector{axialvector correlators in the ground state would be equal,

$$< 0 J^{a}(x) V^{b}(y) J^{b} > = < 0 J^{a}(x) A^{b}(y) J^{b} > :$$
 (2:18)

These correlators can be extracted from decay data, ! + n (n = 1;2;:::) with n even (odd) containing the information about the VV (AA) correlation function. As shown in refs.[2.6], the VV correlator strongly peaks around s ' 0.5 G eV² ' M ² whereas the AA correlator has a broad maximum around s ' 1.5 G eV² ' M ²_{A1}. From that and the approximate avour SU (3) symmetry of the hadron spectrum we conclude that the chiral symmetry is spontaneously broken down to its vectorial subgroup,

$$SU(3)_{L} SU(3)_{R} ! SU(3)_{V}$$
 (2:19)

with the appearance of N $_{\rm f}^2$ 1 = 8 m assless pseudoscalar m esons, the G oldstone bosons [2.7]. These are the analog to the spin waves in a ferrom agnet which underwent spontaneous m agnetization (thus breaking the rotational sym m etry of the m agnet H am iltonian). In nature, how ever, these Goldstone bosons are not exactly massless but acquire a small mass due to the explicit symmetry breaking from the quark masses, M $_{\rm p}^2$ м, where P is a generic symbol for the three pions, the four kaons and the eta. From the sytem at ics of the hadron spectrum, M ' M $_{\rm K}$ M we can immediately conclude that m_d ' m_u since the pions do not contain any strange quarks. These Goldstone m s bosons are in fact the lightest observed hadrons and they saturate the pertinent W ard identities of the strong interactions at low energies. To calculate QCD G reen functions in the non{perturbative regime, one therefore makes use of an elective eld theory (EFT) with the pseudoscalar m esons as the relevant degrees of freedom. The essential feature which makes this EFT amenable to a system atic perturbative expansion is the fact that the interaction between the Goldstone bosons at low energies is weak. To be m ore precise, consider the elastic scattering process + 0! + 0 (for m assless pions) [2.8]

T $(\overset{+}{}^{0} ! \overset{+}{}^{0}) = \frac{t}{F^{2}}$ (2:20)

with t the invariant four{m om entum transfer squared. Indeed, as t approaches zero, the G oldstone boson interaction vanishes. This fact is at the heart of the system atic low energy expansion in terms of sm all m om enta and quark m asses - chiral perturbation theory (CHPT) - as discussed in some detail in the next section. For a m ore detailed account see e.g. the m onograph [2.9], the original papers by G asser and Leutwyler [2.10] or the review [2.3].

II.3. CHIRAL PERTURBATION THEORY

In this section, we brie y review how to construct the elective chiral Lagrangian of the strong interactions at next{to{leading order, following closely the work of G asser and Leutwyler [2.10]. It is most economical to use the external eld technique since it avoids any complication related to the non{linear transform ation properties of the pions. The basic objects to consider are currents and densities with external elds coupled to them [2.11] in accordance with the symmetry requirements. The associated G reen functions autom atically obey the pertinent W ard identities and higher derivative term s can be constructed system atically. The S{m atrix elements for processes involving physical mesons follow then via standard LSZ reduction. To be specific, consider the vacuum {to{vacuum transition amplitude in the presence of external elds

$$e^{iZ [v;a;s;p]} = < 0 \text{ out j} in >_{v;a;s;p}$$
 (2.21)

based on the QCD Lagrangian

$$L = L_{QCD}^{0} + q(v(x) + {}^{5} a(x))q q(s(x) ip(x))q$$
(2.22)

The external vector (v), axial{vector (a), pseudoscalar (p) and scalar (s) elds are herm itean 3 3 m atrices in avor space. The quark mass matrix M (2.12) is contained in the scalar elds (x). The G reen functions of massless QCD are obtained by expanding the generating functional around v = a = s = p = 0. For the real world, one has to expand around v = a = p = 0; s(x) = M. The Lagrangian L is invariant even under local SU (3) SU (3) chiral transform ations if the quark and external elds transform as follow s:

$$q_{R}^{0} = Rq ; q_{L}^{0} = Lq$$

$$v^{0} + a^{0} = R (v + a) R^{y} + iR (R^{y})$$

$$v^{0} \quad a^{0} = L (v \quad a) L^{y} + iL (R^{y})$$

$$s^{0} + ip^{0} = R (s + ip) L^{y}$$
(2:23)

with L;R elements of SU $(3)_{L;R}$ (in general, these are elements of U $(3)_{L;R}$, but we already account for the axial anom aly to be discussed later). The path integral representation of Z reads:

$$e^{i\mathbb{Z} [v;a;s;p]} = \mathbb{D}G \mathbb{D}q\mathbb{D}qe^{id^4 \times L (q;q;G ;v;a;s;p)}$$
(2:24)

It allows one to make contact to the elective meson theory. Since we are interested in processes were the momenta are small (the low energy sector of the theory), we can expand the G reen functions in powers of the external momenta. This amounts to an expansion in derivatives of the external elds. This low energy expansion is not a simple Taylor expansion since the Goldstone bosons generate poles at $q^2 = 0$ (in the chiral lim it) or $q^2 = M^2$ (for nite quark masses). The low energy expansion involves two small parameters, the external momenta q and the quark masses M (or the Goldstone masses M ; M K; M). One expands in powers of these with the ratio M = q^2 xed. The elective meson Lagrangian to carry out this procedure follows from the low energy representation of the generating functional

$$e^{iZ [v;a;s;p]} = DU e^{id^{4}xL_{eff}(U;v;a;s;p)}$$
(2.25)

where the matrix U collects the pseudoscalar G oldstone elds. The low energy expansion is now obtained from a perturbative expansion of the meson EFT,

$$L_e = L_2 + L_4 + :::$$
 (2:26)

where the subscript (n = 2, 4, :::) denotes the low energy dimension (number of derivatives and/or quark mass term s). Let us now discuss the various term s in this expansion. The leading term (called L_2) in the low energy expansion (2.26) can easily be written down in terms of the mesons which are described by a unitary 3 3 matrix in avor space,

$$U^{y}U = 1$$
; det $U = 1$ (2.27)

The matrix U transforms linearly under chiral symmetry, $U^0 = R U L^{\gamma}$. The lowest order Lagrangian consistent with Lorentz invariance, chiral symmetry, parity, G {parity and charge conjugation reads [2.10]

$$L_{2} = \frac{1}{4}F^{2} Tr[r U^{y}r U + {}^{y}U + U^{y}]$$
(2.28)

The covariant derivative r U transforms linearly under chiral SU (3) SU (3) and contains the couplings to the external vector and axial elds,

$$r U = Q U i(v + a)U + iU(v a)$$
 (2:29)

The eld embodies the scalar and pseudoscalar externals,

$$= 2B (s + ip)$$
 (2:30)

There are two constants appearing in eqs.(2.28,2.30). The scale F is related to the axial vector currents, $A^a = F (a^a + ::: and thus can be identified with the pion decay constant in the chiral lim it, <math>F = F f f f + 0$ (M) g, by direct comparison with the matrix {element < $0jA^aj^b > = ip^{ab}F$.* The constant B, which appears in the eld , is related to the explicit chiral symmetry breaking. Consider the symmetry breaking part of the Lagrangian and expand it in powers of the pion elds (with p = 0, s = M so that = 2BM)

$$L_{2}^{SB} = \frac{1}{2}F^{2}BTrM (U + U^{Y})] = (m_{u} + m_{d})BF^{2} - \frac{2}{2} + \frac{4}{24F^{2}} + O(6)] + \dots (2:31)$$

where the ellipsis denotes the contributions for the kaons and the eta. The rst term on the right hand side of eq.(2.31) is obviously related to the vacuum energy,

 $[\]ast$ Strictly speaking the axial-axial correlator in the vacuum has a pion pole term with its residuum given by F .

while the second and third are meson mass and interaction terms, respectively. Since $H_{QCD} = Qm_q = qq$ it follows from (2.31) that

$$< 0juuj0 > = < 0jddj0 > = < 0jssj0 > = F^2Bf1 + 0 (M)g$$
 (2:32)

This shows that the constant B is related to the vev's of the scalar quark densities < 0 jqqj0 >, the order parameter of the spontaneous chiral symmetry breaking. The relation (2.32) is only correct modulo higher order corrections in the quark masses as indicated by the term 0 (M). One can furthermore read o the pseudoscalar mass term s from (2.31). In the case of isospin symmetry (m_u = m_d = m), one nds

$$M^{2} = 2m B f 1 + 0 (M) g = M^{2} f 1 + 0 (M) g$$

$$M_{K}^{2} = (m + m_{s})B f 1 + 0 (M) g = M_{K}^{2} f 1 + 0 (M) g$$

$$M^{2} = \frac{2}{3} (m + 2m_{s})B f 1 + 0 (M) g = M^{2} f 1 + 0 (M) g$$
(2:33)

with M $_{\rm P}$ denoting the leading term in the quark m ass expansion of the pseudoscalar m eson m asses. For the M $_{\rm P}$, the G ell{M ann {O kubo relation is exact, 4M $_{\rm K}^2 = M^2 + 3M^2$. In the case of isospin breaking, which leads to 0 m ixing, these m ass form ulae are som ew hat m ore com plicated (see e.g. ref.[2.10]). Eq.(2.33) exhibits nicely the G oldstone character of the pions { when the quark m asses are set to zero, the pseudoscalars are m assess and SU (3) SU (3) is an exact sym m etry. For sm all sym m etry breaking, the m ass of the pions is proportional to the square root of the sym m etry breaking param eter, i.e. the quark m asses. From eqs.(2.31) and (2.33) one can elim inate the constant B and gets the celebrated G ell{M ann {O akes{R enner [2.12] relations}}

$$F^{2}M^{2} = 2m < 0juj > + 0 (M^{2})$$

$$F_{K}^{2}M_{K}^{2} = (m + m_{s}) < 0juj > + 0 (M^{2})$$

$$F^{2}M^{2} = \frac{2}{3}(m + m_{s}) < 0juj > + 0 (M^{2})$$
(2:34)

where we have used $F_P = Ff1 + 0$ (M)g (P = ,K,), i.e. the di erences in the physical decay constants $F \notin F_K \notin F$ appear in the terms of order M² in eq.(2.34). From this discussion we realize that to leading order the strong interactions are characterized by two scales, namely F and B. Numerically, using the QCD sum rule value $< 0juuj > = (225 \text{ MeV})^3$ one has F'F' = 93 MeV and B' = 1300 MeV. The large value of the ratio B = F' = 14 has triggered some investigations of alternative scenarios concerning the mode of quark condensation [2.13].

O ne can now calculate tree diagram s using the e ective Lagrangian L_2 and derive with ease all so{called current algebra predictions (low energy theorem s). Current algebra is, as should have become evident by now, only the rst term in a system atic

low energy expansion. W orking out tree graphs using L_2 can not be the whole story { tree diagram s are always real and thus unitarity is violated. One has to include higher order corrections to cure this. To do this in a consistent fashion, one needs a counting scheme. The leading term in the low energy expansion of L_{e} (2.26) was denoted L_{2} because it has dimension (chiral power) two. It contains two derivatives or one power of the quark mass matrix. If one assumes the matrix U to be order one, U = O(1), a consistent power counting scheme for local terms containing U, @ U, v, a, s, p, ::: goes as follows. Denote by q a generic smallmomentum (for an exact de nition of 'sm all', see below). Derivatives count as order q and so do the external elds which occur linearly in the covariant derivative r U. For the scalar and pseudoscalar elds, it is most convenient to book them as order q^2 . This can be traced back to the fact that the scalar eld s(x) contains the quark m ass m atrix, thus s(x) М M^2 ở.₩ ith these rules, all terms appearing in (2.28) are of order q^2 , thus the notation L_2 (notice that a term of order one is a constant since $U^{y}U = 1$ and can therefore be disregarded. Odd powers of q clash with parity requirem ents). To sum m arize, the building blocks of all term s containing derivatives and/or quark m asses have the following dimension:

$$\begin{array}{l} (2.35) \\ (2.35$$

where we have introduced the eld strengths F^L;^R for later use. They are de ned via

$$F^{I} = Q F^{I} \quad QF^{I} \quad iF^{I}; F^{I}; I = L; R$$

 $F^{R} = v + a ; F^{L} = v \quad a$
(2:36)

As already mentioned, unitarity calls for pion loop graphs. Weinberg [2.14] made the important observation that diagram swith n (n = 1, 2, :::) meson loops are suppressed by powers of $(q^2)^n$ with respect to the leading term. H is rather elegant argument goes as follows. Consider the S{matrix for a reaction involving N_e external pions

$$S = (p_1 + p_2 + :::+ p_N) M$$
 (2:37)

with M the transition amplitude. Now M depends on the total momentum owing through the amplitude, on the pertinent coupling constants g and the renorm alization scale (the loop diagrams are in general divergent and need to be regularized*),

$$M = M (q; g;) = q^{D} f(q= ; g)$$
 (2:38)

^{*} It is advantegoeus to use dimensional regularization since it that case one avoids the appearance of power{law divergences.

with the total scaling dimension D of M given by [2.14]

$$D = 2 + N_{d} (d 2) + 2N_{L} :$$
 (2:39)

Here, N_L is the number of pion loops and N_d the number of vertices with d derivatives (or quark m ass insertions). The dom inant graphs at low energy carry the sm allest value of D. The leading term s with d = 2 scale like q² at tree level (N_L = 0), like q⁴ at one loop level (N_L = 1) and so on. Higher derivative term s with d = 4 scale as q⁴ at tree level, as q⁶ at one{loop order etc. This power suppression of loop diagram s is at the heart of the low energy expansion in EFT s like e.g. chiral perturbation theory (CHPT).

Up to now, we have been rather casual with the meaning of the word "sm all". By small momentum or small quark mass we mean this with respect to some typical hadronic scale, also called the scale of chiral sym m etry breaking (denoted by).Georgi and M anohar [2.15] have argued that a consistent chiral expansion is possible if 4 F ' 1 G eV. Their argum ent is based on the observation that under a change of the renorm alization scale of order one typical loop contributions (say to the scattering am plitude) will correspond to changes in the elective couplings of the order F 2 = 2 ' $1=(4)^2$. Setting = 4 F and cutting the logarithm ically divergent loop integrals at this scale, quantum corrections are of the same order of magnitude as changes in the renorm alized interaction terms. The factor $(4)^2$ is generic for one {loop integrals (in 3+1 dim ensions). A nother type of argum ent is related to the non {G oldstone spectrum . scattering in the I = J = 1 channel. There, at r = 770 M eV, one hits Consider the {resonance. This is a natural barrier to the derivative expansion of the Goldstone m esons and therefore serves as a cut o . The appearance of the signals the regime of the non {G oldstone particles and describes new physics. It is therefore appropriate to choose ' M ' 770 MeV, which is not terribly dierent from the previous estimate. In summary, small external momenta q and small quark masses M means q=M 1 and M = M1:

We have now assem beld all tools to discuss the generating functional Z at next{to{ leading order, i.e. at 0 (q^4). It consists of three di erent contributions: (1) The anom aly functional is of order q^4 (it contains four derivatives). We denote the corresponding functional by Z_A . The explicit construction was given by W ess and Zum ino [2.16] and can also be found in ref.[2.10]. A geom etric interpretation is provided by W itten [2.17]. (2) The most general electric Lagrangian of order q^4 which is gauge invariant. It leads to the action $Z_2 + Z_4 = d^4xL_2 + d^4xL_4$. (3) One loop graphs associated with the low est order term, L_2 . These also scale as term s of order q^4 .

Let us not discuss the anomaly functional Z_A . It subsumes all interactions which break the intrinsic parity and is responsible e.g. for the decay 0 ! 2. It also generates interactions between ve or more G oldstone bosons [2.17]. In what follows, we will not consider this sector in great detail (for a review, see ref.[2.18]).

W hat is now them ost general Lagrangian at order q^4 ? The building blocks and their low energy dimensions were already discussed { we can have term swith four derivatives

or with two derivatives and one quark m assor with two quark m asses (and, correspondingly, the other external elds). In SU (3), the only invariant tensors are g and , so one is left with (imposing also P,G and gauge invariance) [2.10]

$$L_{4} = \begin{array}{c} X^{10} & X^{2} \\ L_{i}P_{i} + & H_{j}P_{j} \\ & & \\$$

with

$$P_{1} = Tr(r \ U^{Y}r \ U)^{2}; P_{2} = Tr(r \ U^{Y}r \ U)Tr(r \ U^{Y}r \ U)$$

$$P_{3} = Tr(r \ U^{Y}r \ Ur \ U^{Y}r \ U); P_{4} = Tr(r \ U^{Y}r \ U)Tr(\ ^{y}U + \ U^{Y})$$

$$P_{5} = Tr(r \ U^{Y}r \ U)(\ ^{y}U + \ U^{Y}); P_{6} = Tr(\ ^{y}U + \ U^{Y})^{2}$$

$$P_{7} = Tr(\ ^{y}U \ U^{Y})^{2}; P_{8} = Tr(\ ^{y}U \ ^{y}U + \ U^{Y} \ U^{Y})$$

$$P_{9} = iTr(F^{R} r \ Ur \ U^{Y}) \ iTr(F^{L} r \ U^{Y}r \ U)$$

$$P_{10} = Tr(U^{Y}F^{R} \ UF^{L};)$$

$$P_{1} = Tr(F^{R} F^{R}; + F^{L} F^{L};); P_{2} = Tr(\ ^{y})$$

For the two avor case, not all of these terms are independent. The pertinent q^4 e ective Lagrangian is discussed in ref.[2.10]. The rst ten terms of (2.40) are of physical relevance for the low energy sector, the last two are only necessary for the consistent renorm alization procedure discussed below. These terms proportional to P_j (j = 1, 2) do not contain the G oldstone elds and are therefore not directly measurable at low energy constants L_i (i = 1; :::; 10) appearing in (2.40) are the so (called low (energy constants. They are not xed by the symmetry and have the generic structure

$$L_i = L_i^r + L_i^{inf}$$
(2:42)

These constants serve to renormalize the in nities of the pion loops (L_i^{inf}) and the remaining nite pieces (L_i^r) have to be xed phenom enologically or to be estimated by some model (see below). It should be noted that a few of the low (energy constants are in fact nite. At next{to{leading order, the strong interactions dynamics is therefore determ ined in terms of twelve parameters { B, F, L_1 ; :::, L_{10} (remember that we have disregarded the singlet vector and axial currents). In the absence of external elds, only the rst three terms in (2.40) have to be retained.

Finally, we have to consider the bops generated by the lowest order e ective Lagrangian. These are of dimension q^4 (one bop approximation) as mandated by W einberg's scaling rule. To evaluate these bop graphs one considers the neighbourhood of the solution U (x) to the classical equations of motion. In terms of the generating functional, this reads

$$e^{iZ} = e^{i d^{4}x [L_{2}(U) + L_{4}(U)]} DU e^{i d^{4}x [L_{2}(U) + L_{2}(U)]}$$
(2:43)

The bar indicates that the Lagrangian is evaluated at the classical solution. A coording to the chiral counting, in the second factor of (2.43) only the term L_2 is kept. This leads to Z.

$$Z = d^{4}x (L_{2} + L_{4}) + \frac{i}{2} \ln \det D \qquad (2:44)$$

The operator D is singular at short distances. The ultraviolet divergences contained in In detD can be determined via the heat kernel expansion. U sing dimensional regularization, the UV divergences in four dimensions take the form

$$\frac{1}{(4)^2} \frac{1}{d} \frac{1}{4} \operatorname{Sp}\left(\frac{1}{2}^{2} + \frac{1}{12}^{2}\right)$$
(2:45)

The explicit form of the operators ^ and ^ can be found in ref.[2.10]. Using their explicit expressions, the poles in h detD can be absorbed by the following renormalization of the low energy constants:

$$L_{i} = L_{i}^{r} + {}_{i}L; \quad i = 1; :::; 10$$

$$H_{j} = H_{i}^{r} + {}^{\sim}{}_{j}L; \quad j = 1; 2$$
 (2:46)

with

$$L = \frac{1}{16^{2}} d^{4} \frac{1}{d^{4}} \frac{1}{2} [\ln (4) + {}^{0}(1) + 1]$$

$$1 = \frac{3}{32}; \quad 2 = \frac{3}{16}; \quad 3 = 0; \quad 4 = \frac{1}{8};$$

$$5 = \frac{3}{8}; \quad 6 = \frac{11}{144}; \quad 7 = 0; \quad 8 = \frac{5}{48};$$

$$9 = \frac{1}{4}; \quad 10 = -\frac{1}{4}; \quad 1 = -\frac{1}{8}; \quad 2 = \frac{5}{24};$$

$$(2:47)$$

and is the scale of dimensional regularization. The q⁴ contribution $Z_4 + Z_{1 \ loop}$ is nite at d = 4 when expressed in terms of the renormalized coupling constants L_1^r and H_i^r . The next step consists in the expansion of the dimensional operator D in powers of the external elds. This gives the explicit contributions of the one{loop graphs to a given G reen function. The full machinery is spelled out in G asser and Leutwyler [2.10]. In general, one groups the loop contributions into tadpole and unitarity corrections. While the tadpoles contain one vertex and one loop, the unitarity corrections contain one loop and two vertices. The tadpole contributions also depend on the scale of the elective Lagrangian. Both of these loop contributions also depend on the scale of dimensional regularization. In contrast, physical observables are {independent. For actual calculations, however, it is sometimes convenient to choose a particular value of , say, = M or = M . To one{loop order, the generating functional therefore takes the form

$$Z = Z_2 + Z_4 + Z_{one \ loop} + Z_{anom}$$
(2:48)

and what remains to be done is to determ ine the values of the renormalized low energy constants, L_{i}^{r} (i = 1, :::, 10). These are in principle calculable from QCD, they depend on $_{QCD}$ and the heavy quark masses

$$L_{i}^{r} = L_{i}^{r} (Q_{CD}; m_{c}; m_{b}; m_{t})$$
(2:49)

In practice, such a calculation is not feasible. One therefore resorts to phenom enology and determines the L_i^r from data. However, some of these constants are not easily extracted from empirical information. Therefore, one uses constraints from the large $N_{\rm c}$ world. Using this and experimental information from scattering, $F_K = F$, the electrom agnetic radius of the pion and so on, one ends up with the values for the L_{i}^{r} (= M) given in table 1 (large N_c arguments are used to estimate L_{1} ; L_{4} and L_6). For comparison, we also give the values at = M. More accurate data will allow to further pin down this quantity. In the case of SU (2), one can de ne scale{ independent couplings 'i (i = 1;:::;7). These are discussed in ref.[2.10]. Can one now understand the values of the L^r_i from som e underlying principles? A lready in their 1984 paper, Gasser and Leutwyler [2.10] m ade the following observation. They considered an e ective theory of mesons coupled to the pseudoscalars. E lim inating the heavy eld by use of the equations of motion in the region of momenta much smaller than the m ass, one ends up with term s of order q^4 . The values of the corresponding low energy constants are given in term s of M and the {m eson coupling strengths to photons and pions. This leads to a fair description of the SU (2) low energy constants. This method has been generalized by Ecker et al. [2.19] and by D onoghue et al. [2.20]. They consider the lowest order e ective theory of Goldstone bosons coupled to resonance elds (R). These resonances are of vector (V), axial{vector (A), scalar (S) and non{Goldstone pseudoscalar (P) type. For the latter category, only the 0 is of practical importance. The form of the pertinent couplings is dictated by chiral symmetry* in terms of a few coupling constants which can be determined from data (from m eson {m eson and m eson { photon decays). At low momenta, one integrates out the resonance elds. Since their couplings to the G oldstone bosons are of order q^2 , resonance exchange produces term s of order q⁴ and higher. Sym bolically, this reads

So to leading order (q⁴), one only sees the m om entum {independent part of the resonance propagators,

$$\frac{1}{M_{R}^{2}} = \frac{1}{M_{R}^{2}} + \frac{t}{M_{R}^{2}} + \frac{t}{M_{R}^{2}}$$
(2:51)

^{*} For the vectors and axials, this naturally leads to the tensor{ eld form ulation.

and thus the L $_{\rm i}^{\rm r}$ (~' M $_{\rm R}$) can be expanded in term s of the resonance coupling constants and their m asses. This leads to

$$L_{i}^{r}() = \begin{array}{c} X \\ L_{i}^{Res} + \hat{L}_{i}() \\ R = V; A; S; P \end{array}$$
(2:52)

with \hat{L}_i () a remainder. For this scenario to make sense, one has to choose somewhere in the resonance region so that one can neglect the remainder. A preferred choice is = M (as shown in Ref.[2.19], any value of between 500 MeV and 1 GeV does the jbb). In table 1, we show the corresponding values for all low energy constants estimated from resonance exchange. It is apparent that the resonances almost completely saturate the L_i^r , with no need for additional contributions. This method of estimating L_i^r is sometimes called QCD duality or the QCD version of VMD. In fact, it is rather natural that the higher lying hadronic states leave their imprints in the sector of the light pseudoscalars { as already stated, the typical resonance mass is the scale of new physics not described by the G oldstone bosons.

i	Lŗ (M)	L _i r (1)	L ^{res}
1	0:9	0:5	0 : 7	0:5	0.6
2	1:6	0:4	1:2	0:4	1.2
3	3 : 6	1:3	3:	6 1 : 3	3 : 0
4	0:0	0:5	0:3	0:5	0.0
5	22	0:5	1:4	0:5	1.4
6	0:0	0:3	0:2	0:3	0.0
7	0 : 4	0:15	0:4	0:15	0:3
8	1:1	0:3	0:9	0:3	0.9
9	7 : 4	0:2	6 : 9	0:2	6.9
10	5 : 7	0:3	5:	2 0:3	6 : 0

Table 1: Low {energy constants for SU (3)_L SU (3)_R. The rst two columns give the phenom enologically determined values at = M and = M. The L_{i}^{r} (i = 4, :::, 8) are from ref.[2.10], the $L_{9;10}^{r}$ from ref.[2.21] and the $L_{1;2;3}^{r}$ from the recent determination in ref.[2.22]. The ' ' denotes the constants which are not renormalized. The third column shows the estimate based on resonance exchange [2.19].

In CHPT, the structure of any particle is made up by pion bops and higher resonance contributions encoded in the low (energy constants. As an example, consider the pion charge form factor. To lowest (tree) order, it is simply equal to unity as demanded by gauge invariance. At next order in the chiral expansion, loops and counterterms build up the pion radius, with the lions share due to one counterterm (L_9) which is saturated by vector meson exchange. At yet higher orders, one consistently sees more of the energy dependence of the pion form factor. However, in this perturbative approach one does not get the resonance (or similar elects in other channels). That is the reason why we argued that the scale of the resonance masses sets a natural cut o to the range of applicability of CHPT in the meson sector. A more detailed account of this and the many applications of CHPT can be found in the reviews s [2,2,3,2,18,2,22] and the connection of the elective Lagrangian to the QCD W ard identifies is elaborated on in ref.[2,24]. Before considering now the inclusion of matter elds in CHPT, let us brie y discuss the structure of the pion from a quark model point of view.

II.4. MODELLING THE PION

To investigate the form ation of vacuum condensates and the generation of m ass, N am bu and Jona-Lasinio [2.25] proposed a m odel with a H eisenberg {type four nucleon interaction in close analogy to developments in superconductivity. O ne can extend this approach to QCD where in the phase of the spontaneously broken chiral symmetry a scalar quark condensate form s and the quarks acquire a nite constituent m ass of the order of a few hundreds of M eV. The pion as the G oldstone boson appears as a collective quark {antiquark m ode. To discuss these features in som e detail, we follow closely the work of ref.[2.26]. Consider the two{ avor NJL Lagrangian for m assless quarks interacting via a contact force,

$$L_{NJL} = i (0 + G [()^2 (_{5} \sim)^2])$$
 (2:53)

Here, G is a positive coupling strength with the dimension of a squared length. The Lagrangian (2.53) is obviously invariant under chiral SU (2)_L SU (2)_k. It can be thought as a minimal elective Lagrangian mimicking some basic properties of non { perturbative QCD in the long{wavelength limit. One now solves the Dyson equation for the self{energy and identies with the mass M which is dynamically generated by the self{interactions. The resulting self{consistent equation relates M and the coupling G. It has a trivial solution M = 0 which corresponds to the ordinary perturbative

These constituent masses should not be confused with the fundamental mass parameters (current quark masses) entering the QCD Lagrangian. It should be stressed that the notion of a constituent quark is model (dependent but helps to understand qualitatively many features of the hadron properties.

result. However, for G above som e critical value, it has also a non {trivial solution which is determ ined by a self{consistency equation of the form

$$\frac{4N_{f}N_{c}+1}{(2)^{4}}G = d^{4}p\frac{1}{p^{2} - M^{2}+1} = 1$$
(2:54)

with N_f (N_c) the number of avours (colors). The integral in Eq.(2.54) diverges quadratically due to the zero range interaction. One therefore has to regularize the integral. Doing this e.g. by a covariant momentum cut{o $(2 p^2)$, one obtains

$$\frac{4^{2}}{13G^{2}} = 1 \quad \frac{M^{2}}{2} \ln \frac{2}{M^{2}} + 1 \quad : \tag{2:55}$$

Eq.(2.55) clearly exhibits that spontaneous sym m etry breaking only occurs for values of $G_{\rm crit}$. For such values, the mass M starts to deviate from zero and increases with G. The scalar quark condensate acquires a non {vanishing vev which can be interpreted as the probability of noting qq pairs in the vacuum,

< >=
$$\frac{3}{4^2}$$
 M³ $\frac{2}{M^2}$ ln $\frac{2}{M^2}$ + 1 (2:56)

which shows the intim ate relation between the constituent quark mass M and the quark condensate in this schematic model of chiral symmetry violation.



Fig. 2.1: Bethe{Salpeter equation in the Hartree{Fock approximation. The double line represents the pion, the solid lines constituent quarks. The exchange (Fock) diagram is not shown.

In studying the bound{state problem, one nds that the Bethe{Salpeter equation for the vertex function $_5$ (p) in the pseudoscalar isovector channel (shown in g2.1) is equivalent to the condition (2.54) when the total four{m om entum of the quark {antiquark pair is zero, $p^2 = 0$. This m eans that there exists a massless pseudoscalar isovector particle (the pion) related to the spontaneous chiral symmetry breaking. This nicely illustrates the G oldstone theorem in a microscopic picture. A sm and ated by G oldstone theorem, the pion has non{vanishing transition matrix{element into the vacuum via the pion}

axial current which de nes the pion decay constant. In the NJL model, it is related to the constituent quark mass via

$$F = \frac{p}{\frac{3}{2}}M \ln(1+u^2) \frac{u^2}{1+u^2} ; u = \frac{M}{M}$$
(2:57)

which shows that F $\,$ is linked to the collective nature of the pion. The model can, of course, also be treated in the case of explicit chiral sym metry breaking by adding the canonical current quark mass term .

W ith these basic tools, one can now study very di erent problem s related to the physics of the G oldstone bosons and other m esons (if one extends the basic Lagrangian accordingly). Som e of these are:

- The therm odynam ics of the constituent quarks and the pions, i.e. aspects of nite tem perature and density (in the approxim ation that the baryon density is given by three times the constituent quark density). Such character changes of meson properties play an important role in the nuclear equation of state and in hot and dense baryon {rich environments as precursors of the transition to the much discussed (but not yet observed) quark {gluon plasma. For an early reference see [2.27] and the recent reviews [2.28,2.29].
- 2) The extension of the model to the three{ avor case and the study of avorm ixing. This was not addressed in a system atic fashion in the paper by Bernard et al. [2.30] where it was shown that the U (1)_A anom aly forces the inclusion of term swith 2N_f ferm ion elds (within the one-loop approximation to the elds ective potential). Certain aspects of the physics of avorm ixing are reviewed in ref.[2.31].
- 3) The relation of NJL (type m odels to CHPT has been discussed early [2.32]. It has become clear that a direct comparison is hampered by the fact that in the NJL m odel one does not expand in terms of a small parameter. It can nevertheless serve as a guideline to understand the physics behind the low (energy constants (the extended NJL m odel) [2.33] and to get an estimate of p⁶ (and higher order) e ects [2.34].

Let m e brie y elaborate on the last point, i.e. the work of ref.[2.34]. There, a consistent bosonization scheme for the NJL model was developed and the p^2 expansion of certain observables was worked out. In table 2, we show the results for the pion mass, decay constant and the constituent mass. O ne sees that the p^4 approximation is within 1% of the total (Hartree(Fock) result.

0 rder	0 (p²)	0 (p ⁴)	0 (p ⁶)	Total
F	88.6	93.8	93.0	93.1
М	141.5	138.4	139.1	139.0
М	221.2	243.9	241.4	248.1

Table 2: Chiral expansion of the pion decay constant, the pion m ass and the constituent quark m ass to order p^2 , p^4 , p^6 in comparison to the self{consistent result of the bosonized N JL {m odel. A ll num bers are in M eV.

C learly, such results always have to be considered indicative since within the Hartree{Fock (one{loop e ective potential) approximation one does not include pion loops. W hat is most important here is to have a m icroscopic model for the spontaneous chiral symmetry breakdown and its associated G oldstone bosons.

REFERENCES

- 2.1 J.G asser and H. Leutwyler, Phys. Reports C 87 (1982) 77.
- 2.2 H. Leutwyler, in: Proc. XXVI Int. Conf. on High Energy Physics, Dallas, 1992, edited by J.R. Sanford, A IP Conf. Proc. No. 272, 1993.
- 2.3 Ulf-G.Meiner, Rep. Prog. Phys. 56 (1993) 903.
- 2.4 JF.Donoghue, TASI lectures, Boulder, USA, 1993;H.Leutwyler, Bern University preprint BUTP (94/8, 1994.
- 2.5 G. 't Hooft, Nucl. Phys. B 72 (1974) 461.
- 2.6 R.D.Peccei and J.Sola, Nucl. Phys. B 281 (1987) 1; C.A.Dom inguez and J.Sola, Z.Phys. C 40 (1988) 63.
- 2.7 J.Goldstone, Nuovo Cim. 19 (1961) 154.
- 2.8 S.W einberg, Phys. Rev. Lett. 17 (1966) 616.
- 2.9 J.F. Donoghue, E.Golowich and B.R.Holstein, "Dynamics of the Standard Model", Cambridge Univ. Press, Cambridge, 1992.
- 2.10 J.G asser and H.Leutwyler, Ann. Phys. (N.Y.) 158 (1984) 142; Nucl. Phys. B 250 (1985) 465,539.
- 2.11 D.G.Boulware and L.S.Brown, Ann. Phys. (N.Y.) 138 (1982) 392.
- 2.12 M. Gell-M ann, R. J. O akes and B. Renner, Phys. Rev. 175 (1968) 2195.
- 2.13 M.D. Scadron and H.F. Jones, Phys. Rev. D 10 (1974) 967;
 H.Szadijan and J.Stem, Nucl. Phys. B 94 (1975) 163;
 R.J.C rew ther, Phys. Lett. B 176 (1986) 172.
 N.H.Fuchs, H.Szadijan and J.Stem, Phys. Lett. B 238 (1990) 380;
 Phys. Lett. B 269 (1991) 183.
- 2.14 S.W einberg, Physica 96A (1979) 327.
- 2.15 A.M anohar and H.G eorgi, Nucl. Phys. B 234 (1984) 189.
- 2.16 J.W ess and B.Zum ino, Phys. Lett. 37B (1971) 95.
- 2.17 E.W itten, Nucl. Phys. B 223 (1983) 422.
- 2.18 J.Binnens, Int. J.M od. Phys. 8 (1993) 3045.
- 2.19 G. Ecker, J. Gasser, A. Pich and E. de Rafael, Nucl. Phys. B 321 (1989) 311.
- 2.20 J.F.Donoghue, C.Ram irez and G.Valencia, Phys. Rev. D 39 (1989) 1947.
- 2.21 J.Bijnens and F.Comet, Nucl. Phys. B 296 (1988) 557.

- 2.22 J.Bijnens, G.Ecker and J.Gasser, in: The DAFNE Physics Handbook (vol. 1), eds.L.Maiani, G.Pancheri and N.Paver, INFN Frascati, 1992
- 2.23 J.F.D onoghue, in "M edium energy antiprotons and the quark{gluon structure of hadrons", eds. R.Landua, J.M.R ichard and R.K lapish, Plenum Press, New York, 1992;

G.Ecker, in: Quantitative Particle Physics, eds. M. Levy et al., Plenum, New York, 1993;

H.Leutwyler, in \Recent A spects of Quantum Fields", eds. H.M itter and M.G austerer, Springer Verlag, Berlin, 1991.

"E ective eld theories of the standard model", ed. Ulf{G.Meiner, World Scientic, Singapore, 1992;

G.Ecker, \ChiralPerturbation Theory", preprint UW ThPh-94-49, 1995, to app. in Progr. Part. Nucl. Phys., hep-ph/9501357.

- 2.24 H. Leutwyler, Ann. Phys. (N.Y.) 235 (1994) 165; see also preprint BUTP-94/13, 1994.
- 2.25 Y.Nam bu and G.Jona {Lasinio, Phys. Rev. 122 (1961) 345; 124 (1961) 246.
- 226 V.Bemard, Phys. Rev. D 34 (1986) 1601.
- 2.27 V.Bemard, Ulf-G.Meiner and I.Zahed, Phys. Rev. D 36 (1987) 819.
- 2.28 S.K Levansky, Rev. M od. Phys. 64 (1992) 649.
- 2.29 T.Hatsuda and T.Kunihiro, Phys. Reports 247 (1994) 221.
- 2.30 V.Bernard, R.L.Ja e and Ulf-G.Meiner, Nucl. Phys. B 308 (1988) 753.
- 2.31 U.Vogland W.Weise, Prog. Part. Nucl. Phys. 26 (1991) 195.
- 2.32 T.H.Hansson, M.Prakash and I.Zahed, Nucl. Phys. B 335 (1990) 67; V.Bernard and Ulf(G.Meiner, Phys. Lett. B 266 (1991) 403;
 - C.Schuren, E.Ruiz Arriola and K.Goeke, Nucl. Phys. A 547 (1992) 612;
 - S.K levansky and J.M uller, Phys. Rev. C (1994) in print.
- 2.33 J.Bijnens, C.Bruno and E.de Rafael, Nucl. Phys. B 390 (1993) 501.
- 2.34 V.Bernard, A.A.O sipov and Ulf-G.Meiner, Phys. Lett. B 285 (1992) 119.

III. THE PION {NUCLEON SYSTEM

In this section, we will be concerned with the inclusion of baryons in the e ective eld theory. We will specialize to the case of two avors with the pions and nucleons as the asymptotically observed elds. The generalization to the case of three avors will be taken up later. First, we discuss the relativistic formulation. In that case, how ever, the additional mass scale (the nucleon mass in the chiral limit) destroys the one{to{one correspondence between the loop and the small momentum expansion. This can be overcome in the extrement of relativistic limit in which the nucleon is essentially considered as a static source. We will then turn to the system atic renormalization of the e ective pion{nucleon Lagrangian to order p^3 . Finally, we discuss the appearing low { energy constants and the role of the (1232) resonance. As applications, elastic pion{ nucleon scattering and the reaction N ! N at threshold are considered. Reactions involving electroweak probes are relegated to section 4.

III.1. EFFECTIVE LAGRANGIAN

In this section, we will be concerned with the inclusion of baryons in the e ective eld theory. The relativistic form alism dates back to the early days, see e.g. W einberg [3.1], C allan et al. [3.2], Langacker and Pagels [3.3] and others (for a review, see Pagels [3.4]). The connection to QCD G reen functions was performed in a system atic fashion by G asser, Sainio and Svarc [3.5] (from here on referred as GSS) and K rause [3.6]. A s done in the GSS paper, we will outline the form alism in the two{ avor case, i.e. for the pion{nucleon (N) system. The extension to avor SU (3) is spelled out in section 6.

Following GSS, we now discuss them odi cations of the procedure detailed in section 2 to include the nucleons. The starting point is the observation that the time {ordered nucleon matrix elements of the quark currents are generated by the one{nucleon to one{nucleon transition amplitude

$$F (p^{0}; p; v; a; s; p) = < p^{0} \text{ out } p \text{ in } >_{v; a; s; p}^{\text{connected}}; p^{0} \notin p$$
(3:1)

determ ined by the Lagrangian (2.22). Here, \dot{p} in > denotes an incom ing one{nucleon state of m on entum p (and sim ilarly \dot{p}^0 out >). The idea is now to construct in analogy with (2.25) a pion{nucleon eld theory which allows to evaluate the functional F at low energies.

First, we consider the general structure of the elective pion-nucleon Lagrangian L_{N}^{e} . It contains the pions collected in the matrix (valued eld U (x) and we combine the proton (p) and the neutron (n) elds in an isospinor

$$= \begin{array}{c} p \\ n \end{array} \qquad (32)$$

There is a variety of ways to describe the transform ation properties of the spin $\{1/2$ baryons under chiral SU (2) SU (2). All of them lead to the same physics. However,

there is one most convenient choice (this is discussed in detail in Georgi's book [3.7]). In the previous section, we had already seen that the self{interactions of the pions are of derivative nature, i.e. they vanish at zero momentum. This is a feature we also want to keep for the pion {baryon interaction. It calls for a non { linear realization of the chiral symmetry. Following W einberg [3.1] and CCW Z [3.2], we introduce a matrix { valued function K, and the baryon eld transforms as

K not only depends on the group elements $L; R 2 SU(2)_{L;R}$, but also on the pioneld (parametrized in terms of U(x)) in a highly non{linear fashion, K = K(L; R; U). Since U(x) depends on the space{time coordinate x, K implicitly depends on x and therefore the transformations related to K are local. To be more speciec, K is dened via

$$R u = u^{0}K$$
(3:4)

with $u^2(x) = U(x)$ and $U^0(x) = RU(x)L^y = u^{0^2}(x)$.* The transform ation properties of the pion eld induce a well{de ned transform ation of u(x) under SU(2) SU(2). This de nesK as a non{linear function of L, R and (x). K is a realization of SU(2) SU(2),

$$K = {}^{p} \overline{LU \,{}^{y}R \,{}^{y}R} \,{}^{p} \overline{U}$$
(3:5)

The somewhat messy object K 2 SU (2) can be understood most easily in terms of in nitesimal transformations. For $K = \exp(i_{a,a})$, $L = \exp(i_{a,a})\exp(i_{a,a})$ and $R = \exp(i_{a,a})\exp(i_{a,a})$ (with a, a, a real) one nds,

$$\sim = \sim [\sim ~~]=2F + O(\sim^{2}; \sim^{2}; \sim^{2})$$
(3:6)

which means that the nucleon eld is multiplied with a function of the pion eld. This gives some credit to the notion that chiral transform ations are related to the absorption or emission of pions. The covariant derivative of the nucleon eld is given by

$$D = 0 + \frac{1}{2} [u^{Y}; 0 u] = \frac{i}{2} u^{Y} (v + a) u = \frac{i}{2} u (v - a) u^{Y}$$
(3:7)

D transform show ogeneously under chiral transform ations, $D^{0} = K D K^{y}$. The object is the so{called chiral connection. It is a gauge eld for the local transform ations

$$^{0} = K K^{y} + K \Theta K^{y}$$
 (3.8)

^{*} W e adhere here to the notation of [3.5]. The more obvious one with interchanging L and R is e.g. used in [3.7].

The connection contains one derivative. One can also form an object of axial {vector type with one derivative,

$$u = i(u^{Y}r \ u \ ur \ u^{Y}) = ifu^{Y}r \ ug = iu^{Y}r \ Uu^{Y}$$
(3:9)

which transforms how ogeneously, $u^0 = K u K^y$. The covariant derivative D and the axial{vector object u are the basic building blocks for the lowest order e ective theory. Before writing it down, let us take a look at its most general form. It can be written as a string of term swith an even number of external nucleons, $n_{ext} = 0;2;4;:::$. The term with $n_{ext} = 0$ obviously corresponds to the meson Lagrangian (2.28) so that

$$L_{eff}[;;] = L + L + L + :::$$
 (3:10)

Typical processes related to these terms are pion{pion, pion{nucleon and nucleon{ nucleon scattering, in order. In this section, we will only be concerned with processes with two external nucleons and no nucleon loops (in section 5, we will also consider terms with n = 4),

$$L = L_{N} = (x)D(x)(x)$$
 (3:11)

The di erential operator D (x) is subject to a chiral expansion as discussed below. We now wish to construct the generating functional for the vacuum {to{vacuum transitions in the presence of nucleons. For doing that, we add external G rassm ann sources for the nucleon eld to the e ective Lagrangian,

$$L_{eff}[;;] = L + L_N + +$$
 (3:12)

From that, one de nes the vacuum {to{vacuum transition am plitude via

$$expfiZ^{v}[v;a;s;p; ;]g = N [dU][d][d] expi d^{4}x(L + L_{N} + +)$$

$$Z Z Z Z Z$$

$$= N^{0} [dU]expi d^{4}xL + i d^{4}xd^{4}y (x)S(x;y) (y) (detD)$$
(3:13)

where S is the nucleon propagator in the presence of external elds,

$$D S (x;y;U;v;a;s;p) = {}^{(4)} (x y) : (3:14)$$

Evaluating the functional Z at = 0; det D = 1 (i.e. no nucleon loops) one recovers the functional Z, eq.(2.25). Furtherm ore, the leading order terms in the low {energy expansion of F is generated by the tree graphs in Z. How ever, the relation between F and Z beyond leading order is much more complicated due to the fact that the nucleon m ass does not vanish in the chiral lim it as discussed below.

Let us rst consider the e ective pion {nucleon Lagrangian to lowest order. Its explicit form follows simply by combining the connection and the axial {vector u

(which are the objects with the least number of derivatives) with the appropriate baryon bilinears

$$L_{N}^{(1)} = D^{(1)}$$

= (i D m + $\frac{g_{A}}{2}$ 5u) (3:15)

The e ective Lagrangian (3.15) contains two new parameters. These are the baryon mass m and the axial (vector coupling g_A in the chiral lim it,

$$m = m [1 + 0 (m^{2})]$$

$$g_{A} = g_{A} [1 + 0 (m^{2})]$$
(3:16)

Here, m = 939 M eV denotes the physical nucleon m ass and g_A the axial{vector strength m easured in neutron {decay, n ! pe _e, g_A ' 1:26. The fact that m does not vanish in the chiral limit (or is not small on the typical scale ' M) will be discussed below. Furtherm ore, the actual value of m, which has been subject to m uch recent debate, will be discussed in the context of pion{nucleon scattering. The occurence of the constant g_A is all but surprising. W hereas the vectorial (avor) SU (2) is protected at zero m om entum transfer, the axial current is, of course, renorm alized. Together with the Lagrangian L⁽²⁾ (2.28), our low est order pion{nucleon Lagrangian reads:

$$L_{1} = L_{N}^{(1)} + L^{(2)}$$
(3:17)

To understand the low {energy dimension of L $_{\rm N}^{(1)}$, we have to extend the chiral counting rules of section 2 to the various operators and bilinears involving the baryon elds. These are:

$$m = O(1); ; = O(1); D = O(1); = O(1)$$

= O(1); 5 = O(1); = O(1); 5 = O(1); (3:18)
(ib m) = O(p); 5 = O(p)

Here, p denotes a generic nucleon three {m om entum. Since m is of order one, baryon four {m om enta can never be sm all on the typical chiral scale. Stated di erently, any time derivative D₀ acting on the spin {1/2 elds brings down a factor m. However, the operator (ib m) counts as order O (p). The proof of this can be found in ref.[3.6] or in the lectures [3.8]. To lowest order, the Goldberger {Treim an relation is exact, $g_A m = g_N F$, which allows us to write the N coupling in the more fam iliar form $g_N e^{-a}$.

It goes without saying that we have to include pion loops, associated with L_1 given in (3.17). Om itting closed ferm ion loops (det D = 1), the corresponding generating functional reads [3.5]

$$Z Z Z Z expfiZg = N^{0} [dU]exp i dxL^{(2)} + i dx S^{(1)}$$

$$D^{(1)ac}S^{(1)cb} = ab^{(4)}(x y)$$
(3:19)

with S⁽¹⁾ the inverse nucleon propagator related to D⁽¹⁾ (3.15). a, b and c are isospin indices. This generating functional can now be treated by standard methods. The details are spelled out by GSS [3.5]. Let us concentrate on the low {energy structure of the e ective theory which emerges. P ion loops generate divergences, so one has to add counterterm s. This am ounts to

$$L_{1} ! L_{1} + L_{2}$$

$$L_{2} = L_{N}^{(0)} + L_{N}^{(1)} + L_{N}^{(2)} + L_{N}^{(3)} + L^{(4)}$$
(3:20)

The last three terms on the rh.s. of eq.(3.20) are the expected ones. The structure of the N interaction allows for odd powers in p, so starting from $L_N^{(1)}$ to one{loop order one expects conterterms of dimension p^3 . A system atic analysis of all these terms has been given by K rause [3.6]. The rst two terms, $L_N^{(0)}$ and $L_N^{(1)}$, are due to the fact that the lowest order coe cients m and g_A are renormalized (by an in nite amount) when loops are considered. This indicates that the chiral counting is messed up and is completely di erent from the meson sector, the constants B and F are not renormalized in the chiral limit. The origin of this complication lies in the fact that the nucleon mass does not vanish in the chiral limit. To avoid any shift in the values of m and g_A one thus has to add appropriate counter terms

$$L_{N}^{(0)} = m \frac{m}{F}^{2}$$
; $L_{N}^{(1)} = g_{A} \frac{m}{F}^{2} \frac{1}{2}_{5}u$ (3.21)



Fig. 3.1: Chiral expansion for the N scattering amplitude, T_N $p^{N} = D$. Tree graphs contribute at N = 1, 2, 3 :::, n { loop graphs at N = 2,3, ::: (after m ass and coupling constant renorm alization). The contributions from 2,3,::: loops are analytic in the external m om enta at order p^{3} (here, p is a pion four-, nucleon three{m om entum or the pion m ass).

The rst term in (3.2.1) can be easily worked out when one considers the nucleon self(energy N (p) related to the nucleon propagator via S (p) = $[m \quad \beta \quad N(p)]^{1}$ in the one (loop approximation [3.5]). The low (energy structure of the theory in the presence of baryons is much more complicated than in the meson sector. This becomes most and N scattering amplitudes, T transparent when one compares the and T $_{\rm N}$, respectively. W hile $T^{\text{tree}} = p^2$ and $T^{n \text{ loop}}$ $(p^2)^{n+1}$, the corresponding behaviour for T_N is shown in g.3.1 [3.5]. Here, p denotes either a small meson four {m om entum or m ass or a nucleon three (m om entum . Tree graphs for T $_{\rm N}\,$ start out at order p followed by a string of higher order corrections p^2 , p^3 , \dots 0 ne{loop graphs start out at order p^2 (after appropriate m ass and coupling constant renorm alization) and are non {analytic in the external m om enta at order p^3 (in the chiral lim it $\mathfrak{m} = 0$). Higher loops start out at p^2 and are analytic to orders 0 (p^2 ; p^3). This again means that the low (energy constants) associated to L $_{\rm N}^{(2;3)}$ will get renorm alized. Evaluation of one { loop graphs associated with L_1 therefore produces all non {analytic term s in the external m on enta of order p^3 like e.g. leading threshold or branch point singularities. Let us now consider the case m° 🗧 0. Obviously, the Namplitude also contains terms which are non{analytic in the quark masses. A good example is the Adler {W eisberger relation in its di erential form { it contains a factor F 2 and therefore a term which goes like m lnm \cdot D ue to the com plicated low {energy structure of the m eson {baryon system, it has never been strictly proven that one { bop graphs generate all leading infrared singularities, in particular the ones in the quark masses. However, in all calculations performed so far the opposite has never been observed. In any case, the exact one{to{one correspondence between the loop and small momentum expansion is not valid in the meson {baryon system if one treats the baryons fully relativistically. This can be overcome, as will be discussed in the next section, in an extrem e non{relativistic limit. Here, however, we wish to point out that the relativistic form alism has its own advantages. Two of them are the direct relation to dispersion theory and the inclusion of the proper relativistic kinem atics in certain processes. These topics will be discussed later on.

The complete list of the polynom ial counter term s L $_{\rm N}^{(2;3)}$ can be found in ref.[3.6]. Here, let us just list the term s which will be used in the calculations of pion photo{ and electroproduction (see section 4). These are given by

$$L_{N}^{(2)} = c_{1} \quad Tr(+) + c_{6} \quad f^{+} + c_{7} \quad Tr(f^{+}) + ::: \qquad (3.22)$$

$$L_{N}^{(3)} = b_{0} \frac{1}{2F^{2}} \quad _{5} \quad u \quad Tr(+) + b_{11} \frac{g_{A}m}{F^{2}} \quad _{5} \quad \\ + b_{12} \frac{1}{F^{2}} \quad (i \quad D \quad m) \quad Tr(+) + d_{1} \frac{F}{2} \quad Tr(u \quad f^{+}) \quad \\ + d_{2} \frac{F}{2} \quad u \quad Tr(f^{+}) + d_{3}F \quad _{5}i \quad \overset{S}{D} \quad [u ; f^{+}] \quad \\ + d_{4}F \quad i \quad _{5} \quad [u ; f^{+}]$$

$$+ \frac{b_{9}}{F^{2}} D f^{+} + \frac{b_{9}}{F^{2}} Tr(D f^{+})$$

$$+ \frac{g_{A}}{12}b_{13} 5 D ;f] + \frac{i}{2}[u;f^{+}] + ::::$$
(3.23)

where

$$= u^{y} u^{y} u^{y} u^{y} u$$

$$f = u^{y}F^{R} u uF^{L} u^{y}$$

$$F^{L,R} = 0 F^{L,R} 0 F^{L,R} iF^{L,R}; F^{L,R}]$$

$$F^{R} = v + a; F^{L} = v a$$

$$A \overset{\$}{D} B = A (0 +)B A (0)B$$
(3.24)

In the case of having only photons as external elds, f simplies to f = e(0 A) $u^{Y}Qu$) with A the photon eld and Q = diag(1;0) the (nucleon) charge @ A)(uQ u^y matrix. Let us brie y discuss the physical signi cance of the various term s in the pion { nucleon Lagrangian, eqs(3.22,23). L $^{(2)}_{N}$ consists of three terms, the rst (q) is a mass renormalization counterterm and the second and third contribute to the anomalous magnetic moments $_{p,n}$. In GSS [3.5] it was demonstrated that to a high degree of 0, i.e. the isovector anom alous m om ent of the nucleon is given by the accuracy, c_6 loops (in the one (loop approximation). The terms of L $_{\rm N}^{\rm (3)}$ fall into two types. Let us is the term s proportional to b_{10} , b_{11} and b_{12} , in order. The b_{10} (term is needed for the renorm alization of g_A . The b_{11} (term contributes to the renorm alization of g_N and allows to reproduce the empirical deviation of the Goldberger{Treim an relation from unity. In what follows, we will always use g $_{\rm N}$ instead of $g_{\rm A}$ m = F . The term proportional to b_{12} enters the Z {factor which accounts for the renormalization of the external legs. The four terms in (3.23) proportional to d_i (i = 1;:::;4) are nite counterterm s which contribute to pion photo { and electroproduction. The coe cients d_1 ;:::; d_4 are not known a priori. In the N sector, there are three additional terms contributing to pion electroproduction at order q^3 , these are the last three in eq.(3.23). The two terms in eq.(3.23) proportional to b_0 and \tilde{b}_0 are related to the electric mean square charge radii of the proton and the neutron, see section 4.1. The last term in eq.(3.23) is related to the slope of the axial form factor of the nucleon, G_A (k^2) (see section 4.4). O ther term s of order q^2 and q^3 which enter the calculation of pion {nucleon scattering are discussed in ref.[3.5].

To end this section, a few remarks concerning the structure of the nucleons (baryons) at low energies are in order. Starting from a structureless D irac eld, the nucleon is surrounded by a cloud of pions which generate e.g. its anom alous magnetic moment (notice that the lowest order e ective Lagrangian (3.15) only contains the coupling of the photon to the charge). Besides the pion loops, there are also counterterms which encode the traces of meson and baryon excitations contributing to certain properties of the nucleon. Finally, one point which should be very clear by now: O ne can only make a rm statement in any calculation if one takes into account <u>all</u> terms at a given order. For a one{ loop calculation in the meson { baryon system , this amounts to the træe terms of order p, the loop contributions of order p², p³ and the counterterms of order p² and p³. This should be kept in mind in what follows.

III.2. EXTREME NON {RELATIVISTIC LIM IT

As we saw, the fully relativistic treatment of the baryons leads to severe complications in the low {energy structure of the EFT. Intuitively, it is obvious how one can restore the one{to{one correspondence between the loop and the small momentum expansion. If one considers the baryons as extremely heavy, only baryon momenta relative to the rest mass will count and these can be small. The emerging picture is that of a very heavy source surrounded by a cloud of light (alm ost massless) particles. This is exactly the same idea which is used in the so{called heavy quark eld theory methods used in heavy quark physics. Therefore, it appears natural to apply the insight gained from heavy quark EFT's to the pion{nucleon sector. Jenkins and M anohar [3.9,3.10]have given a new formulation of baryon CHPT based on these ideas. It am ounts to taking the extrem e non{relativistic lim it of the fully relativistic theory and expanding in powers of the inverse baryon mass. Notice also that already in the eighties G asser [3.11] and G asser and Leutwyler [3.12] considered a static source model for the baryons in their determ ination of quark mass ratios from the baryon spectrum.

Let us rst spell out the underlying ideas before we come back to the N system . Our starting point is a free D irac eld with mass m

$$L = (i \mathbf{0} \quad m) \quad (3.25)$$

Consider the spin $\{1/2 \text{ particle very heavy. This allow s to write its four {m on entum as}$

$$p = mv + 1$$
 (3.26)

with v the four{velocity satisfying $v^2 = 1$ and 1 a smallo {shellmomentum, v 1 m. One can now construct eigenstates of the velocity projection operator $P_v = (1 + 6)=2$ via

$$= e^{-h(n + h)}$$

6 H = H; 6 h = h (3.27)

which in the nucleon rest{fram ev = (1;0;0;0) leads to the standard non{relativistic reduction of a spinor into upper and lower components. Substituting (3.27) into (3.25) one nds

$$L = H (iv @)H h(iv @ + 2m)h H ig^{2}h + hig^{2}H$$
 (3:28)

with \mathbf{G}^2 the transverse part of the D irac operator, $\mathbf{G} = \mathbf{G}(\mathbf{v} \quad \mathbf{0}) + \mathbf{G}$. From eq.(3.28) it follows that the large component eld H obeys a free D irac equation (m aking use of the equation of m otion for h)

$$v = 0$$
 (3.29)

m odulo corrections which are suppressed by powers of 1=m . The corresponding propagator of H $\,$ reads

$$S(!) = \frac{i}{v + i}; > 0$$
 (3:30)

with ! = v k. The Fourier transform of eq.(3.30) gives the space{time representation of the heavy baryon propagator. Its explicit form $S'(t;r) = (t)^{(3)}(r)$ illustrates very clearly that the eld H represents an (in nitely heavy) static source. The mass{ dependence now resides entirely in new vertices which can be ordered according to their power in 1=m. A more elegant path integral form ulation is given by M annelet al. [3.13]. There is one more point worth noticing. In principle, the eld H should carry a label 'v' since it has a de nite velocity. For the purposes to be discussed we do not need to worry about this label and will therefore drop it.

Let me now return to the N system. The reasoning is completely analogous to the one just discussed. We follow here the system atic analysis of quark currents in avor SU (2) of Bernard et al. [3.14]. We will derive the elective Lagrangian for heavy nucleons in terms of path integrals. In this form ulation, the $1=m_N$ corrections are easily constructed. Consider the generating functional for the chiral Lagrangian of the N { system

where

$$S_{N} = \frac{2}{10} m + \frac{g_{A}}{2} 6_{5} + L_{N}^{(2)} + L_{N}^{(3)} + \cdots$$

$$S_{N} = \frac{2}{10} m + \frac{g_{A}}{2} 6_{5} + L_{N}^{(2)} + L_{N}^{(3)} + \cdots$$

$$S_{N} = \frac{2}{10} m + \frac{g_{A}}{2} 6_{5} + L_{N}^{(2)} + L_{N}^{(3)} + \cdots$$

$$(3:32)$$

The aim is to integrate out the heavy degrees of freedom. To this end the nucleon eld is splitted into upper and lower components with xed four velocity v

$$H_{v} = e^{im v} \frac{x^{2}}{2} (1 + 6)$$

$$h_{v} = e^{im v} \frac{x^{2}}{2} (1 - 6) :$$
(3:33)

In terms of these elds, the action S $_{\rm N}$ m ay be rewritten as

$$S_{N} = d^{4}x H_{v}AH_{v} + h_{v}BH_{v} + H_{v} _{0}B^{Y} _{0}h_{v} h_{v}Ch_{v} :$$
(3:34)

The operators A, B and C have the low energy expansions

$$A = A^{(1)} + A^{(2)} + \dots$$
(3.35)

where A $^{\rm (i)}$ is a quantity of O (q^i) , q denoting a low energy momentum . The explicit expressions read

$$A^{(1)} = i(v \quad D) + g_A (u \quad S)$$

$$A^{(2)} = \frac{m}{F^2} c_1 Tr_{+} + c_2 (v \quad \hat{u}) + c_3 u \quad u + {}_4c[S ; S]u \ u \\ + c_5 (+ \frac{1}{2}Tr_{+}) \quad \frac{i}{4m} [S ; S]((1 + c_6)f^+ + c_7 Trf^+)$$

$$B^{(1)} = iB^2 \frac{1}{2}g_A (v \quad u)_5$$

$$C^{(1)} = i(v \quad D) + n2 + g_A (u \quad S)$$

$$C^{(2)} = A^{(2)} :$$
(3:36)

 $\mathbf{B}^{?} = (g \quad v v)D$ is the transverse part of the covariant derivative which satisf es $\mathbf{fB}^{?}$; $\mathbf{6}g = 0$. Here, we have taken advantage of the simplications for the D irac algebra in the heavy mass formulation. It allows to express any D irac bilinear (= 1, , 5, :::) in terms of the velocity v and the spin{operator 2S = i_5 v. The latter obeys the relations (in d space{time dimensions)

S
$$v = 0; S = \frac{1}{4}; S; S = \frac{1}{2}vv g; S; S = i vS$$
 (3:37)

U sing the convention $^{0123} = 1$, we can rewrite the standard D irac bilinears as:

$$H = v H H; H _{5}H = 0; H _{5}H = 2H S H$$

$$H = 2 v H S H; H _{5} H = 2i(v H S H v H S H)$$
(3:38)

Therefore, the D irac algebra is extrem ely sim ple in the extrem e non{relativistic lim it.

W e return to the discussion of the generating functional. The source term in (3.31) is also rewritten in terms of the $~elds\,H_{\,v}$ and h_{v}

$$Z \qquad Z d4x (+) = d4x (RvHv + HvRv + vhv + hv v) (3:39)$$

with

$$R_{v} = \frac{1}{2} (1 + \mathscr{G}) e^{im v \times}$$

$$_{v} = \frac{1}{2} (1 - \mathscr{G}) e^{im v \times} \qquad (3:40)$$

D i erentiating with respect to the source R $_v$ yields the G reen functions of the projected elds H $_v$. The heavy degrees of freedom, h_v , may now be integrated out. Shifting
variables $h_v^0 = h_v - C^1$ (BH $_v + _v$) and completing the square, the generating functional becomes

$$exp iZ [R_v; R_v; v; v; v; a; s; p] = [dH_v][dH_v][du]_h exp i S^0_N + S + Z$$

$$d^4x (R_vH_v + H_vR_v) + :::$$
(3:41)

where

$$S_{N}^{0} = d^{4}xH_{v}A + (_{0}B_{0}^{Y})C^{1}BH_{v}; \qquad (3:42)$$

and the ellipsis stands for term swith the sources $_v$ and $_v$ [3.17]. In (3.42), $_h$ denotes the determ inant coming from the Gaussian integration over the small component eld, i.e.

$$h = \exp \frac{1}{2} \operatorname{tr} \ln C$$

$$= \operatorname{N} \exp \frac{1}{2} \operatorname{tr} \ln 1 + C^{(1) 1} (i(v D) + g_A (S u) + C^{(1) + \dots + (1) + \dots + (1)$$

A snoted in Ref.[3.13], the space time representation of the h_v propagator, $C^{(1) 1}$, in plies that h_i is just a constant.

The next step consists in expanding the nonlocal functional (3.41) in a series of operators of increasing dimension. This corresponds to an expansion of the matrix C 1 in a power series in 1=m

$$C^{1} = \frac{1}{2m} - \frac{i(v \quad D) + g_{A}(u \quad S)}{(2m)^{2}} + O(q^{2}) :$$
 (3:44)

Thus the electrive heavy nucleon lagrangian up to 0 (q^3) is given as

$$S_{N}^{0} = d_{XH_{v}}^{0} A_{v}^{(1)} + A_{v}^{(2)} + A_{v}^{(3)} + (_{0}B_{v}^{(1)})_{0}\frac{1}{2m}B_{v}^{(1)} + \frac{(_{0}B_{v}^{(1)})_{0}B_{v}^{(2)} + (_{0}B_{v}^{(2)})_{0}B_{v}^{(1)}}{2m} + \frac{(_{0}B_{v}^{(1)})_{0}B_{v}^{(2)} + (_{0}B_{v}^{(2)})_{0}B_{v}^{(1)}}{2m} + \frac{(_{0}B_{v}^{(1)})_{0}B_{v}^{(1)} + (_{0}B_{v}^{(2)})_{0}B_{v}^{(1)}}{2m} + \frac{(_{0}B_{v}^{(1)})_{0}B_{v}^{(1)} + (_{0}B_{v}^{(1)})_{0}B_{v}^{(1)}}{2m} + \frac{(_{0}B_{v}^{(1)})_{0}B_{v}^{(1)} + (_{0}B_{v}^{(1)})_{0}B_{v}^{(1)} + (_{0}B_{v}^{(1)})_{$$

Note that the neglected term s of O (q^4) m ay be suppressed by inverse powers of either m or = 4 F. These two scales are treated on the same footing, the only thing which counts is the power of the low m om entum q. It is important to note that this expansion of the non { local action m akes the closed ferm ion loops disappear from the theory because at any nite order in 1=m, S⁰_N is local (as spelled out in m ore detail in

ref.[3.15]). To complete the expansion of the generating functional up to order q^3 , one has to add the one{loop corrections with vertices from A ⁽¹⁾ only. W orking to order q^4 (which still includes only one{loop diagram s), one also has to include vertices from A ⁽²⁾ and from ($_0B^{(1)y}$)B⁽¹⁾=(2m).

The disappearance of the nucleon m ass term to leading order in 1=m now allows for a consistent chiral power counting. The nucleon propagator is now of the form (3.30), i.e. has chiral power q¹. Consequently, the dimension D of any Feynm an diagram is given by* $_{\rm V}$

$$D = 4L \quad 2I_{\text{M}} \quad I_{\text{B}} + \int_{d}^{A} d(N_{d}^{\text{M}} + N_{d}^{\text{M}\text{B}})$$
(3:46)

with L the number of loops, I_M (I_B) the number of internal meson (baryon) lines and N_d^M , N_d^{MB} the number of vertices of dimension d from the meson, meson (baryon Lagrangian, in order. Consider now the case of a single baryon line running through the diagram [3.15]. In that case, one has

$$N_{d}^{MB} = I_{B} + 1 \qquad (3:47)$$

Together with the general topological relation

$$L = I_{M} + I_{B} \qquad (N_{d}^{M} + N_{d}^{MB}) + 1 \qquad (3:48)$$

we arrive at

$$D = 2L + 1 + (d 2)N_{d}^{M} + (d 1)N_{d}^{MB} : (3:49)$$

C learly, D 2L + 1 so that one has a consistent power counting scheme in analogy to the one in the meson sector. In particular, the coe cients appearing in $L_{N}^{(1)}$ and $L_{N}^{(2)}$ are not renorm alized at any loop order since D 3 for L 1 (if one uses e.g. dimensional regularization). This is in marked contrast to the in nite renorm alization of g_{A} and m in the relativistic approach, see eq.(3.21). As stated before, all mass dependence now resides in the vertices of the local pion (nucleon Lagrangian, i.e. all vertices now consist of a string of operators with increasing powers in 1=m. We have for example

Photon nucleon vertex :
$$ie^{1+3}_{2}$$
 v + 0 (in=)
Pion nucleon vertex : $(g_A = F)^a S q + 0$ (in=)
(3:50)

To sum marize, the elective pion {nucleon Lagrangian takes the form $L_{N}^{e} = L_{N}^{(1)} + L_{N}^{(2)} + L_{N}^{(3)} + L_{N}^{(4)} + \dots$ where the superscript '(i)' denotes the chiral dimension. The

^{*} Since this power counting argument is general, we talk of mesons (M) and baryons (B) for a while (instead of pions and nucleons).

complete list of terms contributing to L $_{\rm N}^{(2)}$ and the corresponding Feynm an rules can be found in appendix A .

Before we turn to the renorm alization of the chiral pion {nucleon EFT, a comment on the heavy ferm ion form alism is necessary. W hile it is an appealing fram ework, one should not forget that the nucleon (baryon) m ass is not extrem ely large. Therefore, one expects signi cant corrections from 1=m suppressed contributions to m any observables. This will become m ore cleareq. in the discussion of threshold pion photo { and electroproduction. It is conceivable that going to one { loop order 0 (q^3) is not su cient to achieve a very accurate calculation. Of course, only explicit and complete calculations can decide upon the quality of the q^3 approximation. This means that higher order calculations should be performed to learn about the convergence of the chiral expansion. For a few selected cases, calculations including part of or all terms of order q⁴ have been performed. We will discuss these in due course. To that accuracy, one has to include the pertinent contact terms from L $_{\rm N}^{\rm (4)}$ and consider one { loop graphs with exactly one insertion from L $_{\rm N}^{(2)}$. Here, let us note that the calculations which include all terms of order q⁴ in the chiral expansion indeed lead to an improvement for the respective theoretical predictions. Ultim ately, one m ight want to include m ore inform ation in the unperturbed Ham iltonian. At present, it is not known how to do that but it should be kept in m ind.

III.3. RENORMALIZATION

In this section, we will be concerned with the renorm alization of the e ective pion { nucleon Lagrangian to order q^3 . In the relativistic case, this problem was addressed for a certain class of divergences in ref.[3.5] and sim ilarly for the heavy mass form alism in refs.[3.14,3.16]. E dker [3.17] has recently given a complete renorm alization prescription of the generating functional at order q^3 as discussed below.

Let us rst consider the nucleon propagator and m ass{shift. The only loop diagram contributing at order q^3 is shown in g.3.2. and leads to [3.14] (we have no tadpole contribution since that involves an odd power of the loop m om entum 1 to be integrated over)

$$loop (!) = 3i \frac{g_A^2}{F^2} \frac{Z}{(2)^d} \frac{d^d l}{v (l k) + \frac{1}{2} M^2 + i} S \quad l(S l)$$

$$= \frac{3g_A^2}{4F^2} M^2 \frac{1}{(2)^d} (!) \quad l \quad (0)$$

$$(3:51)$$

with ! = v k and the loop functions $_{0}J(!)$ and (0) given in appendix B.



Fig. 3.2: One { bop contribution to the nucleon self{energy to order q^3 . The solid and dashed lines denote nucleons and pions, in order.

Putting pieces together, we arrive at

$$(!) = \frac{3g_{A}^{2}}{4F^{2}} 2L! (2!^{2} 3M^{2}) + \frac{!}{8^{2}} (2!^{2} 3M^{2}) \ln \frac{M}{m} + \frac{!}{8^{2}} (m^{2} !^{2})$$

$$\frac{1}{4^{2}} (M^{2} !^{2})^{3=2} \arccos \frac{!}{M} 4M^{2} c_{1} + \frac{B_{20}}{8^{2}} \frac{!}{F^{2}} + \frac{B_{15}!^{3}}{(4 F)^{2}} \cdots$$

$$(3:52)$$

making use of dimensional regularization and separating the in nite from the nite pieces as in eq.(2.47). The last three terms in eq.(3.52) stem from three contact terms of order q^2 and q^3 , respectively, cf. appendix A and eqs(3.60),(3.63). The coe cient of the rst contact term is obviously nite whereas the other two low {energy constants are needed to renorm alize the nucleon self{energy}. The ellipsis in (3.52) stands for terms which do not contribute to the mass shift and Z{factor of the nucleon. The nucleon propagator now takes the form

$$S(!) = \frac{i}{p \quad v m \quad (!)} = \frac{i}{! \quad (!)} :$$
 (3:53)

The propagator develops a pole at $p = m_N v w$ ith m_N the renorm alized nucleon m ass,

$$m_N = m + (0)$$

 $(0) = 4c_1 M^2 - \frac{3g_A^2 M^3}{32 F^2}$: (3:54)

As stated in the previous section, the mass shift (0) is nite and vanishes in the chiral lim it, quite in contrast to the relativistic approach (cf. eq.(3.21)). Notice also that the mass(shift contains the non{analytic piece of order $m^{3=2}$ already found in ref.[3]. The nucleon wave{function renorm alization (Z { factor) is determined by the residue of the propagator at the physical mass pole and given by

$$S(!) = \frac{iZ_{N}}{p \quad v \quad m}$$

$$Z_{N} = 1 + {}^{0}(0) = 1 \quad \frac{3g_{A}^{2} M^{2}}{32 \, {}^{2}F^{2}} \quad 3 \ln \frac{M}{m} + 1 \quad \frac{M^{2}}{2 \, {}^{2}F^{2}} B_{20}^{r}() \quad : \qquad (3:55)$$

Here, the low energy constant B $_{20}$ has eaten up the in nity in the loop contribution via the renorm alization prescription

$$B_{20} = B_{20}^{r}$$
 () + $\frac{20}{4^{2}}L$; $20 = \frac{9}{16}g_{A}^{2}$: (3:56)

In a sim ilar fashion, one can renorm alize all divergences appearing in the various G reen functions. How ever, there exists a more system atic method which we will now turn to.

The starting point for a consistent renorm alization scheme is the generating functional in the presence of the external sources. In the approximations described in section 32, the ferm ion determinant is trivial to any nite order in 1=m and the integration over H_v reduces to completing a square. This leads to:

where the ellipsis stands for term s linear and quadratic in $_{\rm v}$ ($_{\rm v}$) which we will not need in what follows, and U = u². From here on, the standard CHPT procedure as outlined in ref.[3.18] can be applied. One expands the action in the functional integral around the classical solution $U_{\rm cl}[v;a;s;p] = (u_{\rm cl}[v;a;s;p])^2$ of the lowest oder equations of motion. To calculate the loop functional to order p³ one has to expand L⁽²⁾ + L⁽⁴⁾ R_v (A⁽¹⁾) ¹ R_v in the functional integral (3.57) around the classical solution. The divergences are entirely given by the irreducible diagram s (cf. Fig.1 of ref.[3.17]) corresponding to the generating functional

$$Z_{irr}[v;a;s;p;R_{v}] = d^{4}xd^{4}x^{0}d^{4}yd^{4}y^{0}R_{v}(x)(A^{(1)})^{1}(x;y)[_{1}(y;y^{0})^{(4)}(y \quad y^{0}) + _{2}(y;y^{0})](A^{(1)})^{1}(y^{0};x^{0})R_{v}(x^{0})$$
(3:58)

with $(A^{(1)})^{1}$ the propagator of H_{v} in the presence of external eds. The explicit form of the self(energy functional _{1;2} can be found in ref.[3.17]. Here, it is important to note that these diverge as $y ! y^{0}$. The divergences can be extracted in a chiral invariant manner by making use of the heat kernel representation of the propagators in d dimensional Euclidean space. These divergences will then appear as simple poles in

=
$$(4 \quad d)=2.$$
 A fiter som e lengthy algebra as detailed in ref.[3.17] one arrives at

$$\begin{bmatrix} 1 (y; y^{0})^{(4)} (y - y^{0}) + 2 (y; y^{0}) \end{bmatrix} = \begin{bmatrix} 1^{n} (y; y^{0}; -)^{(4)} (y - y^{0}) + 2^{n} (y; y^{0}; -) \end{bmatrix}$$

$$\frac{2L}{F^{2}}^{(4)} (y - y^{0}) \begin{bmatrix} 1 (y) + 2 (y) \end{bmatrix}$$
(3:59)

The generating functional Z $[v;a;s;p;R_v]$ can now be renorm alized by introducing the local counterterm Lagrangian

$$L_{N}^{(3)}(x) = \frac{1}{(4 + F)^{2}} X_{i} B_{i}H_{v}(x) O_{i}(x) H_{v}(x)$$
(3:60)

where the coupling constants B_i are dimensionless and the eld monom ials $O_i(x)$ are of order p^3 . A minimal set consisting of 22 counterterms has been given in ref.[3.17].* In complete analogy to eq.(2.46), one decomposes the low {energy constants B_i as

$$B_{i} = B_{i}^{r} () + (4)^{2}_{i} L$$
 (3:61)

^{*} For on{shell nucleons, one can further reduce this number by using the equations of motion for the nucleons.

The $_{\rm i}$ depend only on g_{A} (strictly speaking on g_{A}) and the corresponding operators $O_{i}(\mathbf{x})$ are given by

(3:62)

The sum of the irreducible one { loop functional (3.58) and the counterterm functional derived from the Lagrangian (3.60) is nite and scale{independent. The renorm alized low (energy constants B_i^r () are measurable (i.e. they can be determined from a t to som e observables) and subject to the follow oing renorm alization group behaviour under scale changes

$$B_{i}^{r}(_{2}) = B_{i}^{r}(_{1}) \qquad \lim_{i} \log \frac{2}{_{1}} :$$
 (3:63)

This completes the formalism necessary to renormalize the pion {nucleon (or meson { baryon) Lagrangian to order q^3 in heavy ferm ion CHPT. In what follows, we will see these renorm alization prescriptions being operative for various physical processes.

III.4. LOW {ENERGY CONSTANTS AND THE ROLE OF THE (1232)

As noted in section 2, in the meson sector the low (energy constants L_1 could all be xed from phenom enological constraints (within a certain accuracy). Furtherm ore, the actual values of these coe cients could be understood from a hadronic duality in terms of resonance exchange. We note, however, that for the non{leptonic weak interactions (which contains 80 new contact term s) this generalized vectorm eson dom inance principle is not that successful [3.19]. In the nucleon sector, the situation is som ewhat sim ilar to the case of non { leptonic weak interactions of the mesons. At present, only a subset of the coe cients in L $_{\rm N}^{\rm (2)}$ and L $_{\rm N}^{\rm (3)}$ (and also in L $_{\rm N}^{\rm (4)}$) have been xed from phenomenology. We will discuss one example below. In most other cases, one resorts to resonance saturation which besides meson resonances involves the nucleon excitations, in particular the (1232) P { wave resonance. The plays a particular role for two reasons. First, its excitation energy is only 300 M eV and second, its coupling to the N system is very strong, q_N / $2q_N$. For these reasons and the degeneracy of the with the nucleon in the lim it of in nite colours, it has been suggested to include the from the start in the e ective theory [3.20]. We will discuss this below. O by joursly, if one does not want to build in the in the EFT, it will feature prom inently in the estimation of certain low {energy constants. We will detail one example which we need for the discussion of elastic N scattering in the threshold region later on.

	occurs in	determ ined from	
C1	m _N , _N	phen.	
C ₂ ;C ₃ ;C ₄	N ! N	res.exch.+ phen.	
C5	m _N , _N (m _u €m _d)	phen.	
C ₆	p;n	phen.	
C7	p;n	phen.	
B ₁ ;B ₂ ;B ₃ ;B ₈	N ! N	res.exch.	
B ₁₀	< $r^2 > 1^{V}$	phen.	
B ₂₀	Z_N	unknow n	

Table 3.1: O courence of low (energy constants and their determ inations from phenom enological (phen.) constraints or estimation based on resonance exchange (res. exch.). Note that c_5 is only contributing for $m_u \notin m_d$. For the de nition of the corresponding elective Lagrangian see (3.36) and appendix A1.

First, let us tabulate the various low {energy constants from $L_{N}^{(2;3)}$ which we will encounter in the following sections and discuss in which process they can be probed (or determ ined). As noted before, while the list for the term s of order q^2 is complete, for the term $s of O(q^3)$ we only exhibit the term s which we will use later on. As becomes clear from table 3.1, certain low {energy constants can only be probed in the presence of external elds. These are, in turn, the best determ ined ones since the nucleon radii and m agnetic m om ents are accurately known (cf. $c_6; c_7$ or B_{10}). The constants related directly to the N interactions have not yet been determined from a global t to N scattering data as it was done in the relativistic case. In view of the present discussion about the low {energy N scattering data such a program has to be performed with adequate care and is not yet available (see section 3.5). As noted in table 3.1, the low { energy constant c_i can be xed from phenom enology. Consider rst c_1 . It is related to the much discussed pion {nucleon {term, N (t) < $p^0 jn^2$ (uu + dd) jp > (t = $(p^0 p^2)$, via [3.14]

$$c_1 = \frac{1}{4M^2}$$
 _N (0) + $\frac{9g_A^2 M^3}{64 F^2}$: (3:64)

U sing the empirical values for F , M and g_A together with the recent determ ination 8 M eV [3,21], this amounts to $_{\rm N}$ (0) = 45

$$c_1 = 0.87 \quad 0.11 \,\mathrm{GeV}^1 \quad (3.65)$$

The two constants c_2 and c_3 are related to the so{called axial polarizability A and the isopin (even N S (wave scattering length a^+ (for the de nitions and discussion see section 3.5)

$$c_{3} = \frac{F^{2}}{2} + \frac{g_{A}^{2}M}{8F^{4}} + \frac{77}{48} + g_{A}^{2} = 525 \quad 0.22 \,\text{GeV}^{1}$$

$$c_{2} = \frac{F^{2}}{2M^{2}} + (1 + \frac{M}{m})a^{+} + \frac{3g_{A}^{2}M^{3}}{64F^{4}} + 2c_{1} + 2c_{1} + \frac{g_{A}^{2}}{8m} = 3.34 \quad 0.27 \,\text{GeV}^{1}$$
(3:66)

using the empirical values $_{\rm A}$ = 2.28 0:10M⁻³ and a⁺ = 0.83 10M ¹ 0:38 (for references, see section 3.5). Note, how ever, that these observables m ight not form the best set to determ ine the constants $c_{1,2,3}$ since the scattering length a^+ is extremely sensitive to the counter term combination $c_2 + c_3$ 2q and, furtherm ore, there are correlations between the S{wave scattering lengths and the N {term . The constants c_6 , c_7 and B $_{10}$ can be determ ined from the isovector and isoscalar anom alous m agnetic moment of the nucleon and its isovector charge radius, respectively [3.5,3.14]. The num erical values of the seven low (energy constants in L $_{\rm N}^{(2)}$ are sum m arized in table 3.2. The constants c₂ and c₃ have also been estim ated making use of the resonance saturation hypothesis [3.22]. Consider c_3 . In that case, the dom inant contribution com es from the

(1232) and there is a small correction due to the N (1440) resonance. In addition,

there is a sizeable contribution due to scalar meson exchange. The pertinent Lagrangians for the coupling of the mesonic and the nucleon excitations to the N system read

$$L_{N} = \frac{3g_{A}}{2^{P} \cdot \frac{1}{2}} a_{B} (Z + \frac{1}{2}) \qquad]u_{a} + hx:$$

$$L_{N} = \frac{1}{4}g_{A} R N = \frac{1}{5} + hx:$$

$$L_{S} = c_{m} Tr(+) + c_{d} Tr(u = u)$$

$$L_{SN} = g_{S}$$
(3:67)

where denotes the Rarita {Schwinger eld and Z param etrizes the o {shellbehaviour of the spin {3/2 eld. This param eter is not well known, the most recent analysis of ref. [3.23] gives 0.8 Z 0.3. We should stress here that it is mandatory to consider these nucleon excitations in the relativistic fram ework. The basic idea is that one starts from a fully relativistic theory of pions coupled to nucleons and nucleon resonances chirally coupled. One then integrates out these excitations from the elective theory which produces a string of pion {nucleon interactions whose coe cients are given in terms of resonance parameters. Finally, one de nes velocity {dependent nucleon elds elim inating the 'lower component' h(x). U sing now the large N_c coupling constant relation g N = 3g N = $\overline{2}$ = 28:42 (close to the empirical value of 28.37) and the phenom enological value g N N = (1=2:::1=4)g N [3.24] (which de nes a parameter R = 1:::1=4), we nd

$$c_{3} = \frac{g_{A}^{2}}{8m^{2}} \frac{m m 4m^{2} m^{2}}{m m} + 4Z [m (2Z + 1) + m (Z + 1)]$$

$$= 2:54 ::: 3:18 \, \text{GeV}^{1}$$

$$c_{3}^{N} = \frac{g_{A}^{2} R}{16 (m m)} = 0:06 ::: 0:22 \, \text{GeV}^{1}$$

$$c_{3}^{S} = 2c_{1} \frac{c_{d}}{c_{m}} = 1:33 \, \text{GeV}^{1} :$$
(3:68)

using $j_{cd} j = 32 \text{ M}$ eV and $j_{cm} j = 42 \text{ M}$ eV [3.31]. In addition, we have assumed that the value of c_1 is saturated by scalar exchange which allows to eliminate the coupling g_s . However, a strongly coupled scalar (isoscalar with $M_s = p \overline{g_s}$ 220 M eV is needed to saturate c_1 this way. A logether, we nd that $c_3^{\text{R}es} = c_3 + c_3^{\text{N}} + c_3^{\text{S}}$ varies between -3.6 and -5.0 G eV ¹, somewhat sm aller than the empirical value discussed above. This demonstrates that the resonance saturation hypothesis can not yet be considered established (as it is in the case of the meson sector). However, in the absence of su ciently many accurate low (energy data in the meson (baryon sector and a system atic evaluation of all counterterms up{to{and{including order q^3}, it is legitim ate to use resonance exchange to estim ate the low {energy constants which appear in the processes one considers. The introduction of this unwanted model {dependence should be considered as a transitional stage until a complete analysis of the various coupling constants based on the to data becomes available.

c_1^0	c_{2}^{0}	c_3^0	c_4^0	c_{5}^{0}	c_6^0	C_{7}^{0}
-1.63	6.20	-9. 86	7.73	0.17	11.22	-2.03

Table 3.2: Numerical values of the dimensionless low (energy constants $c_i^0 = 2m_N c_i$ (i = 1;:::;5) and $c_{6;7}^0 = 2c_{6;7}$ with $m_N = (m_p + m_n)=2 = 938:92 \text{ M eV}$ the nucleon mass. c_4 is determined from the P (wave N scattering volumes and c_5 follows from the strong np mass di erence, $(m_n m_p)_{str} = 2M^2 c_5 (m_d m_u)=16 \cdot c_{6;7}$ are determined from the nucleon isovector and isoscalar anom alous magnetic moments as described in section 4.1.

One particular advantage of the heavy mass formulation is the fact that it is very easy to include the baryon decuplet, i.e. the spin $\{3/2 \text{ states. This has been done in full}$ detailby Jenkins and M anohar [3.10,3.20]. The inclusion of the (1232) is m otivated by the argum ents given in the beginning of this section, in particular the fact that the N mass{splitting m m_N is only about thrice as much as the pion decay constant, * so that one expects signi cant contributions from this close { by resonance (the same holds true for the full decuplet in relation to the octet, see section 6). This expectation is bome out in many phenom enological models and we had also seen in the discussion of the low {energy constants the prominent role of the . How ever, it should be stressed that if one chooses to include this baryon resonance (or the full decuplet), one again has to account for all term s of the given accuracy one aim s at, say 0 (q^3) in a one{loop calculation. This has not been done in the presently available literature. Furtherm ore, m_N does not vanish in the chiral lim it thus destroying the the mass dierence m consistent power counting (as it is the case with the baryon mass in the relativistic form alism discussed in section 3.1). We will come back to this below. In the extreme non {relativistic limit, the is described by Rarita {Schwinger spinor _a with a 2 f1, 2, 3g. This spinor contains both spin $\{1/2 \text{ and spin} \{3/2 \text{ com ponents. The spin} \{1/2 \text{ pieces }$ are projected out by use of the constraint a = 0. One then de nes a velocity { dependent eld via

$$_{a} = e^{im v x} (T + t)_{a}$$
 (3:69)

In terms of the physical states we have

$$T^{1} = \frac{1}{\frac{p}{2}} + \frac{p}{\frac{q}{3}} = \frac{p}{\frac{q}{3}} ; T^{2} = \frac{p}{\frac{p}{2}} + \frac{p}{\frac{q}{3}} = \frac{p}{\frac{q}{3}} ; T^{3} = \frac{1}{\frac{q}{3}} + \frac{p}{\frac{q}{3}} = \frac{p}$$

^{*} O fien it is stated that m m_N ' 2M . W hile that is num erically true, the behaviour of these quantities in the chiral lim it is very di erent. W hile the form er stays constant as \hat{m} ! 0, the latter vanishes.

The e ective non { relativistic N Lagrangian to leading order reads

$$L_{N}^{(1)} = I a v D^{b}T^{b} + T a T^{a} + \frac{3g_{A}}{2 2} (T a^{a}u^{a}H + H u^{a}T^{a})$$
(3:71)

with = m m_N and $u^a = (i=2) \operatorname{Tr} ({}^a u^y r U u^y)$. Clearly one is left with some residual mass dependence. In the language of ref.[3.20] we have set $C = 3g_A = 2 = 1.89$ which is nothing but the SU (4) coupling constant relation discussed before. From the width of the decay ! N one has C = 1.8 [3.10], consistent with the value given before (if one uses the full decuplet the value of C reduces to 1.5). The propagator of the spin{3/2 elds reads} = 1.80 + 1.

S (!) =
$$i \frac{v \ v \ g \ \frac{4}{3}S \ S}{!}$$
 (3:72)

For all practical purposes, it is most convenient to work in the rest { fram ev = (1;0;0;0). In that case, one deals with the well{known non{relativistic isobar model which is discussed in detail in the monograph by Ericson and W eise [3.24]. Consider now the nucleon self{energy (i.e. a diagram like in g.3.2. but with an intermediate state). Its contribution is non{vanishing in the chiral limit. Therefore, a counter term of the following form has to be added [3.25] (like in the relativistic theory of the nucleon alone)

$$L_{N}^{(0)} = m_{0} \operatorname{Tr}(H H)$$

$$m_{0} = \frac{10}{3} \frac{C^{2}}{F^{2}} L + \frac{1}{16^{2}} \ln \frac{2}{6} \frac{5}{6} : \qquad (3:73)$$

C learly, such a contribution destroys the consistent power counting. However, from phenom enological arguments, one might want to consider the quantity as a small parameter. W hile this is not rooted in QCD, it might be worth to be explored in a system atic fashion. Such an analysis is, however, not available at present. Our point of view is that one should not include the as a dynamical degree of freedom in the EFT but rather use it to estimate certain low {energy constants. W hile this might narrow the range of applicability of the approach, it at least allows for a consistent power counting.

III.5. A SPECTS OF PION {NUCLEON SCATTER ING

E lastic pion {nucleon scattering in the threshold region can be considered the most basic process to which the CHPT methods can be applied. This is underlined by the W einberg's very successful current algebra prediction [3.26] for the S{wave pion {nucleon scattering lengths,

$$a_{1=2} = \frac{M}{4 F^2} = 2a_{3=2} = 0:175 M^{-1}$$
 (3:74)

Tom ozawa [327] also derived the sum rule $a_{1=2}$ $a_{3=2} = 3M = 8 F^2 = 0.263 M^{-1}$. Empirically, the combination $(2a_{1=2} + a_{3=2})=3$ is best determined from pion {proton scattering. The Karlsruhe{Helsinki group gives 0.083 0.004 M⁻¹ [3.28] consistent with the pionic atom measurement [3.29] of 0.086 0.004 M⁻¹. The value of $a_{1=2}$ $a_{3=2}$ is more uncertain. The KH analysis leads to 0.274 0.005 M⁻¹ [3.30]. In what follows, we will use the central values from the work of K och [3.28], namely $a_{1=2} = 0.175 M^{-1}$ and $a_{3=2} = 0.100 M^{-1}$. The agreement of the current algebra predictions with these numbers is rather spectacular. Therefore, one would like to know what the next{to{ leading order corrections to the original predictions are. This question was addressed in ref.[3.22]. To be specified, consider the on{shell N forward scattering amplitude for a nucleon at rest. D enoting by b and a the isospin of the outgoing and incoming pion, in order, the scattering amplitude takes the form

$$T^{ba} = T^{+}(!)^{ba} + T(!)^{i}^{bac c}$$
 (3:75)

with q the pion four{m on entum and ! = v q. Under crossing (a \$ b; q ! q) the functions T^+ and T are even and odd, respectively, T (!) = T (!). At threshold one has q = 0 and the pertinent scattering lengths are de ned by

$$a = \frac{1}{4} 1 + \frac{M}{m} T (M)$$
 (3:76)

The S{wave scattering lengths for the total N isospin 1/2 and 3/2 are related to a via $a_{1=2} = a^+ + 2a$; $a_{3=2} = a^+$ a. The abovem entioned central empirical values translate into $a^+ = 0.83 \quad 10 \text{ M}^{-1}$ and $a = 9.17 \quad 10 \text{ M}^{-1}$. In what follows, we will not exhibit the canonical units of 10 2 M $^{-1}$. The benchm ark values are therefore $a^+ = 0.83 \quad 0.38$ and $a = 9.17 \quad 0.17$ compared to the current algebra predictions of $a^+ = 0$ and a = 8.76 (using M = 138 M eV and F = 93 M eV). The empirical values for the forward amplitudes at threshold follow to be T⁺ (M) = 0.17 0.08 fm and T (M) = 1.87 \quad 0.03 fm. The four novel counter terms from $L_N^{(3)}$ which contribute to N scattering are $0_1; 0_2; 0_3$ and 0_8 given in eq.(3.62). Due to crossing symmetry, $L_N^{(2)}$ (these are the term s proportional to $c_{1;2;3}$, cf eq.(3.36)) contributes only to T⁺ (!) whereas $L_N^{(3)}$ solely enters T (!). For the isospin even/odd threshold amplitude we derive the follow ing chiral expansion

$$T^{+}(M) = \frac{2M^{2}}{F^{2}} c_{2} + c_{3} 2q \frac{q_{A}^{2}}{8m} + \frac{3q_{A}^{2}M^{3}}{64 F^{4}} + O(M^{4})$$
 (3:77)

$$T (M) = \frac{M}{2F^{2}} + \frac{M^{3}}{16^{2}F^{4}} + \frac{M^{3}}{16^{2}F^{4}} + \frac{g^{2}}{8m^{4}} + \frac{g^{2}}{8m^{4}} - 4b'()\frac{M^{3}}{F^{2}} + O(M^{4})$$
(3:78)

with $b^{r}() = (B_{1}^{r}() + B_{2}^{r}() + B_{3}^{r}() + 2B_{8}^{r}()) = (16^{2}F^{2})$. b has to be renorm alized as follows to render the isospin (odd scattering am plitude T (!) nite,

$$b = b^{r}$$
 () $\frac{L}{2F^{2}}$; (3:79)

since 1 + 2 + 3 + 2 = 1=2 (cf. eq.(3.62)). It is remarkable that there are no corrections of order M 2 and M 4 in T (M). The order M 2 has to be zero since crossing symmetry forbids any such counter term contribution which also must be analytic in the quark masses. For the loop contribution at order M 4 such an argument does not hold (loops can lead to non (analyticities), but an explicit calculation of all q⁴ loop diagram s shows indeed that they all add up to zero. The various term s in eq.(3.78) are the current algebra prediction, the expansion of the nucleon pole term, the one{loop and the counterterm contribution from $L_{N}^{(3)}$, respectively. is the scale introduced in dim ensional regularization. In what follows, we will use = m = 1:232 G eV, m otivated by the resonance saturation principle. Notice that the contact term contributions are suppressed by a factor M 2 with respect to the leading current algebra term . M atters are di erent for the isospin {even scattering am plitude T + . It consists of contributions of order M 2 and M 3 . From the form of eq.(3.77) it is obvious that the contact term s play a more important role in the determination of T⁺ than for T . The most dicult task is to pin down the various low -energy constants appearing in eqs. (3.77) and (3.78). In ref. [3.22], c_1 was xed as in eq. (3.64). The coe cients c_2 and c_3 where estimated from resonance exchange. This induces a dependence on the o {shell parameter Z as discussed in section 3.4. From the meson sector, scalar meson exchange can contribute to c_1 and c_3 ,

$$c_1 = \frac{1}{2}c_3 = c_1 = q \frac{c_d}{c_m}$$
 (3.80)

with $2c_d=c_m = L_5=L_8$. The central values for the parameters c_d and c_m given in ref.[3.31] lead to $2c_d=c_m = 1:56$. However, within the uncertainty of L_5 and L_8 , this ratio can vary between 0.75 and 2.25. The and the N (1440) contribute to $c_2 + c_3$ and to b()

$$c_{2} + c_{3} = \frac{g_{A}^{2}}{2m^{2}} \left(\frac{1}{2} \quad Z\right) 2m \quad (1 + Z) + m \left(\frac{1}{2} \quad Z\right)$$

$$c_{2} + c_{3}_{N} = \frac{g_{A}^{2} R}{16 (m + m)} \qquad (3:81)$$

$$b^{r} ()_{=m} = q_{A}^{2} \quad \frac{(Z \quad \frac{1}{2})^{2}}{8m^{2}} + \frac{R}{32 (m + m)^{2}}$$

O ther baryon resonances have been neglected since their couplings are either very sm all or poorly (not) known.* C learly, the contribution of the N (1440) is only a sm all

^{*} A remark on the {m eson is in order. The chiral power counting enforces a vertex of order q^2 of the form $L^{(2)} = g$ Tr([u;u]) [3.31]. In forward direction the contraction of the {m eson propagator with the corresponding matrix element vanishes. Therefore, one has no explicit {m eson induced contributions to T of order q^2 and q^3 .

correction to the {contribution. The num erical results are as follows. Consider rst the amplitude T . Using M = 138 MeV, F = 93 MeV, m = 938:9 MeV, Z = 1=4 and R = 1, one has

$$T (M) = (1.57 + 0.24 + 0.08 + 0.02) \text{ fm} = 1.91 \text{ fm}$$
 (3.82)

where we have explicitely shown the contributions from the current algebra, the one loop, the nucleon pole and the counter term s. The total result is in good agreement with the empirical value. The largest part of the M³ term comes from the pion loop diagram s. We should stress that only this loop contribution can close the gap between the Weinberg-Tom ozaw a prediction of 1.57 fm and the empirical value. As stated before, the uncertainties in b are completely masked by the sm all prefactor. If one chooses e.g.

= m, the loop contribution drops to 0.22 fm. The two{loop contribution carries an explicit factor M 5 and is therefore expected to be much smaller. In the case of the isospin{even scattering am plitude T⁺, the situation is much less satisfactory. There are large cancellations between the loop contribution and the 1=m suppressed kinem atical term s of order M 2 and M 3 . Therefore, the role of the contact term s is even further m agni ed. The total result for T⁺ is very sensitive to some of the resonance param eters, the empirical value of T⁺ can, how ever, be obtained by reasonable choices of these (cf. gs. 1 and 2 in [3.22] for the scattering length a⁺). A better understanding of the coe cients of the contact term s appearing at order q² (and higher) is necessary to

further pin down the prediction for T^+ (M).

A nother quantity of interest is the so{called nucleon axial polarisability $_{\rm A}$. It is related to the quenching of the axial vector coupling $g_{\rm A}$ in the nuclear medium as discussed in detail in ref.[3.32]. Consider the standard helicity non{spin{ ip am plitude C = A + B (1 t=4m)¹ with = (s u)=4m and the conventional N am plitude is written as T $_{\rm N}$ = A + ${\bf 6}_{\rm B}$. Here, A and B are functions of and the invariant momentum transfer squared t. The axial polarisability is then de ned as

$$_{A} = 2c_{01}^{+} = 2\frac{\theta}{\theta t}A^{+} (m^{2} + M^{2} t=2;m^{2} + M^{2} t=2)$$
 (3:83)

where the barm eans that the nucleon Born term has been subtracted and we have also indicated the standard notation which refers to the expansion of C (;t) around = 0. Empirically, one has $_{\rm A} = 2.28 \quad 0.04 \,{\rm M}^{-3}$ [3.30]. To get $_{\rm A}$, we calculate the on (shell N scattering amplitude in the cm s and subtract the Born term,

$$T^{+}(!;q^{0};q) = t_{0}(!) + q^{0} \quad qt(!) + \dots \quad (3:84)$$

with the kinematics $v = v^0 = q!'$ and $t = (q - q)^2 = 2 (M^2 - q^2 + q^0 - q)$ (here, q and q^0 are the momenta of the incoming and outgoing pion, respectively). The axial polarizability is then simply given by:

$$a_{A} = t_{1}(0)$$
 : (3.85)

At order q^2 , we have the counterterm contribution proportional to c_3 and at order q^3 , only loops contribute. The possible counterterm of order q^3 proportional to $! q^0$ q gives a vanishing contribution to $_A$. The nalexpression of this calculation was already given in eq.(3.66). Estimating the value of c_3 as discussed above, we nd $_A = 1.3:::1.8 M^{-3}$, som ew hat below the empirical value. It is in portant to stress (see also refs.[3.30,3.32]) that the alone is not su cient to get the empirical value but that one needs additional scalar exchange (as provided here through the resonance saturation).

For the later discussion in section 6, we will have to consider the N amplitude for o {shell pions. For doing that, we choose the pseudoscalar density P^a = iq $_5$ ^aq as the interpolating pion eld. The pion coupling via the pseudoscalar density is given in term s of G ,

$$\langle 0 \mathcal{P}^{b} j^{a} \rangle = {}^{ab}G ;$$
 (3.86)

where G^2 is given as the residue of the vacuum correlator < $0 \not P^a P^b \not D$ > at the pion pole. The o {shell N amplitude is then de ned via

$$d^{4}xe^{iq_{1}x} < N JT (P^{a}(x)P^{b}(0)) N > = \frac{G^{2}i^{5}}{(q_{1}^{2} M^{2})(q_{2}^{2} M^{2})} T^{ab}(q_{1};q_{2}): \quad (3.87)$$

It is now a straightforward exercise to show that the amplitude calculated in this fashion obeys the Adler conditions,

$$T^{+} (q_{1} = q_{2} = 0) = \frac{N(0)}{F^{2}}$$

$$T^{+} (q_{1}^{2} = 0; q_{2}^{2} = M^{2}) = T^{+} (q_{1}^{2} = M^{2}; q_{2}^{2} = 0) = 0:$$
(3:88)

Finally, we stress that GSS [3.5] have evaluated the full o {shell pion{nucleon am plitude in the framework of relativistic baryon CHPT and discussed the so{called remainder of the N {term derived from it. We will come back to these issues in section 6 because the {term is intimately related to the strangeness content of the nucleon and the baryon mass ratios.

III.6. THE REACTION N ! N

7

A nother reaction involving only pions and nucleons is the single pion production reaction N ! N (for some older references, see [3.33]). The interest in this reaction stem s mostly from the fact that it apparently o ers a possibility of determ ining the low (energy elastic scattering amplitude whose precise know ledge allow s to test our understanding of the chiral sym m etry breaking of QCD. How ever, at present no calculation based on chiral perturbation theory is available which links the pion production data to the ! am plitude in a model (independent fashion. Consequently, all presently available determ inations of the S{wave scattering lengths from the abovem entioned data should be taken cum grano salis. Over the last years, new experimental data in

the threshold region have become available [3.34-3.38] which allow for a direct comparison with the CHPT predictions. Beringer [3.39] has performed a tree calculation in relativistic baryon CHPT. In ref.[3.40] the chiral expansion of the threshold am plitudes was reanalyzed in terms of the heavy ferm ion formalism at next{to{leading order.

To be speci c, consider the process ${}^{a}N {}^{b} {}^{c}N$, with N denoting the nucleon (proton or neutron) and 'a;b;c' are isospin indices. At threshold, the transition m atrix { element in the ${}^{a}N$ centre{of{m ass frame takes the form

$$T = i \sim \tilde{k} D_1 (b^{ac} + c^{ab}) + D_2^{abc}$$
 (3:89)

where \tilde{K} denotes the three{m om entum of the incoming pion and the amplitudes D $_1$ and D $_2$ will be subject to the chiral expansion as discussed below. They are related to the more commonly used amplitudes A $_{2I;I}$, with I the total isospin of the initial N system and I the isospin of the two{pion system in the nalstate, via

$$A_{32} = \frac{P_{10}}{10}D_1; A_{10} = 2D_1 \quad 3D_2$$
 (3:90)

A ssum ing that the am plitude in the threshold region can be approxim ated by the exact threshold am plitude, the total cross section can be written in a compact form,

$$tot (s) = \frac{m^2}{2s} \frac{p}{(s;m^2;M^2)} \frac{(s;m^2;M^2)}{3} (s) j_1 D_1 + \frac{2}{2} D_2 \frac{j}{2} S$$

$$(s) = \frac{1}{32^3} \frac{p}{1} dT \frac{p}{T (T + 2m) (T_1 - T) (T_2 - T)} \frac{j}{T_3 - T} ;$$

$$T_1 = \frac{1}{2^3} \frac{p}{5} (P - s - m - M_1 - M_2) (P - s - m + M_1 + M_2) ;$$

$$T_2 = \frac{1}{2^3} \frac{p}{5} (P - s - m - M_1 + M_2) (P - s - m + M_1 - M_2) ;$$

$$T_3 = \frac{1}{2^3} (P - s - m - M_1 + M_2) (P - s - m + M_1 - M_2) ;$$

$$T_3 = \frac{1}{2^3} (P - s - m - M_1 + M_2) (P - s - m + M_1 - M_2) ;$$

$$(3:91)$$

$$(x;y;z) = x^2 + y^2 + z^2 - 2(xy + xz + yz)$$

with s the total centre{of{m ass energy squared. $_3$ (s) denotes the conventional integrated three{body phase space where M $_1$ and M $_2$ stand for the m asses of the nal state pions and one has the inequality 0 T_1 $T_{2;3}$. (x;y;z) is the K allen{ function. The $_{1;2}$ are channel-dependent isospin factors and S is a Bose symmetry factor. For p ! + n and $p ! = 0^{\circ} n$ we have $_1 = 2^{\circ} 2; _2 = 0; S = 1=2$ and $_1 = 0; _2 = 2; S = 1=2$, in order. In the threshold region, one can approximate to a high degree of accuracy the three{body phase space and ux factor by analytic expressions as discussed in m ore detail in section 4. The chiral expansion of the am plitude functions D $_1$ and D $_2$ takes the form *

$$D = f_0 + f_1 + f_2^{2} + \dots$$
(3.92)

^{*} Here, D stands as a generic symbol for $D_{1;2}$.

m odub logarithm s. The rst two coe cients of this expansion have been calculated in ref.[3.40]. For that, one needs only $L_e = L_N^{(1)} + L_N^{(2)} + L_N^{(2)}$ and one nds that none of the low (energy constants c_i will contribute (the sum of the corresponding graphs vanishes). Notice that the much debated next{to{leading order interaction does not appear at this order in the chiral expansion. One can therefore write down low (energy theorem s for $D_{1;2}$ which only involve well{known physical (low est order) parameters,

$$D_{1} = \frac{g_{A}}{8F^{3}} + \frac{7M}{2m} + 0 (M^{2})$$

$$D_{2} = \frac{g_{A}}{8F^{3}} + \frac{17M}{2m} + 0 (M^{2})$$
(3:93)

There are potentially large contributions from diagrams with intermediate (1232) states of the type M² = (m m 2M), which numerically would be of the order 10 M. As shown in [3.40], no such terms appear from diagrams involving one or two intermediate resonances. Consequently, the chiral expansion is well behaved but not too rapidly converging. The order M corrections give approximatively 50% of the leading term. However, the calculations of Beringer [3.39] in relativistic baryon chiral perturbation theory indicate that further 1=m suppressed kinematical corrections are small. The numerical evaluation of eqs.(3.93) amounts to D₁ = 2:4 fm³ and D₂ = 6:8 fm³ or using eq.(3.90)

$$A_{32} = 2.7 M^{-3}; A_{10} = 5.5 M^{-3}$$
 (3.94)

which compare fairly with the recent determ inations of Burkhardt and Lowe (see ref.[3.35]), $A_{32} = 2.07$ 0.10M³ and $A_{10} = 6.55$ 0.16M³. As stressed, how ever, in ref.[3.40], one can confront the LET eqs.(3.93) directly with experimental data and, furtherm ore, the global t to the threshold am plitudes of ref.[3.35] has to be reexam ined critically. The cross sections for p! + n and p! = 0.0 n in comparison to the existing data are shown in g.3.3. They compare well to the existing data for the rst 30 M eV above threshold. To get an idea about the higher order corrections, one can calculate the imaginary parts Im D_{1/2}. Corrections to R e D_{1/2} of the same size indeed turn to be such that they can improve the description of the data since the rst/second reaction allows to test D₁/D₂, respectively.

In addition, one nds that a best t to these data leads to $D_1 = 2.26 \text{ fm}^3$ and $D_2 = 9.05 \text{ fm}^3$ as indicated by the dotted lines in g.3.3. U sing eq.(3.90), this leads to $A_{32} = 2.5 \text{ M}^{-3}$ and $A_{10} = 8.0 \text{ M}^{-3}$ som ew hat di erent from the global best t values of ref.[3.35]. We believe that the energy range covered by the t in ref.[3.35] was too large to reliably extract the threshold am plitudes. In that t, the data in the rst 30 M eV above threshold had too little statistical weight.

A nother in portant remark concerns the fashion in which the S (wave scattering lengths are in general extracted from the N ! N data [3.34,3.35]. It is based on



Fig. 3.3: Total cross sections for p! + n and p! = 0 n in comparison to the data. Squares: ref.[3.36], octagon: ref.[3.37] and diam onds: ref.[3.38]. The dashed lines refer to an approximation discussed in ref.[3.40] and the dash{dotted ones show the best t to these data as discussed in the text (the 1 {band is indicated by the dotted lines).

the old O lsson {Turner m odel [3.33] which param etrizes the chiral sym m etry breaking in terms of one param eter called . This is, of course, an pre{QCD artefact since we now know that the breaking via the quark m asses is of the 3 3 form, i.e. = 0. Therefore, one can no m ore accept such a param etrization. The essential question is now, how can one relate the S-wave scattering lengths and the N threshold am plitudes

in a model independent way. This question is quite nontrivial. The Olsson (Turner model with = 0 only contains the tree level scattering lengths. To establish a m relationship between the amplitudes and the N ! N data beyond leading order, one has to perform a complete one { loop calculation. This has not yet been done. As an estimate, however, we can combine the new low energy theorems for the N threshold amplitudes (3.93) with W einberg's low energy theorems for scattering (i.e. the leading term in the chiral expansion). This way we derive

$$A_{10} = 4 \frac{g_N}{m} 1 + \frac{37M}{14m} \frac{a_0^0}{M^2} + 0 (M^2)$$
(3:95)

$$A_{32} = \frac{p}{10} \frac{g_N}{m} + \frac{7M}{2m} - \frac{a_0^2}{M^2} + O(M^2)$$
(3:96)

The corrections of order M are taken care of by the calculation to order p^2 leading to eqs.(3.93) and what remains to be done is to system atically work out the various contributions at 0 (M²). Ignoring these for the moment and inserting on the left hand side the present t value, we extract $a_0^0 = 0.23$ 0.02 and $a_0^2 = 0.042$ 0.002 which are quite close to the CHPT prediction at next-to-leading order. We stress how ever, the a complete calculation of the 0 (M²) corrections to the above relations is mandatory. We conclude that the values of the S{wave scattering lengths can eventually be infered from the threshold N ! N amplitudes. The complete one{loop calculation which would give a sound basis for doing that is, unfortunately, not yet available. At present, it seems that the most accurate fashion of determ ining in particular a_0^0 are K $_4$ decays.

III.7. THE PION {NUCLEON VERTEX

The last topic we want to address in this section is the pion{nucleon vertex, parametrized in terms of a form factor G $_{\rm N}$ (t). It plays a fundamental role in many areas of nuclear physics, in particular in the description of the nucleon {nucleon force via meson{exchange models. Before we discuss the details, let us stress from the beginning that while the strong pion{nucleon coupling constant g $_{\rm N}$ G $_{\rm N}$ (t = M^2) can be unambigously calculated within CHPT, the N form factor depends on the choice of the interpolating pion eld. Furtherm ore, if one writes a dispersion relation for G $_{\rm N}$ (t), one realizes that the absorptive part starts at t₀ = $(3M)^2$. Therefore, within the context of a one{loop calculation, the momentum dependence of the form factor will entirely stem from some contact term s.

A firer these rem arks, consider the B reit fram e m atrix {elem ent of the pseudoscalar density between nucleon states*

< N (
$$p^{0}$$
) jiq 5 ^aqjN (p) > = 2iB $g_{A} \frac{1 + g(t)}{M^{2} t} H S$ (p p) ^aH : (3:97)

^{*} The B reit fram e ism ost convenient for the calculation of such m atrix {elements since it allows for a unique translation of Lorentz {covariant m atrix {elements into non-relativistic ones.

The form factor g(t) is generated by loop and counterterm contributions. In fact, the loop contribution is divergent and t{independent,

$$g(t) = \frac{B_{23}}{8^{2}F^{2}}t + \frac{M^{2}}{F^{2}}C(g_{A}^{2}; B_{9}; B_{20}; :::)$$
(3:98)

where the constant C sums up all t{independent terms. We do not need its explicit form in what follows. B₂₃ is a nite low {energy constant from L $^{(3)}_{N}$,

$$L_{N}^{(3)} = B_{23} \frac{g_{A}}{(4 F)^{2}} H \text{ is } D H :$$
 (3:99)

The form factor g(t) features in the so{called G oldberger{Treim an discrepancy (for a review, see [3.41]). To be specific, let us look at the relation between the divergence of the axial current and the pseudoscalar density between nucleon states,

$$2B m g_{A} [1 + g(0)] = \frac{m_{N} g_{A}}{F} G \qquad : \qquad (3:100)$$

On the other hand, the strong pion {nucleon coupling constant is de ned via the residue of the pole term in (3.97),

$$2B m g_{A} [1 + g(M^{2})] = g_{N} G ; \qquad (3:101)$$

which leads to the Goldberger{Treim an discrepancy

_N 1
$$\frac{m_N g_A}{F g_N} = g(M^2) g(0) = \frac{M^2}{8^2 F^2} B_{23}$$
: (3:102)

Notice that $_N$ is entirely given by the low (energy constant b_{11} . W ith $m_N = 938.27$ M eV, F = 92.5 M eV, $g_A = 1.257$ and $g_N = 13.3*$ we nd

$$_{\rm N} = 0.04$$
; $B_{23} = 1.433$: (3.103)

If one now describes the whole Goldberger{Treim an discrepancy by a form factor e ect, one identi es the nucleon matrix{element of the pseudoscalar density with G G $_N$ (t)=(M 2 t), so that

$$G_{N}(t) = g_{N}[1 + g(t) - g(M^{2})]$$
 : (3:104)

^{*} In general, we use the Karlsruhe{Helsinki value of $g_N = 13.4$ [3.30]. In light of the recent discussion about the actual value of this quantity, we have adopted here the most recent value proposed by Hohler.

A ssum ing furtherm one the standard monopole form , G $_{\rm N}$ (t) = (2 M 2)=(2 t), one can calculate the cut{o ,

$$= \frac{4 \text{ F}}{2B_{23}} = 700 \text{ M eV} ; \qquad (3:105)$$

close to the result found by GSS [3.5] in the relativistic calculation. However, we stress again that this result depends on the choice of the interpolating eld and that it is based on the assumption that the whole Goldberger{Treim an discrepancy is due to a form factor e ect.

REFERENCES

- 3.1 S.W einberg, Phys. Rev. 166 (1968) 1568.
- 3.2 S.Colem an, J.W ess and B.Zum ino, Phys. Rev. 177 (1969) 2239;
 C.G.Callan, S.Colem an, J.W ess and B.Zum ino, Phys. Rev. 177 (1969) 2247.
- 3.3 P. Langacker and H. Pagels, Phys. Rev. D 8 (1971) 4595.
- 3.4 H. Pagels, Phys. Rep. 16 (1975) 219.
- 3.5 J.Gasser, M E.Sainio and A.Svarc, Nucl. Phys. B 307 (1988) 779.
- 3.6 A.K rause, Helv. Phys. Acta 63 (1990) 3.
- 3.7 H. Georgi, \W eak Interactions and M odern Particle Physics", Benjam in / Cum mings, Reading, MA, 1984.
- 3.8 Ulf-G.Meiner, Int. J.Mod. Phys. E1 (1992) 561.
- 3.9 E. Jenkins and A.V. Manohar, Phys. Lett. B 255 (1991) 558.
- 3.10 E.Jenkins and A.V.M anohar, in "E ective eld theories of the standard m odel", ed.Ulf{G.M ei ner, W orld Scienti c, Singapore, 1992.
- 3.11 J.Gasser, Ann. Phys. (N.Y.) 136 (1981) 62.
- 3.12 J.G asser and H. Leutwyler, Phys. Reports C 87 (1982) 77.
- 3.13 T.Mannel, W. Roberts and Z. Ryzak, Nucl. Phys. B 368 (1992) 264.
- 3.14 V.Bernard, N.Kaiser, J.Kambor and Ulf-G.Meiner, Nucl. Phys. B 388 (1992) 315.
- 3.15 G.Ecker, Czech. J. Phys. 44 (1994) 405.
- 3.16 T. (S. Park, D. (P. M in and M. Rho, Phys. Reports 233 (1993) 341.
- 3.17 G.Ecker, Phys. Lett. B 336 (1994) 508.
- 3.18 J.G asser and H. Leutwyler, Ann. Phys. (N.Y.) 158 (1984) 142.
- 3.19 G. Ecker, J. Kambor and D. Wyler, Nucl. Phys. B 394 (1993) 101.
- 3.20 E. Jenkins and A.V. Manohar, Phys. Lett. B 259 (1991) 353.

- 3.21 J.Gasser, H.Leutwyler and M.E.Sainio, Phys. Lett. 253B (1991) 252, 260.
- 3.22 V.Bernard, N.Kaiser and UlfG.Meiner, Phys. Lett. B 309 (1993) 421.
- 323 M. Benmerrouche, R.M. Davidson and N.C. Mukhopadhyay, Phys. Rev. C 39 (1989) 2339.
- 3.24 T. Ericson and W. Weise, "Pions and Nuclei", Clarendon Press, Oxford, 1988.
- 3.25 V.Bernard, N.Kaiser and UlfG.Meiner, Z.Phys.C 60 (1993) 111.
- 3.26 S.W einberg, Phys. Rev. Lett. 17 (1966) 616.
- 3.27 Y. Tom ozawa, Nuovo Cim. 46A (1966) 707.
- 3.28 R.Koch, Nucl. Phys. A 448 (1986) 707.
- 3.29 W .Beer et al, Phys. Lett. B 261 (1991) 16.
- 3.30 G.Hohler, in Landolt {Bornstein, vol.9 b2, ed.H.Schopper (Springer, Berlin, 1983); M.E.Sainio, private communication.
- 3.31 G.Ecker, J.Gasser, A.Pich and E.de Rafael, Nucl. Phys. B 321 (1989) 311.
- 3.32 M. Ericson and A. Figureau, J. Phys. G 7 (1981) 1197.
- 3.33 S.W einberg, Phys. Rev. 166 (1968) 1568;
 - M G.Olsson and L.Tumer, Phys. Rev. Lett. 20 (1968) 1127; Phys. Rev. 181 (1969) 2142;
 - L.{N.Chang, Phys. Rev. 162 (1967) 1497;
 - R.Rockmore, Phys. Rev. Lett. 35 (1975) 1409;
 - M.G.Olsson, E.T.Osypowski and L.Turner, Phys. Rev. Lett. 38 (1977) 296.
- 3.34 D. Pocanic et al., Phys. Rev. Lett. 72 (1994) 1156.
- 3.35 H.Burkhardt and J.Lowe, Phys. Rev. Lett. 67 (1991) 2622.
- 3.36 M E. Sevior et al, Phys. Rev. Lett. 66 (1991) 2569.
- 3.37 G.Kemelet al, Z.Phys.C 48 (1990) 201.
- 3.38 J.Lowe et al, Phys. Rev. C 44 (1991) 956.
- 3.39 J.Beringer, N Newsletter 7 (1993) 33.
- 3.40 V.Bernard, N.Kaiser and UlfG.Meiner, Phys. Lett. B 332 (1994) 415.
- 3.41 C A.Dom inquez, Riv. Nuovo C im. 8 (1985) N.6.

IV.NUCLEON STRUCTURE FROM ELECTROW EAK PROBES

In this section, we will mostly be concerned with the nucleon structure when realor virtual photons are used as probes. This is of particular interest for the physics program of the existing CW electron machines and intense light facilities. Topics included are C om pton scattering (spin-averaged and spin{dependent) and the classical eld of single and double pion production by real or virtual photons (see e.g. the monograph [4.1]). A nother well{understood probe are the W {bosons. Their interactions with the hadrons lead to the axial form factors and can also be used to produce pions. These topics will be discussed at the end of this section.

IV .1. ELECTROM AGNETIC FORM FACTORS OF THE NUCLEON

The coupling of the photon to the nucleon has an isoscalar and an isovector com ponent. The chiral expansion of the electric and the magnetic form factors of the neutron and the proton amounts to a calculation of the corresponding radii, magnetic moments and so on. Evidently, the further one goes in the loop expansion, higher moments of these form factors are tested. Here, we will concentrate on the form factors at smallm omentum transfer. As it was already mentioned in section 3, the existing precise data on these nucleon properties are mostly used to x the values of some low energy constants. How ever, it is important to understand the interplay of the loop and the counter term contributions and also to critically exam ine the absorptive parts of the isovector form factors.

First, let us consider the matrix {element of the isovector {vector quark current,

$$< p^{0}\dot{j}q - \frac{a}{2}q\dot{p} > = u(p^{0}) - F_{1}^{V}(t) + \frac{i-k}{2m}F_{2}^{V}(t) - \frac{a}{2}u(p)$$
 (4.1)

with $k = p^0$ p and $t = k^2$. This de nes the so{called D irac (F_1^V) and the Pauli (F_2^V) form factors. These are related to the proton and neutron form factors $F_{1;2}^p$ and $F_{1;2}^n$ via

$$F_{1}^{V}(k^{2}) = F_{1}^{p}(k^{2}) \qquad F_{1}^{n}(k^{2})$$

$$F_{2}^{V}(k^{2}) = F_{2}^{p}(k^{2}) \qquad F_{2}^{n}(k^{2}):$$
(4.2)

At zero m om entum transfer, we have $F_1^V(0) = 1$ and $F_2^V(0) = p_n = 3:706$. In relativistic baryon CHPT, these form factors have been discussed by G asser et al. [4.2]. Here, we will elaborate on the heavy ferm ion approach following ref.[4.3]. For that, one rew rites eq.(4.1) in the B reit fram e as

< N (p⁰) jg
$$\frac{a}{2}$$
 qN (p) > = F_1^{V}(t) + $\frac{t}{4m_N^2}F_2^{V}(t)$ v H $\frac{a}{2}$ H
+ $\frac{1}{m_N}F_1^{V}(t) + F_2^{V}(t)$ H [S ; S (p p)] $\frac{a}{2}$ H : (4:3)

This corresponds to the standard decomposition into the electric and magnetic form factors G_E (t) = F_1 (t) + F_2 (t) and G_M (t) = F_1 (t) + F_2 (t), with = t=4m_N^2. The D irac form factor F_1^V (t) is readily evaluated,

$$F_{1}^{V}(t) = 1 + \frac{t}{6} < r^{2} > _{1}^{V} + \frac{g_{A}^{2}}{F^{2}} J(t) + \frac{g_{A}^{2}}{F^{2}} t(t) - 2M^{2}$$
 (t) (4:4)

with the loop functions J (t) and (t) given in appendix B, and (t) = (t) $t^0(t)$. It can be shown analytically that the sum of all loop diagrams does not modify the tree level result, $F_1^V(0) = 1$. This is, of course, nothing but the charge non {renorm alization by the strong interactions. The isovector charge radius

$$< r^{2} > V_{1}^{V} = 6 \frac{dF_{1}^{V}(t)}{dt}_{t=0}$$
 (4:5)

diverges logarithm ically in the chiral lim it,

$$< r^{2} > V_{1}^{V} = \frac{5g_{A}^{2} + 1}{8^{2}F^{2}} \ln \frac{M}{16} - \frac{7g_{A}^{2} + 1}{16^{2}F^{2}} + \frac{3}{4^{2}F^{2}}B_{10}^{r}$$
 (4.6)

where the last term stems from a counterterm of order q^3 (cf. eq.(3.62)). It is worth to stress [4.2] that the coe cient of the logarithm in eq.(4.6) is nine times bigger than in the corresponding expression for the pion charge radius and therefore this term contributes signi cantly even for the physical value of the pion mass. This poses a severe constraint on any serious attempt of modelling the nucleon (say from a quark model point of view). To reproduce the empirical value $< r^2 > {}^V_1 = 0.578 \text{ fm}^2$, one has to set B ${}^r_{10}$ (m) = 0.13.* For B ${}^r_{10}$ (m) = 0, one would get $< r^2 > {}^V_1 = 0.62 \text{ fm}^2$, 8% above the empirical value.

The Pauli form factor F_2^V (t) takes the very simple form

$$F_{2}^{V}(t) = c_{6} - \frac{g_{A}^{2}m}{4F^{2}} \int_{0}^{2} dx \frac{p}{M^{2} + tx(x-1)}$$
(4:7)

which involves the low {energy constant c_6 to be identi ed with the isovector anom alous magnetic moment in the chiral lim it, $c_6 = v$. To order q^3 , we not for the isovector anom alous magnetic moment,

$$_{\rm V} = c_6 - \frac{g_{\rm A}^2 M m}{4 F^2}$$
 (4:8)

^{*} We choose here = m because of the matching conditions discussed in ref.[4.3]. Naturally, any other choice of would do as well since physical observables do not depend on the renorm alization scale.

where the second term is the leading non {analytic piece proportional to $\frac{p}{m}$ rst found by C aldi and Pagels [4.5]. Setting $c_6 = 5.62$, one reproduces the empirical value given after eq.(4.2). The loops generate a correction of about 34 %. The value of q' = 6is not quite surprising if one thinks of generating the corresponding contact term via -m eson exchange. The tensor coupling of the to the nucleon is ' 6. How ever, we should point out that such an estim ate depends cucially on how one chooses the N N and couplings. The isovector m agnetic radius,

$$< r^{2} > {}_{2}^{V} = \frac{6}{v} \frac{dF_{2}^{V}(t)}{dt}_{t=0}$$
 (4:9)

explodes like 1=M in the chiral limit [4.4] and is not a ected by any counter term contribution to order q^3 [4.2],

$$< r^{2} > V_{1}^{V} = \frac{g_{A}^{2} m}{8 F^{2} V} \frac{1}{M} = 0.50 \text{ fm}^{2}$$
 (4:10)

to be compared with the empirical value of $\langle r^2 \rangle_2^V = 0.77 \text{ fm}^2$. It is interesting to compare these results to the ones of the relativistic calculation [4.2]. It becomes obvious that the role of the loop versus the counter term contributions is rather di erent. In [4.2] it was shown that one has to choose c_6 ' 0 to get the empirical value of $_V$. This is due to the additional term s generated by the relativistic one loop diagram s beyond the leading non{analytic term in eq.(4.8) at 0 (q²) (compare the discussion about the power counting in section 3.1 and g.3.1). In the heavy ferm ion form alism, one loop diagram s with insertions solely from the lowest order elective Lagrangian only generate the term in the isovector magnetic moment. A similar statement also holds for the isovector radius, the heavy mass calculation to order q³ just gives the leading singularity.



Fig. 4.1: Two{pion cut contribution to the nucleon isovector electrom agnetic form factors.

Let us now take a closer look at the imaginary parts of the isovector form factors $F_{1,2}^{V}$ (t) (for similar discussions, see refs.[42,4.6,4.7]). As rst observed by Frazer and Fulco [4.8] and discussed in detail by Hohler and Pietarinen in connection with the nucleon electrom agnetic radii [4.9], Im $F_{1,2}^{V}$ (t) exhibits a very strong enhancement close

to threshold, $t_0 = 4M^2$ (the two{pion cut). To be speci c, consider F_2^{V} (t). At low momentum transfer, its absorptive part is dom inated by diagram s like the one shown in g.4.1, i.e.

Im
$$F_2^{V}(t) = \frac{2q_t^3}{(1)t}F(t) = \frac{1}{m}f_+^1(t) + \frac{1}{p}f_-^1(t)$$
 (4:11)

with $q_t = \frac{p}{t=4} = \frac{M^2}{M^2}$, F (t) the pion charge form factor and f^1 (t) the P {wave N partial waves in the t{channel. The latter are calculated from the standard N am plitudes A and B via projection involving ordinary Legendre polynom ials [4.10]. In this procedure, the nucleon pole term of the N am plitudes proportional to 1=(s m²) gives rise to Legendre functions of the second kind, Q_L (Z), which have logarithm ic singularities and a cut along 1 < Z < 1. Consequently, if one continues the partial waves f^4 (t) to the second sheet, they have a logarithm ic branch point at $Z = \cos t = m = (p_t q_t) = 1$, with $p_t = t=4$ m² and = (t 2M²)=4m. This translates into

$$t_c = 4M^2 M^4 = m^2 = 3.98M^2$$
 (4:12)

very close to the physical threshold at $t_0 = 4M^2$. The isovector form factors $F_{1;2}^V$ (t) inherit this logarithm ic singularity (branch point) on the second R iem ann sheet. A ctually, the same phenom enon occurs in the scalar form factor of the nucleon (for m ore details, see ref.[4.10]). Naturally, one asks the question whether this phenom enon shows up in the chiral expansion. Let us rst consider Im F_2^V (t) in the relativistic form ulation of baryon CHPT.Follow ing G asser et al. [4.2], one has

$$\operatorname{Im} F_{2}^{V}(t) = \frac{8g_{A}^{2}}{F^{2}} m^{4} 4 \operatorname{Im}_{4}(t) + \operatorname{Im}_{4}(t)$$
(4:13)

where the loop functions and their in aginary parts $_4$ and $_4$ are given in ref.[4.2]. For our purpose, we only need Im $_4$ (t) since its threshold is the two{pion cut whereas Im $_4$ (t) only starts to contribute at t = 4m². The resulting in aginary part for Im F_2^V (t)=t² is shown in g.4.2 (solid line). One sees that the strong increase at threshold is reproduced since the chiral representation of Im $_4$ (t) indeed has the proper analytical structure, i.e. the singularity on the second sheet at t_c [4.2]. The chiral representation of Im F_2^V (t)=t² does not stay constant on the left shoulder of the {resonance but rather drops. This is due to the fact that in the one loop approximation, one is only sensitive to the rst term in the chiral expansion of the pion charge form factor F^V (t). Indeed, if one sets F (t) 1 in eq.(4.11), the one loop result reproduces nicely the empirical curve (as discussed in more detail in ref.[4.7]). This particular example shows that in

In case of the nucleon scalar form factor, this singularity at $t_{\rm c}$ stems from the partial wave amplitude $f_{\scriptscriptstyle +}^0$ (t) .



Fig. 4.2: Chiral representation of Im F_2^{V} (t)=t² (t in units of the pion m ass) in the relativistic form ulation of baryon CHPT [4.2] (solid line) and in the heavy m ass approach (dashed line) [4.3,4.6].

the relativistic version of baryon CHPT the pertinent analytical structures of current and S{m atrix elements are given correctly.

Let us now turn to the heavy mass approach. The corresponding imaginary part follows from ref.[4.3],

$$\operatorname{Im} F_{2}^{V}(t) = \frac{g_{A}^{2}m}{32F^{2}} t 4M^{2} (t 4M^{2})$$
(4:14)

Here, the in aginary part behaves as q_t^2 close to threshold and not as q_t^3 as dem anded by (4.11). One furtherm ore nds that F_2^V (t) goes like $\ln (2M + t)$ and $t = 4M^2$ is a logarithm ic branch point in the heavy mass approach. This incorrect analytic structure is an unavoidable consequence of the heavy mass limit (m = 1), in which the square root branch point t_0 and the logarithm ic branch point t_c (on the second sheet) coincide. Nevertheless this leads to an enhancement of the in aginary part of F_2^V (t)=t² as shown by the dashed line in g.4.2. The enhancement is stronger than in the relativistic case and stronger than the empirical one (for F (t) 1). In order to get the proper separation of the singularities t_0 and t_c one should therefore perform an order q^5 calculation in the heavy mass approach.

W e also would like to discuss the imaginary part of the isovector D irac form factor F $_1^{\rm V}$ (t). In the heavy mass approach, it reads

$$\operatorname{Im} F_{1}^{V}(t) = \frac{P \frac{1}{1 + 4M^{2} = t}}{96 F^{2}} t + 4M^{2} + g_{A}^{2}(5t + 8M^{2})$$
(4:15)

O ne can show, that this form is exactly the m = 1 lim it of the corresponding expression given in [4.2]. The imaginary part Im F_1^V (t) in the heavy mass lim it shows an abnorm al threshold behaviour, close to threshold it grows linear in q_t and not like q_t^3 as dem anded by eq.(4.11). At the moment we are not able to give a precise explanation for this phenom enon, but certainly it must have to do with the coalescence of the two singularities t_0 and t_c and the behaviour of the t-channel amplitudes f^1 in the heavy mass lim it. The main lesson to be learned from this investigation of the imaginary parts is that the heavy mass formulation also has its own disadvantages. In the in nite nucleon mass lim it the analytical structure (poles and cuts) of certain amplitudes may be disturbed and this may be a hindrance in order to make contact to the dispersion theory.

Finally, a few words about the isoscalar form factors are in order. To one loop accuracy, they are determ ined mostly by some contact term s. For example, the isoscalar magnetic moment, s = p + n follows to be $s = 2c_7 + v$ (as de ned in L⁽²⁾_N) and the isoscalar charge radius $< r^2 > s^2_1$ is determined by the nite low energy constant $b_9 + 2b_9$ (cf. eq.(3.23)). To access the three pion{cut, at which the absorptive parts of the isoscalar form factors start, one has to perform a two loop calculation. Such a two loop calculation will also answer the yet unresolved question whether or not in the isoscalar channel there is an enhancement around $t = 9M^2$. State of the art dispersion theoretical analyses of the nucleon form factors assume only a set of poles in the corresponding spectral distributions [4.10]. Finally, we wish to stress that in this context the matching form alism discussed in ref.[4.3] starts to play a role (which allows to relate matrix (elements in the heavy mass and relativistic form ulation of CHPT) since ultim ately one might want to combine the chiral constraints with dispersion theory.

IV .2. NUCLEON COMPTON SCATTER ING

Low energy C om pton scattering of the nucleon is particularly well suited to investigate the structure of the nucleon since the electrom agnetic probe in the initial and in the nal state is well understood. In this section, we will rst discuss the general form alism of C om pton scattering and then elaborate on the nucleon structure as encoded in the so{called electrom agnetic (C om pton) polarizabilities (;) and the spin-dependent polarizability (). W e will only consider C om pton scattering which allows for a unique eld{theoretical de nition and em pirical extraction of these quantities. W e eschew here

the commonly used but theoretically uncertain non{relativistic treatment of these nucleon structure constants.*

The T-m atrix for the process (k) + p(p) ! $(k^0) + p(p^0)$ in the gauge $_0 = 0 = {}_0^0$ for realphotons, $k = 0 = {}^0$ % and in the centre {of {m ass system } $k_0 = k_0^0 = !$ and $t = (k k^0)^2 = 2!^2 (1 \cos)$ takes the form [4.11] $T = e^2 \sim {}^0 \sim {}_{I\!A} + \sim {}^0 \ \& \sim \& {}^0 A_2 + i \sim {}^0 \ \& \sim {}^0 A_3 + i \sim \& {}^0 \ \& {}^0 > {}^0 \ \& A_5 + i \sim {}^0 \ \& {}^0 > {}^0 \ \& {}^0 > {}^0 \ \& A_6$

(4:16)

^{*} We thus drop the overbar which is frequently used to denote the Compton polarizabilities.

using the operator basis of ref.[4.12]. The A_i are realbelow the pion production threshold, ! < M. From these, one can directly calculate physical obervables like the unpolarized di erential cross section as well as a set of asym metries for scattering polarized photons on polarized protons. The unpolarized di erential cross section in the cm system is

$$\frac{d}{d_{cm}} = \frac{^{2}m}{m+2E} \frac{1}{2}A_{1}^{2}(1+\cos^{2}) + \frac{1}{2}A_{3}^{2}(3-\cos^{2}) + !^{2}\sin^{2} 4A_{3}A_{6}$$

$$+ (A_{3}A_{4} + 2A_{3}A_{5} - A_{1}A_{2})\cos + !^{4}\sin^{2} \frac{1}{2}A_{2}^{2}\sin^{2} + \frac{1}{2}A_{4}^{2}(1+\cos^{2}) (4:17)$$

$$+ A_{5}^{2}(1+2\cos^{2}) + 3A_{6}^{2} + 2A_{6}(A_{4} + 3A_{5})\cos + 2A_{4}A_{5}\cos^{2}$$

with $= e^2 = 4$. The asymmetry for scattering circular polarized photons on polarized protons A_k (i.e. the proton spin parallel or antiparallel to the photon direction \tilde{k}) is given by

$$A_{k} = \frac{d_{m}}{d_{cm}} \frac{d_{m}}{d_{cm}} = \frac{2^{2}m}{m+2E} \qquad A_{3}^{2} \sin^{2} \qquad A_{1}A_{3} (1 + \cos^{2})$$

$$!^{2} \sin^{2} A_{6} (A_{1} + 3A_{3}) + (3A_{3}A_{5} - A_{1}A_{5} + A_{3}A_{4} - A_{2}A_{3}) \cos \qquad (4:18)$$

$$!^{4} \sin^{2} A_{5} (A_{2} - A_{4}) \sin^{2} + 4A_{5}A_{6} \cos + 2A_{6}^{2} + 2A_{5}^{2} \cos^{2})$$

Furtherm ore, we de ne the perpendicular asym m etry A $_{?}$ by considering right-circularly polarized photons m oving in the z-direction scattering on protons with their spin pointing in positive or negative x-direction, $\tilde{k}^{0} = !$ (sin $\cos ; \sin \sin ; \cos)$,

$$A_{?} = \frac{d_{"!}}{d_{cm}} \frac{d_{m}}{d_{cm}} = \frac{2^{2}m}{m+2E} A_{3}(A_{3} A_{1})\cos + !^{2}(A_{1}A_{5} + A_{2}A_{3})\sin^{2} + A_{3}A_{4}(1 + \cos^{2}) + A_{3}A_{5}(3\cos^{2} 1) + 2A_{3}A_{6}\cos$$
(4:19)
+ !^{4} sin^{2} A_{6}(A_{2} + A_{4} 2A_{5}) + A_{5}(A_{2} A_{4} 2A_{5})\cos sin \cos

is (in coordinate-free language) the azim uthal angle (around the axis de ned by the photon m om entum) m easured with respect to the plane spanned by the photon m om entum and the nucleon spin. Clearly, if one changes the di erence in eqs.(4.18,4.19) to a sum one gets in both case just twice the unpolarized cross section. Letting the nucleon spin point in y-direction results in $\cos 2$ sin in (4.19). If one uses left circular polarized photons instead of right circular polarized ones, then both asym m etries eqs.(4.18,4.19) change sign. One can also de ne a general asym m etry, which m eans right (circularly polarized photons m oving in z-direction scatter on polarized protons and

the proton spin lies in the xz-plane inclining an angle with the z-axis. We consider the di erence of cross section for this con guration and the one with reversed proton spin. The corresponding asymmetry reads

$$A() = \cos A_k + \sin A_2$$
 : (4.20)

A () gives the asymmetry for the most general spin alignment conguration.

In forward direction, the $\ensuremath{\mathsf{C}}$ om pton scattering am plitude takes the form

$$\frac{1}{4} T (!) = f_1 (!^2) \sim^0 \quad \sim + i !_2 f!^2 \sim \sim^0 \sim \sim)$$
(4.21)

where the spin {non ip $(f_1(!))$ and the spin { ip $(f_2(!))$ amplitudes have the low energy expansions,

$$f_{1}(!^{2}) = f_{1}(0) + (+)!^{2} + O(!^{4})$$

$$f_{2}(!^{2}) = f_{2}(0) + !^{2} + O(!^{4})$$
(4.22)

in terms of the electric (), the magnetic () and the so{called "spin{dependent" () polarizabilities. The Taylor coe cient $f_1(0)$ is given by gauge invariance,

$$f_1(0) = \frac{e^2 Z^2}{4 m}$$
(4.23)

which means that very soft photons only probe global properties like the charge (Z) and the mass m of the spin $\{1/2 \text{ particle they scatter o} \cdot \text{Eq.}(4.23) \text{ constitutes a venerable low {energy theorem (LET). There exists also a LET for the Taylor coe cient f₂(0) due to Low, G ell{M ann and G oldberger [4.13]. U sing gauge invariance, Lorentz invariance and crossing symmetry, they proved that$

$$f_2(0) = \frac{e^{2}}{8 m^2}$$
(4.24)

where denotes the anom alous magnetic moment of the spin $\{1/2 \text{ particle} \text{ the photon scatters o}$. The nucleon structure rst shows up in the the next $\{\text{to}\}$ leading order terms parametrized in term s of the various polarizabilities. U sing the optical theorem, one can derive the following sum rules

$$\operatorname{Ref}_{1}(!) = \frac{e^{2}Z^{2}}{4 \text{ m}} \frac{!^{2}}{2^{2}} P_{!_{0}}^{Z_{1}} d! \frac{0 \operatorname{tot}(!^{0})}{!^{0^{2}} !^{2}} \\ + \frac{1}{4^{2}} \frac{1}{4^{2}} \frac{d!}{!^{2}} [+ (!) + (!)] \\ = \frac{1}{4^{2}} \frac{1}{1^{2}} \frac{d!}{!^{3}} [+ (!) + (!)]$$

$$(4:25)$$

with tot(!) = (+(!) + (!))=2 the total photo {nucleon absorption cross section and (!) the photoabsorption cross section for scattering circularly polarized photons on

polarized nucleons for total N helicity 3/2 and 1/2, respectively. $!_0 = M + M^2 = 2m$ is the single pion production threshold. A ssum ing furtherm ore that the am plitude f_2 (!) satis es an unsubtracted dispersion relation, D rell, H earn and G erasim ov (D H G) derived the sum rule Z_1

$$I(0) = \int_{0}^{2} \frac{d!}{!} [(1) + (1)] = \frac{e^{2}}{2m^{2}} :$$
(4.26)

One can generalize this to the case of virtual photons ($k^2 < 0$) via

$$I(k^{2}) = \int_{0}^{2} \frac{d!}{!} \left[(!;k^{2}) + (!;k^{2}) \right]$$
(4.27)

with $!_0 = M + (M^2 k^2) = 2m$. This extended DHG sum rule will be discussed later on.

We turn now to the calculation of the cm s am plitudes in heavy baryon CHPT (the polarizabilities to this order are discussed for the relativistic approach in ref.[4.15] and for the heavy m ass calculation in [4.3]). For that, one has to consider nucleon-pole graphs (expanded up to $1=m^2$), ⁰ - exchange and pion loop diagram s. It is important to note that to this order the only contact term s entering are the ones which generate the anom alousm agnetic m om ent. The prediction for the various nucleon polarizabilities will therefore be given entirely in term s of low est order param eters. Consequently, one has a particularly sensitive test of the chiral dynam ics in the presence of nucleons. For the invariant functions A_i one nds (we only give the results for the proton)

$$A_{1} = \frac{1}{m} + \frac{g_{A}^{2}}{8 F^{2}} M \qquad p \frac{p}{M^{2}} \frac{1}{!^{2}} + \frac{2M}{p} \frac{2}{t} \frac{t}{t} \frac{1}{2} \arctan \frac{p}{2M} \frac{t}{2M}$$

$$Z_{1} dz \arctan \frac{(1 z)}{2 M^{2}} \frac{t}{!^{2} z^{2}}$$
(4:28a)

$$A_{2} = \frac{1}{m!^{2}} + \frac{g_{A}^{2}}{8 F^{2}} \frac{t}{(t)^{3=2}} \frac{2M^{2}}{0} dz \arctan \frac{(1 z)^{p} - t}{2^{p} M^{2} !^{2} z^{2}} - \frac{2(1 z)^{p} t(!^{2} z^{2} M^{2})}{4M^{2} 4!^{2} z^{2} t(1 z^{3})}$$

$$(4.28b)$$

$$A_{3} = \frac{!}{2m^{2}} 1 + 2 \qquad (1 + \frac{1}{7}\cos + \frac{g_{A}t!}{8^{2}F^{2}(M^{2} - t)} + \frac{g_{A}^{2}}{8^{2}F^{2}} \frac{M^{2}}{!} \arcsin^{2}\frac{!}{M} + \frac{g_{A}^{2}}{4^{2}F^{2}}!^{4}\sin^{2}\int_{0}^{Z} dx \int_{0}^{Z} dz \frac{x(1 - x)z(1 - z^{2})}{W^{3}} \arcsin\frac{!z}{R} + \frac{!zW}{R^{2}}$$

$$(4:28c)$$

$$A_{4} = \frac{(1+)^{2}}{2m^{2}!} + \frac{g_{A}^{2}}{4^{2}F^{2}} \int_{0}^{Z_{1}} dx \int_{0}^{Z_{1}} dz \frac{z(1-z)}{W} \arcsin\frac{!z}{R}$$
(4.28d)

$$A_{5} = \frac{(1+)^{2}}{2m^{2}!} \frac{g_{A}!}{8^{2}F^{2}(M^{2}-t)} + \frac{g_{A}^{2}}{8^{2}F^{2}} \int_{0}^{Z-1} dx \frac{(1-z)^{2}}{W} \arcsin\frac{!z}{R} + 2!^{2} \cos\frac{x(1-x)z(1-z^{2})}{W^{3}} \arcsin\frac{!z}{R} + \frac{!zW}{R^{2}}$$
(4:28e)

$$A_{6} = \frac{1+}{2m^{2}!} + \frac{g_{A}!}{8^{2}F^{2}(M^{2}-t)} + \frac{g_{A}^{2}}{8^{2}F^{2}} \int_{0}^{Z_{1}} dx \int_{0}^{Z_{1}} dz \frac{(1-z)^{2}}{W} \arcsin\frac{!z}{R}$$

$$2!^{2} \frac{x(1-x)z(1-z^{3})}{W^{3}} \arcsin\frac{!z}{R} + \frac{!zW}{R^{2}}$$
(4.28f)

with

$$W = {}^{p} \overline{M^{2} + t(1 + z)^{2} x(x + 1)}; \quad R = {}^{p} \overline{M^{2} + t(1 + z)^{2} x(x + 1)}$$
(4.28g)

From this, one can read o the polarizabilties as [43,4.15]

$$p = n = 10 \quad p = 10 \quad n = \frac{5e^2 g_A^2}{384 \ ^2F \ ^2M}$$

$$p = n = \frac{e^2 g_A^2}{96 \ ^3F \ ^2M \ ^2}$$
(4.29)

since the isospin factors in the diagram s contributing to , and are the same for the proton and the neutron. Before discussing the num erical results for the cross sections, asym m etries and polarizabilities, let us compare to the recent enum eration of third{ order spin polarizabilities by Ragusa [4.16]. Denoting by A_i the C om pton am plitudes with the B om term s subtracted, we can identify

$$a_{1;1} + a_{1;2} + a_3(0) = \frac{e^2}{8} \frac{e^2}{e!^2} A_1(0;0) = p + p = \frac{11e^2g_A^2}{768 e^2F^2M}$$
 (4:30a)

$$a_{1;1} + a_3(0) = \frac{e^2}{2} \frac{\theta}{\theta t} A_1(0;0) = P = \frac{e^2 g_A^2}{768 \ ^2 F^2 M}$$
 (4:30b)

$$a_{g}(0) = \frac{e^{2}}{4} A_{2}(0;0) = p = \frac{e^{2}g_{A}^{2}}{768 {}^{2}F^{2}M}$$
 (4:30c)

$$a_{2;2} + a_{1} + a_{2} + a_{4} = \frac{e^{2}}{24} \frac{e^{3}}{e^{2}} A_{3}(0;0) = \frac{e^{2}g_{A}^{2}}{96^{-3}F^{2}M^{-2}}$$
 (4:30d)

$$a_{2;2}$$
 2 $_{4} = \frac{e^{2}}{2} \frac{e^{2}}{0!} \frac{e^{2}}{2} A_{3}(0;0) = \frac{e^{2}g_{A}}{16^{-3}F^{2}M^{-2}}$ (4:30e)

$$_{3} = \frac{e^{2}}{4} \frac{0}{0!} A_{6}(0;0) = \frac{e^{2}g_{A}(12 + g_{A})}{384^{-3}F^{2}M^{-2}}$$
(4:30f)

$$_{2} = \frac{e^{2}}{4} \frac{0}{0!} \mathbb{A}_{4} (0;0) = \frac{e^{2} g_{A}^{2}}{192^{-3} F^{2} M^{-2}}$$
(4:30g)

$$_{4} = \frac{e^{2}}{4} \frac{0}{0!} A_{5}(0;0) = \frac{e^{2}g_{A}(12 + g_{A})}{384^{3}F^{2}M^{2}}$$
(4:30h)

O fparticular interest is the so-called C om pton am plitude $f_{E\,E}^1$ recently studied in [4.89] (and references therein) which displays a strong unitarity cusp. We like to discuss it brie y here. The set of C om pton functions introduced in (4.16) is com plete, consequently one can project out $f_{E\,E}^1$ via

$$f_{EE}^{1}(!) = \frac{e^{2}m}{32(! + \frac{p}{m^{2} + !^{2}})} \int_{1}^{Z+1} d$$

$$(A_{1} A_{3})(1 + {}^{2}) + !^{2}(2A_{5} + A_{4} A_{2}) (1 {}^{2})$$
(4:31)

The soft photon lim it in this case is $f_{EE}^1(0) = \hat{e} = (12 \text{ m})$. Evidently, the one-loop representation has a branch point at ! = M and therefore also a cusp. The num erical investigation indeed shows this cusp but it turns out to be rather weak at one-loop order $0 (q^3)$.

We now discuss the numerical results based on the one{loop (order q^3) invariant amplitudes, eqs.(4.28). First, if we set $g_A = 0$, only the nucleon Born graphs expanded in powers of 1=m contribute. For energies below the pion production threshold, the corresponding di erential cross section is within a few percent of the exact Powell cross section [4.17],

$$\frac{d}{d_{lab}} = \frac{1}{2} \frac{E^{0}}{mE} = \frac{1}{2} \frac{E^{0}}{mE} + \frac{E^{0}}{E} + \frac{E^{0}}{E} + \frac{1}{2} \frac{1}{m^{2}} + \frac{2EE^{0}}{m^{2}} (1 \cos_{L})^{2} + \frac{2EE^{0}}{m^{2}} 4(1 \cos_{L}) + \frac{1}{2} (1 \cos_{L})^{2} + \frac{3EE^{0}}{m^{2}} 2(1 \cos_{L}) + \frac{1}{2} (1 \cos_{L})^{2} + \frac{4EE^{0}}{2m^{2}} (1 + \frac{1}{2} \sin^{2}_{L})$$

$$E^{0} = \frac{E}{1 + \frac{E}{m} (1 \cos_{L})}$$

$$(4:32)$$



Fig. 4.3: The unpolarized data from ref.[4.18] in comparison to the chiral expansion of the C ompton amplitude for $_{\rm L} = 135$. The dashed, dash{dotted and solid lines represent the B orn, B om + 0 {exchange and B om + 0 {exchange + bop results. If one approximates the bop contribution by the electrom agnetic polarizabilities as described in eq.(4.33), the dotted line results.

where the scattering angle $_{\rm L}$ and the photon energy E in the laboratory system are related to the on squantities and ! via cos = (m + (m + E) (cos $_{\rm L}$ 1))=(m + E (1 cos $_{\rm L}$)) and ! = E = $\frac{P}{1 + 2E} = m$, respectively.

The chiral expansion to order q^3 (solid line) reproduces well the di erential cross section data of Federspiel et al.[4.18] as shown in g.4.3. (the corresponding results for $_L = 60$ look very sim ilar). In this gure it is also shown that the Born graphs together with the 0 (exchange contribution are not su cient to describe the data. However, as indicated by the dotted line in g.4.3, one is not sensitive to nucleon structure e ects beyond the electrom agnetic polarizabilities (the 1=M -term s). In this latter case, the loop contributions to the A_i are given by

$$A_{1}^{\text{loop}} = \frac{5g_{A}^{2}}{96 \text{ } \text{ } \text{ } ^{2}\text{M}} !^{2} 1 + \frac{1}{10} \cos \ \ \textbf{;} A_{2}^{\text{loop}} = \frac{g_{A}^{2}}{192 \text{ } \text{ } \text{ } ^{2}\text{M}} \textbf{;} A_{3;4;5;6}^{\text{loop}} = 0 : \qquad (4:33)$$

W e will discuss these polarizabilities in m ore detail later on. If one adds the contribution from static exchange (which starts at order q^4) the corrections are not dram atic. In

any case, for a truely meaningful comparison one would have to take into account a host of other q^4 terms. The parallel asymmetry generically changes sign around 90 degrees from negative to positive values as shown in g.4.4 for E = 70 MeV. This is not at all evident from eq.(4.18) since there is no overall cos {factor. For the same photon energy, we also show the perpendicular asymmetry in g.4.4. In both cases, the elects from the nucleon structure encoded in the loop contributions of the A₁ are small. Only for energies ! > 100 MeV one is somewhat sensitive to these structure terms. From an experimental point of view, only an extremely precise measurement of such asymmetries could shed light on the nucleon structure. An accuracy as for the unpolarized case [4.17] is certainly insu cient. If one trusts the q³ approximation up to the pion production threshold, one nds agreement with the few Saskatoon data [4.18] in this range. In this energy range, loop elects are more significant and could be detected experimentally. However, the competing contribution from the (and possible other q⁴ elects) becomes appreciable and makes the analysis of such data much less clear{cut.

Let us now take a closer look at the electrom agnetic polarizabilities of the proton and the neutron. These have been rather accurately determ ined over the last years. For the proton, if one combines the Illinois [4.18], M ainz [4.20] and Saskatoon [4.19] m easurements together with the dispersion sum rule, $(+)_p = (14.2 \ 0.3) \ 10 \ \text{fm}^3$ [4.21], one has

$$p_{p} = (10:4 \quad 0:6) \quad 10 \text{ fm}^{3}; \quad p = (3:8 \quad 0:6) \quad 10 \text{ fm}^{3} \quad (4:34a)$$

Similarly, the dispersion sum rule $(+)_n = (15.8 \ 0.5)$ 10 fm³* [4.22] together with the recent O ak R idge [4.23] and M ainz [4.24] m easurem ents leads to

$$n = (12:3 1:3) \quad 10 \text{ fm}^3; \quad n = (3:5 1:3) \quad 10 \text{ fm}^3 \quad : \quad (4:34b)$$

Notice that we have added the system atic and statistical errors of the empirical determ in nations in quadrature. The salient features of these experimental results are that both the proton and the neutron behave essentially as induced electric dipoles and that their respective sums of the electric and magnetic polarizabilities are almost the same. We should also point out that due to the strong magnetic M1N transition, one naively expects a large contribution from the resonance to the magnetic polarizabilities. These empirical features are already well represented by the lowest order (q³) results (4.29), i.e. $p = n = 13.6 \quad 10^{4} \text{ fm}^{-3}$ and $p = n = 1.4 \quad 10^{4} \text{ fm}^{-3}$. It is also worth to stress that the electrom agnetic polarizabilities explode as 1=M in the chiral limit since the photon sees the long {ranged pion cloud (com pare the chiral limit behaviour of the isovector form - factors discussed in section 4.1). However, at this order one has no isospin breaking (if one works in avor SU (2)) and solely nucleon interm ediate states can contribute in the

^{*} Notice that the uncertainty on the sum rule value for the neutron is presumably underestimated since one has to use deuteron data to extract the photon-neutron cross section.



Fig. 4.4: The parallel and perpendicular asymmetries for E = 70 M eV. For notations, see g.4.3. In case of A₂ we have set $\cos = 1$.

pertinent graphs. In ref.[425], a system atic analysis of all O (q⁴) e ects was presented. In addition to the one{loop diagram swith insertions from L $_{\rm N}^{(1)}$, one also has to include one loop graphs with exactly one insertion from L $_{\rm N}^{(2)}$ and contact terms from L $_{\rm N}^{(4)}$,

$$L_{N}^{(4)} = \frac{1}{4e^{2}} \begin{pmatrix} p & n \end{pmatrix} H f^{+} f_{+} H + \frac{1}{4e^{2}} & n H H Trf^{+} f_{+} \\ + \frac{1}{2e^{2}} \begin{pmatrix} n + n & p & p \end{pmatrix} H f^{+} f_{+} H v v$$

$$\frac{1}{2e^{2}} \begin{pmatrix} n + n \end{pmatrix} H H Tr(f^{+} f_{+}) v v$$
(4:35)

The unknown coe cients we have to determ ine are the four low-energy constants $_{p}$, $_{n}$, $_{p}$ and $_{n}$ (see below). The contact terms of order q² entering (c_{1;2;3}) have
already been determ ined (see section 3). From that, one derives the following form ulae for the electric and magnetic polarizabilities of the proton and the neutron (i = p;n)

with

$$C = \frac{e^{2}}{96^{-2}F^{-2}} = 4:36 \quad 10^{6} \text{ fm}^{-2};$$

$$x_{p} = 9; \quad x_{n} = 3; \quad y_{p} = 71; \quad y_{n} = 39;$$

$$x_{p}^{0} = 3 + s; \quad x_{n}^{0} = 1 \quad s; \quad y_{p}^{0} = \frac{37}{2} + 6s; \quad y_{n}^{0} = \frac{13}{2} \quad 6s;$$

$$c^{+} = 8q + 4c_{2} + 4c_{3} \quad \frac{g_{A}^{2}}{2m};$$

$$(4:37)$$

At this order, the loop contributions to the polarizabilities contain divergences which can be absorbed in the q^4 counter term s. The corresponding renorm alization prescription reads:

$$i = \frac{e^{2}L}{6 F^{2}} c_{2} \frac{x_{i}g_{A}^{2}}{m} + i(); \qquad i = \frac{e^{2}L}{6 F^{2}} c_{2} \frac{3x_{i}^{0}g_{A}^{2}}{m} + i() \quad (4:38)$$

The results shown in eqs.(4.37) have the following structure. Besides the leading 1=M term [4.3,4.15], 0 (q⁴) contributions from the loops have a lnM and a constant piece

M⁰. As a check one can recover the coe cient of the ln M term form the relativistic calculation [14.15] if one sets the new low energy constants c_i and $s_{iv} = 0$. In that case only the 1=m corrections of the relativistic D irac form ulation are treated and one necessarily reproduces the corresponding non {analytic (logarithm ic) term of this approach. The term proportional to $c_2 \ln M$ in eqs.(4.37) represents the e ect of (pion) loops with interm ediate (1232) states [4.26] consistently truncated at order q⁴. We should stress that the decomposition of the loop and counter term pieces at order q^4 has, of course, no deeper physical meaning but will serve us to separate the uncertainties stem ming from the coe cients accompanying the various contact terms. Notice that from now r_i^r () and r_i^r () appearing in eqs. (4.36, 4.38). on we will om it the superscript 'r' on The estimation of these low energy constants is discussed in great detail in ref.[4.27]. The (1232) enters prom inently in the determ ination of the four low-energy constants from L $_{\rm N}^{\rm (4)}$. Therefore, we will determ ine these coe cients at the scale = m (this induces som e spurious scale{dependence since we do not treat the rem ainders as e.g. in eq.(2.52)). In particular, one gets a sizeable contribution to the magnetic polarizabilities due to the strong N M 1 transition. A crude estimate of this has been given in

ref.[4.28] by integrating the M 1 part of the total photoproduction cross section for single pion photoproduction over the resonance region. However, this number is a icted with a large uncertainty. If one simply uses the Born diagram s with an interm ediate point{ like , one nds a result which strongly depends on the strength of the N coupling and on the o {shell parameter Y which is related to the electrom agnetic interaction L^{1}_{N} (we also give L^{2}_{N} used below) [4.32]

$$L^{1}_{N} = \frac{ieg_{1}}{2m} \qquad (Y) \quad {}_{5}T_{3} F + hc:$$

$$L^{2}_{N} = \frac{eg_{2}}{4m^{2}} \qquad (X) \quad {}_{5}T_{3} (0)F + hc: \qquad (4:39)$$

$$(I) = g + \frac{1}{2} (1 + 4I)A + I ; I = X;Y$$

where denotes the Rarita {Schwinger (spin {3/2) spinor, the T⁰s are the isospin 1=2 ! 3=2 transition operators. The parameter A does not appear in any physical observable and can therefore be chosen to be A = 1. These parameters (g;g₂;X;Y) are not very well determined. A conservative estimate therefore is

$$_{p}$$
 (m) = $_{n}$ (m) ' (7:0 7:0) 10 fm³ (4:40)

invoking isospin symmetry. Clearly, the large range in the value for is unsatisfactory and induces a mapr uncertainty in the determ ination of the corresponding counterterm s. We choose the central value of eq. (4.40) as our best determ ination [4.25, 4.27]. In ref.[4.29], the (1232) was included in the elective eld theory as a dynamical degree of freedom and treated non {relativistically (like the nucleon). There, it was argued that the Born graphs have to be calculated at the \circ {shell point ! = 0. This e ect can reduce the large by alm ost an order of m agnitude. This is rem in iscent of the o { shell dependence discussed before. Furtherm ore, as already pointed out in ref.[4.15], the relativistic treatment of the (1232) also induces a nite electric polarizability at order q^4 . This contribution depends strongly on the N couplings q_1 and q_2 as well as the twoo {shellparameters X; Y, cf. eq.(4.39). We thus assign an uncertainty of 2 10 fm³ to the theoretical predictions for the electric polarizabilities. A nother contribution to the coe cients $_{i}$ () and $_{i}$ () comes from loops involving charged kaons [4.30]. Since we are working in SU (2), the kaons and etas are frozen out and e ectively give some nite contact terms. These have been estimated in refs.[425,427]. One nds $_{p}^{K}$ (m) = 1:31 10⁴ fm³ and $_{n}^{K}$ (m) = 0:13 10⁴ fm³. The corresponding numbers for the kaon contributions to the magnetic polarizabilities are a factor 0:12 sm aller. These values might, how ever, considerably overestim ate the kaon loop contribution. Integrating e.g. the data from ref.[4.31] for $p \mid K ; K^{0}$, one gets a much smaller contribution since the typical cross sections are of the order of a few barn. This points towards the importance of a better understanding of SU (3) breaking e ects. At present,

this discrepancy remains to be resolved. Adding the various theoretical uncertainties in quadrature, one ends up with

$$p = (10:5 2:0) \quad 10 \text{ fm}^3 \quad p = (3:5 3:6) \quad 10 \text{ fm}^3$$

$$n = (13:4 1:5) \quad 10 \text{ fm}^3 \quad n = (7:8 3:6) \quad 10 \text{ fm}^3$$
(4:41)

which with the exception of n agree well with the empirical data (4.33,4.34). The in portant new e ect is that the loops of order q^4 generate a ln M term with a large coe cient (for p) which cancels most of the big contribution from the encoded in the coe cients of the L $\frac{(4)}{N}$ contact term s. In case of the neutron, the coe cient in front of the ln M {term is smaller. This points towards the possible in portance of isospin { couplings or in the o {shell parameter Y (for which at breaking in the p and n present we have no empirical indication). Clearly, an independent determ ination of the electric and magnetic nucleon polarizabilities would be needed to further tighten the em pirical bounds on these fundam ental quantities. This was also stressed in ref.[4.29]. It is worth to point out that the uncertainties given in (4.41) do not include e ects of two (and higher) loops which start out at order q^5 . We do not expect these to alter the prediction for the electric polarizabilities signi cantly [4.25]. Notice also that at present the theoretical uncertainties are larger than the experim ental ones (if one im poses the sum rules for (+)). That there is more spread in the empirical numbers when the dispersion sum rules are not used can e.g. be seen in the paper of Federspiel et al. in ref.[4.18]. The role of dispersion theory and its interplay to the chiral perturbation theory results is dicussed in refs.[4.33]

In ref.[4.27], the spin (averaged forward C om pton am plitude $A_{p;n}$ (!) = 4 f (!) was compared with the available data [4.21,4.34]. To lowest order q³ in the chiral expansion, the expressions for $A_{p;n}$ (!) diverge at ! = M . This is an artefact of the heavy mass expansion. The realistic branch point coincides with the opening of the one{pion channel as given after eq.(4.25). To cure this, one introduces the variable $= \frac{z}{1+M} = \frac{1}{2m} = \frac{1}{M} \cdot \frac{1}{(1+M} = 2m) = \frac{1}{10}$. In terms of , the branch point sits at its proper location and $A_{p;n}$ (= 1) is nite. We have

$$A_{p;n}(!) = \frac{e^{2}}{2m}(1 - 1) - 4 + p_{jn}!^{2} + \frac{e^{2}g_{A}^{2}M}{8 + r^{2}} - \frac{3}{2} - \frac{1}{2} + 1 + \frac{1}{2} + \frac{p}{1 - 2} + \frac{1}{4} \arcsin + \frac{11}{24} + \frac{e^{2}g_{A}^{2}M}{8 + r^{2}} - 1 + \frac{10}{3} + \frac{1}{4} + \frac{1}{1 - 2} + \frac{p}{1 - 2} \arcsin + \frac{11}{24} + \frac{1}{2} + \frac{1}{2}$$

where the '+/-' sign refers to the proton/neutron, respectively. The proper analytic continuation above the branch point = 1 is obtained through the substitutions 2 = $\frac{1}{2}$ 1 and arcsin = =2 + iln (+ $\frac{1}{2}$ 1). We should stress that in 1 ī the relativistic form ulation of baryon CHPT such problem s concerning the branch point do not occur [4.15]. In the heavy m ass form ulation one encounters this problem since the branch point ! 0 itself has an expansion in 1=m and is thus di erent in CHPT at order q^3 and q^4 . As shown in ref.[427], the spin {averaged forward C ompton amplitude for the proton is in agreem ent with the data up to photon energies ! ' $\,{\rm M}\,$. It is dominated by the quadratic contribution, i.e. to order q^4 in the chiral expansion the term s of order !⁴ (and higher) are sm all. Sim ilar trends are found for the neutron with the exception of a too strong curvature at the origin. For the proton, the real and im aginary parts of A (!) for > 1 have also been calculated. The imaginary part starts out negative as it should but becomes positive at ! ' 180 MeV. This is due to the truncation of the chiral expansion and can only be overcom e by a more accurate higher order calculation.

The spin (dependent polarizability has not yet been measured. To lowest order in the chiral expansion, explodes like $1=M^2$ in the chiral lim it [4.3],

$$_{\rm p} = _{\rm n} = \frac{e^2 g_{\rm A}^2}{96^{-3} {\rm F}^2 {\rm M}^2} = 4.4 \quad 10^4 {\rm fm}^4 \quad : \qquad (4:43)$$

The q⁴ and q⁵ corrections to this result have not yet been investigated in a system atic fashion. In ref.[4.3], the contribution from the was added (which starts at order q⁵) using the o {shell parameters of ref.[4.35] leading to $_{pm} = 3.66 \quad 10 \text{ fm}^4$, so that

$$p = \frac{1}{p} \frac{100p}{p} + p = 1:50 \quad 10 \text{ fm}^{4};$$

$$n = \frac{1}{n} \frac{100p}{p} + n = 0:46 \quad 10 \text{ fm}^{4};$$
(4:44)

which is rather di erent from the lowest order result, eq.(4.43). One can get an estim ate on the empirical values by use of the dispersion relation (4.25). Using the latest pion photoproduction multipoles from the SAID data basis, one arrives at [4.36]

$$_{\rm p} = 1:34 \quad 10 \text{ fm}^4; \quad _{\rm n} = 0:38 \quad 10 \text{ fm}^4: \quad (4:45)$$

The numbers given in (4.45) di er from the ones in ref.[4.3] because there an older version of the multipoles from the SAID program was used. The theoretical estimates (4.44) are an azingly close to the empirical ones, eq.(4.45). In ref.[4.3], the spin {dependent polarizabilities were also calculated in the relativistic approach. In that case, even on the level of avor SU (2), one nds some isospin {breaking from the one{loop diagram s.

Finally, we turn to a short discussion of the generalized DHG sum rule (4.27). Models for the photoabsorption cross sections [4.37,4.38,4.39] seem to indicate the validity of the DHG sum rule (4.26) (on the qualitative level). The direct experimental test of this sum rule has not yet been performed. Furthermore, the recent EMC measurements

in the scaling region $k^2 j'$ 10 G eV² indicate that the sum rule behaves a 1=k² as $k^2 j$ becomes large and that the sign is opposite to the value at the photon point, $k^2 = 0$. In ref.[4.41], baryon CHPT was used to investigate the slope of I (k²) around the origin. It was found to be negative and of sim ilar size to the recently proposed value by So er and Teryaev [4.42], but opposite to the one of ref.[4.39] (which is due to the contribution). At present, experimental data as well as more detailed theoretical investigation are lacking and we refer the interested reader to [4.41] for more details.

IV .3. AXIAL PROPERTIES OF THE NUCLEON

In the previous sections, we were concerned with the coupling of photons to the nucleon, i.e. pure electrom agnetic processes. W ithin the fram ework of the standard m odel, there also axial currents which can be used as probes. The structure of the nucleon as probed by charged axial currents is encoded in two form factors, the axial and the induced pseudoscalar ones. To be speci c, consider the m atrix {elem ent of the isovector axial quark current, $A^a = q = \frac{1}{5} (a=2)q$, between nucleon states

< N (
$$p^{0}$$
) jA^{a} (p) > = u (p^{0}) G_A (t) + $\frac{(p^{0} p)}{2m}$ G_P (t) $_{5}\frac{a}{2}$ u (p) (4:46)

with $t = (p^0 p)^2$ the invariant m om entum transfer squared. The form of eq.(4.46) follows from Lorentz invariance, isospin conservation and the discrete symmetries C, P and T.* G_A (t) is called the nucleon axial form factor and G_P (t) the induced pseudoscalar form factor. We rst discuss the axial form factor, which probes the spin { isospin distribution of the nucleon (since in a non{relativistic language, this is nothing but the m atrix {element of the G am ov{Teller operator ~ a). The sm all m om entum expansion of the axial form factor takes the form

$$G_A(t) = g_A + \frac{t}{6} < r_A^2 > + \dots$$
 (4:47)

with g_A the axial{vector coupling constant, $g_A = 1.2573$ 0.0028 [4.42], $r_A = \langle r_A^2 \rangle^{1=2}$ the axial root{m ean-square (m s) radius and the ellipsis stands for term s quadratic (and higher) in t. The axial m s radius can be determ ined from elastic (anti) {neutrino{proton scattering or from charged pion electroproduction. W hile the form er m ethod gives $r_A = 0.65$ 0.03 fm [4.43], the latter leads to som ewhat sm aller values $r_A = 0.59$ 0.05 fm [4.1,4.44,4.49]. This apparent discrepancy will be discussed in the next section. In any case, we note that the typical size of the nucleon when probed with the weak charged currents is sm aller than the typical electrom agnetic size, $r_{em} = 0.8$ fm. There is, of course, no a priori reason why these sizes should coincide. This hierachy of nucleon

^{*} We do not consider operators related to so { called second class currents since these are not observed in nature.

radii nds a natural explanation in the topological soliton model of the nucleon [4.45] since there the electrom agnetic size is proportional to $r_B^2 + 6=M_{!}^2$ (0.8)² fm² whereas the axial radius is roughly given by $r_B^2 + 6=(M_{!} + M_{!})^2$ (0.7)² fm², with r_B '0.5 fm the size related to the distribution of topological charge (baryon number). Empirically, the axial form factor can be rather accurately parametrized by a dipole form

$$G_{A}(t) = \frac{g_{A}}{(1 \quad t=M_{A}^{2})^{2}}$$
 (4:48)

and the cut{o $m ass M_A$ is thus related to the axial m s radius via

$$< r_{A}^{2} > = \frac{12}{M_{A}^{2}}$$
 : (4:49)

In heavy baryon CHPT and to one{loop accuracy q^3 , the momentum dependence of the axial form factor is essentially given by some contact terms (sim ilar to the isoscalar form factors discussed in section 4.1) since the absorptive part of G_A (t) starts at the three{pion cut, $t_0 = 9M^2$ (accessible rst at two{bop order),

$$g_{A} = g_{A} Z_{N} + \frac{M^{2}}{32^{2}F^{2}} g_{A}^{2} 4 \ln \frac{M}{2} + g_{A}^{2} + \frac{M^{2}}{4^{2}F^{2}} B_{9}^{r} ()$$

$$< r_{A}^{2} > = \frac{6}{g_{A}} B_{24}$$
(4:50)

with Z_N the nucleon Z {factor (3.55) and the pertinent counter terms in L⁽³⁾_N are O_9 of eq.(3.62) and $B_{24}HS$ [D]; f]H. Stated di erently, to order q⁴ in heavy baryon CHPT, G_A (t) is linear in t since the cut starting at $t_0 = (3M)^2$ rst shows up in the chiral expansion at two{loop order O (q⁵). Therefore, the contribution to order q⁴ m ust be polynom ial in t. The one{loop expression for G_A (t) in the relavistic form ulation can be found in ref.[4.46].

Concerning the induced pseudoscalar form factor, it is generally believed to be understood well in terms of pion pole dominance as indicated from ordinary muon capture experiments, +p! + n (see e.g. refs.[4.47,4.48,4.49]). However, it now seems feasible to measure the induced pseudoscalar coupling constant (the form factor evaluated at $t = 0.38M^2$) within a few percent accuracy via new techniques which allow to minimize the uncertainty in the neutron detection [4.50]. In fact, one is able to calculate this fundamental quantity within a few percent accuracy by making use of the chiral W and identities of QCD. The pseudoscalar coupling constant as measured in ordinary muon capture is dened via

$$g_P = \frac{M}{2m} G_P (t = 0.88M^2)$$
 : (4.51)

The value oft can be understood as follows. If the muon and the proton are initially at rest, energy and momentum conservation lead to

$$t = M^{2} 1 \frac{(M + m_{p})^{2} m_{n}^{2}}{M (M + m_{p})} = 0.88 M^{2} :$$
 (4:52)

To accurately predict g_P in terms of well{known physical parameters, we exploit the chiralW and identity of QCD,

$$[q _{5} \frac{a}{2}q] = m qi_{5} aq :$$
 (4:53)

Sandwiching this between nucleon states, one obtains [42, 43]

$$m G_A (t) + \frac{t}{4m} G_P (t) = 2m B m g_A \frac{1 + g(t)}{M^2 t}$$
 (4:54)

The pion pole in eq.(4.54) originates from the direct coupling of the pseudoscalar density to the pion, eq.(3.86). The residue at the pion pole t = M² is m² G g _N = g _N F M² as discussed in section 3.7. Furtherm ore, to order q⁴, G_A (t) as well as g(t) are linear in t as discussed above. Therefore,

$$m g_{A} + m g_{A} \frac{r_{A}^{2}}{6} t + \frac{t}{4m} G_{P} (t) = \frac{g_{N} F}{M^{2} t} t + g_{N} F + \frac{2B_{23} M^{2} g_{N}}{F}$$
(4:55)

where we have used $2mBg_Am = M^2(g_NF + O(M^2))$. At t = 0, eq.(4.55) reduces to the Goldberger{Treim an discrepancy discussed in section 3.7. G_P (t) can now be isolated,

$$G_{P}(t) = \frac{4m g_{N} F}{M^{2} t} - \frac{2}{3} g_{A} m^{2} r_{A}^{2} + O(t; M^{2})$$
(4:56)

A few remarks are in order. First, notice that only physical and well{determ ined param – eters enter in eq.(4.56). Second, while the rst term on the right{hand{side of eq.(4.56)} is of order q², the second one is 0 (q⁰) and the corrections not calculated are of order q². For g_P, this leads to [4.51]

$$g_{\rm P} = \frac{2M \ g_{\rm N} F}{M^2 + 0.88M^2} - \frac{1}{3} g_{\rm A} M \ m \ r_{\rm A}^2$$
 : (4:57)

Indeed, this relation has been derived long time ago by Adler and Dothan [4.52] with the help of PCAC and by W olfenstein [4.53] using a once{subtracted dispersion relation for the right{hand{side of eq.(4.54) (weak PCAC). It is gratifying that the result of refs.[4.52,4.53] can be m by based on the system atic chiral expansion of low energy QCD G reen functions. In chiral perturbation theory, one could in principle calculate the

corrections to (4.57) by perform ing a two{loop calculation while in A dler and D othan's or W olfenstein's m ethod these either depend (com pletely) on the PCAC assumption or could only be estimated. To stress it again, the main ingredient to arrive at eq.(4.57) is the linear t{dependence of G_A (t) and g(t). Since we are interested here in a very sm all m om entum transfer, t = $0.88M^2$ ' $0.5M^2$, curvature terms of order t² have to be negligible. If one uses for example the dipole param etrization for the axial form factor, eq.(4.48), the t² {term amounts to a 1.3% correction to the one linear in t. Consequently, our results can also be used in radiative m uon capture o hydrogen where the four{ m om entum transfer varies between $M^2 : :: + M^2$. U sing now the PDG values [4.42] for m, M , M = M +, F and g_A together with g_N = 13:31 0:34* and r_A from the (anti)neutrino{nucleon scattering experiments, we arrive at

$$g_{\rm P} = (8:89 \ 0.23) \quad (0:45 \ 0.04) = 8:44 \ 0.23 : (4:58)$$

The uncertainties in eq.(4.58) stem from the range of q_N and from the one for r_A for the rst and second term, in order. For the nal result on $q_{\rm P}$, we have added these uncertainties in quadrature. A measurement with a 2% accuracy of q_P could therefore cleanly separate between the pion pole contribution and the improved CHPT result. This would mean a signi cant progress in our understanding of this fundam ental low (energy parameter since the presently available determ inations have too large error bars to disentangle these values (see e.g. refs.[4.47,4.49]). In fact, one might turn the argum ent around and eventually use a precise determ ination of gp to get an additional determ ination of the strong pion {nucleon coupling constant which has been at the center of much controversy over the last years. The momentum dependence of G_P (t) for t $0:18 \text{ GeV}^2$ has recently been measured [4.49]. The error bars are, 0**:**07 and between however, too large to disentangle between the pion pole prediction and the one given in eq.(4.56). A more accurate determination of the induced pseudoscalar form factor would therefore help to clarify our understanding of the low {energy structure of QCD.

IV .4. THRESHOLD PION PHOTO { AND ELECTROPRODUCTION

In this section, we will be concerned with reactions involving photons, nucleons and pions, i.e. the interplay between vector and axial{vector currents. This has been a topical eld in particle physics in the late sixties and early seventies before the advent of scaling in deep inelastic scattering, see e.g. ref.[4.1]. However, over the last few years renewed interest in the production of pions by real or virtual photons in the threshold region has emerged. This was rst triggered through precise new data on neutral pion photoproduction [4.54,4.55], which lead to a controversy about their theoretical interpretations. Furthermore, new precise data on ⁰ electroproduction [4.56] have given further constraints on the understanding of these fundam ental processes in the non{ perturbative regin e of QCD. In fact, as we will demonstrate, chiral perturbation theory m ethods are best suited to analyze these reactions in the threshold region.

^{*} See the discussion on this in ref.[4.51]

First, we have to supply some basic de nitions. For more details, we refer to refs.[4.1,4.46,4.57,4.58]. Consider the process $(k) + N(p_1) ! = a(q) + N(p_2)$, with N denoting a nucleon (proton or neutron), $a = a pion with an isospin index a and the virtual photon with <math>k^2 < 0$. In the case of photoproduction (real photons), we have $k^2 = 0$ and k = 0. A detailed exposition of the corresponding kinematics can e.g. be found in ref.[4.59]. The pertinent M andelstam variables are $s = (p_1 + k)^2$; $t = (p_1 - p_2)^2$ and $u = (p_1 - q)^2$ subject to the constraint $s + t + u = 2m^2 + M^2 + k^2$. U sing Lorentz invariance and the discrete symmetries P, C and T, the transition current m atrix element can be expressed in term s of six independent invariant functions, conventionally denoted by $A_i(s;u)$; (i = 1; :::; 6), when one m akes use of gauge invariance,

$$J = iu_{2} \int_{i=1}^{X^{6}} M_{i} A_{i}(s; u) u_{1}$$
(4:59)

with

$$M_{1} = \frac{1}{2} (k_{1} + k_{2}); \quad M_{2} = P (2q + k^{2}); \quad P + k(2q + k);$$

$$M_{3} = q + k_{2} + k_{3} + m_{4} = 2 P + k_{2} + k_{3} + m_{5} + m_{5}; \quad (4:60)$$

$$M_{5} = k q + k_{3} + k_{3}^{2}; \quad M_{6} = k_{5} + k_{5} + k_{2}^{2}$$

and $P = (p_1 + p_2)=2$. The amplitudes $A_i(s;u)$ have the conventional isospin decomposition (to rst order in electrom agnetism),

$$A_{i}(s;u) = A_{i}^{(+)}(s;u)_{a3} + A_{i}^{()}(s;u) \frac{1}{2}[_{a};_{3}] + A_{i}^{(0)}(s;u)_{a}:$$
(4:61)

Under (s $\$ u) crossing the am plitudes $A_{1;2;4}^{(+;0)}$; $A_{3;5;6}^{()}$ are even, while $A_{3;5;6}^{(+;0)}$; $A_{1;2;4}^{()}$ are odd. For photoproduction, the num ber of independent am plitudes is further reduced to four. In term s of the isospin components, the physical channels under consideration are

$$J (p! ^{0}p) = J^{(0)} + J^{(+)}$$

$$J (n! ^{0}n) = J^{(+)} J^{(0)}$$

$$J (p! ^{+}n) = \stackrel{p}{2} [J^{(0)} + J^{(-)}]$$

$$J (n! p) = \stackrel{p}{2} [J^{(0)} J^{(-)}]:$$
(4:62)

Having constructed the current transition matrix element J it is then straightforward to calculate observables. The pertinent kinematics and de nitions are outlined in refs. [4.57, 4.58].

For the discussion of the low energy theorem s, we have to spellout the corresponding multipole decomposition of the transition current matrix element at threshold. In the N center of m ass system at threshold i:e:q = (M ; 0;0;0) one can express the current m atrix element in terms of the two S{wave multipole amplitudes, called E₀₊ and L₀₊,

$$\mathcal{J} = 4 \text{ i}(1 +) \stackrel{\text{y}}{_{\text{f}}} E_{0+}(;) \sim + L_{0+}(;) E_{0+}(;) \hat{k} \sim \hat{k}_{i} \qquad (4:63)$$

with $_{i;f}$ two component Pauli-spinors for the nucleon and we chose the C oulom b gauge $_{0} = 0$. For the later discussion we have introduced the dimensionless quantities

$$= \frac{M}{m}; = \frac{k^2}{m^2}:$$
 (4:64)

The multipole E_{0+} characterizes the transverse and L_{0+} the longitudinal coupling of the virtual photon to the nucleon spin. A lternatively to L_{0+} , one also uses the scalar multipole S_{0+} de ned via, S_{0+} (s;k²) = ($\mathfrak{K} \neq k_0$) L_{0+} (s;k²). At threshold, we can express E_{0+} and L_{0+} through the invariant amplitudes A_i (s;u) via (suppressing the isospin indices)

$$E_{0+} = \frac{m^{p}}{8} \frac{(2+)^{2}}{(1+)^{3=2}} \qquad A_{1} + m \frac{(2+)+}{2(1+)} A_{3} + m \frac{(2^{2})}{2(1+)} A_{4} \qquad m A_{6} ;$$

$$L_{0+} = E_{0+} + \frac{m^{p}}{16} \frac{(2+)^{2}}{(1+)^{5=2}} (2^{2}) A_{1} + m^{2} \frac{(2^{2})((2+)+)}{4(1+)} A_{2} \qquad m A_{4} + m^{2} \frac{(2+)+}{2(1+)} A_{5} \qquad m (2+) A_{6} \qquad (4:65)$$

with the A_i (s;u) evaluated at threshold $s_{th} = m^2 (1 +)^2$ and $u_{th} = m^2 (1)^2$

)=(1+). In case of photoproduction, only the electric dipole amplitude $E_{0+}\,$ survives. Finally, one can de ne the S{wave cross section,

$$a_{0} = E_{0+} \hat{J} \qquad \frac{k^{2}}{k_{0}^{2}} L_{0+} \hat{J} \qquad (4.66)$$

where and $k_0 = (s m^2 + k^2) = 2^{p} \overline{s}$ represent, respectively, a measure of the transverse linear polarization and the energy of the virtual photon in the N rest frame. For $k^2 = 0$, this means in particular that $(\tilde{\chi} = \tilde{\eta} \eta) (d = d) = (E_{0+})^2$ as q tends to zero. This completes the necessary form alism.

We discuss now the electric dipole amplitude E_{0+} as measured in neutral pion photoproduction, + p ! ⁰ + p. This multipole is of particular interest since in the early seventies a low {energy theorem (LET) for neutral pion production was derived [4.60].* The recent measurements at Saclay and Mainz [4.54,4.55] seem ed to indicate

^{*} For the sake of brevity, we denote E $_{0+}^{^{0}p}$ by E $_{0+}$.

a gross violation of this LET, which predicts $E_{0+} = 2.3 \quad 10^{+} + at$ threshold. However, the LET was reconsidered (and rederived) and the data were reexam ined, leading to $E_{0+} = (2.0 \quad 0.2) \quad 1^{2}0 + M + in$ agreement with the LET prediction. These developments have been subject of a recent review by D rechsel and T iator [4.61]. Therefore, we will focus here on the additional insight gained from CHPT calculations. In fact, the \LET " derived in [4.60] for neutral pion photoproduction at threshold is an expansion in powers of $= M = m \quad 1=7$ and predicts the coe cients of the rst two terms in this series, which are of order and 2 , respectively, in terms of m easurable quantities like the pion (nucleon coupling constant g_N , the nucleon m ass m and the anom alous m agnetic m om ent of the proton, $_{p}$,

$$E_{0+}(s_{thr}) = \frac{eg_N}{8m} - 1 - \frac{1}{2}(3+p) + O(2)$$
 : (4:67)

In ref.[4.62] it was, however, shown that a certain class of loop diagram s m odi es the LET at next{to{leading order O (2). It is instructive to rederive this result in heavy baryon CHPT [4.3]. Insertions from L $^{(2)}_{N}$ and L $^{(3)}_{N}$ lead to eq.(4.67) when the corresponding quantities are given by their chiral lim it values. However, to order q³, one also has to consider one{loop graphs. The standard derivation of eq.(4.67) is based on the consideration of nucleon pole graphs (supplemented by form factors). We stress that such considerations are not based on a system atic chiral counting. In the threshold region, only the so{called triangle and rescattering diagram s are non-vanishing (com pare the detailed discussion of selection rules in ref.[4.3]) leading to

$$E_{0+}^{\text{loop}} = \frac{eg_{A}}{8 F^{3}} v \quad kJ_{0} (v \quad k) + {}_{0}J(v \quad k) + 2 (v \quad k) + 2 (v \quad k) + 2 (v \quad k)$$
(4:68)

with the loop functions J_0 and $_3$ given in appendix B.At threshold, v = M, so that $J_0(M) + J_0(M) = 0$ and $_3(M) + _3(M) = M$ =32, i.e. only the triangle diagram and its crossed partner contribute. Therefore, these particular one loop diagram s contribute at order 2 ,

$$E_{0+}^{loop} = \frac{eg_{A} M^{2}}{128 F^{3}} = \frac{eg_{A} m^{2}}{128 F^{3}}^{2}$$
(4:69)

Let us look closer at the origin of the nite contribution proportional to M 2 . We follow the argument of ref.[4.62]. In the relativistic calculation, the loops lead to an expression of the form

$$E_{0+}^{loop} = \frac{eg_A M^2}{(4 F)^3} f() f() \qquad (4:70)$$

with

$$f() = \int_{0}^{Z_{1}} dx \int_{0}^{Z_{1}} dy \frac{2xy}{2 + y^{2} + y^{2}} : \qquad (4.70a)$$

Naively, one would argue E_{0+}^{loop} M³, since the form ula (4.70) is manifestly odd under M ! M and this would forbid an even term proportional to M². However, in this argum entation one has already made an assumption, namely that the function f() is analytic at = 0. The explicit form (4.70a) shows that this is not true, f() has a logarithm ic singularity at = 0 and the correct expansion around this point reads

$$f() = \ln + \frac{3^{2}}{8} + \frac{1}{2} + O(); \quad f() = \ln - \frac{2}{8} + \frac{1}{2} + O(): \quad (4:70b)$$

Consequently, we not that the odd combination f() f() has a nonvanishing lim it lim $_{! 0^{+}} [f() f()] = {}^{2}=2 \notin 0$, which is quite astonishing. Of course, without explicit know ledge of f() from the complete loop-calculation in CHPT one would hardly not the peculiar properties of f(). In the standard derivation (and all rederivations) of the incomplete LET one tacitely assumes (without better know ledge) lim $_{! 0^{+}} [f() f()] = 0$. The additional term is non-analytic in the quark mass m since $A_{1,thr}^{loop}$ m [4.62]. Consequently, the correct expansion of E_{0+} in QCD takes the form

$$E_{0+}(s_{thr}) = \frac{eg_N}{8m} = 1 = \frac{1}{2}(3+p) + \frac{m}{4F}^2 + O(2^2) : (4:71)$$

One in mediately notices that the term of order 2 is even bigger (+ 4.35) than the leading order one (-3.46) and of opposite sign. This makes the $\{expansion of E_{0+} truncated at \}$ order ² practically useless for a direct com parison with the data. We also stress that the closeness of the prediction based on the incom plete expansion (4.67) and the reexam ined data has to be considered accidental and is devoid of any physical signi cance. In fact, in ref.[4.59] it was argued that the {expansion of E_{0+} is slow by converging. This has been even further quantied in a calculation [4.63] in the fram ework of heavy baryon CHPT including all terms of order q^4 and including isospin {breaking by di erentiating between the charged and neutral pion masses as proposed in ref.[4.64]. Furthermore, the theoretical prediction of the electric dipole amplitude in 0 production o protons is a icted with some uncertainty related to the contribution to estimate the appearing contact terms. At present, it does not seem to serve as a stringent test of the chiral pion (nucleon dynam ics. M ore accurate data close to threshold are needed to clarify these questions and also the energy {dependence of E_{0+} from the ⁰p threshold (E = 144:68) MeV) to the +n threshold at E = 151:44 MeV. While below this threshold the multipoles are real, above it they in general become complex. To one { loop accuracy, one expects a cusp at the + n threshold with E_{0+} (E = 151:44 MeV) E₀₊ (E = 144:68 MeV) ' 0:7 $1\hat{\theta} = M + [4.63]$. The various analysis of the Mainz data [4.55] give very di erent results, e.g. while Bergstrom [4.65] nds a very steep energy dependence, the analysis of Bernstein leads to an essentially at E_{0+} (E) [4.66]. The situation is

The meaning of low (energy theorems in the framework of the Standard M odel is discussed in ref.[4.90].

di erent for the P {waves. From the two magnetic multipoles M $_{1+}$ and M $_1$ and the electric E $_{1+}$ one form s the combinations

$$P_{1} = 3E_{1+} + M_{1+} \qquad M_{1}$$

$$P_{2} = 3E_{1+} \qquad M_{1+} + M_{1} \qquad (4:72)$$

$$P_{3} = 2M_{1+} + M_{1}$$

To order q^3 and including the pion m ass di erence in the loops, one nds [4.63]

$$P_{1} = \begin{pmatrix} q \\ 1^{2} \\ M^{2}_{0} \\ R \\ \frac{eg_{N}}{8 \\ m^{2}} \\ \frac{eg_{N}}{8 \\ m^{2}} \\ 1 \\ \frac{eg_{A}^{3}}{64 \\ 2F^{3}} \\ \frac{2}{3!^{2}} \\ M^{3}_{+} \\ M^{2}_{+} \\ \frac{!^{2}}{1^{2}} \\ M^{2}_{+} \\ \frac{eg_{A}}{1} \\ \frac{eg_{N}}{8 \\ m^{2}} \\ \frac{eg_{N}}{8 \\ m^{2}} \\ 1 \\ \frac{13!}{10m} \\ \frac{1}{5m!} \\ \frac{M^{2}_{0}}{5m!} \\ \frac{eg_{N}}{1 \\ \frac{1}{2m}} \\ \frac{1}{2m} \\ \frac{eg_{A}^{3}}{64 \\ 2F^{3}} \\ \frac{2}{3!^{2}} \\ M^{3}_{+} \\ M^{2}_{+} \\ \frac{!^{2}}{1^{2}} \\ \frac{M^{2}_{0}}{5m!} \\ \frac{1}{2m} \\ \frac{1}{2m}$$

with $! = (s m^2 + M_0^2) = (2^{p} \overline{s})$ the pion energy in the cm system * A closer look at (4.73) reveals that the term s of order q^3 in the threshold region are very sm all com pared to the leading O (q^2) ones, i.e. one can derive the LETs

$$\frac{1}{q}P_{1} = \frac{eg_{N}}{8m^{2}} + p + 1 = \frac{p}{2} + \frac{g_{N}^{2}(10 - 3)}{48}$$

$$\frac{1}{q}P_{2} = \frac{eg_{N}}{8m^{2}} + p + \frac{p}{2} + p = \frac{g_{N}^{2}}{12}$$
(4:73a)

and similarly for the reaction $n! ^{0}n$,

$$\frac{1}{q}P_{1} = \frac{eg_{N}}{8 m^{2}} + \frac{1}{2} n + \frac{g_{N}^{2} (10 - 3)}{24}$$

$$\frac{1}{q}P_{2} = \frac{eg_{N}}{8 m^{2}} + \frac{1}{2} n + \frac{g_{N}^{2}}{12}$$
(4:73b)

with $_n = 1.913$ the anom alous magnetic m om ent of the neutron. These are examples of quickly converging expansions. In fact, the corresponding P_1 and P_2 of the relativistic calculation agree quite nicely with the LET (remember that in the relativistic

^{* !} is related to the frequently used photon energy in the lab system via ! = (2m E + M $_0^2$)=(2 m 2 + 2m E).

form ulation some higher order term s are included). For example, at E = 151 M eV, the LET predicts $P_1 = 2.47$ and $P_2 = 2.48$ while the P {wave multipoles of ref.[4.59] lead to $P_1 = 2.43$ and $P_2 = 2.60$ (all in units of $10^3 = M_+$) (for $p! ^0p$). In contrast, P_3 is completely dominated by the contribution. Its generic form is $P_3 = q!$ const, where the constant can either be tted from the bell{shaped di erential cross sections about the $^+$ n threshold or by using resonance saturation. Both ways lead to essentially the same number. A lternatively, one could t the coe cient in P_3 from the total cross section data if one excludes the very rst few M eV above threshold. For m ore details on this, we refer the reader to ref.[4.63].

The total cross section for $p ! ^{0}p [4.53,4.54]$ is only sensitive to the value of E_{0+} (s_{thr}) very close to threshold and then dom inated by the P-wave combination $(p_{1}f + p_{2}f + p_{3}f)=3$. This is shown in gure 4.5 for the calculation of refs.[4.59,4.64]. In fact, the estimate of the low {energy constant d₄ in [4.59] should be considered as a best t to these total cross section data. The corresponding di erential cross sections calculated in ref.[4.59] do not agree well with the data since E_{0+} was essentially energy independent 1:3, too large in magnitude to what is needed to produce the bell{ shaped angular distributions. A more detailed account of these topics will be given in ref.[4.63]. Finally, we point out that new data for $p ! ^{0}p$ have been taken at MAM I (M ainz) and SAL (Saskatoon). These are presently in the process of being analyzed.



Fig. 4.5: Total cross section for $p ! ^{0}p$. The solid and dashed curve represent the one{bop CHPT predictions in the isospin lim it and with isospin{breaking in the class I diagram s, respectively [4.59,4.64]. The dotted line is the tree level predictions. The Mainz [4.54] and the Saclay [4.53] data are represented by diam onds and squares, respectively.

The situation is di erent in the case of charged pion production, p! ⁺ n and n! p. Here, there exists a fam ous LET due to K roll and Ruderm an [4.67], which states that the corresponding electric dipole am plitudes do not vanish in the chiral lim it and, furtherm ore, that this leading term of order ⁰ is dom inant. The chiral corrections do not a ect this result as shown in ref.[4.59]. In fact, the quark m ass expansion of E₀₊ for these two channels takes the form (at threshold)

$$E_{0+}^{+n} = {}^{p} \overline{2} \frac{eg_{N}}{8m} 1 \frac{3}{2} + O({}^{2}; {}^{2}\ln) = 26:6 10^{3} = M + ;$$

$$E_{0+}^{-p} = {}^{p} \overline{2} \frac{eg_{N}}{8m} 1 + \frac{1}{2} + O({}^{2}; {}^{2}\ln) = 31:5 10^{3} = M + ;$$
(4:74)

The full one { loop corrections (i.e. no expansion in $\$) have been worked out with the result [4.59]

$$E_{0+}^{+n} = 28:4 \quad 10^{3} = M_{+}; E_{0+}^{p} = 31:1 \quad 10^{3} = M_{+}; \quad (4:75)$$

which com pare, naturally, well with the empirical data $E_{0+}^{+n} = 27.9 \quad 0.5 [4.68], E_{0+}^{+n} = 28.8 \quad 0.7 [4.69], E_{0+}^{-p} = 31.4 \quad 1.3 [4.68] \text{ and } E_{0+}^{-p} = 32.2 \quad 1.2 [4.70]$ (all in canonical units). The numbers in (4.75) should not be considered as rigorous predictions of CHPT since they depend to some extent on the assumptions made on the unknown counter term s. One should perform a similar calculation in HBCHPT to order q^4 . A more accurate determ ination of these threshold multipoles would give a further constraint on the pion (nucleon coupling constant via the Goldberger (M iyazawa (O ehm e sum nule [4.10,4.71])

$$J + \frac{g_{N}^{2}}{2m^{2} M^{2}=2} = \frac{m + M}{m M} a$$
(4:76)

with a the isospin{odd N scattering length and J a dispersion integral over the hadronic p total cross sections. The integral J can be calculated either from the pertinent K arlsruhe{H elsinki cross sections or the ones from the SA ID data basis. One possibility of obtaining the di erence $a_{1=2} = a_{3=2}$, which is most uncertain at present, is via the Panofsky ratio P = (p! n⁰) = (p! n) [4.10],

$$(a_{1=2} a_{3=2})^2 = (9k=q)P \pm 0^{+} f = (9k=q)P R \pm 0^{+} f^2$$
 (4:77)

with $R = (n! p) = (p! n^+)$. To make use of eqs.(4.76,4.77), one needs a very accurate understanding and determ ination of the electric dipole am plitude at threshold for charged pion photoproduction (for further details, see e.g. refs.[4.10,4.72,4.73]). This concludes our discussion of threshold photoproduction.

W e now turn to a short discussion of som e topics related to pion electroproduction. A much more detailed account of these topics can be found in the recent review [4.46]. There, one can ind a thorough discussion of the pertinent low (energy theorems in the various channels. In particular, it is stressed (see also ref.[4.74]) that in a system atic chiral expansion one is only sensitive to the institive few moments of the pertinent nucleon form factors, in contrast to the commonly used practise of supplementing the photon { nucleon and pion {nucleon vertices with the corresponding full form factors. A lso, in the loop expansion there is no need for equating the pion charge form factor and the isovector nucleon charge form factor to maintain gauge invariance as it is offen done. The chiral expansion keeps gauge invariance at any step of the calculation and thus allows naturally for F (t) \in F^V₁ (t). Here, let us brie y discuss the axial m s radius of the nucleon as measured in charged pion electroproduction. The starting point is the venerable LET due to N am bu, Lurie and Shrauner [4.75] for the isospin{odd electric dipole am plitude E $_{0+}^{(i)}$ in the chiral lim it,

$$E_{0+}^{()}(M) = 0; k^{2} = \frac{eg_{A}}{8 F} + \frac{k^{2}}{6}r_{A}^{2} + \frac{k^{2}}{4m^{2}}(v + \frac{1}{2}) + 0 (k^{3})$$
(4:78)

Therefore, m easuring the reactions ${}^{2}p!$ ${}^{+}n$ and ${}^{2}n!$ p allows to extract E $_{0+}^{()}$ and one can determ ine the axial radius of the nucleon, r_{A} . In section 4.3, we had pointed out that the determ inations of the axial radius from electroproduction data and from (anti)neutrino{nucleon scattering show a small discrepancy. This discrepancy is usually not taken seriously since the values overlap within the error bars. How ever, it was show n in ref.[4.76] that pion bops modify the LET (4.78) at order k² for nite pion mass. In the heavy m ass form alism, the coe cient of the k² term reads

$$\frac{1}{6}r_{\rm A}^2 + \frac{1}{4m^2}\left({}_{\rm V} + \frac{1}{2} \right) + \frac{1}{128F^2}\left(1 - \frac{12}{2} \right)$$
(4:79)

where the last term in (4.79) is the new one. This means that previously one had extracted a modi ed radius, the correction being $3(1 \quad 12=2)=64F^2$ ' 0:046 fm². This closes the gap between the values of r_A extracted from electroproduction and neutrino data. As detailed in appendix C, the 1=m suppressed terms (i.e. of order q⁴) modifying the result (4.79) are small [4.91].

A nother interesting quantity is the S{wave cross section de ned in eq.(4.66). The most precise measurement of it for neutral pion production of the proton close to the photon point was presented in ref.[4.56]. In g.4.6 we show the data of ref.[4.56] at $k^2 = 0.042$; 0.0501 and 0.0995 G eV in comparison to the one{loop CHPT result and the corresponding tree level prediction [4.46,4.77]. The most interesting feature of the data is the atness of a_0 as jk^2j increases.

This trend is also exhibited by the one {loop CHPT result but not by the tree graphs (or by tree graphs supplem ented with form factors). Chiral loops are required to explain the trend of the data. We should stress that the calculation of a_0 to one loop accuracy did not involve any new adjustable counter term s (all low {energy constants were previously



Fig. 4.6: The S {wave component of the neutral pion electroproduction cross section, calculated from one{loop CHPT (solid line) and tree graphs (dotted line). The kinematics is W = 1074 MeV and = 0.58 [4.77]. The data extracted in ref.[4.56] are also shown.

tted in photoproduction [4.59] or via nucleon radii). C learly, to have a better test of the chiral dynam ics, one should measure at smaller values of k^2 j since there the loop corrections are sizeable but not as large as at $k^2 = 0.1 \text{ GeV}^2$. In ref.[4.77], it was further stressed that to test the chiral predictions, one should investigate in m ore detail the angular distributions. The m ost striking feature is that the CHPT predictions for the transverse di erential cross sections become a forward peaked as k^2 j becomes larger than 0.04 GeV². Indeed, a group at MAM I (M ainz) has measured this di erential cross section at $k^2 = 0.1 \text{ GeV}^2$ and the shape agrees very nicely with the CHPT prediction [4.78]. C learly, the wide eld of single pion electroproduction in the threshold region is a good testing ground of the chiral dynam ics and just begins to play again an important role. For a more detailed account of the existing predictions and limitations within CHPT, we refer the reader to ref.[4.46].

IV .5. TW O {PION PRODUCTION

In the previous section, we considered single pion photo { and electroproduction. C om plem entary inform ation can be gained from the two pion production process N !

N, with a real or virtual photon. The two pions in the nal state can both be charged, both neutral or one charged and one neutral. Here, we will be concerned with the threshold region, i.e. the photon in the initial state has just enough energy to produce the two pions (and the outgoing nucleon) at rest. This energy is very close to the rst strong resonance excitation of the nucleon, the (1232). In fact, presently available data focus on the resonance region and above. In that case, a two{step reaction m echanism of the form N ! ! N is appropriate to describe these data as detailed in refs.[4.1,4.79,4.80]. A swe will argue, there is how ever a narrow window above threshold which is particularly sensitive to chiral bops, i.e. to the strictures of the spontaneously

broken chiral sym m etry. First m easurem ents of two{pion production at low energies have been perform ed at MAM I and we expect that the theoretical predictions discussed below will give additionalm otivation to perform yet m ore detailed m easurem ents of this particular reaction. The CHPT calculation presented in ref.[4.81] extends the one of D ahm and D rechsel [4.82] who discussed certain aspects of two{pion photoproduction in the fram ework of W einberg's chiral pion {nucleon Lagrangian [4.83].

First, we must outline the formalism necessary to treat two-pion photo- and electroproduction in the threshold region. We will only be concerned with the kinematics close to or at threshold, the corresponding am plitudes and the total cross sections. For a more general discussion we refer the reader to ref.[4.82]. To be specific, consider the process $(k) + N(p_1)! = a(q_1) + b(q_2) + N(p_2)$. The corresponding current transition matrix element is

$$T = <^{a}(q_{1}); \ ^{b}(q_{2}); N (p_{2}) \text{ out j}^{em} (0) \ N (p_{1}) \text{ in } > (4:80)$$

with J^{em} the electrom agnetic current operator and the polarization vector of the photon. From the two initial states p and n we can form in total six nal states

$$p! + p; p! + 0n;$$

$$p! + 0p; n! + n; \qquad (4:81)$$

$$n! + 0p; n! + 0n; \qquad (4:81)$$

In what follows we will concentrate on the channels with a proton in the initial state. To rst order in the electrom agnetic coupling e the threshold am plitudes for p ! + 0 n and n ! - 0 p are equal. In general, one can form ve/six M andelstam variables for the two-pion photo/electroproduction process from the independent fourm on enta. For our purpose, it is most convenient to work in the photon-nucleon center-ofm ass fram e. At threshold, the real or virtual photon has just enough energy to produce the two pions at rest. The threshold center-ofm ass energy squared is $s_{thr} = (p_1 + k)_{thr}^2 = (m + 2M)^2 = m^2 (1 + 4 + 4^2)$. The photon center-ofm ass energy can be expressed in term s of s and the photon virtuality k^2 as $k_0 = (s - m^2 + k^2) = (2 + s) w$ ith its threshold value

$$k_0^{\text{thr}} = \frac{2m}{1+2} + \frac{2}{4} + \frac{4}{4}$$
 (4:82)

in term s of the small parameters and . In the lab system, the threshold value for two pion-photoproduction is given by E $^{\rm thr} = 2M$ (1 +). The kinematics above threshold is discussed in m ore detail in ref.[4.81].

At threshold in the center-ofm ass frame (i.e. $q_1 = q_2 = 0$), the two-pion electroproduction current m atrix element can be decomposed into amplitudes as follows if we work to rst order in the electrom agnetic coupling e,

$$T = {}_{f}^{y} i \sim (\sim \tilde{\kappa}) M_{1}^{ab} + M_{2}^{ab} + M_{3}(a^{3}b + b^{3}a) + \sim \tilde{\kappa} N_{1}^{ab} + N_{2}^{ab} + N_{3}(a^{3}b + b^{3}a) i$$

$$(4:83)$$

in the gauge $_0 = 0$. C learly, for real photons only the M $_{1;2;3}$ can contribute. For virtual photons, gauge invariance T k = 0 allows to reconstruct T as $T_0 = T$ $k=k_0$. The amplitudes M $_{1;2;3}$ and N $_{1;2;3}$ encode the inform ation about the structure of the nucleon as probed in threshold two pion photo- and electroproduction. The physical channels listed in eq.(4.81) give rise to the following linear combination of M $_{1;2;3}$ (and N $_{1;2;3}$ for $k^2 < 0$).

$$p! + p: M_{1} + M_{2};$$

$$p! + 0n: \frac{p}{2}M_{3};$$

$$p! = 0 + 0p: M_{1} + M_{2} + 2M_{3};$$

$$n! + n: M_{1} + M_{2};$$

$$n! = 0 + p: \frac{p}{2}M_{3};$$

$$n! = 0 + p: \frac{p}{2}M_{3};$$

$$n! = 0 + 0n: M_{1} + M_{2} + 2M_{3}$$
(4:84)

C lose to threshold, the invariant m atrix elem ent squared averaged over nucleon spins and photon polarizations takes the form $M_{fij}^2 = \kappa^2 j_1 M_1 + _2 M_2 + _3 M_3 j^2$ with the isospin factors $_{1;2;3}$ given in eq.(4.84). The m ain dynam ical assumption in this relation is that the two-pion photoproduction amplitude in the threshold region can be approximated by the amplitude at threshold. Expressing κ^2 in terms of s and supplementing M_{fij}^2 by the photon ux factor $m^2 = p_1$ $k = 2m^2 = (s m^2)$, we not for the unpolarized total cross section

$$\int_{tot}^{N !} (s) = \frac{m^2}{2s} (s m^2) _3 (s) j_1 M_1 + _2 M_2 + _3 M_3 j_2 S :$$
(4:85)

Here, $_3$ (s) is the integrated three-body phase space, eq.(3.91), and S a Bose symmetry factor, S = 1=2 for the 0 nal state and S = 1 otherwise. For equal pion m asses an excellent approximation to the integrated three (body phase space is given by [4.81]

₃ (s)
$$\frac{M m^{5=2}}{64^{-2} (m + 2M)^{7=2}} [E - 2M (1 +)]^2$$
: (4:86)

O f course, an analogous approximation can be derived for unequalpion masses. Consequently, the unpolarized total cross section can be approximated within a few percent by the handy formula

$$\int_{tot}^{N} N (E) = \frac{M^2 (1 + 1)}{32^2 (1 + 2)^{11 = 2}} j_1 M_1 + {}_2 M_2 + {}_3 M_3 j^2 S (E E^{thr})^2 : (4.87)$$

For electroproduction, the prefactor in eq.(4.87) has to be modiled slightly to account for the virtual photon ux normalization and then it gives the transverse total electroproduction cross section. In general above threshold the total cross section is given by a four{dimensional integral over e.g. the two pion energies and two angle variables (for details, see ref.[4.81]). One remark on isospin breaking is in order. To one{loop accuracy O (q³), it is legitim ate to work with one nucleon and one pion mass. However, the pion mass di erence M M $_{0}$ = 4:6 M eV in reality leaves a 11.9 M eV gap between the production threshold of two neutral versus two charged pions. W hile we are not in position of performing a calculation including all possible isospin-breaking e ects, a m inim alprocedure to account for the mass di erence of the physical particles is to put in these by hand in the pertinent kinematics, such that the thresholds open indeed at the correct energy value. To be speci c, for the ⁺ p nal state the threshold photon energy is E ^{thr} (⁺ p) = 320:66 M eV whereas for ⁰ ⁰ p it is E ^{thr} (⁰ ⁰ p) = 308:77 M eV. Therefore, in the pertinent three(body phase space integrals we will di erentiate between neutral and charged pion mass when we present results incorporating the correct opening of the thresholds.

Consider now the chiral expansion of the threshold am plitudes M $_{1;2;3}$ and N $_{1;2;3}$. In each case we will give two complete chiral powers, the leading and next-to-leading term. It is worth to elaborate a bit on the chiral counting here. The S{m atrix elements are calculated up {to {and {including order q3. This means that the threshold am plitudes are given to order g since two chiral powers are factored out, ĩ q. Due to the various selection rules which apply for heavy baryon CHPT and additional ones due to the threshold kinem atics, only a few diagram s are contributing. These are discussed in detail in [4.81]. Here, we just mention that the leading nonzero contributions com es from tree graphs with one insertion from $L_{N}^{(2)}$. At next order, one has a plethora of possible contributions. Four loop graphs (plus their crossed partners) remain and the only contact terms which survive are the ones with one insertion from $L_{M}^{(3)}$.* The corresponding low {energy constants are estim ated via resonance saturation, i.e. the contribution. One expects sizeable e ects from the (1232) since rst it is quite close to threshold and second its couplings to the N system are very large (about twice the nucleon couplings). On rst sight the distance of only 14:6 M eV of the (1232) from threshold seem s to give rise to overw helm ing contributions since one naively expects that the very sm all denom inator 1= (m²) $s_{hr} = 1 = m$ m 2M) 1 = (m + m + 2M)enters the result. However, as shown in ref.[4.81], this dangerous denom inator always gets cancelled by exactly the same term in the num erator in the corresponding diagram s. Therefore, the expansion in M is not a priori useless. For the transverse threshold am plitude, the resulting chiral expansion takes the form

$$M_{1} = \frac{eg_{A}^{2}M}{4m^{2}F^{2}} + O(q^{2})$$
(4:88a)

^{*} W ithin our approximation, the contribution from $L^{(4)}$ containing the W ess-Zum ino term incorporating the anomalous (natural parity violating) vertex ! 3 vanishes.

$$M_{2} = \frac{e}{4m F^{2}} (2g_{A}^{2} - 1 - v) + \frac{eM}{4m^{2}F^{2}} (g_{A}^{2} - v) + \frac{eg_{A}^{2}M}{8m m^{2}F^{2}}B + \frac{eg_{A}^{2}M}{64 F^{4}} \frac{8 + 4r}{1 + r} \arctan^{p} \frac{1 + r}{1 + r} \frac{r}{1 + r} \frac{1 + r + r^{2}}{(1 + r)^{3 = 2}} \frac{1 + r + r^{2}}{2} + \arctan \frac{r}{1 + r} + i \frac{p}{3} \frac{(2 + r)}{1 + r} \frac{1 + r + r^{2}}{(1 + r)^{3 = 2}} \ln \frac{2 + r + p}{1 + r + r^{2}} + 0 (q^{2});$$
(4.88b)

$$M_{3} = \frac{e}{8m F^{2}} (1 + v) + \frac{eM}{8m^{2}F^{2}} + \frac{eg_{A}^{2}M}{16m m^{2}F^{2}} B$$

$$+ \frac{eg_{A}^{2}M}{256 F^{4}} - 6 + \frac{4 + 2r}{1 + r} \arctan^{p} \frac{1 + r}{1 + r} + \frac{r}{1 + r} + \frac{1 + r + r^{2}}{(1 + r)^{3 - 2}} + \arctan^{p} \frac{r}{1 + r}$$

$$+ i \frac{p}{3}(2 + r) + \frac{1 + r + r^{2}}{(1 + r)^{3 - 2}} \ln \frac{2 + r + p}{1 + r + r^{2}} + O(q^{2})$$

$$(4.88c)$$

with the ratio $r = k^2 = 4M^2$ and

$$B = \frac{2m^2 + m m m^2}{m m} + 4Z [m (1 + 2Z) + m (1 + Z)]$$
(4:88d)

which involves the o -shell parameter Z of the N vertex. In fact, taking the allowed range of Z given in ref.[4.43], one nds a weak Z dependence, i.e. 9.9 G eV < B < 15:1 GeV.Furthermore, from the isospin factors of eq.(4.88) we see that to order M the contributions are absent in the 0 channels. We also note that to low est order, M₁ = 0 and M $_2$ = $2M_3$ so that the production of two neutral pions is strictly suppressed. A nother point worth m entioning is that the transverse am plitudes M $_{2;3}$ are k^2 -dependent only through their loop contribution. This can be understood from the fact that the tree graphs have to be polynomial in both M and k^2 and that a term linear in k^2 is already of higher order in the chiral expansion. It is also not possible to further expand the r-dependent functions since $r = k^2 = 4M^2$ counts as order one and all term s have to be kept. We also notice that the amplitudes M_{2:3} have a sm ooth behaviour in the chiral lim it. Finally, we note that the loop contribution to the transverse amplitudes of two pion production as given in eq.(4.88) have a nonzero im aginary part even at threshold. This comes from the rescattering type graphs. Due to unitarity the pertinent loop functions have a right hand cut starting at $s = (m + M)^2$ (the single pion production threshold) and these functions are here to be evaluated at $s = (m + 2M)^2$ (the twopion production threshold). In the electroproduction case, we also have the longitudinal threshold am plitudes N_{1:2:3}. Since we can no more exploit the condition k = 0 the photon coupling to an out-going pion line is non-vanishing and therefore we obtain a nonzero contribution already at leading order 0 (g) involving a pion propagator (for charged pions). Adding up all term s which arise at order q and q^2 we nd the following

results

$$N_{1} = O(q);$$

$$N_{2} = \frac{eM(1+)}{F^{2}(4M^{2}-k^{2})} + \frac{e(2g_{A}^{2}-1)}{4mF^{2}} + O(q);$$

$$N_{3} = \frac{1}{2}N_{2} + O(q):$$
(4:89)

It is interesting to note that none of the low energy constants $c_1; c_2; c_3; c_4$ and the anom abus magnetic moments which enter $L_N^{(2)}$ show up in the nalresult.

We now turn to the numerical results for the threshold am plitudes and total cross sections. The isospin symmetric case is discussed in great detail in ref.[4.81]. To connect to the experimental situation, consider the three{body phase space with the physical masses for the corresponding pions. This automatically takes care of the various threshold energies. In the loops we work, how ever, with one pion mass. This elect is small as discussed in ref.[4.81]. In g.4.7 we show the calculations with the correct phase{space and using the threshold matrix{elements.



Fig. 4.7: Total cross sections (in nb) for the p initial state ($k^2 = 0$) with the correct three{body phase space. The dotted, dashed and dashed{dotted lines refer to the ⁺ p, ^{+ 0}n, and the ^{0 0}p nal state, in order. For p ! ⁺ p we show the rst correction as discussed in ref.[4.81]. The various thresholds are indicated.

For p! + p, we show the rst correction above threshold from $L_{N}^{(1)}$ which is bigger than the matrix {element calculated with the threshold amplitudes already 3 MeV above threshold in this particular channel (for details, see [4.81]). At $E = 320 \,\mathrm{MeV}$, the total cross section for 0 production is 0.5 nb w hereas the competing 0 + n nal state has tot = 0.013 nb. D ouble neutral pion production reaches tot = 1.0 nb at E = 324.3 MeV in comparison to $tot(p!^{0+n}) < 0.05$ nb and $tot(p!^{+})$ p) = 0:4 nb.This means that for the rst 10...12 MeV above 0 threshold, one has a fairly clean signal and much more neutrals than expected. We stress that the enhancem ent of the total cross section in the ⁰ ⁰ channel is a chiral loop e ect. Consider the corresponding threshold matrix element M₁ + M₂ + 2M₃. W hereas the Born graphs contribute only 4.9 GeV 3 , the loop contribution at the same chiral power is much larger (26.6 + 16.0 i) GeV 3 and enhances the total cross section by a factor 50. A lready the know legde of the order of magnitude of the experimental 0 cross sections allows to test the enhancement e ect of the chiral loops. Remember that to leading order in the chiral expansion, the production of two neutral pions is com pletely suppressed. Of course, the above threshold correction for this channel, which comes from $L_{N}^{(2)}$ (and higher orders) should be calculated system atically. The rst correction, which vanishes proportional to $\dot{\mathbf{y}}_i$ j (i = 1;2) at threshold, has been calculated and found to be very small. The corresponding cross section at E = 320, 325 and 330 MeV is $\frac{\text{rst corr}}{\text{tot}} = 0.009; 0.026$ and 0:056 nb, i.e a few percent of the leading order result. It is, therefore, conceivable that the qualitative features described above will not change if even higher order corrections are taken into account. A m ore detailed account of these topics can be found in ref.[4.81].

IV .6. W EAK PION PRODUCTION

A s discussed in the preceeding sections, single and double pion production o nucleons by real or virtual photons gives in portant inform ation about the structure of the nucleon. A s stressed in particular by A dler [4.84], weak pion production involves the isovector axial am plitudes and a uni ed treatm ent of pion photo-, electro- and weak production allows to relate inform ation from electron {nucleon and neutrino{nucleon scattering experiments. In this spirit, we will consider here pion production through the isovector axial current in the threshold region, extending the classical work of A dler [4.84], A dler and D othan [4.52] and N am bu, Lurie and Shrauner [4.75], who have considered soft pion em ission induced by weak interactions making use of PCAC and gauge invariance, relating certain electrow eak form factors of the nucleon to particular threshold multipole am plitudes. The corrections beyond this were considered in ref.[4.85], were novel relations between various axial threshold multipole am plitudes and physical observables like electrow eak form factors, S { wave pion {nucleon scattering lengths and, in particular, the nucleon scalar form factor, (t) < N jf (uu + dd) jN > are given.

We have been informed by Th. W alcher that a rst analysis of double neutral pion production making use of the TAPS detector seem to indicate an even stronger increase of the p⁰ cross section at threshold as indicated by the calculation of ref.[4.81].

We consider processes of the type $(k_1) + N(p_1) ! l(k_2) + N(p_2) + {}^{a}(q)$, which involve the isovector vector and axial{vector currents. Here, we will focus on the pion production induced by the axial current, $A^{b} = q_{5}({}^{b}=2)q$ in terms of the u and d quark elds. Denoting by $k = k_1$ k the four{momentum of the axial current, the pertinent M and elstam variables are $s = (p_1 + k)^2$, $t = (q k)^2$ and $u = (p_1 q)^2$ subject to the constraint $s + t + u = 2m^2 + M^2 + k^2$. The pertinent m atrix element decom poses into an isospin{even and an isospin{odd part (analogous to the N scattering am plitude),

< N (p₂); ^a (q) out
$$A^{b}$$
 N_{1} (p in > = i ^{ab} T ⁽⁺⁾ abc ^c T ⁽⁾ (4:90)

 $u_1 = {}_5u$. Notice that one can use is the axial polarization vector, where the Dirac equation to transform terms of the type k into lepton mass terms via k $_1$ for $_5 u = m_1 u_1 _5 u$. This means that in the approximation of zero lepton m ass, one has k = 0 and all diagram s where the axial source couples directly to a pion line vanish. The general D irac structure for the transition current involves the eight operators $O_1 = (6 \, \mathbf{\hat{q}} \, \mathbf{\hat{q}} \, 6) = 2; \, Q =$ $q_{i} O = 6; O_4 =$ $_{1}(p+p_{2})=2; O_{5} = K_{5}$ **q;**₽= $_{1}(p+p_{2})=2; 0_{7} = k; 0 = k k which are accompanied by invariant functions$ 6 denoted $A_i^{()}$ (s;u) (i = 1;:::;8) [4.84]. At threshold in the N center of mass frame, one can express the pertinent matrix element in terms of six S{wave multipoles, called $L_{0+}^{()}$, $M_{0+}^{()}$ and $H_{0+}^{()}$,

$$T^{()} = {}_{2}u = {}_{0}u = {}_{1}A_{i}^{()} (s_{th}; u_{th})u_{1} = 4 (1+) {}_{2}v = {}_{0}L_{0+}^{()} + k_{0+}^{(1)} + i \sim \hat{k}(\sim)M_{0+}^{()} {}_{1}:$$
(4:91)

At threshold, one can express L_{0+} , M_{0+} and H_{0+} (suppressing isospin indices) through the invariant amplitudes A_i (s;u) via

$$M_{0+} = \frac{p}{\frac{2}{(2+)^{2}}} M_{1} A_{3}$$

$$L_{0+} = \frac{p}{\frac{(1+)^{3=2}}{(2+)^{2}}} \frac{A_{1}}{(2+)^{2}} M_{1} + m_{1} A_{2}$$

$$+ \frac{2(1+)(2+)}{(2+)^{2}} A_{3} + m_{1} + \frac{2}{2} A_{4} + \frac{2}{m}^{2} A_{5} + m^{2} + \frac{1+2}{2} A_{6}$$

$$H_{0+} = \frac{p}{\frac{(2+)^{2}}{(2+)^{2}}} \frac{2(1+)}{(2+)^{2}} A_{1} \frac{1}{m} A_{3} \frac{1}{2} A_{4} \frac{m}{2} A_{6} + A_{7} + mA_{8}$$

$$(4:92)$$

where the $A_i(s;u)$ are evaluated at threshold.

We seek an expansion of these threshold multipoles in powers of and up to and including order O (2 ;) (m odulo logarithm s). To work out the corrections at order O (q^3), it it mandatory to perform a complete one{loop calculation with insertions from

 $L_{N}^{(1)}$ and the tree diagrams with exactly one insertion from $L_{N}^{(3)}$. One also has to consider tree graphs with two insertions from $L_{N}^{(2)}$ with a nucleon propagator, which scales as 1=q, in between. The resulting low (energy theorems for the various S{wave multipoles are

$$M_{0+}^{(+)} = \frac{p \frac{1}{M^2 - k^2}}{\frac{16 \text{ m F}}{M^2 - k^2}} g_A^2 + C_M^{(+)}M + O(q^3)$$

$$M_{0+}^{(-)} = \frac{p \frac{1}{M^2 - k^2}}{16 \text{ m F}} G_M^V (k^2 - M^2) - q_A^2 + C_M^{(-)}M + O(q^3)$$
(4:93a)

$$L_{0+}^{(+)} = \frac{1}{3 \text{ M F}} \quad (k^2 \quad M^2) \quad \frac{1}{4} \quad (0) \quad \frac{a^+ F}{M} \quad \frac{g_A^2 M}{16 \text{ m F}} + C_L^{(+)} M^2 + O \quad (q^3)$$

$$L_{0+}^{(-)} = \frac{1}{8 \text{ F}} \quad G_E^V \quad (k^2 \quad M^2) + \frac{M}{2m} \quad (g_A^2 + 1) \quad \frac{k^2}{8m^2} \quad + C_L^{(-)} M^2 + O \quad (q^3)$$

$$(4.93b)$$

$$H_{0+}^{(+)} = \frac{a^{+}F}{k^{2} M^{2}} + \frac{(0) (k^{2} M^{2})}{12 F (k^{2} M^{2})} + C_{H}^{(+)}M + O(q^{2})$$

$$H_{0+}^{()} = \frac{a F}{k^{2} M^{2}} + \frac{M [G_{E}^{V} (k^{2} M^{2}) 1]}{8 F (k^{2} M^{2})} + \frac{1}{16 m F} + C_{H}^{()}M + O(q^{2})$$
(4:93c)

with a the isopin (even and odd S { wave N scattering lengths. The form of the pion pole term in H $_{0+}^{(\)}$ can easily be understood from the fact that as k ! q one picks up as a residue the forward N scattering am plitude which at threshold is expressed in term s of the two S { wave scattering lengths. The relation between axial pion production and the N scattering am plitude has also been elucidated by A dler in his sem in al work [4.84]. The constants C $_{\rm H\ ;L\ ,M}^{(\)}$ subsum e num erous k^2 { independent kinem atical, loop and counterterm corrections (the latter ones stem m ainly from L $_{\rm N}^{(3)}$) which we do not need for the following discussion and which are di cult to pin down exactly. There is, how ever, one exception to this. The chiral W and identity ($A^{\rm b}$ = m qi $^{\rm b}$ $_5q$ M 2 dem ands that k_0L_{0+} + k^2 H $_{0+}$ M 2 and thus with k_0 = m (2 + 2 +)=2(1 +) we have

$$C_{H}^{(+)} = \frac{a^{+}F}{2m M^{2}} \frac{(0)}{8 M^{2}m F} + \frac{g_{A}^{2}}{32 m^{2}F} = \frac{c_{2} + c_{3}}{4 m F}$$

$$C_{H}^{()} = \frac{2g_{A}^{2} + 5}{64 m^{2}F} :$$
(4:94)

The num erical values of the constants are $C_{H}^{(+)} = 1:0 \text{ GeV}^{3}$ and $C_{H}^{(-)} = 0:5 \text{ GeV}^{3}$. The argument of the various nucleon form factors in (4.94) is the threshold value of the invariant momentum transfer squared $t_{thr} = (q + k_{thr}^{2}) = (k^{2} - M^{2}) = (1 + 1) = 1$ $k^2 = M^2 + 0$ (q³). Of particular interest is the low {energy theorem for $L_{0+}^{(+)}$ where one has the following slope at the photon point $k^2 = 0$

$$\frac{\mathcal{Q}L_{0+}^{(+)}}{\mathcal{Q}k^2}_{k^2=0} = \frac{\mathcal{Q}(M^2)}{3MF} + \mathcal{Q}(M) = \frac{g_A^2}{128F^3} \frac{6}{5} \quad \arctan\frac{1}{2} + \mathcal{Q}(M) : \quad (4:95)$$

It is very interesting to note that although $L_{0+}^{(+)}$ vanishes identically in the chiral lim it M = 0 the slope at $k^2 = 0$ stays nite. The form al reason for this behaviour is the non {analytic dependence of $L_{0+}^{(+)}$ on M which does not allow to interchange the order of taking the derivative with respect to k^2 at $k^2 = 0$ and the chiral limit. Notice also that for k^2 ' 0 and assuming that C $_{\rm L}^{~(+~)}$ of the order of 1 G eV $^{-3}$, the term proportional to the scalar form factor (M^2) (0)=4 dom in tes the behaviour of $I_{h_+}^{(+)}$ using the num bers from the recent analysis of Gasser, Leutwyler and Sainio [4.86]. In principle, an accurate m easurem ent of this particular multipole in weak pion production allows for a new determ ination of the elusive nucleon scalar form factor and the N {term. This m ight open the possibility of another determ ination of this fundam ental quantity. In the standard m odel, the axial part of the weak neutral current is the third component of the isovector axial current. To see this most interesting correction, one should therefore consider neutral neutrino reactions like $p! p^0$ (in that case the zero lepton mass approximation is justied). First, however, a complete calculation involving also the isovector vector current has to be performed to nd out how cleanly one can separate this multipole in the analysis of neutrino {induced single pion production. For that, it will be mandatory to include the resonance since the presently available data are concentrated around this mass region [4.87]. In parity (violating electron scattering, the interference of this axial current with the vector one is suppressed by the factor $4\sin^2$ w), with \sin^2 v 0:23 the W einberg (weak mixing) angle. (1

In contrast to this, the behaviour of H $_{0+}^{(+)}$, which also contains the scalar form factor, is dominated by the pion pole term proportional to a^+ . At $k^2 = 0$, one nds H $_{0+}^{(+)} = 32.8 \,\text{GeV}^2 + (0) (M^2)$ 14:1 G eV 3 0:14 G eV 2 . The uncertainty in a^+ , $a^+ = 0:38$ 10=M, gives as large a contribution as the term proportional to the scalar form factor.

F inally, we point out that A dler's relation between weak single pion production and the elastic neutrino (nucleon cross section at low energies [4.88] is also modi ed by the novel term proportional to the scalar form factor of the nucleon.

REFERENCES

- 4.1 E.Amaldi, S.Fubiniand G.Furlan, Pion Electroproduction, Springer Verlag, Berlin, 1979.
- 4.2 J.Gasser, M.E.Sainio and A.Svarc, Nucl. Phys. B 307 (1988) 779.

- 4.3 V.Bernard, N.Kaiser, J.Kambor and UlfG.Meiner, Nucl. Phys. B 388 (1992) 315.
- 4.4 M A B.Beg and A.Zepeda, Phys. Rev. D 6 (1972) 2912.
- 4.5 D.G. Caldiand H. Pagels, Phys. Rev. D 10 (1974) 3739.
- 4.6 Ulf-G.Meiner, Int. J.Mod. Phys. E1 (1992) 561.
- 4.7 Ulf-G.Meiner, Review talk at WHEPP (III, Madras, India, 1994, preprint CRN (94/04.
- 4.8 W R. Frazer and J. Fulco, Phys. Rev. 117 (1960) 1603, 1609.
- 4.9 G. Hohler and E. Pietarinen, Phys. Lett. B 53 (1975) 471.
- 4.10 G.Hohler, in Landolt {Bornstein, vol.9 b2, ed. H.Schopper (Springer, Berlin, 1983).
- 4.11 R E. Prange, Phys. Rev. 110 (1958) 240.
- 4.12 A.C. Hearn and E. Leader, Phys. Rev. 126 (1962) 789.
- 4.13 F.Low, Phys. Rev. 96 (1954) 1428; M.Gell-M ann and M.L.Goldberger, Phys. Rev. 96 (1954) 1433.
- 4.14 S.D rell and A C. Heam, Phys. Rev. Lett. 16 (1966) 908; S.B. Gerasim ov, Sov. J. Nucl. Phys. 2 (1966) 430.
- 4.15 V.Bernard, N.Kaiser and Ulf-G.Meiner, Phys. Rev. Lett. 67 (1991) 1515; Nucl. Phys. B 373 (1992) 364.
- 4.16 S.Ragusa, Phys. Rev. D 49 (1994) 3157.
- 4.17 JL. Powell, Phys. Rev. 75 (1949) 32.
- 4.18 F J. Federspielet al, Phys. Rev. Lett. 67 (1991) 1511.
- 4.19 E L. Hallin et al, Phys. Rev. C 48 (1993) 1497.
- 4.20 A.Zieger et al, Phys. Lett. B 278 (1992) 34.
- 4.21 M.Dam ashek and F.Gilm an, Phys. Rev. D 1 (1970) 1319;
- 4.22 V A. Petrunkin, Sov. J. Nucl. Phys. 12 (1981) 278.
- 4.23 J.Schm iedm ayer et al, Phys. Rev. Lett. 66 (1991) 1015.
- 4.24 K.W. Rose et al, Phys. Lett. B 234 (1990) 460.
- 4.25 V.Bernard, N.Kaiser, A.Schmidt and UlfG.Meiner, Phys.Lett. B 319 (1993) 269.
- 4.26 M.N.Butler and M.J.Savage, Phys. Lett. B 294 (1992) 369.
- 4.27 V.Bernard, N.Kaiser, A.Schmidt and UlfG.Meiner, Z.Phys. A 348 (1994) 317.
- 4.28 N.C.Mukhopadhyay, A.M. Nathan and L. Zhang, Phys. Rev. D 47 (1993) R7.
- 4.29 N M . Butler, M J. Savage and R. Springer, Nucl. Phys. B 399 (1993) 69.
- 4.30 V.Bernard, N.Kaiser, J.Kambor and Ulf-G.Meiner, Phys. Rev. D 46 (1992) 2756.
- 4.31 R.Erbe et al, Phys. Rev. 188 (1969) 2060.

- 4.32 M. Benmerrouche, R.M. Davidson and N.C. Mukhopadhyay, Phys. Rev. C 39 (1989) 2339.
- 4.33 A.L'vov, Phys. Lett. B 304 (1993) 29; B.R.Holstein and A.M.Nathan, Phys. Rev. D 49 (1994) 6101.
- 4.34 T A . A m strong et al., Phys. Rev. D 5 (1970) 1640; T A . A m strong et al., Nucl. Phys. B 41 (1972) 445.
- 4.35 R D. Peccei, Phys. Rev. 181 (1969) 1902.
- 4.36 A M .Sandor et al, Phys. Rev. D 50 (1994) R 6681.
- 4.37 I.Karliner, Phys. Rev. D 7 (1973) 2717.
- 4.38 R L.W orkm an and R A.A mdt, Phys. Rev. D 45 (1992) 1789.
- 4.39 V.Burkert and Z.Li, Phys. Rev. D 47 (1993) 46.
- 4.40 J.Ashm an et al, Nucl. Phys. B 328 (1989) 1.
- 4.41 V.Bernard, N.Kaiser and UlfG.Meiner, Phys. Rev. D 48 (1993) 3062.
- 4.42 Particle Data Group, Phys. Rev. D 45 (1993) S1.
- 4.43 T.K itagakiet al, Phys. Rev. D 28 (1983) 436;
 LA.Ahrens et al, Phys. Rev. D 35 (1987) 785;
 LA.Ahrens et al, Phys. Lett. B 202 (1988) 284.
- 4.44 A. delGuerra et al, Nucl. Phys. B 107 (1976) 65;
 M.G.Olsson, E.T. O sypow ski and E.H. Monsay, Phys. Rev. D 17 (1978) 2938.
- 4.45 Ulf-G.Meiner, Phys. Reports 161 (1989) 213.
- 4.46 V.Bernard, N.Kaiser, T.-S.H.Lee and UlfG.Meiner, Phys. Reports 246 (1994) 315.
- 4.47 G.Bardin et al, Phys.Lett. B104 (1981) 320.
- 4.48 J.Bemabeu, Nucl. Phys. A 374 (1982) 593c.
- 4.49 S.Choiet al, Phys. Rev. Lett. 71 (1993) 3927.
- 4.50 D.Taqqu, contribution presented at the InternationalW orkshop on "Large Experim ents at Low Energy Hadron M achines", PSI, Sw itzerland, April 1994; and private communication.
- 4.51 V.Bernard, N.Kaiser and UlfG.Meiner, Phys. Rev. D 50 (1994) 6899.
- 4.52 S.L.Adler and Y.Dothan, Phys. Rev. 151 (1966) 1267.
- 4.53 L.W olfenstein, in: High-Energy Physics and Nuclear Structure, ed. S.Devons (Plenum, New York, 1970) p.661.
- 4.54 E.M azzucato et al, Phys. Rev. Lett. 57 (1986) 3144.
- 4.55 R.Beck et al, Phys. Rev. Lett. 65 (1990) 1841.
- 4.56 T.P.W elch et al, Phys. Rev. Lett. 69 (1992) 2761.
- 4.57 S.Nozawa and T.-S.H.Lee, Nucl. Phys. A 513 (1990) 511, 544.

- 4.58 F A. Berends, A. Donnachie and D. L. Weaver, Nucl. Phys. B 4 (1967) 1.
- 4.59 V.Bernard, N.Kaiser and UlfG.Meiner, Nucl. Phys. B 383 (1992) 442.
- 4.60 IA .Vainshtein and V.I.Zakharov, Sov. J.Nucl. Phys. 12 (1971) 333; Nucl. Phys. B 36 (1972) 589;
 - P.de Baenst, Nucl. Phys. B 24 (1970) 633.
- 4.61 D.D rechsel and L.Tiator, J.Phys. G: Nucl. Part. Phys. 18 (1992) 449.
- 4.62 V.Bernard, J.Gasser, N.Kaiser and UlfG.Meiner, Phys. Lett. B 268 (1991) 291.
- 4.63 V. Bernard, N. Kaiser and Ulf-G. Mei ner, \Neutral Pion Photoproduction o Nucleons Revisited", preprint CRN 94-62 and TK 94 18, 1994, hep-ph/9411287.
- 4.64 V.Bernard, N.Kaiser and Ulf-G.Meiner, N Newsletter 7 (1992) 62.
- 4.65 J.C. Bergstrom, Phys. Rev. C 44 (1991) 1768.
- 4.66 A M . Bernstein, private communication.
- 4.67 N M . K roll and M A . Ruderm an, Phys. Rev. 93 (1954) 233.
- 4.68 J.P. Burg, Ann. Phys. (Paris) 10 (1965) 363.
- 4.69 M J.A dam ovitch et al, Sov. J. Nucl. Phys. 2 (1966) 95.
- 4.70 E L.Goldwasser et al, Proc. X II Int. Conf. on High Energy Physics, Dubna, 1964, ed. Ya.-A. Smorodinsky (Atom izdat, Moscow, 1966).
- 4.71 M L.Goldberger, H.M iyazawa and R.Oehme, Phys. Rev. 99 (1955) 986.
- 4.72 R L.W orkm an, R A . A mdt and M M . Pavan, Phys. Rev. Lett. 68 (1992) 1653.
- 4.73 G. Hohler, Karlsruhe University preprint TTP 92{21, 1992.
- 4.74 V.Bernard, N.Kaiser and UlfG.Meiner, Phys. Lett. B 282 (1992) 448.
- 4.75 Y.Nambu and D.Lurie, Phys. Rev. 125 (1962) 1429; Y.Nambu and E.Shrauner, Phys. Rev. 128 (1962) 862.
- 4.76 V.Bernard, N.Kaiser and Ulf-G.Meiner, Phys. Rev. Lett. 69 (1992) 1877.
- 4.77 V.Bernard, N.Kaiser, T.{S.H.Lee and UlfG.Meiner, Phys. Rev. Lett. 70 (1993) 387.
- 4.78 M.D istler and Th.W alcher, private communication.
- 4.79 JM . Laget, Phys. Rep. 69 (1981) 1.
- 4.80 P.W. .Carruthers and H.W. .Huang, Phys. Lett. B 24 (1967) 464;
 H.W. .Huang, Phys. Rev. 174 (1968) 1799;
 S.C. .Bhargava, Phys. Rev. 171 (1968) 969.
- 4.81 V.Bernard, N.Kaiser, UlfG.Meiner and A.Schmidt, Nucl. Phys. A 580 (1994) 475
- 4.82 R.Dahm and D.Drechsel, in Proc. Seventh Amsterdam Mini{Conference, eds. H.P.Blok, J.H.Koch and H.DeVries, Amsterdam, 1991.

- 4.83 S.W einberg, Phys. Rev. 166 (1968) 1568.
- 4.84 S.L.Adler, Ann. Phys. (N.Y.) 50 (1968) 189.
- 4.85 V.Bernard, N.Kaiser, and Ulf-G.Meiner, Phys. Lett. B 331 (1994) 137.
- 4.86 J.Gasser, H. Leutwyler and M.E. Sainio, Phys. Lett. B 253 (1991) 252,260.
- 4.87 S.J.Barish et al, Phys. Rev. D 19 (1979) 2521;
 M.Pohlet al, Lett. Nuovo C im ento 24 (1979) 540;
 N.J.Baker et al, Phys. Rev. D 23 (1981) 2495.
- 4.88 SLAdler, Phys. Rev. Lett. 33 (1974) 1511; Phys. Rev. D 12 (1975) 2644.
- 4.89 J.C. Bergstrom and E.L. Hallin, Phys. Rev. C 48 (1994) 1508.
- 4.90 G.Ecker and UlfG.Meiner, \W hat is a Low {Energy Theorem ?", Comments. Nucl.Part.Phys. (1995) in print.
- 4.91 A.Schmidt, Thesis TU Munchen, 1995 (unpublished).

V.THE NUCLEON {NUCLEON INTERACTION

One of the best studied objects in nuclear physics is the interaction between two nucleons. It is well known that to a high degree of accuracy one can consider nuclei as m ade of nucleons which behave non { relativistically and interact pair { w ise. Furtherm ore, three{ and m any {body forces are believed to be sm all. This has lead to the construction of sem i{phenom enological boson exchange potentials. These describe accurately deuteron properties and low {energy nucleon {nucleon phase shifts. The salient feature of these potentials $[5.1\{5.9\}]$ can be sum marized as follows. At large separation, there is one{pion exchange rst introduced by Yukawa [5.1]. The interm ediate{range attraction between two nucleons can be understood in term s of a ctitious scalar { isoscalar { m eson with a mass of approximatively 550 MeV.! {meson exchange gives rise to part of the short{range repulsion and the features prom inently in the isovector{tensor channel, where it cuts down most of the pion tensor potential. There are, of course, di erences in the various potentials but these will not be discussed here. As we will show in what follows, the elective chiral Lagrangian approach of QCD can be used to gain some insight into the question why these potentials work after all. One can also extend these considerations to many {nucleon forces as well as meson {exchange currents. The latter are the cleanest signal of non {nucleonic degrees of freedom in nuclei, in particular of pions [5.10,5.11]. First, however, we have to discuss som e technical subtleties related to the appearance of di erent energy scales in two (and m any) nucleon systems.

V.1. GENERAL CONSIDERATIONS

The consequences of the spontaneous chiral sym metry breakdown for the problem of the forces between nucleons were rst discussed by W einberg [5.12,5.13]. Since in his papers and the subsequent ones of the Texas/Seattle group [5.14,5.15,5.16,5.17,5.18]another language than the previously discussed one is used, we rst have to review the construction of the chiral Lagrangian and the power counting in this scheme. W e will then address the problem of sm all energy scales (sm all energy denom inators) related to the nuclear binding. Since SU (2) SU (2) is locally isom orphic to SO (4) and SU (2) SO (3), one can use stereographic coordinates to describe the G oldstone bosons living on the three sphere S³ SO (4)/SO (3). The covariant derivative of the pions is

$$D' = \frac{\theta \sim}{2D F}$$
; $D = 1 + \frac{\sim^2}{4F^2}$: (5:1)

Notice that we use F = 93 M eV in contrast to the conventions of refs.[5.12-5.18] which have F = 186 M eV. Nucleons are decribed by a D irac spinor N, which is also a Pauli spinor in isospace. The elds D irac formula constructed out of the elds D, N and their covariant derivatives,

$$D \vec{D} = (0 \vec{D} + i\vec{E} - \vec{D})$$

$$D N = (0 + \vec{\tau} \cdot \vec{E} -)N$$
(5.2)

where E = i(~ D) = F and t = ~=2 the isospin generators in the $\frac{1}{2}$ representation (for more details, see appendix C. There it is also shown how to include the resonance in this fram ework). The most general elective chiral Lagrangian follows by considering all possible isoscalar terms and imposing proper Lorentz, parity and time {reversal invariance and herm iticity. The explicit chiral symmetry breaking is due to the fourth component of a chiral four vector with coel cient (m $_u + m_d$)=2. The construction of these terms is also discussed in appendix C.

Consider now the S{m atrix for the scattering process of N incoming and N outgoing nucleons, all with m omenta smaller than some scale Q, say Q M'm. This means that the nucleons are non{relativistic and it thus is appropriate to use old{fashioned time{ordered perturbation theory. In that case, one deals with energy denominators for the intermediate states instead of the usual particle propagators. The idea is now to order all contributions in powers of Q=m. This is, however, not straightforward. In fact, as will become clear later, one is dealing with a three scale problem,

$$m \quad Q \quad \frac{Q^2}{2m} \quad : \tag{5:3}$$

The appearance of the nucleon m assm is obvious and the related scale can be rem oved by either de ning velocity {dependent elds (cf. section 3.3) or using the equation of m otion to elim inate the large time derivatives, $@_0N mN$, as described below [5.13]. The occurence of the third (sm all) scale $Q^2=2m$ is related to the presence of shallow nuclear bound states. In fact, consider a time {ordered diagram with only N nucleons in the interm ediate state. The energy denom inator associated to such a diagram is of order $Q^2=2m$, whereas all other diagram s contain at least one pion in the interm ediate state and have energy denom inators of order Q. The appearance of this sm all scale causes the perturbation theory to diverge and leads to the form ation of nuclear bound states. It is instructive to understand this in m ore detail from conventional Feynm an diagram techniques. For that, consider the box graph (called I) shown in g.5.1 for static nucleons. For a nucleon at rest, the propagator takes the form

$$S_{\rm N} (q) = \frac{1}{q^0 + 1}$$
 (5:4)

with the projection matrix onto positive energy, zero momentum Dirac wave functions.



Fig. 5.1: The box diagram for NN scattering discussed in the text. Solid (dashed) lines give nucleons (pions) and the pertinent momenta are exhibited.

0 ne nds for I

where the polynom ial P (q) includes term s that are non {vanishing as q^0 goes to zero. Consequently, the integral over q^0 in I has an infrared (IR) divergence. The contour of integration is pinched between the two poles at $q^0 =$ i and it is therefore in possible to distort it in such a way to avoid these singularities. This is distinctively di erent from the single { nucleon case discussed in the previous sections. Of course, this IR divergence is an artefact of the approximation (5.4), i.e. treating the nucleons as static. Indeed, if one includes the nucleon kinetic energy, $L_{kin} = N r^2 N = 2m$, the poles are shifted i] and the $\frac{1}{9}$ integral has the nite value 2i m = $\frac{1}{9}$ $\frac{1}{2}$. However, to q^0 ' [ef =2m from the counting of sm all m om enta one would expect this integral to scale as Q^{-1} (since each propagator scales as Q^{-1}), i.e. it is enhanced by a large factor m = Q. This enhancem ent is at the heart of the nuclear binding. Such small scales can only com e from reducible diagram s and to avoid these, one de nes an e ective potential as the sum of tim e{ordered perturbation theory graphs for the T {m atrix excluding those with pure nucleon intermediate states [5.12, 5.13]. The full machinery of expanding in powers of Q is therefore only applied to the reducible diagram s and the full S {m atrix is obtained by solving a Lippm ann {Schwinger or Schrodinger equation with the elective potential. This will be discussed in more detail when we consider the NN {potential. At this point we should stress that this separation of reducible versus irreducible diagram s is unavoidable but still poses som e concern to the purist since in the process of solving such bound state problem s, one can not completely exclude som e large m om entum components.

To remove the scale m from the problem, one can make use of a eld rede nition and replace the time derivative of the nucleon eld by the nucleon eld equations [5.13]

$$i\theta_0 = \frac{1}{2D F^2} t$$
 (~ $_0 \theta$) N = m + $\frac{g_A}{D F} t$ (~ \tilde{r})~ + ::: N (5:6)

So the chiral invariant time derivative of the nucleon eld in the interaction Lagrangian simply changes the coe cients of other term sallowed (and required) by chiral symmetry. Therefore, one can simply adopt a de nition of the elds and the constants in L_e such that no time derivatives appear. A lternatively, one could use the methods described in section 3.3.

In sum m ary, once the scales m and $Q^2=2m$ are removed by considering irreducible diagrams and using appropriate eld de nitions, one can order all remaining contributions to the N (nucleon forces in powers of Q=m Q=M. To do that, we have to extend the power counting scheme discussed in section 3.2 (since there it was assumed that only one nucleon line runs through a given diagram). Let us do that in time (ordered perturbation theory. Derivatives are counting as order Q, pion elds as $Q^{1=2}$ (using

the conventional norm alization $1 = \frac{p}{2!}$ for pion elds), interm ediate nucleons or 's* as Q¹ and loop integrals as d³k Q³. The chiral dimension of a graph with E_n external nucleon lines, D interm ediate states, L loops, C connected pieces and V_i vertices of type i (with d_i derivatives or pion m assess and n_i (p_i) nucleon (pion) elds) follows to be (we set C = 1 for the moment) [5.12,5.13,5.14]

= 3L D +
$$V_{i}(d_{i} = \frac{p_{i}}{2})$$
 (5:7)

and using the topological identities

$$D = \bigvee_{i}^{X} V_{i} \quad 1 \tag{5:7a}$$

$$L = I \qquad \begin{array}{c} X \\ V_i + 1 \end{array} \tag{5:7b}$$

$$2I + E_{n} = \bigvee_{i}^{X} V_{i} (p_{i} + n_{i})$$
(5:7c)

with I the total number of internal lines, one arrives at

$$= 2L + 2 \frac{1}{2}E_{n} + V_{i i}$$

$$I = d_{i} + \frac{1}{2}n_{i} 2 :$$
(5:8)

In case of C > 1, this generalizes to

= 2(L C) + 4
$$\frac{1}{2}E_n + V_{i i}$$
 (5:9)

It is now in portant to notice that chiral sym metry dem ands

This can easily be understood. Operators involving pions only have at least two derivatives or two powers of M and nucleon bilinears have at least one derivative. A s before, to lowest order one calculates tree diagram s with $_{i} = 0$. Loop diagram s are suppressed by powers of Q². We have now assembled all tools to take a closer look at the nucleon {nucleon potential and the problem of m any {nucleon forces.

^{*} We include here the since that has also been done in ref.[5.17] which reported rst full scale num erical results. We rem ind the reader here of the reservations made in section 3.4.



Fig. 5.2: Lowest order diagram s contributing to the NN interaction. (a) is a set of four{nucleon contact term s and (a) is the one{pion exchange.

V.2. THE NUCLEON {NUCLEON POTENTIAL

From eq.(5.8) it follows that for $E_n = 4$ (C = 1), the minimum value of is given by L = 0 (tree diagrams) with $_i = 0$. The latter condition can either be fullled having diagrams with $d_i = 1$ and $n_i = 2$ (one{pion exchange}) or having $d_i = 0$ and $n_i = 4$ (four{nucleon contact terms) (see g.5.2).

The corresponding Lagrangian reads

$$L^{(0)} = L + L_{N} + L_{NN} = \frac{1}{2D^{2}} (@ ~)^{2} \frac{M^{2}}{2D} ~^{2}$$

$$N i@_{0} m \frac{g_{A}}{2DF} t ~(~\tilde{r}) ~ \frac{1}{2DF^{2}} t ~(~_{0}@) N ~(5:11)$$

$$\frac{1}{2}C_{S} (NN) (NN) + \frac{1}{2}C_{T} (N~N) N(~N)$$

where C_s and C_T are new low (energy constants related to L of eq.(3.10). Notice that because of Ferm i statistics (Fierz rearrangement) one can rewrite a non (derivative four{nucleon contact term involving t as a combination of the last two terms in (5.11). It is straightforward to construct the interaction H am iltonian related to L⁽⁰⁾ as detailed in refs.[5.12,5.13]. For the two{nucleon case, the elective potential derived then is simply the sum of one{pion exchange and a contact interaction arising from the two last terms in eq.(5.11). One nds in coordinate space

with Y (r) = exp(M r)=4 r the standard Yukawa function. C learly, the potential (5.12) is only a crude approximation to the NN forces. In particular, the correlated J = T = 0 pion pair exchange that is believed to furnish the intermediate range attraction is hidden in the constant C_s . As stressed by W einberg [5.12,5.13], the constant C_s has to be "unnaturally" large to lead to shallow nuclear bound states. If one considers e.g. the L = 0 spin singlet state and approximates the potential by C ⁽³⁾ (r_1 r_2), with

 $C = C_S = 3C_T + g_A^2 = 4F^2$, one can solve the Lippm ann {Schwinger equation in m om entum space and nds after renorm alization ($C \ ! \ C_R$) a bound state with binding energy

$$B = \frac{16^{-2}}{m^{3}C_{R}^{2}} \qquad : \qquad (5:13)$$

A coording to naive dimensional analysis [5.19] one expects $C_R = 2^2 = 1 = 2^2 = 1 \text{ GeV}^2$. However, to get the deuteron binding energy of B = 2.22 MeV, $C_R = 2^2 \text{ must}$ have the large value of (260 MeV)². This then suggests that Q = m ($Q = -)^2$ and one adopts the rule that a pure nucleon intermediate state counts as if it contributes two more powers of 1=Q than any other intermediate state. Notice also that the lowest order NN potential leaves no room for the short-range repulsion or the spin (orbit forces and alike. On the other hand, one (pion exchange is known to describe well the higher partial waves in NN scattering at low energies.

In the work of refs.[5.16-5.18], the was also put in the elective theory based on the closeness of this resonance to the nucleon ground state [5.20], i.e. m M. In that case, one has additional lowest order terms, collected in L⁽⁰⁾

$$L^{(0)} = [i\theta_{0} \frac{1}{2F^{2}D}t^{(3=2)} (\sim_{0}\theta) m]$$

$$\frac{h_{A}}{2FD} [NT S(r) \sim +hc:] = N \sim tN \qquad NS[T + hc:] + :::$$
(5:14)

where the ellipsis stands for terms involving more 's which are irrelevant for the NN potential. The constant h_A can be calculated e.g. from the decay width (! N), h_A ' 2:7. D_T is a new low {energy constant and only enters the calculation of 3N (or more) forces.

To calculate corrections, one also has to consider the term swith $_i = 1$ and $_i = 2$. These are discussed in detail in refs.[5.14,5.16,5.17]. We only give a short outline of the pertinent elective Lagrangians here. For L⁽¹⁾, one nds

$$L^{(1)} = \frac{B_1}{4F^2D^2} N N [(r \sim)^2 \quad (Q \sim)^2] \quad \frac{B_2}{4F^2D^2} N t \sim N \quad (r \sim r \sim) \frac{B_3 M^2}{4F^2D} N N \sim^2$$
$$\frac{D_1}{2F D} N N N (t \sim r \sim) N \frac{D_2}{2F D} (N t \sim N N t \sim N) \quad r \sim$$
$$\frac{1}{2}E_1 N N N t N N t N \frac{1}{2}E_2 N N N t \sim N N t \sim N$$
$$\frac{1}{2}E_3 (N t \sim N N t \sim N) N t \sim N + :::$$

(5**:**15)

where the B_i , D_i and E_i are new parameters. B_3 is obviously related to the N {term and $B_{1;2}$ could be determined from N scattering (cf. section 4.3). At present, this has
not been done but all B_i , D_i and E_i are left free. The six{ferm ion term s proportional to $E_{1;2;3}$ do not enter the NN potential. A loo, term s with explicit 's are not shown. The term s with $_i = 2$ take the form

$$L^{(2)} = \frac{1}{2m} N r^{2} N \frac{A_{1}^{0}}{2F} N (t \sim r^{2}) t + \overline{r^{2}N} (t \sim r^{2}) N$$

$$\frac{A_{2}^{0}}{2F} \overline{rN} (t \sim r^{2}) r N^{0} [(NCrN)^{2} + (\overline{rNN})^{2}] C_{2}^{0} (NrN) (rNN) + :::$$
(5.16)

where the A_i^0 and C_i^0 are undetermined coecients and the ellipsis denotes other terms with two derivatives or more pionelds. Because only the term proportional to B_1 contains a time derivative, the corresponding interaction Hamiltonian can be taken as $L^{(1)} = L^{(2)}$ up to term s with more pionelds.

We are now in the position to system atically discuss the corrections to the lowest order potential $V_{12}^{(0)}$, eq.(5.12). As already noted in [5.13], the rst corrections arise from the same graphs as in g.5.2 with exactly one insertion from L ⁽¹⁾. However, since all time derivatives have been eliminated, one would have to construct a vertex with an odd number of three{m om enta. This clashes with parity and one therefore concludes that

$$V_{12}^{(1)} = 0$$
 : (5:17)

The second corrections fall into two classes. The $\;$ rst one are tree graphs with exactly one vertex or kinetic energy insertion from L $^{(2)}$, leading to

$$V_{12}^{(2)}(\mathbf{q};\mathbf{\tilde{k}}) = \frac{2g_{A}}{F}\mathbf{\tilde{t}}_{1} \quad \mathbf{\tilde{t}}_{2}\frac{\sim_{1}}{\mathbf{q}^{2} + M^{2}}\mathbf{q}_{1}\mathbf{q}^{2} + A_{2}\mathbf{\tilde{k}}^{2} + \frac{2g_{A}}{(\mathbf{q}^{2} + M^{2})^{1-2}} = \frac{\mathbf{q}^{2}}{4\mathbf{m}} \quad \frac{\mathbf{\tilde{k}}^{2}}{\mathbf{m}} + C_{1}\mathbf{q}^{2} + C_{2}\mathbf{\tilde{k}}^{2} + (C_{3}\mathbf{q}^{2} + C_{4}\mathbf{\tilde{k}}^{2})\sim_{1} \quad 2^{*} + i\frac{C_{5}}{2}(\sim_{1} + \sim_{2}) \quad (\mathbf{q}\;\mathbf{\tilde{k}}) + C_{6}\sim_{1} \quad \mathbf{q}_{2^{*}} \quad \mathbf{q} + \mathcal{C}\sim_{1} \quad \mathbf{\tilde{k}}\sim_{2} \quad \mathbf{\tilde{k}}$$

$$(5:19)$$

with q = p p^0 the transferred m on entum, $\tilde{k} = (p + p^0) = 2, 2m + E$ the energy in the center of m ass and p $\binom{0}{i}$ is the initial (nal) on m on entum. The A_i;C_i are combinations of the A_i;C_i⁰ in (5.16). Second, there are the one{loop contributions, i.e. the two{pion exchange, of the form

$$V_{12;loop}^{(2)} = V_{12;no}^{(2)} + V_{12;one}^{(2)} + V_{12;two}^{(2)}$$
(5.20)

corresponding to no, one and two isobars in the intermediate states. The rst term on the rh.s. of (5.20) reads [5.14]

$$V_{12,no}^{(2)} = \frac{1}{32F^4} t_1 t_2^2 \frac{d^3 l}{(2)^3} \frac{1}{!_+!} \frac{(!_+!_!)^2}{!_++!} \frac{g_A}{2F^2} t_1 t_2^2 \frac{d^3 l}{(2)^3} \frac{1}{!_+!} \frac{q^2 l^2}{!_++!} \frac{q^2}{!_++!} \frac{q^2}{!_++!$$

 $q = (q - 1)^2 + 4M^2$ and the explicit expressions for the contributions with one or two interm ediate isobars can be found in ref.[5.16]. Finally, the third corrections have also been evaluated. For the same reason as discussed above, one nds

$$V_{12;tree}^{(3)} = 0$$
 ; (5:21)

and in the one { loop graphs one has exactly one insertion from eq.(5.15) leading to

$$V_{12;loop}^{(3)} = V_{12;no}^{(3)} + V_{12;one}^{(3)}$$
 (5.22a)

with

$$V_{12;no}^{(3)} = \frac{1}{4} \frac{g_{A}}{2F^{2}} \sum_{i=1}^{2} \frac{d^{3}l}{(2)^{3}} \frac{1}{!_{+}^{2}!^{2}} - 3(q^{2} \tilde{l}^{2})^{2} 4M^{2}B_{3} = B_{1}(q^{2} \tilde{l}^{2}) + 16B_{2} \sum_{i=1}^{2} (q^{2})^{2} 4M^{2}B_{3} = B_{1}(q^{2}) \sum_{i=1}^{2} (q^{2})^{2} (q^{2})^{2} + 16B_{2} \sum_{i=1}^{2} (q^{2})^{2} q^{2} q^{2}$$

and the corresponding expression with one intermediate is given in ref.[5.16]. We are now at the point to discuss the structure of the momentum space potentials. The term proportional to A_1 in (5.19) can be considered as coming from the expansion of the pion {nucleon form factor in powers of momenta over the cut{o. Indeed, a typical monopole form factor F_M (q^2) = $^2 = (^2 + q^2)$ would amount to $A_1 = 2 = ^2 = 2 \text{ GeV}^2$ for = 1 GeV. The A₂ {term is a so {called non-adiabatic correction and the last term in the square brackets in (5.19) is the energy {dependent recoil correction. Thism odied one (pion exchange gives the long (range part of the potential. At interm ediate distances, the two {pion exchange generated from the one { loop diagram s com es in. M any of the box and crossed box diagram s are well-known from the work of Bruckner and W atson [5.21], Sugawara and von Hippel [522] and Sugawara and Okubo [523]. However, the diagram s {vertex have coe cients which are either xed by chiral sym m etry (like with one N N e.g. g_{A}^{4} or $(g_{A} h_{A})^{2}$), or are in principle determ ined from N scattering (the $B_{1:2:3}$). We com e back to these later on. Furtherm ore, at this order there is no correlated two{pion exchange, it only shows up at order $(Q = M)^4$ and higher. This is consistent with the analysis of the interm ediate (range attraction m ade in ref. 5.24) based on the spectral analysis of the scalar pion form factor. All physics of shorter ranges is buried in the various contact terms, i.e. the coe cients C_i. The various loop integrals like (5.20a) or (5.22b) are all divergent. At present, this is treated by a momentum space cut{o. The form of the cut{o function is chosen to be gaussian as in the N im egen approach [5.9]. Speci cally, all bop m om enta lare cut o by exp ($1^2 = 2^2$). Furtherm ore, since as argued before all m om enta should be sm aller than som e scale , the transferred m om entum q is also damped with the same type of $cut\{o, exp(q^2 = 2^2)\}$. In practise, = M is chosen. Of course, one would like to see a more elegant regularization employed such

as dimensional regularization. The potential is then transformed into coordinate space, where it is energy-dependent and takes the form

$$V = \sum_{p=1}^{X^0} V_p (r; \frac{\theta}{\theta r}; \frac{\theta^2}{\theta r^2}; E) O^p ; \qquad (5.23)$$

with

$$V_{p}(\mathbf{r};\frac{\theta}{\theta r};\frac{\theta^{2}}{\theta r^{2}};E) = V_{p}^{0}(\mathbf{r};E) + V_{p}^{1}(\mathbf{r};E)\frac{\theta}{\theta r} + V_{p}^{2}(\mathbf{r};E)\frac{\theta^{2}}{\theta r^{2}}; \qquad (5:23a)$$

and the O^{p=1;:::;20} are a complete basis of operators made of the \sim_i , the \sim_i (i = 1;2), the tensor operator S₁₂, the total spin operator $S = (\sim_1 + \sim_2)=2$ as well as the angular momentum operator L = ir \tilde{r} . Alltogether, the potential contains 26 parameters, but it should be stressed that some of these are indeed not free but given by constraints from N scattering. Let us brie y dicuss the connection between the B_i (i = 1;2;3) and the various c_i discussed in section 3.4.0 ne nds [5.25]

$$B_{1} = 4c_{3} = 13.6 \,\text{GeV}^{1}; B_{2} = 8c_{1} = 7.0 \,\text{GeV}^{1}; B_{3} = 4q \frac{1}{m} = 17.5 \,\text{GeV}^{1};$$
(5.24)

for the central values of the c_i from section 3.4. These constraints have not yet been in plemented in the numerical calculations. Furthermore, in the tting procedure of ref.[5.17], even the fundamental parameters F, g_A and h_A were left free.

The parameters are xed from a best t to deuteron properties (binding energy, m agnetic m om ent and electric quadrupole m om ent) and the np and pp phase shifts with J 2 and T_{lab} 100 M eV. The higher partial waves are supposedly dom inated by one pion exchange and were therefore not used to constrain the t [5.17]. The results for the deuteron properties are sum marized in table 5.1 and some typical phases are shown in

g.5.3 (m ore of these can be found in ref.[5.17]). The resulting values for F , g_A and h_A are 86 M eV, 1.33 and 2.03, respectively, not too far from their empirical values. U sing the G oldberger{Treim an relation, this corresponds to a pion{nucleon coupling constant of 14.5.

0 bærvable	F盳	Exp.
B [M eV]	2.18	2,224579 (9)
_d [n.m.]	0.851	0.857406(1)
Q [fm ²]	0.231	0.2859 (3)
	0.0239	0.0271(4)

Table 1: Deuteron properties: binding energy B, magnetic moment $_d$, quadrupole moment Q and the asymptotic D/S ratio [5.17]. The data are from ref.[5.26].



Fig. 5.3: Best tto some partial waves. Phase shifts in degrees versus T_{lab} in MeV.W e thank L.Ray and U.van Kolck for supplying us with the pertinent numbers.

The calculated D {state probability is 5%, i.e. of comparable size to what one gets from the Bonn or Paris potential. The L = 0 singlet and triplet scattering lengths are predicted to be -15.0 fm and 5.46 fm, in fair agreem ent with the empirical values of -16.4(1.9) fm and 5.396(11) fm [5.27], respectively. A close look at table 1 and the phase shifts reveals that the t is not too satisfactory, in particular the deviation in the deuteron quadrupole m om ent as well as in certain P {waves are quite substantial for the accuracy one is used from the sem i-phenom enological potentials. Holinde has argued [5.28] that some of these discrepancies reside in the asymmetric treatment of the pions and the mesons. In particular, the ne cancellations between the tensor forces from the are unbalanced here. This in turn leads to an overall unsatisfactory the and the tensor force which mostly shows up in the before mentioned observables. At present, it is not clear how one has to go about these problem s. Clearly, more detailed ts allowing also for variations in the cut-o function and its associated cut o are called for and as already stressed a few times, the strictures from the single nucleon sector on some of the parameters should be enforced. On the positive side, it is worth pointing out that such

a straightforward potential based solely on chiral symmetry constraints can describe the low {energy NN phases and deuteron properties within some accuracy.

V.3. MORE THAN TWO NUCLEONS

Nucleons interact m ainly via two{body forces. However, there is some indication of sm all three{nucleon forces (for a recent review see ref.[5.29]). Standard two{nucleon potentials when employed to 3 H and 4 H e tend to lead to an underbinding of typically 0.5 to 5 M eV. This in turn m eans that if this discrepancy is due to a three{nucleon force, it has to be sm all, typically a few percent of the 2N force. However, recent calculations of the Bochum group [5.30] have indicated that a ne{tuning of the NN {potential can lead to a satisfactory description of the three{nucleon system. This, however, involves rather large charge{sym m etry breaking e ects. The chiralLagrangian analysis can shed som e light about the size of three (and m any) nucleons forces to be expected as w ill be discussed in this section. The role of chiral sym m etry, i.e. the use of the pseudovector N coupling leading to sm all 3N forces has long been conjectured [5.31,5.32]. A s w ill be show n, this can now be put on mer grounds.

To leading order, the potential between A nucleons is simply given by a pair{w ise sum of the lowest order two{nucleon potential (5.12) since decreasing the number of connected pieces C costs powers of Q, cf. eq.(5.9). Therefore, $m_{in} = 6$ 3A leading to

$$V^{(0)}(\mathbf{r}_{1}; :::; \mathbf{r}_{A}) = \bigvee_{\substack{(ij)}} V^{(2)}(\mathbf{r}_{1} + \mathbf{r}_{J});$$
 (5:25)

where the sum runs over all nucleon pairs. This is a rather crude approximation. Therefore, one has to consider corrections [5.13,5.18]. We follow here the recent analysis by van Kolck [5.18]. To order $= m_{in} + 2$, one has the following form for the potential between A nucleons:

At this order, the 2N potential contains the one{pion exchange recoil (5.19) am ong other terms. The 3N potential consists of the three types of terms shown in g.5.4. and the double pair potential $V_{2;2}$ is made of two sets of diagrams, the rst being two disconnected OPE graphs and the second one one OPE graph separated from a lowest order two{nucleon contact term. It was rst shown by W einberg [5.12] that all diagrams s containing the non{linear N vertex add up to zero to lowest order. Furtherm ore, as detailed in [5.18], the remaining three{body forces and double pair forces are canceled by the energy-dependence of the two{body potential when the latter is iterated in the



Fig. 5.4: Tree graphs contributing to the three{nucleon potential. For each class of contributions, one typical diagram is shown. All other irreducible time orderings have to be considered.

Lippm ann-Schwinger equation. Such kind of cancellation had been noticed before [5.33] and it means that if one chooses to work with an energy {dependent N N {potential, one has to include at the same time 3N and double{2N forces calculated consistently in the same framework.

The corrections at next order, $= m_{in} + 3$ are discussed in [5.18]. The correction to the double{pair potential vanishes for the same reasons discussed before (5.17) and the remaining 3N potential takes the form

$$V_{3}^{(3)}(\mathbf{q}_{ij};\mathbf{q}_{jk}) = E_{1}\mathbf{t}_{i} \ \mathbf{t}_{k} + E_{2}\sim_{i} \ \kappa^{*}\mathbf{t}_{i} \ \mathbf{t}_{k} + E_{3}\sim_{j} \ (\kappa \ \gamma_{k})\mathbf{t}_{j} \ \mathbf{t}_{i}(\mathbf{t}_{k})$$

$$\frac{g_{A}}{2F^{2}} \frac{1}{!_{ij}}\sim_{k} \ \mathcal{B}_{1} \ D_{1}(\mathbf{t}_{i} \ \mathbf{t}_{k}\sim_{i} + \mathbf{t}_{j} \ \mathbf{t}_{k}\sim_{j}) \ 2D_{2}\mathbf{t}_{j} \ \mathbf{t}_{i}(\mathbf{t}_{k})\sim_{i} \ \gamma_{j} \ \mathcal{B}_{1}$$

$$+ 2 \ \frac{g_{A}}{2F^{2}} \ \frac{2}{!_{ij}!_{jk}^{2}}\sim_{i} \ \mathcal{B}_{1}^{*}\mathbf{t}_{i} \ \mathbf{t}_{k} \ (B_{1}\mathbf{q}_{ij} \ \mathcal{B}_{1} + B_{3}M^{2})$$

$$B_{2}\mathbf{t}_{j} \ \mathbf{t}_{i}(\mathbf{t}_{k})\sim_{j} \ (\mathbf{q}_{i} \ \mathbf{q}_{k}) + \text{two cyclic permutations of (ijk) ; (5:27)}$$

which contains 8 parameters, three of which are in principle xed by N scattering, cf. (5.24), and the D_i could be determined form {deuteron scattering or pion production on two{nucleon systems. The three E_i can only be xed from data on 3N systems. Such an analysis is not yet available. A simplication arises if one includes the (1232) in the e ective Lagrangian. In that case, one has an additional 3N force of order = m in + 2 which has the form (5.27) and the corresponding low {energy constants can be expressed in terms of properties,

$$E_{1} ! 0; E_{2} ! \frac{1}{9} \frac{D_{T}^{2}}{m m_{N}}; E_{3} ! \frac{1}{18} \frac{D_{T}^{2}}{m m_{N}};$$

$$D_{1} ! \frac{4}{9} \frac{D_{T} h_{A}}{m m_{N}}; D_{2} ! \frac{2}{9} \frac{D_{T} h_{A}}{m m_{N}};$$

$$B_{1} ! \frac{4}{9} \frac{h_{A}^{2}}{m m_{N}}; B_{2} ! \frac{2}{9} \frac{h_{A}^{2}}{m m_{N}}; B_{3} ! 0$$
(5:28)

with D_T a new low {energy constant. The terms proportional to h_A^2 , i.e. the two{ pion exchange pieces, are nothing but the old Fujita{M iyazawa force [5.34] which is

accompanied here by a shorter{range contribution proportional to the parameter D_T . The relation of these results to existing three{nucleon force models is discussed in [5.18]. No explicit 3N calculation has yet been performed in the framework outlined here.

To sum marize, the chiral Lagrangian approach in plies that few {nucleon forces are generically smaller than the dom inant two{nucleon forces. There are strong cancellations between the leading (static) 3N force, the double{pair forces and the iterated leading energy{dependence of the two{nucleon force. The remaining 3N force is expected to be dom inated by the Fujita{M iyazawa force plus a shorter{range term involving one new parameter, D_T . These 3N forces are expected to be of order 0 (M ²=M ²), i.e. som e 5% of the N N contribution. By a sim ilar argument, one expects even smaller 4N forces of order 0 (M ⁴=M ⁴) (less than 1% com pared to the N N contribution). Consequently, this analysis leads one to expect that four{nucleon system s are underbound by roughly four times the triton underbinding when pure N N forces are used. These dimensional arguments have yet to be substantiated by a quantitative calculation in the fram ework outlined here.

V.4. THREE (BODY INTERACTIONS BETW EEN NUCLEONS, PIONS AND PHOTONS

In the previous sections we saw that the calculation of the two{body interactions between nucleons involves a large number of free parameters and that the resulting potential does not yet have the accuracy of the standard sem i{phenom enological ones. It was therefore proposed by W einberg [5.15] to use the empirical know ledge about the two (body interactions between nucleons as well as pions and nucleons and combine these with the remaining contributions from the potential of the same power in small m om enta, which are graphs with three particles (or two pairs of particles) interacting. Stated di erently, if one looks at any nuclear process like elastic pion scattering, pion photoproduction and so on, the calculation of the S{matrix element $< A_{\rm J}$] $A_{\rm J} > A_{\rm J}$ is split into two parts. On one hand, chiral perturbation theory is used to calculate the irreducible kernel I to a certain power in Q = M and on the other hand, one uses phenom enological input to construct the nuclear wave{function A. The virtue of this method lies in the fact that it orders the relevant contributions to I in a systematic fashion and thus can explain the dom inance of certain digram s contributing to a certain process (which is often already known from models but also often not fully understood). One thus encompasses many of the problem swhich arise in the CHPT calculation of the NN interaction. However, one also looses a certain degree of consistency since one does not calculate nuclear wave{functions and operators in the same fram ework. This has to be kept in m ind in what follows. How much this could be improved by system atically generating also the nuclear wave {functions from a potential solely derived from chiral symmetry as described in section 5.2 is not yet clear.

In ref.[5.15], this method is applied to pion scattering on complex nuclei. To be more speci c, the calculation is simplied by considering the corresponding scattering length, i.e. the reaction with the in-com ing and out-going pion having vanishing three{ m om entum. In this process, we have $N_n = A$ external nucleons and N = 1 external pions. The leading irreducible graphs are those in which the pion scatters o a single nucleon, evaluated using the lowest order vertices with $_i = 0$ in the tree approximation (this is what is called the in pulse approximation). To second order in smallmomenta, the two{body interactions involving loop graphs and tree diagrams with $_i = 1;2$ are taken from phenomenological models of N scattering and the remaining three{body interactions between two nucleons and the pion (calculated from tree diagrams with $_i = 0$ vertices) are shown in g.5.5 (these are the ones that contribute to the pion{ nucleus scattering length).



Fig. 5.5: Irreducible connected graphs for the interaction of a pion with a pair of nucleons that contribute to the pion {nucleus scattering length. Only one time {ordered diagram per class is shown.

The details of the calculation are found in ref.[5.15]. The pion{nucleus scattering length takes the form

$$a_{ab} = \frac{1 + M = m_{N}}{1 + M = Am_{N}} X_{ab} + a_{ab}^{(r)} + a_{ab}^{(r)} ; \qquad (5.29)$$

where a, b are the pion isovector indices, $a_{ab}^{(r)}$ is the pion scattering length of the rth nucleon and m_N is the nucleon mass. The three{body contribution stemming from diagram s 5.5a{ftakes the form

$$a_{ab}^{three \ body} = \frac{M^2}{32 \ ^4F^4 (1 + M = m_d)} \frac{X}{r < s} \frac{1}{q_{rs}^2} 2t^{(r)} \ t^{(s)} ab \ t^{(r)}_{a} t^{(s)}_{b} \ t^{(s)}_{a} t^{(s)}_{b}$$

$$\frac{g_{A}^{2}}{32^{-4}F^{-4}(1+M^{-1}m_{d})} \sum_{r

$$+ \frac{g_{A}^{2}}{32^{-4}F^{-4}(1+M^{-1}m_{d})} \sum_{r(5:30)$$$$

where r;s label individual nucleons and $(t_c^{(\)})_{ab} = i_{abc}$ is the pion isospin vector. Notice that there is some cancellation between the second and third term in eq.(5.30) as q_{rs} ! 1 so that the result is less sensitive to the nuclear wave function at sm all separation (see also ref.[5.35]). For an isoscalar nucleus like the deuteron, the expressions in (5.30) simplify considerably since the last term vanishes and $t_a^{(r)}t_b^{(s)} + t_a^{(s)}t_b^{(r)}$ can be replaced by (2=3) $_{ab} \tau^{(r)} \tau^{(s)}$. Even more in portant, for an isoscalar nucleus the nom inally leading term in eq.(5.29) are vanishing since they involve an expectation value of $_{r} \tau^{(r)} \tau^{()}$. So one is left with sm all O (M²) contributions of the {term type to the impulse approximation. This is the reason why it makes sense to compare the corrections calculated in CHPT directly with the empirical values of the d scattering length and nds [5.36,5.37]

$$\frac{1 + M = m_N}{1 + M = m_d} (a_p + a_n) = (0.021 \quad 0.006) M^1 ;$$
 (5.31)

with m_d the deuteron mass. The rst term in (5.30) gives the well{known and large rescattering contribution. It is much bigger than the remaining three{body terms due to the anom alously large radius of the deuteron. Using empirical information on N scattering to calculate the rescattering contribution (for details, see e.g. [5.37]), and the Bonn potential to produce the deuteron wave function for the calculation of the remaining three{body contributions [5.15], one nds

$$a^{\text{three body}} = (0.026 \ 0.001) M^{1} \ 0.0005 M^{1}$$
; (5:32)

where the rst number refers to the rst term in (5.30) and the second one to the remaining three{body contributions. The latter ones are very small, well within the uncertainties of the other dom inant terms. This justi as the naltheoretical result of

 $(0.047 \ 0.006)$ M¹ in good agreem ent with the empiral value of $(0.056 \ 0.009)$ M¹ [5.37]. This is a good example how the chiral Lagrangian machinery can be used to explain why one is allowed to take only certain graphs like the rescattering contribution but neglect the others which are of the same order in the expansion in sm all m om enta.

A similar calculation has been been performed by Beane et al. [5.38] for pion photoproduction on nuclei. They have considered all corrections which are suppressed

by two powers in sm allm on enta as compared to the lowest order in pulse approximation using $_i = 0$ vertices. Again, there is a single scattering contribution taken from phenom enology plus some three{body interactions (and disconnected graphs involving pairs of nucleons for A > 2). In the case of the deuteron and considering neutral pion photoproduction, the calculation simpli es enorm eously. At threshold, the invariant matrix{element takes the form

$$M = 2iJ \sim E ; \qquad (5:33)$$

with ${\tt J}$ the spin of the deuteron. The single scattering contribution can be compactly written as

$$E^{ss} = \frac{1 + M}{1 + M} = m_{d} E_{0+}^{0} + E_{0+}^{0} S (k=2) = 1:33 \quad 10^{3} M^{-1}$$
(5:34)

with S (k=2) the deuteron form factor evaluated for the threshold kinem atics (k = 0:685 fm⁻¹) and using the currently accepted value of $E_{0+}^{0}{}^{p} = 2:0 \quad 10^{\circ} M^{-1}$ and taking $E_{0+}^{0}{}^{n} = 0:5 \quad 10^{\circ} M^{-1}$ from the incomplete \LET " as discussed in section 4.4. C learly, this prediction hinges on these particular values for the elementary pion photoproduction electric dipole am plitudes. A lso, the one for the n! 0 n is not taken from experiment. The three(body graphs which contribute are of the exchange current type (see also the next section). In class (a), the photon couples to the pion which is interchanged between the two nucleons and the second class (b) involves the diagram swith exactly one N N and one N N low est order vertex. Their contributions to E_d take the form

$$E^{(a)} = \frac{eg_{A} M m_{N}}{8^{2} (M + m_{d})F^{3}k} F^{(kr)} dr$$

$$E^{(b)} = \frac{eg_{A} M m_{N}}{16^{2} (M + m_{d})F^{3}} \int_{0}^{2} dz G^{(zkr)} dr$$
(5:35)

with the integrands F (kr) and G (zkr) given in [5.38]. Using again the Bonn potential to generate the deuteron wave function, one nds

$$E^{(a)} = 224 \quad 10^{\circ} M^{-1}; \quad E^{(b)} = 0242 \quad 10^{\circ} M^{-1}:$$
 (5:36)

Sum m ing up (5.34) and (5.36), one nds $E_d = 3:99 \ 10^{\circ} M^{-1}$ in good agreem ent with the empirical value of (3:74 0:25) $10^{\circ} M^{-1}$ [5.39]. How ever, as already stressed, the single scattering contribution (5.34) is a icted by large uncertainties and it remains to be seen which value for $E_{0+}^{\circ p}$ the new data from M ainz and Saskatoon will favor and how accurate the guess for the electric dipole am plitude $E_{0+}^{\circ n}$ will turn out to be (once m easured).



Fig. 5.6: The lowest order and dom inant pion exchange current diagrams. The wiggly line denotes an electroweak probe.

V.5. EXCHANGE CURRENTS

M eson exchange currents arise naturally in the m eson {exchange picture of the nuclear forces. An external electrom agnetic or axial probe does not only couple to the nucleons (im pulse approxim ation, one{body operators) but also to the m esons in ight or leads to nonlinear seagull-type vertices (these are typical two{body operators), cf. g.5.6.

A lso known since many decades [5.40], the rst compelling evidence for meson exchange currents came from the calculation of neutron radiative capture at threshold and deuteron photodisintegration [5.10,5.11] (for reviews, see [5.19,5.20]). By now, the existence of these two{body operators can be considered veri ed experim entally [5.41]. M ore than 10 years ago, the so{called "chiral lter hypothesis" was introduced [5.42]. It states that the response of a nucleus to a long{wavelength electroweak probe is given solely by the soft{pion exchange terms dictated by chiral symmetry. Consider any exchange current contribution X, this means

$$X = X + X_2 + X_R + X_N = X_{soft pion} (1 + C)$$
 (5:37)

where C is a generally small correction to the leading one (C 1), R denotes the e ects of heavier m eson exchange and N the excitation of nucleon resonances. Stated di erently, all the heavier m esons and nucleon excitations, multi{pion exchanges and form factor e ects are not seen, even up to energies of the order of 1 G eV (although individual contributions can be large). W hy this holds true at such energies has not yet been explained.

Rho [5.43] has given a simple argument how the "chiral lter" can occur in nuclei for small and moderate momentum transfer. H is lowest order analysis follows closely the one of W einberg [5.12]. Any matrix {element M E of the electrice potential V or of a current J has the form M E Q F (Q = m), as discussed before. In the presence of a slow ly varying external electrom agnetic eld A (or a weak one), the H am iltonian takes the form

and this additional term H_{ext} modi es the power counting. Since one derivative is replaced by the external current, the tree graphs (L = 0) with the lowest power must full

$$d_i + \frac{1}{2}n_i \quad 2 = 1 \tag{5:39}$$

which leads to $d_i = 0$ and $n_i = 2$ or $d_i = 1$ and $n_i = 0$. In contrast to the case of the nuclear forces, to leading order no four{nucleon contact terms contribute. This means that there is no short{ranged two{body current (to lowest order), the exchange current is entirely given in terms of the soft{pion component derived from (5.38). This justi es the chiral lter hypothesis at tree level.

Park et al.[5.44] have also investigated one {loop corrections to the axial {charge operator, the st correction to the pertinent soft { pion m atrix { elem ent is suppressed by $(Q = m)^2$. The authors of ref.[5.44] use the heavy mass form alism which simplies the calculation considerably. They argue that {function type contact term s are suppressed by the short { range nuclear correlations. Stated di erently, since the chiral counting is only meaningful as long as $Q^2 = m^2$ 1, one can not describe processes that involve energy or momentum scales exceeding this criterion. Short{distance interactions are therefore not accessible by chiral perturbation theory. This is di erent in philosophy from the calculation of the nuclear forces by van Kolck et al. [5.14,5.17] where it is arqued that the four {nucleon contact term s are sm eared out over a distance 1=M .In ref.[5.44], it is shown that the loop corrections are small for distances r 0:6 fm, which m eans that the low est order argum ent of R ho [5.43] is robust to one { loop order. To be m ore precise, the results of ref. [3.44,5.45] can be sum m arized as follows. One writes the nuclear m atrix element of the axial (charge operator as

$$M^{axial} = M_1 + M_2; \quad M_2 = M_2^{tree} (1+)$$
 (5:40)

where the subscript n = 1;2 refers to one{ and two{body operators and 'tree' correponds to the diagram s shown in g.5.6 (with renormalized couplings). One nds almost independently of the mass number that < 0:1, i.e. the tree contribution dom inates. In this particular case, the two{body operator is of comparable size to the one{body operator, $M_2^{\text{tree}}=M_1$ 0:7 [5.45] which is su cient to explain the empirical value of $M_2^{\text{exp}}=M_1$ 1:6:::2:0 [5.46].

The therm all np capture has also been considered by Park et al. [5.47] including terms to order $(Q = m)^3$. A part from the dominant one pion exchange diagrams (g.5.6), there are additional graphs corresponding to two {pion exchange as well as counterterm contributions saturated by and ! m eson exchange. While in impulse approximation one nds (np! d) = 305:6 mb, the exchange currents calculated in CHPT together with a short{range correlation cut{o} 0 < r_c < 0:7 fm lead to (np! d) = 334 3 mb, in nice agreement with the experimental value, exp (np! d) = 334:2 0:5 mb [5.48]. A gain, the soft pion contribution gives the dominant part of the two {body enhancement.

The calculations presented here seem to lend credit to the chiral lter hypothesis and dem onstrate once m one the importance of chiral symmetry in nuclear phenomena. Furtherm one, in nuclei it appears natural to make use of the short{range correlations to suppress operators of the contact term type and alike. Form one details on the calculation of exchange currents from chiral Lagrangians we refer the reader to refs.[5.45,5.47,5.49]. W hat remains mysterious is why the chiral lter hypothesis works up to so high energies { the answer to this lies certainly outside the realm of baryon CHPT.

REFERENCES

- 5.1 H.Yukawa, Proc. Phys.-M ath. Soc. Jpn. 17 (1935) 48.
- 5.2 R.Machleidt, K.Holinde and Ch.Elster, Phys. Rep. 149 (1987) 1.
- 5.3 W N.Cottingham et al., Phys. Rev. D 8 (1973) 800:
 M.Lacom be et al., Phys. Rev. C 21 (1980) 861.
- 5.4 M M. Nagels, T.A. Riken and J.J. de Swart, Phys. Rev. D 17 (1978) 768.
- 5.5 S.Deister et al, Few {Body System s 10 (1991) 1.
- 5.6 IE.Lagaris and V R.Phandharipande, Nucl. Phys. A 359 (1981) 331: R B.W iringa, V G J.Stoks and R.Schiavilla, \An accurate Nucleon {Nucleon Potential", ANL preprint 1994, nucl-th/9408016.
- 5.7 S.O. Backman, G.E. Brown and J.A. Niskanen, Phys. Reports 124 (1984) 1.
- 5.8 G.E. Brown and A.D. Jackson, \The Nucleon {Nucleon Interaction", North { Holland, Amsterdam, 1976.
- 5.9 T A.Rijken, Ann. Phys. 164 (1985) 1, 23.
- 5.10 D.O.Riska and G.E.Brown, Phys. Lett. B 38 (1972) 193.
- 5.11 J.Hockert, D.O.Riska, M.Gariand A.Hu mann, Nucl. Phys. A 217 (1973) 19.
- 5.12 S.W einberg, Phys. Lett. B 251 (1990) 288.
- 5.13 S.W einberg, Nucl. Phys. B 363 (1991) 3.
- 5.14 C.Ordonez and U.van Kolck, Phys. Lett. B 291 (1992) 459.
- 5.15 S.W einberg, Phys. Lett. B 295 (1992) 114.
- 5.16 U. van Kolck, Thesis, University of Texas at Austin, 1992
- 5.17 C.Ordonez, L.Ray and U.van Kolck, Phys. Rev. Lett. 72 (1994) 1982.
- 5.18 U. van Kolck, Phys. Rev. C 49 (1994) 2932.
- 5.19 JF.M athiot, Phys. Reports 173 (1989) 63.
- 5.20 D.O.Riska, Phys. Reports 181 (1989) 207.
- 5.21 K A. Brueckner and K M. W atson, Phys. Rev. 92 (1953) 1023.
- 5.22 H. Sugawara and F. von Hippel, Phys. Rev. 172 (1968) 1764.
- 5.23 H. Sugawara and S. Okubo, Phys. Rev. 117 (1960) 605, 611.

- 5.24 Ulf-G.Meiner, Comments Nucl. Part. Phys. 20 (1991) 119.
- 5.25 Ulf-G.Meiner, \Baryon Chiral Perturbation Theory A.D. 1994", Bonn preprint TK 94 17, 1994, Czech. J. Phys., in print.
- 5.26 T E O . Ericson, Nucl. Phys. A 416 (1984) 281c.
- 5.27 R P.Haddock et al, Phys. Rev. Lett. 14 (1985) 318; H P.Noyes, Phys. Rev. 130 (1963) 2025.
- 5.28 K. Holinde, \Hadron {Hadron Interactions", Julich preprint, 1994.
- 5.29 J.L. Friar, Czech. J. Phys. 43 (1993) 259.
- 5.30 H.W itala, W.G lockle and Th. Cornelius, Few Body Systems 6 (1989) 79; Nucl. Phys. A 496 (1989) 446; Phys. Rev. C 39 (1989) 384;
 I.Slaus, R.M achleidt, W.Tornow, W.G lockle and H.W itala, Comments Nucl. Part. Phys. 20 (1991) 85;
 W.G lockle, private communication.
- 5.31 K A. Brueckner, C A. Levinson and H M. Mahmoud, Phys. Rev. 95 (1954) 217.
- 5.32 G E.Brown, A M.Green and W J.Gerace, Nucl. Phys. A 115 (1968) 435; G E.Brown, Comments Nucl. Part. Phys. 5 (1972) 6.
- 5.33 SN.Yang and W.Glockle, Phys. Rev. C 33 (1986) 1774; SA.Coon and JL.Friar, Phys. Rev. C 34 (1986) 1060.
- 5.34 J.Fujita and H.Miyazawa, Prog. Theor. Phys. 17 (1957) 360.
- 5.35 M R.Robilotta and C.W ilkin, J.Phys.G 4 (1978) L115.
- 5.36 JM. Eisenberg and D.S. Koltun, \Theory of Meson Interactions with Nuclei", Wiley{Interscience, New York, 1980.
- 5.37 T. Ericson and W W eise, \Pions and Nuclei", Oxford University Press, Oxford, 1988.
- 5.38 S.R. Beane, C.Y. Lee and U. van Kolck, Duke preprint, 1995.
- 5.39 P.Argan et al, Phys. Rev. Lett. 41 (1978) 629; Phys. Rev. C 21 (1980) 1416.
- 5.40 F.V illars, Helv. Phys. Acta 20 (1947) 476; H.M iyazawa, Prog. Theor. Phys. 6 (1951) 801.
- 5.41 B.Frois and J.{F.M athiot, Comments Part. Nucl. Phys. 18 (1989) 291.
- 5.42 K.Kubodera, J.Delorm e and M.Rho, Phys. Rev. Lett. 40 (1978) 755 M.Rho and G.E.Brown, Comments Part. Nucl. Phys. 10 (1981) 201.
- 5.43 M.Rho, Phys. Rev. Lett. 66 (1991) 1275.
- 5.44 T.-S.Park, D.-P.M in and M.Rho, Phys. Reports 233 (1993) 341.
- 5.45 T.-S. Park, I.S. Towner and K. Kubodera, Nucl. Phys. A 579 (1994) 381.
- 5.46 E K.W arburton and I.S. Towner, Phys. Reports 242 (1994) 103.
- 5.47 T.-S.Park, D.-P.M in and M.Rho, preprint SNUTP 94{124, 1992, nucl-th/9412025.
- 5.48 A E.Cox, S.W ynchank and C.H.Collie, Nucl. Phys. 74 (1965) 497.
- 5.49 M. Rho, Phys. Reports 240 (1994) 1.

VI.THREE FLAVORS, DENSE MATTER AND ALL THAT

In this section, we will rst be concerned with the extension to the case of three avors and discuss the calculation of baryon masses and {term s. Then, we will turn to the topic of kaon {nucleon scattering and the behaviour of pions in dense matter. This latter topic (also extended to kaons) is a rather new and rapidly developing eld, so we can only provide a state of the art sum mary. Finally, various om issions are brie y touched upon.

VI.1. FLAVOR SU (3), BARYON MASSES AND {TERMS

Let us rst provide the necessary de nitions for the three avor m eson {baryon system. It is most convenient to write the eight m eson and baryon elds in term s of SU (3) m atrices and B, respectively,

$$B = \begin{bmatrix} B \\ 0 \end{bmatrix} \xrightarrow{p_1} 0 + p_1 = 0 \\ 0 \\ 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p_1 = 0 \\ p_2 = 0 \\ p$$

with

$$U() = u^{2}() = expfi = F_{p}g$$
 (62)

with F_p the pseudoscalar decay constant in the chiral limit. Of course, beyond leading order, one has to account for the fact that $F \in F_K \in F$ [6.1]. The covariant derivative acting on B reads

$$D B = @ B + [;B]$$

= $\frac{1}{2} u^{y} @ i(v + a)]u + u @ i(v a)]u^{y}$ (6:3)

with v and a external vector and axial(vector sources. Under SU $(3)_{\rm L}$ SU $(3)_{\rm R}$, B and D B transform as

$$B^{0} = KBK^{Y}$$
; (D B)⁰ = K (D B)K^Y (6:4)

It is now straightforward to construct the lowest { order 0 (p) m eson { baryon Lagrangian,

$$L_{MB}^{(1)} = Tr iB D B m_0 BB + \frac{1}{2}DB _5 fu ; Bg + \frac{1}{2}FB _5 [u ; B]$$
 (6:5)

where m₀ stands for the (average) octet m ass in the chiral limit. The trace in (6.5) runs over the avor indices. Notice that in contrast to the SU (2) case, one has two possibilities of coupling the axial u to baryon bilinears. These are the conventional F and D couplings subject to the constraint $F + D = g_A = 1.26$. At order O (p²) the baryon m ass degeneracy is lifted by the term s discussed below. However, there are m any other term s at this order. If one works in the one{loop approximation, one also needs the term s of order O (p³) (or eventually from O (p⁴)). The complete locale ective Lagrangians $L_{MB}^{(2)}$ and $L_{MB}^{(3)}$ are given by K rause [6.2]. The extension of this to the heavy m ass form alism is straightforward, as spelled out in detail in the review article by Jenkins and M anohar [6.3]. For our purpose, we only give the low est order Lagrangian and the three term s of order q² which account for quark m ass insertions (in the isospin limit m_u = m_d),

$$L_{MB}^{(1)} = Tr(Biv DB) + DTB(Sfu;Bg) + FTr(BS[u;B])$$
 (6:6)

with the baryons considered as static sources and equivalently theirm on enta decompose as $p = m_0 v + 1$, v = 1 m. Beyond leading order and in the present context we consider only counter terms of chiral power q^2 which account for quark m ass insertions,

(2)

$$L_{MB}^{(2)} = b_{D} \operatorname{Tr}(Bf_{+};Bg) + b_{F} \operatorname{Tr}(B[_{+};B]) + b_{0} \operatorname{Tr}(BB) \operatorname{Tr}(_{+})$$
(6:7)

with $_{+} = u^{y} u^{y} + u^{y} u$ and $= 2B_{0} (M + S)$ where S denotes the nonet of external scalar sources. As we will see later on, the constants b_{D} , b_{F} and b_{0} can be xed from the know ledge of the baryon masses and the N -term (or one of the K N -term s). The constant b_{0} can not be determined from the baryon mass spectrum alone since it contributes to all octet members in the same way.

We now consider the calculation of baryon m asses in CHPT.G asser [6.4] and G asser and Leutwyler [6.5] were the st to system atically investigate the baryon m asses at next{to{bading order. The quark m ass expansion of the baryon m asses takes the form

$$m_{\rm B} = m_0 + M + M^{3=2} + M^2 + \dots$$
 (6:8)

The non {analytic piece proportional to M³⁼² was rst observed by Langacker and Pagels [6.6]. If one retains only the term s linear in the quark masses, one obtains the Gell-M ann {O kubo relation $m + 3m = 2 (m_N + m)$ (which is full led within 0.6% in nature) for the octet and the equal spacing rule for the decuplet, m m = m m = (experim entally, one has 142:145:153 M eV). However, to extract quark m ass m ratios > from the expansion (6.8), one has to work harder. This was done in refs. [6.4, 6.5]. The non{analytic terms were modelled by considering the baryons as static sources surrounded by a cloud of mesons and photons { truely the rst calculation in the spirit of the heavy m ass form alism. The most important result of this analysis was the fact that m_d) com es out consistent with the value obtained from the ratio R = (m) $m_{s}) = (m_{11})$

the m eson spectrum. Jenkins [6.7] has recently repeated this calculation using the heavy ferm ion EFT of refs. [6.3, 6.8], including also the spin {3/2 decuplet elds in the EFT. She concludes that the success of the octet and decuplet m ass relations is consistent with baryon CHPT as long as one includes the decuplet. Its contributions tend to cancel the large corrections from the kaon loops like m $_{\rm S}M$ $_{\rm K}^2$ ln M $_{\rm K}^2$. The calculation was done in the isospin lim it $m_u = m_d = 0$ so that nothing could be said about the quark m ass ratio R. This latter question was recently addressed by Lebed and Luty [6.9] who arrive at a negative conclusion concerning the possibility of extracting current quark m ass ratios > from the baryon spectrum. We follow here ref.[6.10] in which the whole scalar sector, ie the baryon masses and {term s, are considered and which sheds som e doubt on the results obtained so far when the decuplet is included in the EFT. Following [6.10], a com plete calculation up to order q^3 involves only intermediate octet states. At this order (one-loop approximation) one has three counterterms with a priori unknown but nite coe cients. These can be xed from the octet masses (m $_{\rm N}$;m ;m ;m) and the value $_{\rm N}$ (0) since one of the counter term s appears in the baryon mass form ulae in such a way that it always can be lum ped together with the average octet m ass in the chiral limit. This allows to predict the two K N -term s, $\binom{(1)}{K N}$ (0) and $\binom{(2)}{K N}$ (0) as well as the -term shifts to the respective C heng-D ashen points and the matrix element $m_s < p_{jss} p_j > .$ To this order in the chiral expansion, any baryon m ass takes the form

$$m_B = m_0 \frac{1}{24 F_p^2} B^{3} + B^{K} M_{K}^{3} + B^{M} B^{3} + B^{D} b_{D} + B^{F} b_{F} 2b_{0} (M^{2} + 2M_{K}^{2})$$
 (6:9)

The second term on the right hand side comprises the Goldstone boson loop contributions and the third term stems from the counter term seq.(6.7). Notice that the loop contribution is ultraviolet nite and non-analytic in the quark masses since M $_{,K}^3$, M $^{3=2}$. The constants b_D , b_F and b_D are therefore nite. The numerical factors read

$${}_{N} = \frac{9}{4} (D + F)^{2}; {}_{N}^{K} = \frac{1}{2} (5D^{2} \quad 6D F + 9F^{2}); {}_{N} = \frac{1}{4} (D \quad 3F)^{2};$$

$$= D^{2} + 6F^{2}; {}_{K}^{K} = 3 (D^{2} + F^{2}); {}_{E}^{M} = D^{2};$$

$$= 3D^{2}; {}_{K}^{K} = D^{2} + 9F^{2}; {}_{E}^{M} = D^{2};$$

$$= \frac{9}{4} (D \quad F)^{2}; {}_{K}^{K} = \frac{1}{2} (5D^{2} + 6D F + 9F^{2}); {}_{E}^{M} = \frac{1}{4} (D + 3F)^{2};$$

$${}_{N}^{D} = 4M_{K}^{2}; {}_{N}^{F} = 4M_{K}^{2} 4M^{2}; {}_{E}^{M} = 4M^{2}; {}_{E}^{F} = 0;$$

$${}_{D}^{D} = \frac{16}{3}M_{K}^{2} + \frac{4}{3}M^{2}; {}_{E}^{F} = 0; {}_{D}^{M} = 4M_{K}^{2}; {}_{E}^{F} = 4M_{K}^{2} + 4M^{2}:$$

$${}_{C}^{M} = 4M_{K}^{2}; {}_{E}^{F} = 0; {}_{E}^{M} = 4M_{K}^{2}; {}_{E}^{F} = 0; {}_{E}^{M} = 4M_{K}^{2}; {}_{E}^{F} = 4M_{K}^{2} + 4M^{2}:$$

At this order, the deviation from the Gell-M ann-O kubo form ula is given by

$$\frac{1}{4} 3m + m \qquad 2m_{N} \qquad 2m = \frac{3F^{2} D^{2}}{96 F_{p}^{2}} M^{3} \qquad 4M_{K}^{3} + 3M^{3}$$

$$= \frac{3F^{2} D^{2}}{96 F_{p}^{2}} M^{3} \qquad 4M_{K}^{3} + \frac{1}{P - \frac{1}{3}} (4M_{K}^{2} M^{2})^{3=2}$$
(6:10)

where in the second line we have used the GMO relation for the -m eson m ass, which is legitim ate if one works at next-to-leading order.

Further inform ation on the scalar sector is given by the scalar form factors or – term s which measure the strength of the various matrix-elements m $_q$ qq in the proton. One de nes:*

$${}_{N}(t) = \hat{m} < p^{0}juu + dd\dot{p} >$$

$${}_{K N}^{(1)}(t) = \frac{1}{2}(\hat{m} + m_{s}) < p^{0}juu + ss\dot{p} >$$

$${}_{K N}^{(2)}(t) = \frac{1}{2}(\hat{m} + m_{s}) < p^{0}j uu + 2dd + ss\dot{p} >$$
(6:11)

with $t = (p^0 \quad p)^2$ the invariant m on entum transfer squared and $m = (m_u + m_d)=2$ the average light quark m ass. At zero m on entum transfer, the strange quark contribution to the nucleon m ass is given by

$$m_{s} < pjsjp > = \frac{1}{2} - \frac{M^{2}}{4M_{K}^{2}} - 3_{KN}^{(1)}(0) + \frac{M^{2}}{KN}(0) + \frac{1}{2} - \frac{M_{K}^{2}}{M^{2}} - N(0)$$
 (6:12)

making use of the leading order m eson mass form ulae M 2 = 2m B $_0$ and M $_K^2$ = (m + m_s)B $_0$ which are su ciently accurate to the order we are working. The chiral expansion at next-to-leading order for the -term s reads

$$_{\rm N} (0) = \frac{{\rm M}^2}{64 {\rm F}_{\rm p}^2} - 4 {\rm N} {\rm M} - 2 {\rm N}_{\rm N} {\rm M}_{\rm K} - \frac{4}{3} {\rm N} {\rm M} - 2 {\rm M}^2 (b_{\rm D} + b_{\rm F} + 2b_{\rm O})$$
 (6:13a)

for j = 1;2 with the coe cients

$${}^{(1)}_{K} = \frac{7}{3}D^{2} \quad 2D F + 5F^{2}; \quad {}^{(2)}_{K} = 3 (D F)^{2}; \quad {}^{(1)} = 1; \quad {}^{(2)} = 3;$$

$${}^{(1)}_{D} = 1; \quad {}^{(2)}_{D} = 0; \quad {}^{(1)}_{F} = 0; \quad {}^{(2)}_{F} = 1; \quad {}_{N} = \frac{1}{3} (D + F) (3F D):$$

$$(6:13c)$$

This completely determ ines the scalar sector at next-to-leading order. Note that the N-term is given as $_N(0) = m(m_N = 0m)$ according to the Feynman-Hellm an theorem.

^{*} These quantities are renorm alization-group invariant in a mass-independent subtraction scheme, which is what one usually employs.

The shifts of the $-\text{term} \ s$ from t = 0 to the respective C heng-D ashen points do not involve any contact term s_r

$$(2M^{2}) \qquad _{N} (0) = \frac{M^{2}}{64} \frac{4}{F_{p}^{2}} \frac{4}{3} _{N} M$$

$$+ \frac{2}{3} _{N}^{K} \frac{M^{2}}{P} \frac{M_{K}^{2}}{2M} \ln \frac{P}{P} \frac{2M_{K} + M}{2M_{K}} + M_{K}$$

$$+ \frac{4}{9} _{N} \frac{M^{2} M^{2}}{P} \frac{M^{2}}{2M} \ln \frac{P}{P} \frac{2M_{K} + M}{2M_{K}} + M$$

$$(6:14a)$$

(6:14b)

Notice that the shifts of the two K N -term s acquire an imaginary part since the pion loop has a branch cut starting at t = 4M ² which is below the kaon Cheng{D ashen point t = 2M $_{K}^{2}$ *. In the lim it of large kaon and eta m ass the result eq.(6.14a) agrees, evidently, with the ancient calculation of Pagels and Pardee [6.11] once one accounts for the num erical error of a factor 2 in that paper. Clearly, the -term shifts are nonanalytic in the quark m asses since they scale with the third power of the pseudoscalar m eson m asses. Our strategy will be the following: W e use the empirically known baryon m asses and the recently determined value of $_{N}$ (0) [6.12] to x the unknown parameters m $_{0}$; b_{F} and b_{0} . This allows us to predict the two K N -term s $_{K N}^{(j)}$ (0). The shifts of the -term s are independent of this t.

Since we use N(0) as input in what follows, let us brie y review the status of this much debated quantity. The quantity N(0) can be calculated from the baryon spectrum. To leading order in the quark masses, one nds

$$y = \frac{m}{m_{s}} \frac{m + m}{m_{s}} \frac{2m_{N}}{1} + 0 \text{ (M}^{3=2})$$

$$y = \frac{2 < pjssjp >}{< pjuu + ddjp >}$$
(6:15)

^{*} Since we choose the GMO value for the mass, $M + M > 2M_{K}$, the loop does not contribute to the imaginary part in eq.(6.14b). For the physical value of the mass this contribution is tiny compared to the pion loop.

where y is a measure of the strange quark content of the proton. Setting y = 0 as suggested by the OZI rule, one nds $_{\rm N}$ (0) = 26 M eV. However, from the baryon mass analysis it is obvious that one has to include the O (M $^{3=2}$) contributions and estimate the O (M 2) ones. This was done by G asser [6.4] leading to

N (0) =
$$\frac{35 \quad 5M \text{ eV}}{1 \quad y} = \frac{0}{1 \quad y}$$
 (6:16)

However, in N scattering one does not measure $_{\rm N}$ (0), but a quantity called $_{\rm N}$ de ned via

$$_{\rm N} = {\rm F}^2 {\rm D}^+ (=0;t=2{\rm M}^2)$$
 (6:17)

with the bar on D denoting that the pseudovector Born terms have been subtracted, D = DD_{DV}. The amplitudes D are related to the more conventional N scattering amplitudes A and B via D = A + B, with = (s u)=4m. The superscript '' denotes the isospin (even or odd). D is useful since it is related to the total cross section via the optical theorem. The kinematical choice = $0;t = 2M^2$ (which lies in the unphysical region) is called the Cheng {Dashen point [6.13]. The relation between Ν and the N scattering data at low energies is rather com plex, see e.g. Hohler [6.14] for a discussion or G asser [6.15] for an instructive pictorial (given also in ref.[6.16]). B ased on dispersion theory, Koch [6.17] found $_{\rm N}$ = 64 8 M eV (notice that the error only re ects the uncertainty of the method, not the one of the underlying data). Gasser et al. [6.12] have recently repeated this analysis and found N = 60 MeV (for a discussion of the errors, see that paper). There is still som e debate about this value, but in what follows we will use the central result of ref.[6.12]. Finally, we have to relate $_{\rm N}$ (0) and

 $_{\rm N}$. The relation of these two quantities is based on the LET of B rown et al. [6.18] and takes the following form at the Cheng{D ashen point:

$$_{\rm N} = _{\rm N} (0) + _{\rm N} + R$$

 $_{\rm N} = _{\rm N} (2M^2) _{\rm N} (0)$ (6:18)

 $_{\rm N}$ is the shift due to the scalar form factor of the nucleon, and R is related to a remainder not xed by chiral symmetry. The latter was found to be very small by GSS [6.19], R = 0:4 M eV. In this case, one is dealing with strong S {wave and N interactions. Therefore, the suspicion arises that the one{loop approximation is not su cient to give an accurate description of the scalar form factor (com pare e.g. ref.[6.20]). Therefore, G asser et al. [6.12] have performed a dispersion{theoretical analysis tied together with CHPT constraints for the scalar form factor $_{\rm N}$ (t). The resulting num erical value is

$$_{\rm N} = (15 \quad 0.5) \,{\rm M \, eV}$$
 (6.19)

which is a stunningly large correction to the one{loop result (see below). If one parametrizes the scalar form factor as $_N(t) = 1 + \langle r_S^2 \rangle t + O(t^2)$, this leads to

 $< r_s^2 > = 1:6 \text{ fm}^2$, substantially larger than the typical electrom agnetic size. This means that the scalar operator m (uu + dd) sees a m ore extended pion cloud. Notice that for the pion, a similar enhancem ent of the scalar radius was already observed, $(r_s^2 = r_{em}^2)$ / 1:4 [6.20]. Putting pieces together, one ends up with $_{\rm N}$ (0) = 45 8 M eV [6.12] to be compared with 0 = (1) $y = (35 \quad 5) M eV = (1 \quad y)$. This means that the strange quarks contribute approximately 10 MeV to the N {term and thus the mass shift induced by the strange quarks is $m_s < p_{jss} p > ' (m_s = 2m^\circ) = 10 \text{ MeV} ' 130 \text{ MeV}$. This is consistent with the estimate made in ref. [6.21] based on the heavy mass formalism including the decuplet elds. The e ect of the strange quarks is not dram atic. All speculations starting from the rst order form ula (6.15) should thus be laid at rest. The lesson to be learned here is that m any sm alle ects can add up constructively to explain a seem ingly large discrepancy like N 0 0. W hat is clearly needed are more accurate and reliable low {energy pion {nucleon scattering data to further pin down the uncertainties. We now return to the order q^3 calculation in CHPT. We use an average value $F_p = (F + F_K) = 2'$ 100 M eV, together with F = 0.5 and D = 0.75, which leads to $q_A = 1.25$. The uncertainties in these parameters and how they a ect the results are discussed in [6.10]. The four unknowns, which are the three low energy constants b_D ; b_F and b_0 and the average octet mass (in the chiral limit) m₀ are obtained from a least square t to the physical baryon m asses (N;;;) and the value of $_{\rm N}$ (0) ′ 45 M eV . This allows to predict $\binom{(1)}{KN}$ (0) and $\binom{(2)}{KN}$ (0) and the much discussed matrix element $m_s < p_{jss} p >$, is: the contribution of the strange quarks to the nucleon m ass. Typical results are [6.10]: (a) The strangeness m atrix element in most cases is negative and of reasonable m agnitude of about 200 M eV, (b) within the accuracy of the calculation, the ΚN -tem s tum out to be

$$_{\rm K N}^{(1)}$$
 (0) ' 200 50 M eV ; $_{\rm K N}^{(2)}$ (0) ' 140 40 M eV (6.20)

which is comparable to the rst order perturbation theory analysis having no strange quarks, $\binom{(1)}{KN}(0) = 205 \text{ MeV}$ and $\binom{(2)}{KN}(0) = 63 \text{ MeV}$ [6.22]. These results are indeed most sensitive to the value of N (0). The -term shifts are given by [6.11,6.23]

$$_{\rm N}$$
 (2M²) $_{\rm N}$ (0) = 7:4M eV (6:21)

which is half of the empirical value discussed above, eq.(6.19). Furtherm ore, one nds

$$\begin{pmatrix} (1) \\ K N \end{pmatrix} \begin{pmatrix} (2M \\ K \end{pmatrix} \begin{pmatrix} 2 \end{pmatrix} \begin{pmatrix} (1) \\ K N \end{pmatrix} \begin{pmatrix} (0) \\ K \end{pmatrix} = (271 + i303) M eV$$

$$\begin{pmatrix} (2) \\ K N \end{pmatrix} \begin{pmatrix} (2M \\ K \end{pmatrix} \begin{pmatrix} (2) \\ K \end{pmatrix} \begin{pmatrix} (2) \\ K \end{pmatrix} \begin{pmatrix} (2) \\ K \end{pmatrix} \begin{pmatrix} (0) \\ K \end{pmatrix} = (21 + i303) M eV$$

$$(6.22)$$

The realpart of the rst {term can be estimated simply via Re($_{K N}^{(1)}$ (2M $_{K}^{2}$) $_{K N}^{(1)}$ (0)) ' [$_{N}$ (2M 2) $_{N}$ (0)] (M $_{K}$ = M) 3 = 7:4 46:2 MeV = 340 MeV. The rather small real part in $_{K N}^{(2)}$ stems from the large negative contribution of the {loop which leads

to strong cancellations. Notice the large in aginary parts in $\binom{(j)}{KN} (2M_K^2) = \binom{(j)}{KN} (0)$ due to the two{pion cut. The situation concerning the empirical determinations of the kaon{nucleon {terms is rather uncertain, see e.g. refs.[6.24,6.25]. Since most of the phase shift data stem from kaon{nucleus scattering, it is advantegeous to de ne the KN {terms but means of the pucker isomin

{term s by m eans of the nuclear isospin,

$${}^{0}_{KN} = \frac{1}{4} 3 {}^{(2)}_{KN} + {}^{(1)}_{KN} ; {}^{00}_{KN} = \frac{1}{2} {}^{(2)}_{KN} {}^{(1)}_{KN} :$$
(6:23)

The best available determ inations give ${}^{0}_{K N}$ (0) = 599 377 M eV and ${}^{00}_{K N}$ (0) = 87 66 M eV which translates into ${}^{(1)}_{K N}$ (0) = 469 390 M eV and ${}^{(2)}_{K N}$ (0) = 643 378 M eV.

As has been argued e.g. in refs.[6.3,6.7,6.8], one should account for the spin-3/2 decuplet in the EFT since it leads to a natural cancellation of the large kaon cloud contributions of the type m $_{\rm S}M_{\rm K}^{\ 2}$ ln M $_{\rm K}^{\ 2}$. However, as shown in section 3.4, the inclusion of these elds spoils the power counting much like the nucleon mass term in the relativistic formulation of baryon CHPT. For the case at hand, one also has an in nite renorm alization of the three constants $b_{\rm D}$, $b_{\rm F}$ and $b_{\rm D}$ [6.10],

$$b_{\rm D} = b_{\rm D}^{\rm r}$$
 () $\frac{{\rm C}^2}{2{\rm F}_{\rm p}^2} {\rm L}$; $b_{\rm F} = b_{\rm F}^{\rm r}$ () $+ \frac{5 {\rm C}^2}{12{\rm F}_{\rm p}^2} {\rm L}$; $b_0 = b_0^{\rm r}$ () $+ \frac{7 {\rm C}^2}{6{\rm F}_{\rm p}^2} {\rm L}$: (6:24)

the scale of renorm alization, = 231 M eV the average octet{decuplet m ass with splitting, C the Goldstone{octet{decuplet coupling as discussed after (3.71) and L is de ned in (2.47). The graphs with interm ediate decuplet (states start to contribute at order q⁴ (explicit form ulae can e.g. be found in ref.[6.10]). If one now assumes that these contributions dom inate at this order, one does not nd a consistent description of the scalar sector, as long as one keeps F and D close to their physical values. In {term s turn out to be very sm all and the strange m atrix {elem ent that case, the K N $m_s < p_{jss} p_j > very large$. In ref.[6.7], some additional tadpole diagram s with one insertion from $L_{M B}^{(2)}$ were considered, but that does not alter these conclusions. As stressed before, a complete and system atic q^4 calculation has to be perform ed before one can draw a de nite conclusion on the role of the interm ediate decuplet states. For another critical discussion, see e.g. ref. [6.26]. How ever, there is one curious result in the two{ avor sector we would like to mention [6.10, 6.27]. If one considers the shift of the {term calculated with interm ediate nucleon and (1232) states and uses the large Ν N_c coupling constant relation, one nds

$$(2M^{2}) \qquad _{N} (0) = \frac{3g_{A}^{2}M^{2}}{64^{2}F^{2}} \qquad M + (4) \qquad 4^{p} \frac{p}{2} M^{2} \ln \frac{1}{M} + \frac{1}{M} 1 \\ + \frac{8}{M}^{2} \frac{p}{2Y^{2} + M^{2} 2^{2}} \operatorname{arctan} \frac{p}{2Y^{2} + M^{2} 2^{2}} = 14 \text{ M eV}$$

$$(6.25)$$

very close to the empirical value (6.19) (with = m m = 293 M eV). This means that the graph with the interm ediate (1232) contributes alm ost as much as the low est order diagram with a nucleon as interm ediate state. However, the spectral distribution Im_{N} (t)=t² is much less pronounced around t = 4M² as in the analysis of ref.[6.12] but has a longer tail leading to the same result for the integral. The {contribution m ocks up the higher loop corrections of the dispersive analysis [6.12]. This is sim ilar to the discussion of the spectral function related to the interm ediate range attraction in the nucleon {nucleon interaction (cf. section 5.2). It remains to be seen how the result (6.25) will be a ected when all q⁴ (and possibly higher order) term s will be accounted for. The essential di erence to the baryon m ass calculation and the shifts of the K N

{term s is that the kaon and eta contributions to (6.25) are essentially negligible (they contribute approximately one extra M eV to (6.25)), i.e. we are dealing with an SU (2) statem ent so that one does not expect higher loop diagram s to contribute signi cantly.

VI.2. KAON {NUCLEON SCATTER ING

A topic of current interest is the dynam ics of the kaon {nucleon system based on SU (3) extensions of chiral e ective Lagrangians. Such investigation were in particular triggered by K aplan and N elson [6.28] who proposed a mechanism for kaon condensation in dense nuclear matter using a (how ever incom plete) chiral Lagrangian. B esides this, kaon {nucleon scattering at low energies is of its own interest as a testing ground for threeavor chiral dynam ics. In comparison to the SU (2) sector of pion {nucleon interaction (discussed to some extend in sect.III.5) the kaon {nucleon dynam ics involves several com plications. First, the pertinent expansion parameter for the chiral expansion is much larger, nam ely

$$\frac{M_{K}}{m}$$
 / 0:53 (6:26)

in comparison to = M = m' 0:14 for the N system. Therefore one expects the next-to-leading order corrections to be num erically less suppressed in comparison to the leading term s. The K N system with strangeness S = +1 is physically still quite simple at low energies since it is a purely elastic scattering process with a dom inant S {wave contribution. The analogous K N system with strangeness S =1 how ever greatly di ers, mainly because there are a number of baryons and baryon resonances with 1 but none with S = +1. For the K p reaction there exist inelastic channels dow n S = to threshold involving a pion and a hyperon, $K p ! ^{0} ; ; ^{0}$. Furtherm ore, there is a subthreshold resonance in the K p system, the (1405) of isospin zero. It may be considered as a kind of a K p bound state which can decay into the kinem atically channel and thus receives its physical width. The dynam ical di erences in open K N versus K N naturally show up in the values of the corresponding S-wave scattering lengths [6.29]. The experimental numbers stem from data on kaon {nucleon scattering and K -atom ic level shifts and we give them here without error bars,

$$a^{K^{+}p} = 0:31 \text{ fm}; \qquad a^{K^{+}n} = 0:20 \text{ fm}; \qquad (6:27)$$
$$a^{K^{-}p} = (0:67 + i0:63) \text{ fm}; \qquad a^{K^{-}n} = (+0:37 + i0:57) \text{ fm}:$$

The experimental values for K⁺N are comparable in magnitude to N scattering lengths, a ⁺p = $a_{3=2} = a^+$ a ' 0:14 fm. The negative signs of the a^{K^+N} signal that the K⁺-nucleon interaction is repulsive. Characteristic for the K N scattering lengths is their very large imaginary part, which originates from the open inelastic; channels and the subthreshold (1405)-resonance.

Such inelastic channels are not a problem for CHPT of kaon {nucleon interaction, since they re ect them selves simply as unitarity cuts in the amplitudes which extend below the physical threshold. They are mainly of kinematical origin. On the other hand, the existence of a strong subthreshold resonance (like (1405) in K p) poses a problem to the expansion scheme of chiral perturbation theory, since bound states and (subthreshold) resonances can not be obtained at any nite order of perturbation theory. They require in nitely many orders and are out of the scope of pertubation theory. Consequently, the (1405) will have to be added by hand (com pare the discussion concerning the decuplet states in the EFT in section 3.4). In ref.[6.30] a model has been proposed to generate the (1405) dynamically. For that one solves a Schrodinger equation for the coupled K N; and channels with local or separable meson-baryon potentials linked to an SU (3) chiral Lagrangian. The latter means that the relative strengths of the transition potentials are xed by the Clebsch {G ordan coe cients of a chiral m eson-baryon vertex. A few nite range parameters for the potentials and a coupling stength are then adjusted to the mass and width of the (1405) and several m easured branching ratios. It turns out that such a few parameter t is quite successful in describing the low {energy K N data. W e will not pursue this approach further here, but outline some aspects of KN and KN scattering in CHPT.

The leading order Lagrangian is a straightforward generalization of the chiral N – Lagrangian of sect.III to SU (3) as described in section 6.1. To lowest order, all octet baryon m asses equal. The baryon m ass splittings are due to higher orders in the quark m ass expansion. The corresponding $L_{M B}^{(1)}$ is given in (6.6) for the heavy ferm ion approach. From that, one nds for the S{wave scattering lengths:

$$a^{K^+p} = 1 + \frac{M_K}{m} - \frac{1}{4} \frac{M_K}{F_p^2} = 2a^{K^+n} - 0.6 \text{ fm}$$
 (6.28)

This current algebra result has the correct negative sign and the order of magnitude is reasonable. It is about a factor 2, respectively 1.5, too large for K⁺ p and K⁺ n if we use $F_p = F = 93 \text{ MeV}$. However, there is quite some theoretical uncertainty in this leading order result. A coording to the chiral power counting the prefactor $1=(1 + M_K = m)$ could be neglected and F_p could be taken to be $F_K = 1.22 F$. Such am biguities point towards the importance of higher order calculations, at least up to order q^3 . The corresponding K N scattering lengths have the opposite sign of the K⁺ N ones to leading order, i.e. the chiral K N interaction is attractive. This feature was considered as quite in portant for understanding the dynam ics of the (1405)-resonance form ation in ref.[6.30].

At next-to-leading order, O (q^2) , the SU (3) chiralLagrangian contains a host of new term s. We display here only those which contribute to the S{waves [6.10,6.31]

where the rst three term s obviously coincide with the ones given in eq.(6.7). In ref.[6.32] the last two terms have been forgotten. The complete list of terms at order q^2 and q^3 for avor-SU (3) can be found in ref.[6.2] (for the relativistic approach). There are 13 new parameters for chiral SU (3) in comparison to 3 (c_1 ; c_2 ; c_3) in avor SU (2) and some of the d_i contain 1=m corrections from the expansion of the relativistic leading order Lagrangian form ulated in terms of D irac-elds. The coe cients of the rst three terms in eq.(6.29) (often named "sigma {terms") can be xed at this order from the mass splittings in the baryon octet and the empirical value of the N stress.

$$m = \frac{16}{3} b_{d} (M_{K}^{2} = M^{2}) = 79 M \text{ eV}; \quad b_{d} = 0.066 \text{ G eV}^{-1}$$

$$m = m_{N} = 8 b_{f} (M_{K}^{2} = M^{2}) = 383 M \text{ eV}; \quad b_{f} = 0.213 \text{ G eV}^{-1} \quad (6:30)$$

$$m_{N} (0) = -2M^{2} (b_{d} + b_{f} + 2b_{0}) = 45 M \text{ eV}; \quad b_{0} = -0.517 \text{ G eV}^{-1}$$

O f course such a t is som ew hat problem atic, since one neglects all higher order in the quark m asses, com pare the discussion after eq.(6.15). Consequently, the strangeness content of the proton

$$y = \frac{2(b_0 + b_d - b_d)}{2b_0 + b_d + b_f} / 0.4$$
(6:31)

is about twice the value obtained by G asser, Leutwyler and Sainio [6.12]. If one enforces, however, y ' 0, which is possible due to the uncertainties going into the theoretical analysis of y, one nds $b_0 = 0.279 \text{ GeV}^{-1}$. This corresponds to $_N (0) = 26 \text{ MeV}$, the usual estimate at linear order in the quark masses. The K ⁺ N and K N scattering lengths are now given as [6.31]:

$$a^{K} = 1 + \frac{M_{K}}{m} - \frac{1}{4} \frac{M_{K}}{F_{p}^{2}} - 1 + M_{K} (D_{s} + D_{v})$$

$$a^{K} = 1 + \frac{M_{K}}{m} - \frac{1}{4} \frac{M_{K}}{F_{p}^{2}} - \frac{1}{2} + M_{K} (D_{s} - D_{v})$$
(6:32)

with D s and D v some linear combination of the $b_{d;f;0}$ and $d_{1;:::;10}$. O f course, as long as one has not found a reliable way to estimate all new coe cients, the expressions given

above have not much predictive power. A similar situation occurs for the isospin even

N scattering length a^+ . At this order, the K N scattering lengths are still real, since the scattering into the inelastic channels is a loop e ect which rst shows up at order q^3 .

In ref.[6.32] an order q^3 calculation for the K N scattering lengths has been presented. In this work, how ever, the loop contribution has not been separated cleanly from the counterterm s at the same order. The know ledge of the magnitude of the full loop correction (say at a scale ' 1 G eV) would how ever be very important in order to get a feeling for the genuine size of the (non-analytic) corrections in SU (3). We rem ind here that for the isospin odd N scattering length a the loop correction at order M³ just has the right sign and magnitude to 11 the gap between the W einberg-Tom ozaw a prediction and the empirical value. The K N scattering lengths given in ref.[6.32] are still real at order q^3 . How ever, at this order the rescattering processes K N ! ; ! K N into the inelastic channels are possible and they manifest them selves as com plex valued scattering lengths, cf. g.6.1.



Fig. 6.1: Rescattering diagram which leads to the imaginary part of the scattering lengths at order q^3 for K $\,$ N $\,$! K $\,$ N $\,$.

In the heavy m ass lim it one calculates the following nonzero in aginary parts from the rescattering diagram s

$$\operatorname{Im} a^{K} = 1 + \frac{M_{K}}{m} - \frac{1}{2} \frac{M_{K}^{2} P_{K}}{32 P_{p}^{2}} = \frac{4}{5} \operatorname{Im} a^{K} P_{K} - 0.63 \text{ fm}$$
(6:33)

which are surprisingly close to the empirical values. However, one should not put to much emphasis on these numbers since all mass splittings in the baryon octet have been neglected and this a ects the available phase space. It is interesting to observe that such a simple rescattering calculation tends to explain the near equality of the imaginary parts for proton and neutron K -scattering and at least gives the correct order of magnitude without an explicit (1405).

C learly, all what has been discussed here points towards the importance of more system atic calculations using the complete chiral Lagrangian at a given order. M any of the present controversies in the literature (in particular concering the in-m edium behavior of pions and kaons) stem from the use of incom plete Lagrangians. It is also clear that baryon CHPT for avor SU (3) is just at its beginning and a lot more work (com plete higher order calculations) is necessary in order to judge the quality of such an approach.

VI.3. THE PION IN MATTER

In this chapter we will describe how elective chiral Lagrangians can be used to get inform ation on the modil cation of pionic properties in nuclear matter (so-called medium modil cations). The medium modil cations of hadron properties are relevant for a broad class of problems in nuclear physics. Among these are pion and kaon condensation (in neutron stars), chiral symmetry restoration in relativistic heavy ion collisions, the "dropping" of hadron masses in medium, to name a few. In the following we will only touch upon part of these many issues, namely the density dependence of the quark condensate < uu > and the density dependence of the pion decay constant F and of the pion mass M . We will make use of chiral elective Lagrangian techniques and show that such a method indeed leads to the correct linear terms in density. The latter are often called "low-density theorem s" and can be derived from a multiple scattering expansion. We follow here closely the ideas spelled out in ref.[6.33].

If one rem embers the additional com plications one encounters in each step in extending chiral perturbation theory from the pure meson sector (section 2) to single baryon processes (sections 3 and 4) and further to the B = 2 (here, B denotes the baryon number) sector of nucleon-nucleon interaction and exchange currents (section 5) it is not surprising that a rigorous form ulation of a system atic chiral expansion in nuclearm atter (ite: at nite baryon density) has not yet been found. Finite baryon density introduces a new scale parameter, the Ferm imomentum of the nucleons k_F (sections have to be considered. A rigorous (and still predictive) expansion scheme which can account for all of these many-body com plexities as well as the chiral structure of QCD has not yet been form ulated and may even be too dem anding. A recent approach due to Shankar [6.34] appears prom ising but needs further detailed study. For a general approach to non {relativistic e ective theories, see Leutwyler [6.35].

In a rst step, follow ing ref.[6.33] one can simply use the free space chiralLagrangian for the B = 0 and B = 1 sectors developped so far and evaluate the pertinent nucleon operators at the mean eld level. Consequently, one works to linear order in the nuclear m atter density,

Form ally, such a mean eld approximation means that any local term in the elective N Lagrangian of the form H (x)O (x)H (x) is replaced by $\frac{1}{2}$ Tr [D (x)] . The averaging trace here goes over both spin and isospin coordinates since we consider only a hom ogeneous, isospin symmetric and (spin-)unpolarized nuclear matter distribution of density . Of course, such a mean eld approximation may have no rigorous foundation, but intuitively it should be reasonable at least for low densities. Furthermore, since its starting point

is the most general e ective chiral Lagrangian to a given order the information gained this way is more general than model calculations of the in {medium properties.

Let us now apply the simple mean eld approximation to the elective chiral Lagrangian L⁽²⁾ + L⁽¹⁾_N + L⁽²⁾_N. The averaging procedure going along with the mean eld approximation used to decribe spin and isospin symmetric nuclear matter makes the piece L⁽¹⁾_N vanishing identically since iv $\ell = _{0}\ell$ gives zero and the free term m is absent in the heavy mass formulation. The other terms in L⁽¹⁾_N vanish since the isospin traces Tr ; Tru are zero by construction. So we have to consider only L⁽²⁾_N and here not all terms vanish in the mean eld approximation. These are the ones proportional to $c_1; c_2 = q_1^2 = 8m; c_3$ which are of scalar{isoscalar nature. One obtains the following " nite density chiral Lagrangian" (in the isospin limit m_u = m_d)

L() =
$$\frac{F^2}{4} + \frac{c_3}{2}$$
 Tr(uu) + $\frac{c_2}{2} - \frac{g_A^2}{16m}$ Tr(vuvu) + $\frac{F^2}{4} + c_1$ Tr(+) (6:35)

The term s coming from [(v D^2) D D]=2m have been neglected. They either represent nucleon kinetic energies or as the term s Tr (); Tr (v v) start out at order in the expansion in powers of the pion m ass, which are not of interest here. The form of eq.(6.35) is very illustrative when compared with the free space Lagrangian L⁽²⁾. The two parameters F ' F and B₀ = < uu > =F² have become density dependent through the mean eld approximation of the nucleons. One can immediately read o the corresponding medium modi cations. From the last term in eq.(6.35) we get the density dependence of quark condensate (in the absence of pions + is proportional to quark m ass times quark condensate),

$$\frac{\langle uu \rangle ()}{\langle uu \rangle (0)} = 1 + \frac{4c_1}{F^2} = 1 - \frac{N}{F^2M^2}$$
(6:36)

where we used the leading order relation to the N sigma term $_{\rm N}$ (0) = 4GM 2 $(x \in eq.(3.54))$. The result eq.(6.36) for the linear term of the density dependence of the quark condensate has been derived by several authors using quite di erent m ethods [6.36,6.37] and is often called \low -density theorem ". It was also found in calculations using the Nambu-Jona-Lasinio model, see e.g. ref. [6.38]. It it quite interesting that the simple mean eld approximation to the e ective chiral N Lagrangian very naturally leads to this general result. This gives some con dence in the approximations one is using. Putting in the empirical value of $_{\rm N}$ (0) = 45 M eV one nds a 30 % reduction of the quark condensate at norm alnuclear matter density, giving strong hints that the chiral restoration will take place at about several times nuclear matter density. This is an important issue for relativistic heavy ion collisions, where one hopes to reach such high densities. Of course, in order to make a more quantitative statem ent about the actual chiral restoration density one has to know corrections to eq. (6.36) at higher order in the density . There is a certain similarity to the calculation of the temperature

dependence of the quark condensate in CHPT, it allows for an accurate calculations at low T but can not be used to calculate the critical temperature since then the higher order corrections have to completely cancel the leading term, i.e. one has no longer a controlled expansion [6.39].

The rst and second term in eq.(6.35) take the form of a density dependent pion kinetic term . As nite density breaks Lorentz invariance the time and spatial components of the pionic gradients are treated now dimensity. One sees that at nite density the pion decay constant splits up into a "time component" F_t () and a "spatial component" F_s () which behave dimensity. They are given as

$$F_{t}^{2}() = F^{2} + 2c_{2} + 2c_{3} - \frac{g_{A}^{2}}{4m})$$
 (6:37)
 $F_{s}^{2}() = F^{2} + 2c_{3}$

The phenom enon that the breakdown of Lorentz invariance leads to di erent time and space components of the pion decay constant has also been discussed in some model calculations of pion properties [6.40]. Using the most general elective chiral Lagrangian at order q² and the simplem ean eld approximation to describe density such a behaviour, i.e. $F_t \notin F_s$, follows very naturally from the underlying chiral structure. Therefore one should take care of this possibility in model calculations of the in-medium elects of the pion.

Of particular interest is the density dependence of the pion m ass because of the pions G oldstone boson nature. The inverse pion propagator is

$$D^{-1}(!;q;) = !^{2} q^{2} (!;q;); \qquad (6:38)$$

with the self{energy correction due to the interaction with the medium . Performing an expansion of eq.(6.35) to quadratic order in the pion eld one nds

D¹ (!;q;) = 1 +
$$\frac{2c_3}{F^2}$$
 (!² q²) + $\frac{1}{F^2}$ 2c₂ $\frac{q_A^2}{4m}$!² 1 + $\frac{4c_1}{F^2}$ M²: (6:39)

Evaluating the poles of this propagator one $\ nds$ for the e ective pion m ass M 2 () = $!\,^{2}$ (q = 0;)

$$M^{2}() = M^{2} 1 + \frac{4c_{1}}{F^{2}} 1 + \frac{F^{2}}{F^{2}} 2c_{2} + 2c_{3} \frac{g_{A}^{2}}{4m}^{1}$$

$$= M^{2} 1 \frac{1}{F^{2}} 2c_{2} + 2c_{3} 4q \frac{g_{A}^{2}}{4m}$$

$$= M^{2} T^{+}(M) = M^{2} 4 1 + \frac{M}{m} a^{+}$$
(6:40)

with a^+ the isospin even N scattering length calculated to lowest order (cf. eq.(3.66) without the loop contribution M^{3}). In ref.[6.41] it was emphasized that the linear term in the density dependence of the pion mass is proportional to the isoscalar N scattering length a⁺. This fact is a rigorous result from the leading order of a multiple scattering expansion. The correct coe cient proportional to a + is indeed reproduced here using the complete chiral N Lagrangian at order q^2 and the mean eld approxim ation. The statem ent in ref.[6.41] that the chiral Lagrangian techniques can not give this result is therefore wrong. The argum entation of ref.[6.41] was based on an incom plete chiralLagrangian which consists only of the term proportional to c_1 (the "sigm a-term"). We would like to stress here again that the complete chiral Lagrangian up to a given order has to be used to autom atically produce the correct results m odulo corrections of higher order. Since the value of a^+ ' 0.01 M⁻¹ is negative one nds that the pion m ass slightly increases with density. Of course, this statem ent is based exclusively on the know ledge of the very small linear term in density and could be modied by higher orders, 0 (²) and so on. In the absence of calculations including higher orders in the density one even has no control on the range of validity of the linear density approximation, in: to what fraction of nuclear matter density it is reliable.

As a nalissue let us address the validity of the G ell-M ann-O akes-R enner (G M O R) relation at nite density. It is often assumed or found to hold in model calculations, see e.g. ref.[6.42]. At zero density, the G M O R relation is well founded in chiral perturbation theory and takes the form, $F^{2}M^{2} = m^{2} < uu + dd > + 0 (m^{2})$. An extension to nite densities is not immediately obvious since there is not just a single density dependent pion decay constant and, also, other quantities are density (dependent. The combination of eqs.(6.36,6.37,6.40) yields an in-m edium version of the G M O R relation to linear order in the density , but only if one replaces the free space pion decay constant F by its density dependent time component F_{t} (),

$$F_{+}^{2}()M^{2}() = m < uu + dd > ()$$
 (6:41)

This relation only holds modulo corrections which are of higher order in the light quark m asses and the density. It is important to note that for the spatial component F_s () the in-medium GMOR relation does not hold. In ref.[6.33] functional methods have been used to show that the in-medium properties of the pion up to linear order in the density do not depend on the actual choice of the interpolating pion eld. This feature is of course quite important in order to make the concept of density dependent mass, decay constant and so on meaningful at all. The independence from the interpolating

eld becom es also clear if one goes back to eq.(6.35), the "density dependent chiral Lagrangian". A ny param etrization of the chiral matrix U(x) in terms of som e pion eld (exponential, -m odel gauge, stereographic coordinates, :::) gives the sam e result for the expansion truncated at the quadratic order and this is all one needs to read o the density dependent pion m ass and decay constant.

The salient features from this study of the pion in nuclearm atter can be sum m arized as follows. Using the electric chiral Lagrangian up to order q^2 and a simple mean eld approximation to describe the nuclear density, one can easily reproduce the so-called \bw-density theorem s" for the density variation of the quark condensate and the pion mass which follow from a multiple scattering expansion. The pion decay constant F splits up into a time component $F_t()$ and a spatial component $F_s()$, which do not have the same density dependence, nevertheless both actually decrease with density. The corresponding linear coe cients $2c_2 + 2c_3$ $q_1^2 = 4m$ and $2c_3$ are negative. The results presented here could be obtained rather easily from lowest order tree level chiral Lagrangian $L_{N}^{(1;2)}$, but is seem s rather nontrivial to go to higher orders. For example at order q³ loop diagram s give rise to non-local term s of the form $d^4vH(x)O(x;v)H(v)$ and it is not clear how to handle them in mean eld approximation. Furtherm ore, all four-nucleon terms showing up in the B = 2 sector should be considered, since they will give information on ² correction and eventually nuclear correlations. Presently a system atic scheme to account for all these complications is unknown and may only be feasible if one supplies phenom enological inform ation as discussed in section 5.

VI.4. M ISCELLANEOUS OM ISSIONS

In this section, we want to give a list of topics not covered in detail. This list is neither meant to be complete or does the order imply any relevance. The references should allow the interested reader to further study these topics.

Isospin violation: A lthough the down quark is almost twice as heavy as the up quark, the corresponding isospin violations are perfectly masked in almost all observables since (m d m₁₁)= 1. All purely pionic low {energy processes autom atically conserve isospin to order $m_d = m_u$ besides from true electrom agnetic e ects. The reason is that G {parity forbids a term of the type uu dd using only pion elds with no derivatives. Therefore, W einberg [6.43] considered isospin violating e ects in the scattering lengths of neutral pions o nucleons. A spointed out by W einberg [6.43] and later quanti ed by Bernard et al.[6.44], the absolute values of a $\begin{pmatrix} 0 n \\ \end{pmatrix}$ and of a $\begin{pmatrix} 0 p \\ \end{pmatrix}$ are hard to pin down accurately. However, in the di erence m any of the uncertainties cancel and one expects a sizeable isospin violating e ect of the order of 30% [6.43]. In view of this, Bernstein [6.45] has proposed a second (generation experiment to accurately measure the phase of the reaction p! ⁰p below ⁺ n threshold and use the three{channel unitarity to deduce the tiny ⁰p phase. M ore recent discussions of these topics are due to van Kolck [6.46] and W einberg [6.47].

Baryon octet and decuplet properties: The high energy hyperon beams at CERN and Ferm ilab allow to study aspects of the electrom agnetic structure of these particles. For example, one can make use of the Primako e ect to measure the polarizabilities. In the quark model, one expects + since in a system with like{sign charges like the (dds) dipole excitations are strongly suppressed

[6.48]. This was quantized in a CHPT calculation to order q³ in ref.[6.49]. To that + **′** 1:5 . Further studies of hyperon radiative decays accuracy, one expects and an analysis of the octet m agnetic m om ents can be found in refs. [6.50, 6.51, 6.52]. In the EFT with the spin $\frac{3}{2}$ decuplet as active degrees of freedom, one can also address the properties of the decuplet states. Topics considered include the E 2=M 1 m ixing ratio in the decay ! N [6.53], the decay ! N ' + ' (where ' denotes a lepton) [6.54] or the strong and electrom agnetic decays of the decuplet states [6.55]. Furtherm ore, in the large N_c lim it, one needs the spin $\{3/2 \text{ states to restore}\}$ unitarity in N scattering (see [6.56] and references therein). This is often used as a strong support for the inclusion of the decuplet states in the EFT. However, we would like to stress that since the chiral and the large N $_{\rm c}$ lim ites do not com mute, considerable care has to be taken when such aroum ents are employed, see e.g. refs.[6.57,6.58,6.59]).

K aon and pion condensation: The work of K aplan and N elson [6.28] triggered a urry of papers addressing the question whether B ose condensates of charged m esons m ay be found in dense nuclear m atter form ed e.g. in the cores of neutron stars, the collapse of stars or in the collision of heavy ions. The physical picture behind this is the attractive S {wave kaon {nucleon interaction which could lower the e ective m ass of kaons to the extent that the kaons condense at a few times the nuclear m atter density. This question, its consequences for the nuclear equation of state, neutron stars and the related question of S {wave pion condensation are addressed e.g. in refs.[6.31,6.32,6.33,6.41,6.60,6.61,6.62,6.63,6.64] (and references given therein).

Finally, let us m ention that a state of the art update can be found in the workshop proceedings [6.65] from which m any m ore references can be traced back.

REFERENCES

- 6.1 J.G asser and H. Leutwyler, Nucl. Phys. B 250 (1985) 465.
- 6.2 A.Krause, Helv. Phys. Acta 63 (1990) 3.
- 6.3 E.Jenkins and A.V.M anohar, in "E ective eld theories of the standard model", ed.Ulf(G.M ei ner, W orld Scientic, Singapore, 1992.
- 6.4 J.Gasser, Ann. Phys. (N.Y.) 136 (1981) 62.
- 6.5 J.Gasser and H. Leutwyler, Phys. Reports C 87 (1982) 77.
- 6.6 P.Langacker and H.Pagels, Phys. Rev. D 8 (1971) 4595.
- 6.7 E. Jenkins, Nucl. Phys. B 368 (1992) 190.
- 6.8 E. Jenkins and A.V. Manohar, Phys. Lett. B 259 (1991) 353.
- 6.9 R F. Lebed and M A. Luty, Phys. Lett. B 329 (1994) 479.

- 6.10 V.Bernard, N.Kaiser and UlfG.Meiner, Z.Phys.C 60 (1993) 111.
- 6.11 H. Pagels and W. Pardee, Phys. Rev. D 4 (1971) 3225.
- 6.12 J.Gasser, H.Leutwyler and M.E.Sainio, Phys. Lett. 253B (1991) 252, 260.
- 6.13 T P. Cheng and R. Dashen, Phys. Rev. Lett. 26 (1971) 594.
- 6.14 G.Hohler, in Landolt (Bornstein, vol.9 b2, ed. H.Schopper (Springer, Berlin, 1983).
- 6.15 J.G asser, in "Hadrons and Hadronic Matter", eds. D.Vautherin et al., Plenum Press, New York, 1990.
- 6.16 Ulf-G.Meiner, Int. J.Mod. Phys. E1 (1992) 561.
- 6.17 R.Koch, Z.Phys. C 15 (1982) 161.
- 6.18 L.S.Brown, W.J.Pardee and R.D.Peccei, Phys. Rev. D 4 (1971) 2801.
- 6.19 J.Gasser, M E. Sainio and A. Svarc, Nucl. Phys. B 307 (1988) 779.
- 6.20 J.Gasser and Ulf-G.Meiner, Nucl. Phys. B 357 (1991) 90.
- 6.21 E. Jenkins and A.V. Manohar, Phys. Lett. B 281 (1992) 336.
- 6.22 R L. Ja e and C. Korpa, Comm. Nucl. Part. Phys. 17 (1987) 163.
- 6.23 V.Bernard, N.Kaiser, J.Kambor and Ulf-G.Meiner, Nucl. Phys. B 388 (1992) 315.
- 624 P.M. Gensini, J. Phys. G: Nucl. Part. Phys. 7 (1981) 1177.
- 625 PM.Gensini, in N Newsletter no. 6, eds. R.E.Cutkowsky, G.Hohler, W.Kluge and BMK.Nefkens, April 1992.
- 6.26 M A.Luty and A.W hite, Phys. Lett. B 319 (1993) 261.
- 627 I. Jam eson, A W . Thom as and G . Chanfray, J. Phys. G: Nucl. Part. Phys. 18 (1992) L159.
- 6.28 D.B.Kaplan and A.E.Nelson, Phys. Lett. B175 (1986) 57; B192 (1987) 193.
- 6.29 O.Dum brais et al, Nucl. Phys. B 216 (1982) 277.
- 6.30 P.B. Siegeland W. Weise, Phys. Rev. C 38 (1988) 2221.
- 6.31 G E.Brown, C.H.Lee, M. Rho and V. Thorsson, Nucl. Phys. A 567 (1993) 937.
- 6.32 C.H.Lee, H.Jong, D.P.M in and M.Rho, Phys. Lett. B 326 (1994) 14.
- 6.33 V. Thorsson and A.W irzba, \S {wave M eson {N ucleon Interactions and the M eson M ass in N uclear M atter from E ective C hiral Lagrangians", N O R D ITA preprint 1995, in preparation..
- 6.34 R. Shankar, Rev. M od. Phys. 66 (1994) 124.
- 6.35 H. Leutwyler, Phys. Rev. D 49 (1994) 3033.
- 6.36 E.G. D rukarev and E.M. Levin, Nucl. Phys. A 511 (1988) 697.
- 6.37 T.D.Cohen, P.J.Furnstuhland D.K.Griegel, Phys. Rev. C 45 (1992) 1881.
- 6.38 M.Lutz, S.K limt and W.Weise, Nucl. Phys. A 542 (1992) 521.

- 6.39 P.G erber and H. Leutwyler, Nucl. Phys. B 321 (1989) 387.
- 6.40 A. Le Yaouanc, L. Oliver, S. Ono, O. Pene and J.-C. Raynal, Phys. Rev. D 31 (1985) 137;
 - V.Bemard, Phys. Rev. D 34 (1986) 1601.
- 6.41 J.Delome, M. Ericson and T.E.O. Ericson, Phys. Lett. B 291 (1992) 379.
- 6.42 V.Bemard and Ulf(G.Meiner, Nucl. Phys. A 489 (1988) 647.
- 6.43 S.W einberg, Trans. N.Y. Acad. Sci. 38 (1977) 185.
- 6.44 V.Bernard, N.Kaiser and Ulf{G.Meiner, Phys. Lett. B 309 (1993) 421.
- 6.45 A.M. Bernstein, N Newsletter 9 (1993) 55.
- 6.46 U. van Kolck, Thesis, University of Texas at Austin, 1992.
- 6.47 S.W einberg, \Strong Interactions at Low Energies", preprint UTTG-16-94, 1994.
- 6.48 H J. Lipkin and M A . M oinester, Phys. Lett. B 287 (1992) 179.
- 6.49 V.Bernard, N.Kaiser, J.Kambor and Ulf-G.Meiner, Phys. Rev. D 46 (1992) 2756.
- 6.50 E. Jenkins et al., Nucl. Phys. B 397 (1993) 84.
- 6.51 H.Neufeld, Nucl. Phys. B 402 (1993) 166.
- 6.52 E. Jenkins et al, Phys. Lett. B 302 (1993) 482.
- 6.53 M N.Butler, M J.Savage and R P.Springer, Phys. Lett. B 304 (1993) 353.
- 6.54 M N.Butler, M J.Savage and R P.Springer, Phys. Lett. B 312 (1993) 486.
- 6.55 M N.Butler, M J.Savage and R P.Springer, Nucl. Phys. B 399 (1993) 69; Phys. Rev. D 49 (1994) 3459.
- 6.56 R.Dashen, E.Jenkins and A.V.Manohar, Phys. Rev. D 49 (1994) 4713.
- 6.57 J.Gasser and A.Zepeda, Nucl. Phys. B 174 (1980) 45.
- 6.58 E.Jenkins and A.V.M anohar, in \E ective Field Theories of the Standard M odel", ed.Ulf{G.M einer (W orld Scientic, Singapore, 1992).
- 6.59 N.Kaiser, in \Baryons as Skyrm e Solitons", ed. G.Holzwarth (World Scientic, Singapore, 1993).
- 6.60 H D. Politzer and M. B. W ise, Phys. Lett. B 273 (1991) 156.
- 6.61 G E.Brown, K.Kubodera, M.Rho and V.Thorsson, Phys. Lett. B 291 (1992) 355.
- 6.62 V.Thorsson, M.Prakash and J.M.Lattim er, Nucl. Phys. A 572 (1994) 693; Nucl. Phys. A 574 (1994) 851,
- 6.63 C.H. Lee, G.E. Brown and M. Rho, \Kaon condesation in nuclear starm atter", SeculUniversity preprint SNUTP-94-28, hep-ph/9403339, 1994.
- 6.64 H.Yabu, F.M yhrer and K.Kubodera, Phys. Rev. D 50 (1994) 3549 (and references therein).
- 6.65 Proceedings of the workshop on \ChiralD ynam ics: Theory and Experim ent", B R. Holstein and A M. Bernstein (eds), Springer Lecture Notes in Physics, 1995, in print.

APPENDIX A: FEYNMAN-RULES

Here, we wish to collect the pertinent Feynm an rules which are needed to calculate tree and loop diagrams. In order to praram etrize the SU (2) matrix U in terms of pion elds we use the so-called sigm a gauge $U = \begin{bmatrix} 1 & -2 = F^2 + i \\ 1 & -2 = F^2 + i \\ - = F$ which is more convenient than the exponential param etrization $U = \exp[i \\ - = F]$. For elective vertices involving 3 and more pions, the Feynm an rules di er in the two param etrizations. Of course the complete S-matrix for a process with a certain number of on-shell external pions is the same in either param etrization. The param etrization dependence of matrix elements for o -shell pions signals that these are indeed non-unique in CHPT. Physically this is clear, since in order to calculate eg: o -shell pion amplitudes, one has to know the exact pion eld of QCD, not just some interpolating eld.

W e use the following notation:

- 1 M om entum of a pion or nucleon propagator.
- k M om entum of an external vector or axial source.
- q M om entum of an external pion.

Photon polarization vector.

- A Polarization vector of an axial source.
- p M om entum of a nucleon in heavy m ass form ulation.

and pion isopin indices are a;b;c;d;3.Furtherm ore, v is the nucleon four-velocity and S its covariant spin {vector. A llparam eters like $Q = F;g_A;m;:::$ are m eant to be taken in the chiral lim it. W e also give the orientation of m om enta at the vertices, i.e. which are "in"-going or "out"-going.

Vertices from L⁽²⁾

pion propagator:

$$\frac{i^{ab}}{l^2 M^2 + i0}$$
 (A:1)

1 pion, pseudoscalar source:

3 pions, pseudoscalar source:

0 (A :3)

1 pion, axial source (k in):

$$F \stackrel{ab}{}_{A} k$$
 (A:4)

3 pions, axial source (all q's out):

$$\frac{1}{F} = \int_{A}^{bc \ de} (q_2 + q_3 \ q_1) + \int_{bd \ ce}^{bd \ ce} (q_1 + q_3 \ q_1) + \int_{be \ cd}^{bc \ cd} (q_1 + q_2 \ q_1)] \quad (A:5)$$

2 pions, photon $(q_1 \text{ in}, q_2 \text{ out})$:

 e^{a3b} $_{1}(q_{1} q_{2})$ (A:6)

4 pions, photon:

(A :7)

- 2 pions, 2 photons:
- 2ie^{2 0} ^{ab} ^{a3 b3}) (A:8)
- 2 pions, scalar source:

- i^{ab}M² (A:9)
- 4 pion vertex (all q's in):

$$\frac{i}{F^2} f^{ab cd} [(q_1 + q_2)^2 M^2] + {}^{ac bd} [(q_1 + q_3)^2 M^2] + {}^{ad bc} [(q_1 + q_4)^2 M^2] g (A : 10)$$

0

- Vertices from L $_{\rm N}^{\rm (1)}$
- nucleon propagator:

$$\frac{1}{v + i0}$$
 (A:11)

1 pion (q out):

$$\frac{q_{A}}{F}S \quad q^{a} \tag{A.12}$$

photon:

$$\frac{10}{2}(1+3)$$
 v (A.13)

2 pions
$$(q_1 \text{ in}, q_2 \text{ out})$$
:

$$\frac{1}{4F^2} v (q+q_2)^{abc c}$$
 (A:14)

1 pion, 1 photon:

$$\frac{ieg_{A}}{F} \quad \hat{S}^{3b \ b} \qquad (A:15)$$

3 pions (allq's out):

$$\frac{g_{A}}{2F^{3}} \begin{bmatrix} a & bc \\ S & (q+q_{3}) + b & ac \\ S & (q+q_{3}) + c & ab \\ S & (q+q_{2}) \end{bmatrix}$$
 (A:16)

2 pions, photon:

$$\frac{ie}{4F^2} \left({}^{a \ b3} + {}^{b \ a3} \ 2^{3 \ ab} \right) \quad v \tag{A:17}$$
3 pions, photon:

(A :18)

axial source:

$$ig_A S_A$$
^b (A :19)

1 pion, axial source:

$$\frac{i}{2F} A \vec{v}^{bc c} \qquad (A : 20)$$

2 pions, axial source:

$$\frac{ig_A}{2F^2}S_A \left({}^{ab\ c} + {}^{bc\ a} 2^{ac\ b} \right)$$
 (A :21)

Vertices from $L_{N}^{(2)}$

$$L_{N}^{(2)} = H \frac{1}{2m} (v D^{2}) \frac{1}{2m} D D \frac{ig_{A}}{2m} fS D; v ug_{f}Tc_{+}$$

$$+ (c_{2} \frac{g_{A}^{2}}{8m}) (v \hat{u}) + c_{3}u u + 4c + \frac{1}{4m}) [S; S] u u \qquad (A.22)$$

$$+ c_{5} Tr(\sim_{+}) \frac{i}{4m} [S; S] (1 + v)f^{+} + \frac{1}{2} (s v) Tr(f^{+}) H$$

0

with $\sim_{+} = +$ (1=2) Tr + (this term is only non {vanishing form $_{u} \notin m_{d}$). All parameters, g_{A} ; m; c_{i} ; $_{s;v}$ are understood as the ones in the chiral lim it. $f^{+} = u^{Y}F^{R}u + uF^{L}u^{Y}$ with $F^{L,R}$ the eld strength tensor corresponding to external (isovector) left/right vector sources (isovector photon, W and Z boson). The external vector source in f^{+} is understood to have also an isoscalar component (isoscalar photon). Here, we will use the (C oulom b) gauge v = 0 for the photon. The rst three terms in $\frac{f^{2}}{N}$ stem from the 1=m expansion of the chiral nucleon D irac Lagrangian. They have no counter part in the relativistic theory, is: no bilinears form involving {matrices. Their coe cients are

xed in terms of the lowest order parameters and nucleon mass m. The other terms involving the new low (energy constants come from the most general relativistic lagrangian at order q^2 (see ref.[3.6]) after translation into the heavy mass formalism. One observes, that there is some overlap between the two types of terms at order q^2 , namely in the $(v \quad u^2)$ and [S; S]u u terms and the magnetic moment couplings. The low (energy constants c_6 and c_7 which are discussed in sections three and four are related to the anom alous magnetic moments of the nucleon (in the chiral limit) via

$$c_6 = v; c_7 = \frac{1}{2}(s v) :$$
 (A 23)

The pertinent Feynm an insertions read (p_1 is always ingoing and p_2 outgoing):

nucleon propagator:

$$\frac{1}{2m} = 1 - \frac{1^2}{(v + i\theta)}$$
 (A.24)

2 photons:

$$\frac{ie^2}{2m}$$
 (1 + ³) ⁰ (A :25)

1 pion (q out):

$$\frac{q_{\rm A}}{2m \, \rm F} S \quad (p+p_2)v \quad q^2 \qquad (A.26)$$

1 photon (k in):

$$\frac{ie}{4m}(1 + {}^{3}) \quad _{1}(p + p_{2}) + \frac{ie}{2m}[S ; S k](1 + (1 + v))^{3}) \qquad (A.27)$$

1 pion, 1 photon (q out):

$$\frac{eg_{A}}{2m F}S \quad v \quad {}^{a}q + {}^{a3})$$
 (A :28)

2 pions (q in, q out):

$$\frac{i^{ab}}{F^{2}} = 4q 1M^{2} + (2c_{2} \frac{q_{A}^{2}}{4m})v_{1}q_{2}q_{2}+ 2c_{3}q_{1} 2q_{4}$$

$$+ \frac{1}{8m F^{2}} a^{bc} c(p_{1} + p_{2}) (q_{1} + q_{2}) v_{1}(q_{2} + p_{2})v (q_{1} + q_{2}) \qquad (A 29)$$

$$\frac{1}{F^{2}} 2c_{4} + \frac{1}{2m} a^{bc} c[S_{1}q_{1}S_{2}q_{4}]$$

3 pions (allq's out):

$$\frac{ig_{A}}{4m F^{3}} \stackrel{abc}{=} [v_{1}G_{A} (q_{1}) + v_{2}G_{A} (q_{1}) + v_{3}G_{A} (q_{2})]$$

$$\frac{g_{A}}{4m F^{3}}S (p+p_{2})[\stackrel{abc}{=} v_{2}(q+q_{3}) + \stackrel{bac}{=} v_{2}(q+q_{3}) + \stackrel{cab}{=} v_{2}(q+q_{2})]$$
(A:30)

2 pions, 1 photon $(q_1 \text{ in}, q_2 \text{ out})$:

$$\frac{ie}{2m F^{2}} 2(1 + v) [S; S k] + 1 + pp) (a b^{3} + b a^{3} 2^{3} a^{b}) + \frac{e}{8m F^{2}} 1(q q_{2})^{abc} (c + 3c) + 2c_{3} \frac{e}{F^{2}} a^{3b} 1(q q_{2})$$
(A:31)

3 pions, 1 photon (all q's out):

$$\frac{eq_{A}}{4m F^{3}}S = 2^{a3} b^{c}v \quad (q+q_{B} q_{I}) + 2^{b3} a^{c}v \quad (q+q_{B} q_{I}) + 2^{c3} a^{b}v \quad (q+q_{L} q_{I}) + a^{b}c v \quad (q+q_{B}) + b^{b}a^{c}v \quad (q+q_{B}) + c^{c}a^{b}v \quad (q+q_{L})$$

$$(A:32)$$

2 pions, 2 photons:

$$\frac{ie^2}{4m F^2} \circ e^{a[b^3 + b^{a^3} + 2^{a^3 b^3} 2^{ab}(1 + 3)] + 4ic_3 \frac{e^2}{F^2} \circ e^{ab}(1 + 3) = 4ic_3 \frac{e^2}{F^2} + 4ic_3 \frac{e^2}{F^2} +$$

axial source:

$$\frac{i g_A}{2m F} S (p+p_2)_A v^b$$
 (A:34)

1 pion, axial source (q out):

$$\frac{i}{4m F} \stackrel{abc c}{=} [A \quad (p+p_2) \quad A \quad vv \quad 1 \not (p p_2) + 2\frac{c_3}{F} \stackrel{ab}{=} A \quad q \\
+ \frac{1}{F} (2c_2 \quad \frac{q_A^2}{4m}) \stackrel{ab}{=} A \quad vv \quad q \stackrel{i}{+} \frac{i}{2m F} (1 + v) [S \quad A; S \quad K]^{bc \ c} \qquad (A:35) \\
+ i \quad 2c_4 + \frac{1}{2m} \quad [S \quad q; S \quad A] \stackrel{abc \ c}{=}$$

APPENDIX B:LOOP FUNCTIONS

Here, we will de ne many of the loop functions which frequently occur in our calculations and we will give these functions in closed analytical form as far as possible. D ivergent loop functions are regularized via dimensional regularization and expanded around d = 4 space-time dimensions. In the following all propagators are understood to have an in nitesimal negative imaginary part.

$$\frac{1}{i} \frac{d^{d} l}{(2)^{d}} \frac{1}{M^{2} \frac{1}{2}} = M^{d^{2}} (4)^{d=2} (1 \frac{d}{2})$$
(B:1)

$$= 2M^{2} L + \frac{1}{16^{2}} \ln \frac{M}{M} + O (d 4)$$
 (B 2)

with

_

$$L = \frac{d^{4}}{16^{2}} \frac{1}{d^{4}} + \frac{1}{2} (E 1 \ln 4)$$
 (B.3)

containing a pole in d = 4 and $_{\rm E}$ = 0:557215:... The scale is introduced in dimensional regularization.

$$\frac{1}{i} \frac{d^{d}l}{(2)^{d}} \frac{f_{1;l}; l l g}{(v l !)(A^{1} l)} = f_{J_{0}}(!); v J_{1}(!); g J_{2}(!) + v v J_{3}(!)g \quad (B : 4)$$

$$J_0(!) = 4L! + \frac{!}{8^2} 1 2 \ln \frac{M}{4^2} \frac{1}{4^2} \frac{P}{M^2} \frac{1}{M^2} \frac{P}{M^2} + O(d 4)$$
(B:5)

$$J_1(!) = ! J_0(!) + ; J_2(!) = \frac{1}{d - 1} (M^2 - !^2) J_0(!) !$$
 (B:6)

$$J_3 (!) = ! J_1 (!) \quad J_2 (!)$$
 (B :7)

$$\frac{1}{i} \frac{d^{d}l}{(2)^{d}} \frac{fl;l;llg}{v l(v l !)^{2}(M !)} = f^{*}_{0}(!);v^{*}_{1}(!);g^{*}_{2}(!) + vv^{*}_{3}(!)g^{*}(B :8)$$

U sing the identity

$$\frac{1}{v \quad l(v \quad l \quad = \frac{1}{!})!} = \frac{1}{v \quad l \quad v \quad l \quad = \frac{1}{v \quad l}} \frac{1}{v \quad l}$$
(B:9)

$$\sim_{i}(!) = \frac{1}{!} J_{i}(!) J_{i}(0) ; (i = 0;1;2;3)$$
 (B:10)

$$\frac{1}{i} \frac{d^{d}l}{(2)^{d}} \frac{f_{1}l_{1}l_{2}l_{3}l_{4}}{(v l l^{2} l_{1})^{2}} = f_{G_{0}}(!); v G_{1}(!); g G_{2}(!) + v v G_{3}(!)g (B:11)$$

U sing the identity

$$\frac{1}{(v \ l \ l^{2})} = \frac{0}{0!} \frac{1}{v \ l}$$
(B :12)

$$G_{i}(!) = \frac{0}{0!} J_{i}(!); \quad (i = 0;1;2;3)$$
 (B:13)

$$\frac{1}{i} \frac{d^{d}l}{(2)^{d}} \frac{f_{1}l_{1}l_{2}l_{3}l_{4}}{v_{1}(1+k)^{2}} = f_{0}(!)_{k}k_{1}(!) + v_{2}(!)_{k}$$

$$g_{3}(!) + k k_{4}(!) + (k v + k v)_{5}(!) + v v_{6}(!)g$$
(B:14)

where $! = v + k_i^2 k = 0$ since we consider only real photons.

$$_{0}(!) = \frac{1}{16^{2}!} + \arcsin\frac{!}{M} \arcsin\frac{!}{M}$$
 (B:15)

The vector and tensor functions $_{j}(!); j = 1;2;3;4;5;6$ can be obtained by the following procedure. One multiplies the dening equation with v thereby canceling a factor v l or one multiplies by 2k and uses the identity $2k \quad l = (M^{2} \quad \hat{t}) \quad (M^{2} \quad (l+k)^{2}))$ for $k^{2} = 0$. This leads to linear relation among the $_{j}(!); j \in 0$ where the right hand sides

are loop functions with fewer propagators. For illustration of the m ethod, we give the explicit solution for the functions $_{j}(!)$:

$$_{1}(!) = \frac{1}{8^{2}!^{2}} p \frac{1}{M^{2}} !^{2} \arccos \frac{!}{M} \frac{1}{2} M !$$
 (B :16)

$$_{2}(!) = 2L + \frac{1}{16^{2}} + \frac{1}{2\ln M} + \frac{1}{8^{2}!} - \frac{p}{2}M = \frac{p}{M^{2}!} - \frac{1}{2} \operatorname{arccos} \frac{1}{M}$$
 (B:17)

$${}_{3}(!) = L! + \frac{!}{16^{2}} \ln \frac{M}{1} + \frac{1}{16^{2}} \ln \frac{P}{M^{2}} \frac{!}{!^{2}} \arccos \frac{!}{M} + \frac{M^{2}}{32^{2}!} + \arcsin \frac{!}{M} \arcsin \frac{!}{M}$$
(B:18)

$$_{4}(!) = \frac{1}{32^{2}!^{3}} M^{2} + \arcsin \frac{!}{M} \arcsin \frac{!}{M} 2!^{p} \frac{!}{M^{2}!^{2}} \arccos \frac{!}{M} + !^{2} (B : 19)$$

$$_{5}(!) = L + \frac{1}{16^{2}} \ln \frac{M}{M} + \frac{p \frac{M^{2} !^{2}}{M^{2} !^{2}} \operatorname{arccos} \frac{!}{M} \frac{M^{2}}{32^{2} !^{2}} + \operatorname{arcsin} \frac{!}{M} \operatorname{arcsin} \frac{!}{M}$$
(B 20)

(L) = 2L + \frac{1}{16^{2}} \ln \frac{M}{M} + \frac{p \frac{M^{2} !^{2}}{16^{2} !} \operatorname{arccos} \frac{!}{M} \frac{M^{2}}{32^{2} !^{2} !^{2}} + \operatorname{arcsin} \frac{!}{M} \operatorname{arcsin} \frac{!}{M}
(B 20)

$$_{6}(!) = 2L! + \frac{!}{16^{2}} - 1 - 2\ln\frac{M}{2} - \frac{1}{8^{2}} + \frac{1}{M^{2}} - \frac{1}{2} \arccos\frac{!}{M}$$
 (B :21)

From now on we give only the scalar loop functions for d = 4.

$$\frac{1}{1} \frac{d^{4}l}{(2)^{4}} \frac{1}{v l(v l !)^{2} (M^{2} (l k^{3}))} = _{0} (!)$$
 (B :22)

U sing the sam e identity as for $\widetilde{\ }_{i}\left(!\right)$

$$_{0}(!) = \frac{1}{!} _{0}(!) _{0}(!) = \frac{1}{8^{2}!^{2}} \arcsin^{2} \frac{!}{M}$$
 (B.23)

$$\frac{1}{i} \frac{d^{4}l}{(2)^{4}} \frac{1}{v \quad l(M \quad \tilde{f})^{2} (M \quad 2)} = _{0} (!)$$
(B:24)

$${}_{0}(!) = \frac{1}{32 \text{ M}^{2}!^{2}} \text{ M} \qquad {}^{p} \frac{1}{M^{2} !^{2}} + \frac{1}{16 {}^{2}M^{2}!^{2}} ! \qquad {}^{p} \frac{1}{M^{2} !^{2}} \operatorname{arcsin} \frac{!}{M} \quad (B : 25)$$

$$\frac{1}{i} \frac{d^4 l}{(2)^4} \frac{1}{(v - \frac{3}{4})(M^2 - \frac{3}{4})(M^2 - (1 + k)^2)} = _0 (!)$$
 (B.26)

$$_{0}(!) = \frac{1}{8^{2}!} (M^{2} !^{2})^{1=2} \arccos \frac{!}{M} \frac{!}{2M}$$
 (B.27)

$$\frac{1}{i} \frac{d^{4}l}{(2)^{4}} \frac{1}{(v - \frac{3}{2})(M^{2} - \frac{3}{2})^{2}(M^{2} - (1 + k)^{2})} = _{0}(!)$$
 (B.28)

$$_{0}(!) = \frac{1}{16^{2} M^{2}!} (M^{2} !^{2})^{1=2} \arccos \frac{!}{M} \frac{!}{2M} (B : 29)$$

For the calculation of form factor we need i with v = 0 but $k = t \in 0$.

$$\frac{1}{i} \frac{d^{d}l}{(2)^{d}} \frac{1;ll}{v \quad l(M^{2} \quad (l+k)^{2})} = _{0}(t);g \quad _{3}(t) + ::: \qquad (B:30)$$

$$_{0}$$
 (t) = $\frac{1}{8} = \frac{1}{t}$ arctan $\frac{p}{2M}$ (B :31)

$$_{3}(t) = \frac{1}{32}M + \frac{1}{2}\frac{2M^{2}}{t}p_{--}\frac{p_{--}}{tarctan}\frac{t}{2M}$$
 (B:32)

The two loop function which enter the nucleon isovector D irac from factor are:

$$J(t) = \frac{1}{16^2} \frac{t}{6} + \int_{0}^{2} dx M^2 + tx (x - 1) \ln 1 + \frac{t}{M^2} x (x - 1)$$
(B:33)

(t) =
$$\frac{1}{16^2} \int_{0}^{2} dx \ln 1 + \frac{t}{M^2} x (x - 1)$$
 (B:34)

APPENDIX C: THE "AXIAL RADIUS DISCREPANCY"

In the end of sect.IV 4 we discussed pion electroproduction at threshold and found that chiral loops modify the LET of N am bu, Lurie and Shrauner. The conclusion was that an analysis of threshold charged pion electroproduction data in terms of soft pion theory (in order to link the measured cross sections to nucleon electrom agnetic and axial form factors) does not lead to the nucleon mean square axial radius r_A^2 itself but to a modi ed quantity r_A^2 which subsumes the chiral loop corrections. For this "discrepancy", $r_A^2 = r_A^2$, the leading term which survives in the chiral lim it is given in eq.(4.79). Num erically it is a {10% e ect and allows to understand the system atic discrepancies between present (anti)neutrino experiments (which measure the true nucleon axial radius) and charged pion electroproduction experiments [4.76]. (N ote that we are considering here exclusively small values of the momentum transfer k²).

Since the lowest order result of the discrepancy $r_A^2 = r_A^2$ is quite small, one should investigate higher order corrections (in = M = m) in order to see whether the num erical value of the leading order prediction is actually reliable. Such a complete next-to-leading order calculation has been done in ref.[4.91]. For that one has to go to order q^4 in the chiral expansion, which am ounts to an evaluation of all one loop graphs with

a single insertion from $L_{N}^{(2)}$ and possible counter terms. The latter were estimated from resonance exchange contributions, in the present case from (770) and (1232) exchange.

The relevant observable to be studied is the transition matrix element for p! ⁺ n in the center of mass frame at threshold. Only the transverse part is of interest and it takes the form

$$\Gamma \qquad \frac{ieg_{A}}{2F} \sim \sim E^{2}(k = 4 \text{ i}(1 + 1) \sim \sim e^{+}_{0} \text{ f}_{; \text{thr}}^{n} \qquad (C:1)$$

The auxiliary quantity E (k^2) introduced here is proportional to the transverse threshold $S \{w \text{ ave } m \text{ ultipole for } ^+ - \text{electroproduction}$. In the chiral lim it the corresponding current algebra result becomes exact and gives, when expanded in k^2 ,

$$E(k^2) = 1 + \frac{k^2}{6}r_A^2 - \frac{k^2}{2m^2} + \frac{1}{4} + O(k^3)$$
 (C.2)

with

$$r_{A}^{2} = r_{A}^{2} + O(2);$$
 $n = n \frac{g_{A}^{2} m M}{8 F^{2}} + O(2)$ (C:3)

the nucleon m ean square axial radius and neutron m agnetic m om ent in the chiral lim it. Note that the axial m ean square radius has no non-analytic piece m in its quark m ass expansion [4.91] (in contrast to the isovector m agnetic m om ent). The aim is to work out all tree and loop graphs up to order q^4 which contribute to the slope term s $E^0(0) = @E(0) = @E(0) = @k^2$ proportional to M⁰ and/or M¹. The quantity $6E^0(0)$ is the sum of the m ean square axial radius r_A^2 and a host of other term s. Am ong these other term s the contributions from the relativistic B om graphs including electrom agnetic form factors will not be counted for the axial radius discrepancy r_A^2 r_A^2 , since such e ects are taken into account in the standard analysis of the pion electroproduction data. The discrepancy therefore subsum es (per de nition) all additional loop (and counter term) e ects which go beyond the form factors. Stated di erently, the dicrepancy represents all those k²-pieces which are m issing in a tree calculation (with form factors) of E (k²). A fter som e lengthy calculation [4.91] one arrives at

$$\mathbf{r}_{A}^{2} = \frac{3}{64F^{2}} \quad 1 \quad \frac{12}{2} + \frac{3M}{64m F^{2}} + \frac{3c^{+}(4)M}{32 F^{2}} + \frac{3c^{+}(4)M}{32 F^{2}} + \frac{3g_{A}^{2}}{32 F^{2}} + \frac{3g_{A}^{2}}{8^{2}m F^{2}} \quad \ln \frac{M}{2} + \frac{2}{16} + \frac{7}{12} + \frac{1}{4} + 6E^{0}(0) + 6E^{0}(0)$$
(C:4)

The rst term in (C.4) is the leading order result given already in eq.(4.79). The combination of low energy constants $c^+ = 8q + 4c_2 + 4c_3$ $q_1^2 = 2m$ can be related to the isospin even N scattering length a^+ via $c^+ = 8 F^2 a^+ = M^2$. The last two terms in

(C 3) represent the counter term contributions at order q^4 which have been identi ed with (770) and (1232) exchange contributions,

$$6E^{0}(0) = \frac{3(1 + M)}{16^{2}g_{A}mF^{2}}$$

$$6E^{0}(0) = \frac{M}{p \sqrt{2}m^{2}m^{2}} \frac{m^{2} mm + m^{2}}{mm} 2m(Y + Z + 2YZ) 2m(Y + Z + 4YZ)$$

Here, ' 6 stands for the tensor-to-vector ratio of the NN couplings and $= g_1$ ' 5 is the N coupling constant. The second N coupling g₂ of eq.(4.39) does not contribute at order q⁴. The o -shell parameters Y;Z have been estimated roughly in ref.[4.32] as 0:75 Y 1:67 and 0:8 Z 0:3. For a num erical evaluation of (C.5) these ranges are much too large and they should be further constrained. The strategy of ref.[4.91] was to link them to some nucleon structure parameters. The o -shell parameter Y enters the (1232) contribution to the proton magnetic polarizability

(C:5)

$${}_{p}^{()} = \frac{e^{2} {}^{2}}{18 {}_{m} {}^{2}m^{2}} \frac{m^{2} {}_{m} {}^{m} m + m^{2}}{m {}_{m} {}^{m}} 4Y m (Y + 1) + m (2Y + 1)$$
 (C :6)

Experimental determinations of this quantity in ref.[4.28] give a value of $p^{()}_{\rm p}$ ' 7 10^(fm) fm³ corresponding to Y ' 0:12. It is clear that the wide range of Y mentioned above is inacceptably large, since it also leads to (absurd) negative magnetic polarizabilities

⁽⁾ Furtherm one the o -shellparam eter Z of the N -vertex has been constrained. The (1232) gives a large contribution to the P-wave N scattering volume a $_{33}$. In the isobar m odel (an approximation without the o -shell parameter Z) the experimental value $a_{33} = 0.214$ =M ³ is understood to come in equal shares from nucleon pole graphs and from (1232) excitation. U sing a fully relativistic treatment of the (R arita-Schwinger spinors) the maximal value of $a_{33}^{max} = 0.185$ =M ³ is obtained with Z ' 0.3. These values of Y = 0.12 and Z = 0.3 will now be used to evaluate (C.5). Putting all pieces together, one nds for the axial radius discrepancy

$$\mathbf{r}_{A}^{2} = (4:6 + (3:1 + 1:1 + 4:5) + (7:2 + 8:0))^{2} 10 \text{ fm}^{2}$$

= (4:6 + 0:5) 10 fm² (C:7)

The rst term (-4.6) gives the leading order contribution and the others are the order corrections. In (C.6) $a^+ = 0.83 \quad 10=M$ and = 1 GeV has been used. A lthough there is some numerical uncertainty in the order correction, one obverves that the individual loop and counter term contributions cancel each other to a large extent if one makes reasonable assumptions on the parameters involved. We stress that the individual term s at order in (C.7) coming from certain classes of loop diagram s have no physical meaning, only the total sum counts. The latter tends to be quite small, similar to the sum of and contributions. In essence one concludes from this complete order q^4 calculation that no dram atic corrections to the leading order prediction of the discrepancy $r_A^2 = r_A^2$ are to be expected. It is now the task of future precision experiments (like + -electropoduction at low k^2 close to threshold and inverse -decay) to test the prediction of $r_A^2 = r_A^2$ presented here.

In this appendix, we brie y sum marize how one can use stereographic coordinates to parametrize the non{linearly realized pion and matter elds. This form alism is used extensively in section 5. The pions inhabit the three{sphere of radius F,

$$S^{3} = \frac{SO(4)}{SO(3)}$$
 (D:1)

since SU (2)_L SU (2)_R SO (4) and SU (2)_r SO (3) (local isom orphism s). Embedding the sphere in euclidean space E⁴ of cartesian coordinates = f g = f[~]; ₄ = g, the sphere is de ned via

$$X^4$$
 2 = F² : (D 2)

Three pion elds ~ can be obtained by applying e.g. a four{rotation $R (\sim) (R R^T = 1)$ to the north pole,

$$(\sim) = R_{4} (\sim) F$$
 : (D 3)

In stereographic coordinates, one has

$$R [r] = \begin{array}{c} ij \frac{1}{D} \frac{i}{2F^{2}} & \frac{1}{D} \frac{i}{F} \\ \frac{1}{D} \frac{j}{F} & \frac{1}{D} 1 \frac{z^{2}}{4F^{2}} \end{array}; \quad (D:4)$$

with

D
$$1 + \frac{2}{4F^2}$$
 : (D :5)

The corresponding covariant derivative follows to be as given in eq.(5.1). It transforms linearly under the unbroken subgroup SU $(2)_V$ but highly non{linear under SU $(2)_A$. Furtherm ore, it expresses the G oldstone boson character of the pions, i.e. their interactions vanish as the momentum transfer goes to zero.

 $E\,xp\,licit\,sym\,m\,etry\,b\,reak\,ing\,can\,\,be\,included\,\,in\,\,the\,follow\,ing\,w\,ay.\,R\,ew\,rite\,the\,m\,ass$ term in two{ avorQCD as

$$L_{mass} = \frac{1}{2} (m_u + m_d) (qq) = \frac{1}{2} (m_u - m_d) (2qt_3 q) = \frac{u}{d} :$$
 (D:6)

The rst term is the fourth component of the four{vector $S = (2qi_5tq;qq)$ and the second the third component of the SO (4) vector $P = (2qi_3;qi_5q)$ with opposite transform ation properties under parity and time reversal. Both term s break chiral sym m etry,

but only the second one breaks isospin (the invariant SO (3) subgroup does not a ect the fourth component). In terms of the pion elds, one constructs

$$S[r] = \frac{\sim}{DF}; 1 = \frac{\sim^2}{2DF^2};$$
 (D:7)

which leads to the canonical pion m ass term

L
$$_{mass} = \frac{1}{2D}M^{2} \sim^{2} + constant = \frac{1}{2}M^{2} \sim^{2} + \dots;$$
 (D.8)

and the pion mass squared is proportional to $(m_u + m_d)$.

To include the matter elds = $_{\rm N}$; ;::; one has to furnish a representation $t^{()}$ of SO (3). For the nucleon, the corresponding 2 2 matrices aret^(N) $t = \frac{1}{2}$, with the Paulim atrices. It is then most convenient to use a non{linear realization

$$N = \frac{p}{n} = \frac{1}{\frac{p}{D}} 1 + i_{5} \frac{\tau}{F} N; \qquad (D:9)$$

where $_{\rm N}$ transforms linearly under the chiral group. The pertinent covariant derivative is given in eq.(5.2). For the isobar,

the construction is the same, only that the $t^{()} = t^{(3=2)}$ are herm itean 4 d m atrices. The covariant derivative reads

$$D = (0 + \tau^{(3=2)} E) ; \qquad (D : 11)$$

with E de ned after eq.(5.2). Finally, one also needs the 2 4 spin (S_i) and isospin (T_a) $\frac{1}{2}$! $\frac{3}{2}$ transition matrices satisfying

$$S_{i}S_{j}^{+} = \frac{1}{3} (2_{ij} \quad \dot{i}_{ijk \ k});$$

$$T_{a}T_{b}^{+} = \frac{1}{6} (a_{b} \quad \dot{i}_{abc}t_{c}):$$
(D:12)