

Relativistic Bound State Equation for Unstable Fermions and the Toponium Width

Wolfgang Kummer and Wolfgang M. Oeditsch¹

Institut für Theoretische Physik
Technische Universität Wien
Wiedner Hauptstraße 8-10/136, A-1040 Wien
Austria

Abstract

The bound state problem for a fermion-antifermion system is considered taking into account a finite decay width of the constituents. We propose an exactly solvable relativistic zero order equation similar to that of Barbieri and Remiddi, but including a constant width. We focus especially on the $t\bar{t}$ system for which we reconsider our recent calculation of the bound state corrections to the toponium width, which was performed in the narrow width approximation and needed the use of second order Bethe-Salpeter perturbation theory. We show that one obtains the same result already in first order BS perturbation theory if one uses our present approach. Furthermore the large cancellations of gauge dependent terms is demonstrated to be a consequence of a Ward identity. This cancellation mechanism is shown to be valid for general fermion-antifermion systems.

Vienna, January 1995

¹e-mail: wmoedrit@ecxph.tuwien.ac.at, FAX: +43-1-5867760

1. The possible importance of off-shell corrections to the decay width in the calculation of the cross section for top-antitop production by e^+e^- lately has been the subject of detailed analysis [1, 2]. In a recent publication [3] we addressed this problem in the context of Bethe-Salpeter (BS) perturbation theory starting from the Barbieri-Remiddi equation [4]. We were able to show by explicit calculation that large cancellations took place and that the net result can be interpreted as a pure time dilatation effect. This had been conjectured qualitatively before [2] in view of the known results from muonium [5].

In this note we will improve our understanding of this result and derive a general theorem, valid for similar cases. From a theoretical point of view our calculation for toponium has been unsatisfactory since it was obtained in the narrow width approximation. As we will explain below this approximation is not strictly applicable to that system. We now include the "bulk" of the decay width already in the zero order equation to avoid this problem. This offers also the more technical advantage that we then need only to consider first order BS-perturbation theory in contrast to the calculation in the narrow width approximation where second order perturbation theory was needed. Furthermore we are able to show that the cancellation of the gauge dependent terms is a result of a Ward identity. This fact also considerably simplifies the actual calculation.

2. The Bethe-Salpeter (BS) approach for weakly bound systems is based upon the BS-equation

$$G = D + D K G \quad (1)$$

for the tt Green function $G(P; k; k^0)$, where D is the product of the two (full) propagators

and K represents the $2\pi i$ BS-kernel. All four point functions in (1) depend on the total momentum $P = (P_0; \vec{0})$; the incoming (outgoing) lines carry momentum $P = 2k$ ($P = 2k$).

Perturbation theory starts from an equation similar to eq. (1) with $G_0; D_0$ and K_0 chosen in such a way that the exact solution is known. If D_0 consists of the nonrelativistic fermion-propagators only and if K_0 is the Coulomb kernel the zero order equation corresponding to (1) (after integrating out k_0 and k_0^0) simply reduces to the Schrodinger equation. Another very convenient zero order equation which already includes the relativistic free fermion propagators and still remains solvable has been found by Barbieri and Remiddi (BR) some time ago [4]. However, as is the case for the top quark – if the decay width of the constituents becomes comparable to the binding energy, perturbation theory runs into troubles because of the occurrence of terms $(= (\frac{1}{2}m))^n$ in graphs like the one in fig. 1. To circumvent these difficulties one has to include at least a part of the exact selfenergy function in the zero order equation [6].

The key point of the BS approach as generalized to the case of a particle with finite width is the approximation to the free fermion propagator $E_k = \frac{1}{\vec{k}^2 + m^2}$

$$S(\frac{P_0}{2} + k_0) = [(\frac{P_0}{2} + k) - m - (\frac{P_0}{2} + k)]^{-1} \quad (2)$$

$$= \frac{1}{\frac{P_0}{2} + k_0 - E_k + i\frac{\Gamma}{2}} + \frac{1}{\frac{P_0}{2} + k_0 + E_k - i\frac{\Gamma}{2}}; \quad (3)$$

which is surely valid for a slowly moving particle. This choice leaves the relativistic projectors

$$(\not{p}) = \frac{E_p - (\vec{p} + \vec{m})}{2E_p}; \quad (4)$$

undisturbed and thus introduces a minimal number of terms involving the width in the

calculation. By integrating the zero order BS-equation for the Feynman amplitudes (

$$D_0 = S_+ \quad S_-)$$

$$iG_0 = -D_0 + D_0 K_0 G_0 \quad (5)$$

with respect to the zero components of momentum under the assumption that K_0 is instantaneous and with a kernel of the special form ($a = 4=3a_s$)

$$K_0 = [\gamma_0 \gamma^+ \gamma^+ \gamma^0] \int \frac{d^4 q}{q^2} \quad (6)$$

with

$$\begin{aligned} &= 2E_k = (E_k + m) \\ &= \frac{q}{2} \frac{1}{P_0 + 2E_k + i} \\ &= \frac{1}{2} (1 - \gamma_0) \end{aligned}$$

so as to annihilate the second term in (3) we can show that

$$G_0 = i(2)^4 (k_0) D_0 + [\gamma^+ \gamma^+ \gamma^0 \gamma_0] \int \frac{d^4 q}{q^2} \quad (7)$$

$$\frac{2!}{k_0^2} \frac{(2)^3}{2!} \frac{k_0 k_0}{2!} \frac{G_C(E^0; k; k^0)}{m^0} \frac{2!^0}{k_0^0} \frac{1}{2!}$$

with

$$! = E_k \frac{P_0 + i}{2} \quad (8)$$

is the solution of eq. (5). Primes denote the dependence on the corresponding momenta

and $G_C(E^0; k; k^0)$ is the nonrelativistic Coulomb Green function [7] evaluated at

$$E^0 = \frac{1}{4m} [(P_0 + i)^2 - 4m^2]:$$

This solution shows that the width of a particle can also consistently be included for a relativistic zero order equation similar to the nonrelativistic case where the replacement $E \rightarrow E + i$ in the Green function has been proposed first [8].

The corresponding solution for the BS-wave functions [6] ($\epsilon_n = E_p - M_n^{(0)} = 2 - i$)

$$\psi_n(p; \epsilon) = \psi_0(p; \epsilon)_0 = i \frac{S_0}{(p_0^2 - \epsilon_n^2)} \frac{(p) \epsilon_n^{2!}}{(p) \epsilon_{M_n^{(0)}}^{2!}} (p): \quad (9)$$

is identical to the BR wave-function for stable quarks and belongs to the spectrum of bound states $P_n^{(0)} = M_n^{(0)} - i_t = \frac{q}{1 - \epsilon_n^2} - i$. It appears as the real residue of the complex pole. The small imaginary quantity i determines the integration around that pole.

In eqs. (9) S is a constant 4×4 matrix which represents the spin state of the particle-antiparticle system :

$$S = \begin{matrix} \begin{matrix} \uparrow \downarrow \\ \uparrow \downarrow \end{matrix} & \begin{matrix} 5 \\ a_m \sim \end{matrix} & : \text{ singlet} \\ \begin{matrix} \uparrow \uparrow \\ \downarrow \downarrow \end{matrix} & & : \text{ triplet} \end{matrix} \quad (10)$$

is simply the normalized solution of the Schrodinger equation in momentum space, depending on the usual quantum numbers $(n, l, m); a_1, a_0$ in (10) describe the triplet states.

In the following it will sometimes be sufficient to use the nonrelativistic approximations of eqs. (9) ($\epsilon_n - \epsilon_n = (p^2 - m^2 + E_n) = 2 - i$)

$$(p; \epsilon)^r = \frac{p - 2i\epsilon_n}{p_0^2 - \epsilon_n^2} (p) S = \psi_0(p; \epsilon)^r_0 \quad (11)$$

Near a bound state, whose pole now lies in the lower half plane at $P_0 = P_n^{(0)}$, the

Green function of the unperturbed (BR) equation behaves as

$$G_0(P; k; k^0) = \frac{(k)}{P_0} \frac{(k^0)}{P_n^{(0)}} \quad (12)$$

A generic perturbation H changes G_0 into

$$G = \sum_{i=0}^{\infty} G_0 (H G_0)^i : \quad (13)$$

Since $G; G_0; \dots$ denote Feynman amplitudes, H in (13) differs from such an amplitude by a minus sign. It is straightforward to show that [6].

$$H = K + K_0 + iD^{-1} - iD_0^{-1} \quad (14)$$

The lhs of eq. (13) looks like (12) with $P_n^{(0)}$ replaced by P_n and with perturbed wave functions.

Formally expanding both sides of eq. (13) near a certain pole $P_n^{(0)}$

$$\begin{aligned} G_0 &= \sum_{m=0}^{\infty} g_m (P_0 - P_n^{(0)})^{m-1} \\ H &= \sum_{m=0}^{\infty} h_m (P_0 - P_n^{(0)})^m \end{aligned} \quad (15)$$

and using $g_0 =$ allows the determination of the shift of the position of the pole

$P_n = P_n - P_n^{(0)}$ to arbitrary order

$$P_n = M_n - \frac{i}{2} n = \langle h_0 \rangle + \langle h_0 g_1 h_0 \rangle + \langle h_0 i h_1 \rangle + O(n^3) : \quad (16)$$

A shorthand for BS expectation values has been used which incorporates four-integrations

of the relative momenta ($\int^R dk = (2\pi)^{-4} \int^R d^4 k$):

$$\langle h_{ii} \rangle = \int^Z dk dk^0 \int^0_{ij} (k) a_{ii^0 jj^0}(k; k^0) \int^0_{ij^0} (k^0); \quad (17)$$

As already in eq. (1) etc., in composite expressions integrations and summations over internal variables are understood. i, j etc. represent spinor as well as color indices. A similar formula as eq. (16) holds for the perturbed wave functions [9]. It is important to note that eq. (16) is not a power series in s .

3. The correction to the decay width is obtained from the imaginary part of the expectation values, receiving contributions from the graphs in g.2 and g.3. In ref. [3] the problem of the gauge independent calculation of bound state corrections to the toponium decay width had been addressed. It was shown that it is possible to obtain a gauge independent result which can be interpreted as a correction due to time dilatation. But since the top quark decay width is of the same order of magnitude as the binding energy it is obviously preferable to use the equations valid for decaying particles as described above. In our previous approach it was essential that $O(\frac{2}{s})$ corrections arose from all three terms denoted in (16) because the BR equation was used to zero order. Here we will show that within our present formalism we need only to consider first order BS perturbation theory e.g. the first term in (16).

Corrections to the fermion propagator have to be included via the perturbation kernel (see g.2)

$$h_0^{(1)} = (iD^{-1} - iD_0^{-1})_{P_0=M_n^{(0)}-i} : \quad (18)$$

We emphasize that the perturbation kernel has to be evaluated at $P_0 = M_n^{(0)} - i$. With eq. (14) and $D_0^{-1}j_{P_0=M_n^{(0)}-i} = D_0^{-1}j_{=0}$ we find

$$h_0^{(1)} = i[(\not{p} - \frac{i}{2}\not{n}) + \frac{i}{2}\not{0}] (\not{p}^0 - m) + (\not{p} - m) (\not{p} + \frac{i}{2}\not{n} - \frac{i}{2}\not{0}) (2)^4 (k - k) \quad (19)$$

where $n = (1; 0)$ and $p = \frac{M_n^{(0)}}{2} + k$, $p^0 = \frac{M_n^{(0)}}{2}$ k^0 denote the four momentum of the quark and the antiquark, respectively. The direct product refers to the t t spinor space. The factors $(\not{p} - m)$ and $(\not{p}^0 - m)$ compensate the super uous propagator on the line without . We have neglected term s that will contribute to $O(\epsilon^2 - m)$.

Since we are calculating higher order e ects we may also drop the dependence on the mass of the bottom quark although it may be included in principle. For the electroweak theory we use the R -gauge with the gauge xing Lagrangian (M denotes the mass of the W -Boson)

$$L_{gf} = \frac{1}{2} W^+ W^- + \frac{M^2}{2} W^0 W^0$$

because it eliminates mixed W propagators. The gauge parameter will not be fixed in the following. We obtain for the imaginary parts of in (19) ($s = \sin \theta_W$, $P = (1 - s^2) = 2$) for $p^2 > M^2$ and $p^2 > M^2$:

$$\text{Im } \mathcal{M}_W = \frac{e^2}{s^2 16} [P + A^{(W)} + P A^{(')}] \quad (20)$$

where

$$\begin{aligned} A^{(W)} &= (\not{p}^2) + (\not{p}^2;) \\ A^{(')} &= m^2 \left(\frac{(\not{p}^2)}{2M^2} - \frac{(\not{p}^2;)}{p^2} \right) \\ (\not{p}^2) &= \frac{(\not{p}^2 - M^2)^2}{2p^4}; \quad (\not{p}^2;) = \frac{1}{2} \left(1 - \frac{M^2}{2p^2} \right) \end{aligned} \quad (21)$$

Now we observe that the term s in (19) which include have already been calculated in [3] except for the imaginary parts occurring in the trace

$$T = \frac{1}{2} \text{tr} [\not{0} S^Y + \not{0} P \not{p} + S (\not{p} - m)] =$$

$$\frac{1}{2} (p_0 + i) \ln (1 - \frac{p^2}{m^2}) + (p_0 - i) \frac{i}{2} : \quad (22)$$

and in

$$p^2 - m^2 - 2m p_0 - p^2 - m^2 - i : \quad (23)$$

We will be able to neglect these imaginary parts in the following since they either give rise to real corrections (which we are not interested in) or to imaginary ones from $\text{Re} \epsilon$, both of which are of $O(\frac{2}{s_{\text{weak}}})$. Of course, these corrections would be required in a complete calculation of the toponium decay width to this order in order to obtain a gauge independent result. This goes beyond the scope of the present paper. Furthermore, these corrections are state independent, expected to be very small and therefore of minor practical interest, anyhow. As can be seen from (22) the axial part of ϵ does not contribute within the expectation value.

Therefore, the contribution from first order BS perturbation theory to the decay width to $O(\frac{2}{s})$ reads

$$\langle \bar{h} h_0^{(1)} \rangle = \Gamma_1 + \Gamma_3 \quad (24)$$

with

$$\Gamma_1 = 4 \text{Im} \langle \bar{h} h \rangle (p) \langle \bar{p}^0 - m \rangle (2) \langle k - k \rangle = \quad (25)$$

$$= \Gamma_0 - 2 \Gamma_0 \frac{2}{n} + \frac{e^2 m}{16 s^2} [(2M^2 + m^2) \frac{m^2}{m^4} + 2 (m^2;) \text{Re} \frac{p^2}{m^2} i;$$

$$\Gamma_3 = 4 \text{Im} \langle \bar{h} h \rangle_2 \langle \bar{p}^0 - m \rangle (2) \langle k - k \rangle = \Gamma_0 (1 + \frac{2}{n}) \quad (26)$$

A factor 2 arises here and in the following, counting both self energy contributions of the

t and t line:

$$\Gamma_0 = -2\text{Im} \Pi^{(W)}(m) = \frac{e^2 m}{16 s^2} \left(1 + \frac{m^2}{2M^2}\right) \frac{(m^2 - M^2)^2}{m^4} \quad (27)$$

In the sum $\Gamma_1 + \Gamma_3$ the leading Γ_0 terms cancel and the resulting correction is already of $O(\frac{m^2}{s^2})$.

Now we turn to the calculation of the vertex correction (Fig. 3). Performing the color trace this gives rise to the kernel

$$h_0^{(2)} = i\text{Im} \Pi_0 = \frac{4}{g^2} \quad (28)$$

In [3] we calculated $\text{Im} \Pi_0 \equiv \lim_{q^2 \rightarrow 0} p^2 \Gamma^{(2)}(p; q)$ directly. Here we will show that it is possible to use a Ward identity to determine this correction. We observe that

$$\Gamma_0^{(2)}(p; q)_{q^2 \rightarrow 0} = g_{CD} T^a \frac{\partial}{\partial p_0} \Gamma^{(1)}(p) \quad (29)$$

holds at least at the one-loop level. In the graphs shown in Fig. 3 to the required order it proves sufficient to evaluate the leading contribution by putting the fermions on the mass shell and applying the nonrelativistic wave functions (11). Using the explicit expression (20) for $\text{Im} \Pi$ we obtain

$$\begin{aligned} \text{Im} \Pi_0 &= \frac{e^2}{s^2 16} \left[A^{(W)}(m^2) + A^{(V)}(m^2) + 2m^2 \frac{\partial}{\partial p^2} (A^{(W)}(p^2) + A^{(V)}(p^2)) \right]_{p^2=m^2} \\ &= \frac{e^2}{32 s^2} F \\ F &= \left(1 + \frac{m^2}{2M^2}\right) \left(1 - \frac{M^2}{m^2}\right) \left(1 + 3\frac{M^2}{m^2}\right) + 4 \ln\left(\frac{m^2}{M^2}\right) \end{aligned} \quad (30)$$

This result agrees with that of ref. [3] and one obtains a correction to the decay width:

$$\Gamma_2 = \frac{e^2}{16 s^2} h \frac{4}{g^2} iF : \quad (31)$$

The fact that the gauge dependent terms in the sum $\Gamma_1 + \Gamma_2$ cancel is thus traced back to the identity (29). The gauge independent contribution from first order BS-perturbation theory is now different from that of [3] (actually it gives now the net result) in accordance with the different zero order equation, used in our present context.

To complete our calculation we finally check the contributions of second order perturbation theory. It can be shown that due to the fact that the first weak correction now is already of order $O(\frac{2}{s})$, these corrections do not contribute to the required order. Thus our present approach simplifies the calculation of bound state corrections to the decay width considerably.

Summing up all contributions we get the result

$$\Gamma_{\text{boundstate}} = \Gamma_1 + \Gamma_2 + \Gamma_3 = \Gamma_0 \frac{2}{s} \quad (32)$$

in agreement with our previous calculation [3].

4. The above considerations suggest that the result (32) is only a special case of a more fundamental statement, with broader range of applicability, which we will now derive.

Consider a fermion whose decay can be described by the imaginary part of a selfenergy function which will have the general form (in any covariant gauge in the relevant sector of the theory)

$$\Sigma(p) = \Sigma_S(p^2) + \Sigma_V(p^2) + \Sigma_P(p^2) + \Sigma_{SA}(p^2): \quad (33)$$

Within the expectation value (17) the pseudoscalar and axial vector parts vanish. Furthermore we will drop all factors $(p^2 - m^2)$, which is a valid approximation in all cases where $2m \ll m$. If this is not the case, either the whole calculation makes no sense

because then one would have to include the entire rung of Coulomb interactions of the remaining particle with the heavy decay products or the decay width becomes very small, or both. Let us therefore consider the correction to the decay width which results from the remaining parts of the self energy in an on shell renormalization scheme (m is the pole mass of the fermion). We expand around that mass shell:

$$\Gamma(p) = i(\text{Im } \Sigma_S + \not{p} \text{Im } \Sigma_V) + (\not{p}_S + \not{p}_V)(p^2 - m^2): \quad (34)$$

Terms of $O(g_{\text{weak}}^2)$ are understood to be neglected (cf. remarks after eq. (23)) and we denote

$$\begin{aligned} X &:= X(m^2); \quad X = S; V \\ \frac{0}{X} &:= \frac{\partial}{\partial p^2} X(p^2)_{p^2=m^2}: \end{aligned}$$

From eqs. (29) and (34) we can easily calculate the vertex correction, say, for the upper particle line:

$$\Gamma_0^+ = gT^a \Gamma_0 [i\text{Im } \Sigma_V + 2m(\frac{0}{S} + m\frac{0}{V})] + \quad (35)$$

After some algebra we can write to the required order

$$h_0^{(+)} = (\text{Im } \Sigma_S + m \text{Im } \Sigma_V + \frac{1}{2} \Gamma_0) (\not{p}^0 - m) iD_0;^1_{=0} [i\text{Im } \Sigma_V + 2m(\frac{0}{S} + m\frac{0}{V})] \quad (36)$$

$$h_0^{(+)} = \frac{4}{q^2} [i\text{Im } \Sigma_V + 2m(\frac{0}{S} + m\frac{0}{V})] \Gamma_0 \Gamma_0$$

Since

$$= 2u \text{Im } u_{p=(m, \vec{p})} = \text{Im } \Sigma_S(m^2) - m \text{Im } \Sigma_V(m^2) \quad (37)$$

and $(iD_{0;=0}^{-1} - K_{0;=0}) = 0$ we obtain for the bound state correction to the decay width to $O(\frac{2}{s_{\text{weak}}})$

$$\delta\Gamma_{\text{boundstate}} = 4i \text{Im} \left(\langle \psi^0 | m \rangle \right) = \frac{\hbar^2}{2m^2} \quad (38)$$

Now it is clear from the preceding argument that it applies not only to the toponium system but also to all other systems which can be described as weakly bound fermion-antifermion system with unstable components. Another example which is known for a long time is provided by the muonium system where the above result has also been obtained first by explicit calculation [5]. We conclude that the formalism developed here and especially the use of the identity (29) simplifies the problem of the bound state correction to the decay width in a profound way. It is now possible to clearly isolate the underlying cancellation mechanism which automatically gives a gauge independent result which can be interpreted as time dilatation alone. However, because we may now also include the scalar part of the self energy function and since K_0 can be any kernel, the result (38) is found to comprise [3] and [5] as special cases of a more general theorem: The leading bound state corrections for weakly bound systems of unstable fermions (with decays like $t \rightarrow b + W^+; \quad \tau \rightarrow e + \bar{\nu}_e + \dots$) are always of the form (38). Among the consequences we especially note that (38) may be even safely applied to toponium even if one uses e.g. a renormalization group improved potential.

Acknowledgement: This work has been supported by the Austrian Science Foundation (FWF), project P10063-PHY within the framework of the EEC-Program "Human

Capital and Mobility", Network "Physics at High Energy Colliders", contract CHRX-CT93-0357 (DG 12 COM A).

References

- [1] Y .Sum ino, K .Fu jii, H .H agiwa ra, H .M urayam a, C .K .N g,
PhysRev.D 47 (1993) 56.
- [2] M .Jezabek, J.H .K uhn, T .Teubner, Z Phys.C 56 (1992) 653;
M .Jezabek, T .Teubner, Z Phys.C 59 (1993) 669.
- [3] W .M odritsch, W K um m er, Nucl.Phys.B 430 (1994) 3.
- [4] R .Barbieri, E .Rem iddi, NuclPhysB 141 (1978) 413;
W .Buchm ullaer, E .Rem iddi, NuclPhysB 162 (1980) 250.
- [5] H .U berall, PhysRev.119 (1960) 365.
- [6] W K um m er, W .M odritsch, "R igorous Q CD -Potential for the tt-System at
threshold", TUW 94-14, to appear in Z Phys.C .
- [7] J.Schw inger, JM ath Phys 5 (1964) 1606.
- [8] V .S.Fadin, V .A .Khoze, Sov.J.NuclPhys.48 (1988) 309;
M .J.Strassler, M .E .Peskin, PhysRevD 43 (1991) 1500.
- [9] G .P .Lepage, Phys.Rev.A 16 (1977) 863;
W .K um m er, Nucl.PhysB 179 (1981) 365.




Fig. 1: Multi-decay contribution




Fig. 2: Weak off-shell corrections (D^1)




Fig. 3: Coulomb enhancement ($\sim h_0^{(2)}$)

This figure "fig1-1.png" is available in "png" format from:

<http://arxiv.org/ps/hep-ph/9501406v1>