

DPNU-94-56

INS-~~rep.~~-1075,

Nov.1994

Can the Nambu-Goldstone Boson Live on the Light-Front?

Yoonbai Kim¹, Sho Tsujimaru² and Koichi Yamawaki³

Department of Physics, Nagoya University, Nagoya 464-01, Japan

Institute for Nuclear Study, University of Tokyo, Tanashi, Tokyo 188, Japan

Abstract

We show that the Nambu-Goldstone (NG) boson restricted on the light-front (LF) can only exist if we regularize the theory by introducing the explicit breaking NG-boson mass m . The NG-boson zero mode, when integrated over the LF, must have a singular behavior $\sim 1/m^2$ in the symmetric limit of $m^2 \rightarrow 0$. In the discretized LF quantization this peculiarity is clarified in terms of the zero-mode constraints in the linear model. The LF charge annihilates the vacuum, while it is not conserved in the symmetric limit in the NG phase.

¹E-mail address: yoonbai@eken.phys.nagoya-u.ac.jp

²E-mail address: sho@insu-tokyo.ac.jp

³E-mail address: yamawaki@eken.phys.nagoya-u.ac.jp

Recently there has been renewed interest in the light-front (LF) quantization [1] as a promising approach to solve the nonperturbative dynamics [2, 3]. Based on the trivial vacuum structure, the LF quantization with a Tamm-Dancoff truncation has successfully described the bound state spectra and their wave functions in several old theoretical models in $(1+1)$ dimensions, particularly within the framework of the discretized LF quantization (D LFQ) [4, 5]. However, realistic theories like QCD in $(3+1)$ dimensions include rich structures such as confinement, spontaneous symmetry breaking (SSB), etc., which are basically on account of the nontrivial vacuum in the conventional equal-time quantization. How can one reconcile such a nontrivial structure of the theory with the trivial vacuum of the LF quantization? It seems to be now a general consensus that the zero mode [4] plays an essential role to realize the spontaneous symmetry breaking on the LF [6, 7, 3]. Problem of the zero mode in the LF vacuum was first addressed back in 1976 by Maskawa and Yamawaki [4] who discovered, within the canonical theory of D LFQ, the second class constraint so-called zero mode constraint, through which the zero mode is not an independent degree of freedom but a complicated operator-valued function of all other modes. One may thus expect that solving the vacuum state in the ordinary equal-time quantization is traded for solving the operator zero mode in the LF quantization. Actually, several authors have recently argued in $(1+1)$ dimensional models that the zero-mode solution might induce the spontaneous breaking of discrete symmetries [7]. However, the most outstanding feature of the spontaneous symmetry breaking is the existence of the Nambu-Goldstone (NG) boson for the continuous symmetry breaking. Thus the real question to be addressed is whether or not the zero-mode solution automatically produces the NG phase, particularly in $(3+1)$ dimensions.

In this paper we shall show, in the context of D LFQ, how the NG phenomenon is realized due to the zero modes in $(3+1)$ dimensions while the vacuum remains in the trivial LF vacuum. We encounter a striking feature of the zero mode of the NG boson: Naïve use of the zero-mode constraints does not lead to the NG phase at all

("no-go theorem") in contrast to the current expectation mentioned above. Within the DLFQ, it is inevitable to introduce an infrared regularization by explicit breaking mass of the NG boson m . The NG phase can only be realized via peculiar behavior of the zero mode of the NG-boson fields: The NG-boson zero mode, when integrated over the LF, must have a singular behavior $\sim 1/m^2$ in the symmetric limit $m^2 \rightarrow 0$. This we demonstrate both in a general framework of the LSZ reduction formula and in a concrete field theoretical model, the linear model, within a framework of DLFQ. The NG phase is in fact realized in such a way that the vacuum is trivial while the LF charge is not conserved in the symmetric limit $m^2 \rightarrow 0$.

Let us first prove a "no-go theorem" that the naive LF restriction of the NG-boson field leads to vanishing of both the NG-boson emission vertex and the corresponding current vertex; namely, the NG phase is not realized in the LF quantization.

Based on the LSZ reduction formula, the NG-boson emission vertex $A \rightarrow B +$ may be written as

$$\begin{aligned} \langle B(q) | \mathcal{A} | i \rangle &= i \int d^4x e^{iqx} \langle B | \mathcal{J}(x) | i \rangle \\ &= i(2\pi)^4 (\delta_A - p_B - q) \langle \mathcal{P}_A - p_B - q | \langle B | j(0) | i \rangle; \end{aligned} \quad (1)$$

where $\mathcal{J}(x)$ and $j(x) = 2\mathcal{Q}(x) = (2\mathcal{Q}_+ \mathcal{Q}_- - \mathcal{Q}_?^2)(x)$ are the interpolating field and the source function of the NG boson, respectively, and $q = p_A - p_B$ are the NG-boson four momenta and $q = (q^+, q^2) [8]$. It is customary [9] to take the collinear momentum, $q = 0$ and $q \neq 0$ (not a soft momentum), for the emission vertex of the exactly massless NG boson with $q^2 = 0$. Here we adopt the DLFQ, $x \in [-L; L]$, with a periodic boundary condition [10] in the x direction and take the continuum limit $L \rightarrow \infty$ in the end of the whole calculation [4]. Without specifying the boundary condition, we would not be able to formulate consistently the LF quantization anyway even in the continuum theory [11]. Then the NG-boson emission vertex should vanish on the LF due to the periodic boundary condition:

$$(2\pi)^3 \langle \mathcal{P}_A - p_B | \langle B | j(0) | i \rangle$$

$$= \int d^2x \lim_{L \rightarrow \infty} \int_L^{\infty} dx \, 2\partial_+ \partial_- \mathcal{A} = 0: \quad (2)$$

Another symptom of this disease is the vanishing of the current vertex (analogue of g_A in the nucleon matrix element). When the continuous symmetry is spontaneously broken, the NG theorem requires that the corresponding current J contains an interpolating field of the NG boson (ϕ), that is, $J = f \partial_- \phi + \mathcal{J}$, where f is the "decay constant" of the NG boson and \mathcal{J} denotes the non-pole term. Then the current conservation $\partial_- J = 0$ leads to

$$\begin{aligned} 0 &= \int d^3x \partial_- \mathcal{J}(\mathbf{x}) \mathcal{A}|_{x^0=0} \\ &= i(2\pi)^3 \int d^3x \frac{m_A^2 - m_B^2}{2p_A^+} \mathcal{J}^{p^+}(0) \mathcal{A}|_{x^0=0} \end{aligned} \quad (3)$$

where $\int d^3x \lim_{L \rightarrow \infty} \int_L^{\infty} dx \, d^2x$ and the integral of the NG-boson sector 2 has no contribution on the LF because of the periodic boundary condition as we mentioned before. Thus the current vertex $\mathcal{J}^{p^+}(0) \mathcal{A}|_{x^0=0}$ should vanish at $q^2 = 0$ as far as $m_A^2 \neq m_B^2$ [12].

This is actually a manifestation of the conservation of a charge $\mathcal{Q} = \int d^3x \mathcal{J}^{p^+}$ which is constructed only from the non-pole term. Note that \mathcal{Q} is equivalent to the full LF charge $Q = \int d^3x J^+$, since the pole part always drops out of Q due to the integration on the LF, i.e., $Q = \mathcal{Q}$. Therefore the conservation of \mathcal{Q} inevitably follows from the conservation of Q : $[\mathcal{Q}, P_-] = [Q, P_-] = 0$, which in fact implies vanishing current vertex mentioned above. This is in sharp contrast to the charge integrated over usual space $\mathbf{x} = (x^1, x^2, x^3)$ in the equal-time quantization: $Q^{\text{et}} = \int d^3x J^0$ is conserved while $\mathcal{Q}^{\text{et}} = \int d^3x \mathcal{J}^0$ is not.

Thus the NG bosons are totally decoupled, i.e., the NG phase is not realized on the LF. Note that this is a direct consequence of the periodic boundary condition and the first-order form of $2 = 2\partial_+ \partial_- \phi + \mathcal{J}^2$ in ∂_- in contrast to the second order form in ∂_0 in the equal-time quantization.

Now, we propose to regularize the theory by introducing explicit breaking mass of the NG boson m . The essence of the NG phase with a small explicit symmetry

breaking can well be described by the old notion of the PCAC hypothesis: $\partial J(x) = f m^2 \phi(x)$, with $\phi(x)$ being the interpolating field of the (pseudo-) NG boson. From the PCAC relation the current divergence of the non-pole term $\mathcal{J}(x)$ reads $\partial \mathcal{J}(x) = f(2 + m^2) \phi(x) = f j(x)$. Then we obtain

$$\begin{aligned} \int d^3x \partial \mathcal{J}(x) \langle \mathcal{A} | &= f m^2 \int d^3x \phi(x) \langle \mathcal{A} | \\ &= \int d^3x f j(x) \langle \mathcal{A} |; \end{aligned} \quad (4)$$

where the integration of the pole term $2 \phi(x)$ is dropped out as before. The second expression of (4) is nothing but the matrix element of the LF integration of the zero mode (with $P^+ = 0$)! $\frac{1}{2L} \int_L^R dx \phi(x)$. Suppose that $\int d^3x \phi(x) = \int d^3x \phi(x)$ is regular when $m^2 \rightarrow 0$. Then this leads to the "no-go theorem" again. Thus in order to have the non-zero NG-boson emission vertex (R.H.S. of (4)) as well as the non-zero current vertex (L.H.S.) at $q^2 = 0$, the zero mode $\phi(x)$ must behave as

$$\int d^3x \phi(x) \sim \frac{1}{m^2} (m^2 \rightarrow 0): \quad (5)$$

This situation may be clarified when the PCAC relation is written in the momentum space:

$$\frac{m^2 f j(q^2)}{m^2 - q^2} = \partial J(q) = \frac{q^2 f j(q^2)}{m^2 - q^2} + \partial \mathcal{J}(q): \quad (6)$$

What we have done when we reached the "no-go theorem" can be summarized as follows. We first set L.H.S. of (6) to zero (or equivalently, assumed implicitly the regular behavior of $\int d^3x \phi(x)$) in the symmetric limit in accord with the current conservation $\partial J = 0$. Then in the LF formalism with $\epsilon = 0$ ($q^2 = 0$), the first term (NG-boson pole term) of R.H.S. was also zero due to the periodic boundary condition or the zero-mode constraint. Thus we arrived at $\partial \mathcal{J}(q) = 0$. However, this procedure is equivalent to playing a nonsense game: $\lim_{m^2, q^2 \rightarrow 0} (\frac{m^2 - q^2}{m^2 - q^2}) = 0$ as far as $f j \neq 0$ (NG phase). Therefore the " $m^2 = 0$ " theory with vanishing L.H.S. is ill-defined on the LF, namely, the "no-go theorem" is false. The correct procedure should be to take the symmetric limit $m^2 \rightarrow 0$ after the LF restriction $\epsilon = 0$ ($q^2 = 0$)

[13], although (6) itself yields the same result $f_j = \partial_j \Phi$, irrespectively of the order of the two limits $q^2 \rightarrow 0$ and $m^2 \rightarrow 0$. Then (5) does follow. This implies that at quantum level the LF charge $Q = \Phi$ is not conserved, or the current conservation does not hold for a particular Fourier component with $q = 0$ even in the symmetric limit:

$$\frac{1}{i} [Q; P] = \partial_j J_j|_{q=0} = f \lim_{m^2 \rightarrow 0} m^2 \int d^3x \neq 0; \quad (7)$$

Let us now demonstrate that (5) and (7) indeed take place as the solution of the constrained zero-modes in the NG phase of the $O(2)$ linear model:

$$L = \frac{1}{2} (\partial_- \phi)^2 + \frac{1}{2} (\partial_+ \phi)^2 - \frac{1}{2} m^2 (\phi^2 + \chi^2) - \frac{1}{4} (\phi^2 + \chi^2)^2 + c; \quad (8)$$

where the last term is the explicit breaking which regularizes the NG-boson zero mode.

In the DLFQ we can clearly separate the zero modes (with $P^+ = 0$), ϕ_0 , χ_0 ($\frac{1}{2L} \int_{-L}^L dx \phi(x)$ (similarly for χ_0), from other oscillating modes (with $P^+ \neq 0$), ϕ'_i , χ'_i (similarly for χ'). Through the Dirac quantization of the constrained system the canonical commutation relation for the oscillating modes reads [4]

$$[\phi'_i(x); \chi'_j(y)] = \frac{i}{4} \epsilon_{ij}^{(2)} (x-y) \frac{x-y}{L} \epsilon_{ij}^{(2)} (x^2 - y^2); \quad (9)$$

where each index stands for ϕ or χ , and the $\epsilon(x)$ is the sign function. By use of (9) we can introduce creation and annihilation operators simply defined by the Fourier expansion of ϕ'_i with respect to x even when the interaction is included. Thus the physical Fock space is constructed upon the LF vacuum ("trivial vacuum") which is defined to be annihilated by the annihilation operators without recourse to the dynamics.

On the other hand, the zero modes are not independent degrees of freedom but are implicitly determined by ϕ' and χ' through the second class constraints so-called zero-mode constraints [4]:

$$\frac{1}{2L} \int_{-L}^L dx \left(\phi^2 - \chi^2 \right) + \left(\phi^2 + \chi^2 \right) = 0; \quad (10)$$

and similarly, $\frac{1}{2L} \int_L^R dx f[\phi] = 0$. Note that through the equation of motion these constraints are equivalent to the characteristic of the DLFQ with periodic boundary condition: $\frac{1}{2L} \int_L^R dx \partial_+ \phi = 0$; (similarly for ∂_-) which we have used to prove the "no-go theorem" for the case of $m^2 = 0$.

Actually, in the NG phase ($m^2 < 0$) the equation of motion of ϕ reads $(\partial^2 + m^2)\phi(x) = (\phi^3 + \phi^2 + 2v\phi)j(x)$, with $\phi^0 = v$ and $m^2 = -v^2 + v^2 = -v^2$, where v is the classical vacuum solution determined by $v^2 + v^3 = c$. Integrating the above equation of motion over x , we have

$$\int_L^R d^3x j(x) - m^2 \int_L^R d^3x \phi(x) = \int_L^R d^3x \phi^2(x) = \int_L^R d^3x \phi = 0; \quad (11)$$

where $\int_L^R d^3x \phi(x) = \int_L^R d^3x \phi^0(x)$. Were it not for the singular behavior (5) for the zero mode ϕ , we would have concluded $(\partial^2 + m^2)\phi(x) = 0$ which $\int_L^R d^3x \phi(x) = 0$ in the symmetric limit $m^2 \rightarrow 0$. Namely, the NG-boson vertex at $q^2 = 0$ would have vanished, which is exactly what we called "no-go theorem" now related to the zero-mode constraint. On the contrary, direct evaluation of the matrix element of $j =$

$(\phi^3 + \phi^2 + 2v\phi)$ in the lowest order perturbation yields non-zero result even in the symmetric limit $m^2 \rightarrow 0$: $\langle j \rangle = 2v \langle \phi \rangle = 2v \neq 0$ ($\phi = 0$);

which is in agreement with the usual equal-time formulation. Thus we have seen that naive use of the zero-mode constraints by setting $m^2 = 0$ leads to the internal inconsistency in the NG phase. The "no-go theorem" is again false.

We now study the solution of the zero-mode constraints in the perturbation around the classical (tree level) SSB vacuum, since we need to formulate the NG phase on the LF at least for the theory whose SSB is already described at the tree level in the equal-time quantization. It is convenient to divide the zero modes ϕ_0 (or ϕ) into classical constant piece v (or v) and operator part ϕ_0 (or ϕ), and also do the zero-mode constraints. The classical part of the zero-mode constraints is nothing but the condition that determines the minimum of the classical potential and we have chosen a solution that $v = 0$ and $v = v$; i.e., $\phi_0 = \phi$, $\phi = \phi + v$. The operator zero modes are solved perturbatively by substituting the expansion $\phi_i = \sum_{k=1}^P \phi_i^{(k)}$

under the Weyl ordering.

The lowest order solution of the zero-mode constraints and for ϕ takes the form :

$$(\phi^2 + \phi_0^2) \phi = \frac{1}{2L} \int_0^L dx (\phi'^3 + \phi' \phi'^2 + 2v \phi' \phi) : \quad (12)$$

Then (5) immediately follows [14]:

$$\lim_{m^2 \rightarrow 0} m^2 \int_0^L dx \phi^3 = \int_0^L dx \phi^3 (\phi'^3 + \phi' \phi'^2 + 2v \phi' \phi) \neq 0 : \quad (13)$$

This is our main result. This actually ensures non-zero ϕ vertex through (11): $\langle \phi(0) \phi^3 \rangle = 2v$; which agrees with the previous direct evaluation as it should.

Let us next discuss the LF charge operator. The $O(2)$ current in this model is given by $J = \phi \partial \phi$. As was noted in Ref.[4], the corresponding LF charge $Q = \oint \phi^R d^3x (\phi' \phi' - \phi' \phi')$ contains no zero-modes including the pole term which was dropped by the integration due to the periodic boundary condition and the ϕ , so that Q is well-defined even in the NG phase and hence annihilates the vacuum simply by the P^+ conservation:

$$Q |0\rangle = 0 : \quad (14)$$

This is also consistent with explicit computation of the commutators: $\langle \phi | Q ; \phi' | \rangle = i \phi' | \rangle = 0$ and $\langle \phi | Q ; \phi' | \rangle = i \phi' | \rangle = 0$ [15], which are contrasted to those in the usual equal-time case where the spontaneously broken charge does not annihilate the vacuum: $\langle \phi | Q^{et} ; \phi | \rangle = i \phi | \rangle = 0$; $\langle \phi | Q^{et} ; \phi | \rangle = i \phi \neq 0$.

Since the PCAC relation is now an operator relation for the canonical field $\phi(x)$ with $f = v$ in this model, (13) ensures $\langle \phi | P^- | \rangle \neq 0$ or a non-zero current vertex $\langle \phi^+ | j | \rangle \neq 0$ ($q^2 = 0$) in the symmetric limit. Noting that $Q = \oint \phi$, we conclude that the regularized zero-mode constraints indeed lead to non-conservation of the LF charge in the symmetric limit $m^2 \rightarrow 0$:

$$Q |P^- \rangle = i v \lim_{m^2 \rightarrow 0} m^2 \int_0^L dx \phi^3 \neq 0 : \quad (15)$$

This can also be confirmed by direct computation of $[Q; P]$ through the canonical commutator and explicit use of the regularized zero-mode constraints. At first sight there seems to be no distinction between the spontaneous and the explicit symmetry breakings on the LF. However, the singular behavior of the NG-boson zero mode (5) or (13) may be understood as a characterization of the spontaneous symmetry breaking.

Our result implies that solving the zero-mode constraints without regularization would not lead to the NG phase at all in contradiction to the naive expectation [7]. Our treatment of the zero modes in the canonical DLFQ is quite different from that proposed recently by Wilson et al. [3] who eliminate the zero modes by hand in the continuum theory instead of solving the zero-mode constraints. They also arrived at the non-conservation of the LF charge without zero mode, while still claiming the conservation of the full LF charge in contrast with our result. The relationship between these two approaches are not clear at the moment. Finally, it should be noted that there exists another no-go theorem that forbids any LF field theory (even the free theory) satisfying the Wightman axioms [16]. This no-go theorem is also related to the zero modes but has not yet been overcome by the DLFQ or any other existing approach and is beyond the scope of this paper.

We would like to thank T. Kugo, Y. Ohnuki and I. Tutsui for useful discussions. Y.K. is a JSPS Postdoctoral Fellow (No.93033). This work was supported in part by a Grant-in-Aid for Scientific Research from the Ministry of Education, Science and Culture (No.05640339), the Ishida Foundation and the Sumitomo Foundation. Part of this work was done while K.Y. was staying at Institute for Theoretical Physics at U.C. Santa Barbara in May, 1994, which was supported in part by the Yoshida Foundation for Science and Technology and by the U.S. National Science Foundation under Grant No. PHY 89-04035.

References

- [1] P. A. M. Dirac, Rev. Mod. Phys. 21, 392 (1949).
- [2] For a review, see S. J. Brodsky, G. McCartor, H. C. Pauli and S. S. Pinsky, Particle World 3, 109 (1993) and references therein.
- [3] K. G. Wilson, T. S. Walhout, A. Harindranath, W. Zhang, R. J. Perry and S. D. Glazek, Phys. Rev. D 49, 6720 (1994).
- [4] T. Maskawa and K. Yamawaki, Prog. Theor. Phys. 56, 270 (1976).
- [5] H. C. Pauli and S. J. Brodsky, Phys. Rev. D 32, 1993 (1985); *ibid* 2001 (1985).
- [6] T. Heinzl, S. K rushe and E. W emer, Phys. Lett. B 277, 54 (1991).
- [7] T. Heinzl, S. K rushe and E. W emer, Z. Phys. C 56, 415 (1992); D. G. Robertson, Phys. Rev. D 47, 2549 (1993); C. M. Bender, S. Pinsky, and B. Van de Sande, Phys. Rev. D 48, 816 (1993).
- [8] We choose the LF "time" as $x^+ = x \frac{1}{2}(x^0 + x^3)$ and the longitudinal and transverse coordinates are denoted by $x \equiv (x^-; x^\perp)$ with $x^- = \frac{1}{2}(x^0 - x^3)$ and $x^\perp = (x^1; x^2)$, respectively.
- [9] S. Weinberg, Phys. Rev. 150. 1313 (1966)
- [10] A consistent DLFQ can also be constructed with antiperiodic boundary condition, which, however, requires each zero mode to be identically zero and hence no vacuum expectation value of the field, namely, no spontaneous symmetry breaking.
- [11] P. Steinhardt, Ann. Phys. 32, 425 (1980).
- [12] For the nucleon matrix element with $m_A = m_B = m_N$, (3) should read $0 = u(p_B) \gamma^+ \gamma_5 u(p_A) g(q^2 = 0)$, with $u \gamma^+ \gamma_5 u = 2m_N$ and $u \not{q} u = 0$ ($q = 0; q \neq 0$), which implies $g_A(0) = 0$.

- [13] The role of the explicit mass m in defining LF charge was discussed by R. Carlitz, D. Heckathorn, J. Kaur and W. K. Tung, Phys. Rev. 11, 1234 (1975), however without consistent boundary condition and zero-mode constraints.
- [14] As is seen from the canonical commutator, $\phi(x)$ scales like a dimensionless field under the scale transformation in the x direction and hence the zero mode ϕ_0 is independent of L . Thus there is no subtlety between the two limits, $L \rightarrow 1$ and $m^2 \rightarrow 0$.
- [15] By explicit calculation with a careful treatment of the zero-modes contribution we can also show that $\langle \phi_0^2 \rangle = \langle \phi_0^2 \rangle = 0$.
- [16] N. Nakanishi and K. Yamawaki, Nucl. Phys. B 122, 15 (1977).